


Chapter 4:

Euclidean Geometry



Greek Math

Greek Math

- Pythagoras died ~ 500BC. Euclid born ~325BC.
Between them lived Socrates, Plato and Alexander
the Great.

Greek Math

- Pythagoras died ~ 500BC. Euclid born ~325BC. Between them lived Socrates, Plato and Alexander the Great.
- Plato established the Academy in Athens ~387BC, giving scholars time, resources and collaborators. Legend says that above the entrance was “Let no man ignorant of geometry enter here.”


Greek Math

- Pythagoras died ~ 500BC. Euclid born ~325BC. Between them lived Socrates, Plato and Alexander the Great.
- Plato established the Academy in Athens ~387BC, giving scholars time, resources and collaborators. Legend says that above the entrance was “Let no man ignorant of geometry enter here.”
- Many great Greek mathematicians studied at the Academy. Greatest of them was Eudoxus of Cnidus.


Academy in Athens



Life in Ancient Athens



Life in Ancient Athens



The Elements

The Elements

- Euclid wrote the *Elements*, which is the most important book in math history. Much of it is not his original work—it is compiled from other sources.



The Elements

- Euclid wrote the *Elements*, which is the most important book in math history. Much of it is not his original work—it is compiled from other sources.
- The major sources were likely the Pythagoreans, Hippocrates and Eudoxus. Many of the proofs are believed to be original to Euclid.



The Elements

- Euclid wrote the *Elements*, which is the most important book in math history. Much of it is not his original work—it is compiled from other sources.
- The major sources were likely the Pythagoreans, Hippocrates and Eudoxus. Many of the proofs are believed to be original to Euclid.
- Its success is due to its logical and axiomatic presentation. This deductive style is the central approach to mathematics today.



The Elements

The Elements

- The text was the primary math textbook for *two thousand years*.

The Elements

- The text was the primary math textbook for *two thousand years*.
- It is divided into 13 “books” on topics from geometry, algebra and number theory.

The Elements

- The text was the primary math textbook for *two thousand years*.
- It is divided into 13 “books” on topics from geometry, algebra and number theory.
- Book I is devoted to propositions from planar geometry.


The Elements

- The text was the primary math textbook for *two thousand years*.
- It is divided into 13 “books” on topics from geometry, algebra and number theory.
- Book I is devoted to propositions from planar geometry.
- Euclid’s postulates allow him to use a straightedge and compass, which play a central role.

Definitions, Common Notions and Postulates


Proving Propositions

Triangle Angle Sum




[Link](#)

Triangle Angle Sum




[Link](#)



A diagram illustrating the Pythagorean theorem. It features a large square divided into four right triangles. The two triangles on the left have legs labeled a and b , and the hypotenuse of the top-left triangle is labeled c . The bottom-left triangle has legs a and b , and its hypotenuse is also labeled c . The two triangles on the right are congruent to the ones on the left, with legs a and b , and hypotenuse c . The total area of the large square is equal to the sum of the areas of the four triangles plus the area of the central red square.

$$a^2 + b^2 = c^2$$

PYTHAGOREAN THEOREM



A diagram illustrating the Pythagorean theorem. It features a large square divided into four right triangles. The two triangles on the left have legs labeled a and b , and the hypotenuse of the top-left triangle is labeled c . The bottom-left triangle has legs a and b , and its hypotenuse is also labeled c . The two triangles on the right are congruent to the ones on the left, with legs a and b , and hypotenuse c . The total area of the large square is equal to the sum of the areas of the four triangles plus the area of the central red square.

$$a^2 + b^2 = c^2$$

PYTHAGOREAN THEOREM

The Aftermath

The Aftermath

- Book I showed how to construct an equilateral triangle and a square.

The Aftermath

- Book I showed how to construct an equilateral triangle and a square.
- Ancient Greeks also know how to construct a regular pentagon, and given a regular polygon with m sides, they could construct a regular polygon with $2m$ sides.

The Aftermath

- Book I showed how to construct an equilateral triangle and a square.
- Ancient Greeks also know how to construct a regular pentagon, and given a regular polygon with m sides, they could construct a regular polygon with $2m$ sides.
- Major open question: For which values of n is the regular n -gon constructible with a straightedge and compass?

The Aftermath

The Aftermath

- There was no progress made until 1796, when a 19-year-old named Carl Friedrich Gauss used some beautiful algebra to find a construction of the regular 17-gon.



The Aftermath

- There was no progress made until 1796, which a 19-year-old named Carl Friedrich Gauss used some beautiful algebra to find a construction of the regular 17-gon.
- Then, at 24, he entirely classified which regular polygons are constructible.



The Aftermath

- There was no progress made until 1796, which a 19-year-old named Carl Friedrich Gauss used some beautiful algebra to find a construction of the regular 17-gon.
- Then, at 24, he entirely classified which regular polygons are constructible.
- He did not prove one direction of his theorem, though. That was done by Pierre Wantzel.



The Aftermath



The Aftermath

- The Gauss-Wantzel theorem says this:



The Aftermath

- The Gauss-Wantzel theorem says this:
- A regular n -gon can be constructed with straightedge and compass if and only if n is a power of 2 or the product of a power of 2 and any number of distinct Fermat primes.



The Aftermath

- The Gauss-Wantzel theorem says this:
- A regular n -gon can be constructed with straightedge and compass if and only if n is a power of 2 or the product of a power of 2 and any number of distinct Fermat primes.
- A Fermat prime is a prime of the form $2^{2^k} + 1$.



The Elements

The Elements

- Book I: Planar geometry

The Elements

- Book I: Planar geometry
- Book II: Algebraic geometry

The Elements

- Book I: Planar geometry
- Book II: Algebraic geometry
- Book III: Circles

The Elements

- Book I: Planar geometry
- Book II: Algebraic geometry
- Book III: Circles
- Book IV: Inscribing and circumscribing

The Elements

- Book I: Planar geometry
- Book II: Algebraic geometry
- Book III: Circles
- Book IV: Inscribing and circumscribing
- Book V: Magnitudes, ratios and rules of arithmetic

The Elements

- Book I: Planar geometry
- Book II: Algebraic geometry
- Book III: Circles
- Book IV: Inscribing and circumscribing
- Book V: Magnitudes, ratios and rules of arithmetic
- Book VI: Proportions between lines and shapes

The Elements

The Elements

- Book VII, VIII and IX: Number theory

The Elements

- Book VII, VIII and IX: Number theory
- Book X: Square roots

The Elements

- Book VII, VIII and IX: Number theory
- Book X: Square roots
- Book XI: Solid figures

The Elements


- Book VII, VIII and IX: Number theory
- Book X: Square roots
- Book XI: Solid figures
- Book XII: Volume

The Elements


- Book VII, VIII and IX: Number theory
- Book X: Square roots
- Book XI: Solid figures
- Book XII: Volume
- Book XIII: Platonic solids




Platonic solids



Platonic solids




Hypatia



THE
MURDER
OF ANCIENT ALEXANDRIA'S
GREATEST
SCHOLAR



Hypatia



THE
MURDER
OF ANCIENT ALEXANDRIA'S
GREATEST
SCHOLAR



Hypatia



Hypatia

- Hypatia is the first woman in math history that we know a lot about.



Hypatia

- Hypatia is the first woman in math history that we know a lot about.
- She was a first-class thinker and highly respected philosopher, mathematician and teacher.





Hypatia

- Hypatia is the first woman in math history that we know a lot about.
- She was a first-class thinker and highly respected philosopher, mathematician and teacher.
- Her life ended in tragedy, as she was brutally murdered for defending religious freedom.



The Legacy of Axiomatic Thinking





The Aftermath

The Aftermath

- Recall that Euclidean geometry was built from 5 axioms, the final of which was *the parallel postulate*. This one was slightly controversial.

The Aftermath

- Recall that Euclidean geometry was built from 5 axioms, the final of which was *the parallel postulate*. This one was slightly controversial.
- Can that postulate be deduced from the other four? Unfortunately, no.

The Aftermath

- Recall that Euclidean geometry was built from 5 axioms, the final of which was *the parallel postulate*. This one was slightly controversial.
- Can that postulate be deduced from the other four? Unfortunately, no.
- In this way, the parallel postulate is *undecidable* if your axioms are only the first four.


The Aftermath

The Aftermath


- This implies that there exist non-Euclidean geometries. Instead of the parallel postulate, you can have an axiom saying that if ℓ is a line and P is a point off of ℓ , then there are 2+ lines through P that are parallel to ℓ . Or no lines that are parallel.

The Aftermath


- This implies that there exist non-Euclidean geometries. Instead of the parallel postulate, you can have an axiom saying that if ℓ is a line and P is a point off of ℓ , then there are 2+ lines through P that are parallel to ℓ . Or no lines that are parallel.



Hyperbolic



Euclidean



Elliptic

The Aftermath

The Aftermath

- Our set theory today is built from the *ZFC axioms*. These are 8 completely basic axioms about sets, plus the slightly-controversial *axiom of choice*.

The Aftermath

- Our set theory today is built from the *ZFC axioms*. These are 8 completely basic axioms about sets, plus the slightly-controversial *axiom of choice*.
- From these, not only is set theory built, but number theory, too.

The Aftermath



- Our set theory today is built from the *ZFC axioms*. These are 8 completely basic axioms about sets, plus the slightly-controversial *axiom of choice*.
- From these, not only is set theory built, but number theory, too.
- Kurt Gödel proved that every set of axioms that leads to number theory must have undecidable statements. There must be statements that can neither be proven nor disproven.

Approximating $\sqrt{2}$

Approximating $\sqrt{2}$

- This chapter is on geometry and the next chapter is on number theory. A bridge between these topics is using geometry to approximate $\sqrt{2}$.

Approximating $\sqrt{2}$


- This chapter is on geometry and the next chapter is on number theory. A bridge between these topics is using geometry to approximate $\sqrt{2}$.
- We previously learned about the Indian geometry from their *sulbasutras*. These also contained an approximation:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Approximating $\sqrt{2}$

- This chapter is on geometry and the next chapter is on number theory. A bridge between these topics is using geometry to approximate $\sqrt{2}$.
- We previously learned about the Indian geometry from their *sulbasutras*. These also contained an approximation:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$




1.414213...

Approximating $\sqrt{2}$


- This chapter is on geometry and the next chapter is on number theory. A bridge between these topics is using geometry to approximate $\sqrt{2}$.
- We previously learned about the Indian geometry from their *sulbasutras*. These also contained an approximation:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$



Approximating $\sqrt{2}$

Suppose you have two 1×1 squares



1


1

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1

1


Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

If you could combine them into a single square, it would be a $\sqrt{2} \times \sqrt{2}$ square

$\sqrt{2}$




$\sqrt{2}$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1


1

Approximating $\sqrt{2}$


- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1




$\frac{1}{2}$ $\frac{1}{2}$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1


1

Approximating $\sqrt{2}$


- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1



1


$\frac{1}{3}$ $\frac{1}{3}$ $\frac{1}{3}$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares




Approximating $\sqrt{2}$


- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1 $\frac{1}{3}$




$\frac{1}{3}$

Approximating $\sqrt{2}$


- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares



1 $\frac{1}{3}$




Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Suppose you have two 1×1 squares





Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Divide into 8 pieces





Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Place them on
the diagram




$$\frac{1}{3 \cdot 4}$$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

Place them on
the diagram




1 $\frac{1}{3}$

$\frac{1}{3}$ $\frac{1}{3 \cdot 4}$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$




Place them on
the diagram

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$




This is not a square, but it's close. And it does use all of the original two squares.

Approximating $\sqrt{2}$

- Goal:


$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$



Approximating $\sqrt{2}$

- Goal:


$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$



Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$



Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

$$2 \cdot W \times \left(1 + \frac{1}{3} + \frac{1}{3 \cdot 4} \right) - W^2 = \frac{1}{3 \cdot 4} \times \frac{1}{3 \cdot 4}$$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

$$2 \cdot W \times \left(1 + \frac{1}{3} + \frac{1}{3 \cdot 4} \right) = \frac{1}{3 \cdot 4} \times \frac{1}{3 \cdot 4}$$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

$$2 \cdot W \times \left(1 + \frac{1}{3} + \frac{1}{3 \cdot 4} \right) = \frac{1}{3 \cdot 4} \times \frac{1}{3 \cdot 4}$$
$$\frac{2W(12 + 4 + 1)}{3 \cdot 4} = \frac{1}{3 \cdot 4 \cdot 3 \cdot 4}$$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

$$2 \cdot W \times \left(1 + \frac{1}{3} + \frac{1}{3 \cdot 4} \right) = \frac{1}{3 \cdot 4} \times \frac{1}{3 \cdot 4}$$
$$\frac{2W(12 + 4 + 1)}{3 \cdot 4} = \frac{1}{3 \cdot 4 \cdot 3 \cdot 4}$$
$$34W = \frac{1}{3 \cdot 4}$$

Approximating $\sqrt{2}$

- Goal:

$$\sqrt{2} \approx 1 + \frac{1}{3} + \frac{1}{3 \cdot 4} - \frac{1}{3 \cdot 4 \cdot 34}.$$

$$2 \cdot W \times \left(1 + \frac{1}{3} + \frac{1}{3 \cdot 4} \right) = \frac{1}{3 \cdot 4} \times \frac{1}{3 \cdot 4}$$

$$\frac{2W(12 + 4 + 1)}{3 \cdot 4} = \frac{1}{3 \cdot 4 \cdot 3 \cdot 4}$$

$$34W = \frac{1}{3 \cdot 4}$$

$$W = \frac{1}{3 \cdot 4 \cdot 34}.$$



Shout-outs!



Shout-outs!

- Apollonius lived ~262-190 BC, modern-day Turkey.



Shout-outs!

- Apollonius lived ~262-190 BC, modern-day Turkey.
- Wrote the text *Conic*. A major study in ellipses, parabolas, hyperbolas, and tangent lines.







Shout-outs!

- Apollonius lived ~262-190 BC, modern-day Turkey.
- Wrote the text *Conic*. A major study in ellipses, parabolas, hyperbolas, and tangent lines.
- Apollonius problem: Given 3 circles, find a 4th circle tangent to all 3.
(Solved in 1956.)




Shout-outs!

- Apollonius lived ~262-190 BC, modern-day Turkey.
- Wrote the text *Conic*. A major study in ellipses, parabolas, hyperbolas, and tangent lines.
- Apollonius problem: Given 3 circles, find a 4th circle tangent to all 3.
(Solved in 1956.)





Shout-outs!

- A related problem is one of my favorite in all of math.




• A_r





Shout-outs!

- A related problem is one of my favorite in all of math.




• A_r





Shout-outs!

- A related problem is one of my favorite in all of math.




• A
r




Shout-outs!

- A related problem is one of my favorite in all of math.




• A_r




Shout-outs!

- A related problem is one of my favorite in all of math.




• A
m



Shout-outs!

- A related problem is one of my favorite in all of math.



Shout-outs!

Shout-outs!

- Packing problems are still important areas of study.

Shout-outs!

- Packing problems are still important areas of study.
- In the 17th century, Johannes Kepler conjectured that the most efficient way to pack spheres will result in about 75% of the volume being filled.

Shout-outs!

- Packing problems are still important areas of study.
- In the 17th century, Johannes Kepler conjectured that the most efficient way to pack spheres will result in about 75% of the volume being filled.
- This was proved in 1998 by Thomas Hales.

Shout-outs!

Shout-outs!

- In 2017, Maryna Viazovska generalized this to higher dimensions.



Shout-outs!

- In 2017, Maryna Viazovska generalized this to higher dimensions.
- Example: in 8 dimensions, only 25% of your space can be filled with hyperspheres. In 24 dimensions, only 0.1% of your space can be filled with hyperspheres.



Shout-outs!

- In 2017, Maryna Viazovska generalized this to higher dimensions.
- Example: in 8 dimensions, only 25% of your space can be filled with hyperspheres. In 24 dimensions, only 0.1% of your space can be filled with hyperspheres.
- For her work, Viazovska won the 2022 Fields medal. She was the second woman in history to win it.



School of Athens



School of Athens



People's History

People's History of Geometries

