




Chapter 8:

Modern Calculus



Two Early Contributors To The Mathematization of Science

Kepler

- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:




Kepler


- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.




Kepler




- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.



Kepler



- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.



Kepler

- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.



Kepler

- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.
 2. A line segment joining a planet and the Sun sweeps out equal areas during equal intervals of time.



KEPLER'S LAW OF PLANETARY MOTION

Kepler's Second Law

www.scientutts.com

SAMPLE USE ONLY

This is also known as the law of equal areas.

one

2. A line segment joining a planet and the Sun sweeps out equal areas during equal intervals of time.

KEPLER'S LAW OF PLANETARY MOTION

Kepler's Second Law

www.scientutts.com

SAMPLE USE ONLY

This is also known as the law of equal areas.



one


2. A line segment joining a planet and the Sun sweeps out equal areas during equal intervals of time.

Kepler

- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.
 2. A line segment joining a planet and the Sun sweeps out equal areas during equal intervals of time.





Kepler





- Following Copernicus' heliocentric theory of the solar system, Johannes Kepler devised three *laws of planetary motion*:
 1. The orbit of a planet is an ellipse with the Sun at one of the two foci.
 2. A line segment joining a planet and the Sun sweeps out equal areas during equal intervals of time.
 3. The square of the orbital period of a planet is directly proportional to the cube of the semi-major axis of its orbit.

Galileo




Galileo


- Galileo Galilei ran experiments to investigate objects in motion.




Galileo




- Galileo Galilei ran experiments to investigate objects in motion.
- Example: He rolled objects down an incline.




Galileo




- Galileo Galilei ran experiments to investigate objects in motion.
- Example: He rolled objects down an incline.




Galileo





- Galileo Galilei ran experiments to investigate objects in motion.
- Example: He rolled objects down an incline.



Galileo



- Galileo Galilei ran experiments to investigate objects in motion.
- Example: He rolled objects down an incline.
- He discovered that if it travels d inches in the first second, then it travels $3d$ inches in the next second, $5d$ inches in the third second, and so on.



Isaac
Newton



Gottfried
Wilhelm Leibniz



Newton

Newton

- Grew up in Woolsthorpe, Lincolnshire, England.

Newton

- Grew up in Woolsthorpe, Lincolnshire, England.
- Thought scientifically early on. Was destined to be a farmer, but lucky for us he was terrible at it.

Newton

- Grew up in Woolsthorpe, Lincolnshire, England.
- Thought scientifically early on. Was destined to be a farmer, but lucky for us he was terrible at it.



Newton

- Grew up in Woolsthorpe, Lincolnshire, England.
- Thought scientifically early on. Was destined to be a farmer, but lucky for us he was terrible at it.




Apple Tree!

Newton




Newton

- Went to Cambridge. Began pondering scientific foundations.




Newton

- Went to Cambridge. Began pondering scientific foundations.
- Cambridge hired Issac Burrow, their first math professor. Burrow examined Newton on Euclid's *Elements*.



Newton

- Went to Cambridge. Began pondering scientific foundations.
- Cambridge hired Issac Burrow, their first math professor. Burrow examined Newton on Euclid's *Elements*.
- Newton took the chance to study advanced math, and invented more. Spent his money on math books, and borrowed more.



Newton

Newton

- Newton made his first big math discovery. He generalized *the binomial theorem* to rational exponents.

Newton

- Newton made his first big math discovery. He generalized *the binomial theorem* to rational exponents.
- After his second year, Cambridge was closed due to the plague, for 20 months. His time at home was among the most productive in scientific history. This is called his “miracle year.”



Newton

- Newton made his first big math discovery. He generalized *the binomial theorem* to rational exponents.
- After his second year, Cambridge was closed due to the plague, for 20 months. His time at home was among the most productive in scientific history. This is called his “miracle year.”
- The 24 year old discovered his fundamental theories on gravity, motion and optics. He invented differential and integral calculus. He told no one.



Newton

Newton

- Newton recorded his work in a small notebook which he called his “waste book.”

Newton

- Newton recorded his work in a small notebook which he called his “waste book.”



Newton

- Newton recorded his work in a small notebook which he called his “waste book.”



Newton


- Newton recorded his work in a small notebook which he called his “waste book.”



Link to scan


On the Board: Isaac Newton's Fluxions and Fluents

Newton vs. Leibniz




Newton vs. Leibniz

- **Theorem (Newton).** Suppose $y = x^r$ where r is a rational number. Then, $\frac{\dot{y}}{\dot{x}} = rx^{r-1}$.



Newton vs. Leibniz


- **Theorem (Newton).** Suppose $y = x^r$ where r is a rational number. Then, $\frac{\dot{y}}{\dot{x}} = rx^{r-1}$.



- **Theorem (Leibniz).** Suppose u and v are fluents which depend on a variable t , and $w = uv$. Then, $\dot{w} = u\dot{v} + v\dot{u}$.


Newton vs. Leibniz

- **Theorem (Newton).** Suppose $y = x^r$ where r is a rational number. Then, $\frac{\dot{y}}{\dot{x}} = rx^{r-1}$.




- **Theorem (Leibniz).** Suppose u and v are fluents which depend on a variable t , and $w = uv$. Then, $\dot{w} = u\dot{v} + v\dot{u}$.

Newton vs. Leibniz




Newton vs. Leibniz

- It is likely that Leibniz independently invented the calculus. Their work was fundamentally the same, but they developed it in different ways. Their work *looked* quite different. Today, we use Leibniz's notation.



Newton vs. Leibniz


- It is likely that Leibniz independently invented the calculus. Their work was fundamentally the same, but they developed it in different ways. Their work *looked* quite different. Today, we use Leibniz's notation.



- Their countrymen defended them. A bitter rivalry ensued.

Newton vs. Leibniz

- It is likely that Leibniz independently invented the calculus. Their work was fundamentally the same, but they developed it in different ways. Their work *looked* quite different. Today, we use Leibniz's notation.



- Their countrymen defended them. A bitter rivalry ensued.
- Newton's work was better; Liebniz's notation was better. Both were superb.

Isaac Newton



Isaac Newton

- In the late 1660s and early 1670s, Newton slowly revealed some of his work. First on infinite series and then his reflecting telescope. He regretted it.



Isaac Newton

- In the late 1660s and early 1670s, Newton slowly revealed some of his work. First on infinite series and then his reflecting telescope. He regretted it.
- In 1684, Halley asked Newton how the planets moved. This set off his second intense period of inquiry, leading to his masterpiece: *Principia*.



Isaac Newton

Isaac Newton

- In the *Principia*, Newton:

Isaac Newton

- In the *Principia*, Newton:

-Develops calculus

Isaac Newton

- In the *Principia*, Newton:

- Develops calculus

- Laws of gravity

Isaac Newton

- In the *Principia*, Newton:

- Develops calculus

- Laws of gravity

- Derives Kepler's laws

Isaac Newton

- In the *Principia*, Newton:

- Develops calculus

- Laws of gravity

- Derives Kepler's laws

- Shows inverse law

Isaac Newton

- In the *Principia*, Newton:

- Develops calculus
- Laws of gravity
- Derives Kepler's laws
- Shows inverse law
- Estimates relative mass
of Sun and planets

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion
 - Explains irregularities of Moon's motion

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion
 - Explains irregularities of Moon's motion
 - IDs equilateral bulge

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion
 - Explains irregularities of Moon's motion
 - IDs equilateral bulge
 - Explains the tides

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion
 - Explains irregularities of Moon's motion
 - IDs equilateral bulge
 - Explains the tides
 - Explains comets paths

Isaac Newton

- In the *Principia*, Newton:
 - Develops calculus
 - Laws of gravity
 - Derives Kepler's laws
 - Shows inverse law
 - Estimates relative mass of Sun and planets
 - Explains Sun's motion
 - Explains irregularities of Moon's motion
 - IDs equilateral bulge
 - Explains the tides
 - Explains comets paths

Albert Einstein regarded Newton with highest regard.

Isaac Newton

Isaac Newton

- Random fact: His biographer says Newton's associates only ever saw him laugh once.

Isaac Newton

- Random fact: His biographer says Newton's associates only ever saw him laugh once.
- William Stukeley wrote a biography on Newton, published in 1752. He wrote (minor translation changes):

Isaac Newton

- Random fact: His biographer says Newton's associates only ever saw him laugh once.
- William Stukeley wrote a biography on Newton, published in 1752. He wrote (minor translation changes):

All the time he was with him, he never observed him to laugh; but once. He remembers, it was on this occasion.

He asked a friend, to whom he had lent a copy of Euclid's *Elements* to read, what progress he had made and how he liked it. [The friend] answered by desiring to know what use and benefit in life is that kind of study to him? Upon which Sr. Isaac was very merry.

Isaac Newton

Isaac Newton

- Newton also spent much time studying alchemy (transform metals into other metals), largely by collecting others' work and annotating it, or by running experiments. May have suffered from Mercury poisoning. He also studied theology. Almost all of this was only known after his death.

Isaac Newton

- Newton also spent much time studying alchemy (transform metals into other metals), largely by collecting others' work and annotating it, or by running experiments. May have suffered from Mercury poisoning. He also studied theology. Almost all of this was only known after his death.
- First scientist to receive a ceremonial funeral in Westminster Abbey. Inscription read:

Isaac Newton

- Newton also spent much time studying alchemy (transform metals into other metals), largely by collecting others' work and annotating it, or by running experiments. May have suffered from Mercury poisoning. He also studied theology. Almost all of this was only known after his death.
- First scientist to receive a ceremonial funeral in Westminster Abbey. Inscription read:

Here lies that which
was mortal of Isaac Newton

Poem:
Isaac Newton Alone

Maria Agnesi




Maria Agnesi

- The calculus now needed to be explained in an accessible way to the scientific community. This was challenging and important.




Maria Agnesi

- The calculus now needed to be explained in an accessible way to the scientific community. This was challenging and important.
- In 1748, Italian mathematician Maria Agnesi published the first unified treatment of algebra, coordinate geometry, differential calculus and integral calculus. Her book was widely praised.



Maria Agnesi

- The calculus now needed to be explained in an accessible way to the scientific community. This was challenging and important.
- In 1748, Italian mathematician Maria Agnesi published the first unified treatment of algebra, coordinate geometry, differential calculus and integral calculus. Her book was widely praised.
- She was the first woman to ever be appointed to a professorship in math.



Rigorizing The Calculus


Rigorizing The Calculus

- Turning “calculus” into “real analysis” is largely due to Austin-Luis Cauchy and Karl Weierstrass.



Rigorizing The Calculus


- Turning “calculus” into “real analysis” is largely due to Austin-Luis Cauchy and Karl Weierstrass.
- Cauchy thought in terms of limits. For example, he expressed $\lim_{x \rightarrow a} f(x) = L$ as



Rigorizing The Calculus

- Turning “calculus” into “real analysis” is largely due to Austin-Luis Cauchy and Karl Weierstrass.
- Cauchy thought in terms of limits. For example, he expressed $\lim_{x \rightarrow a} f(x) = L$ as


When the successively attributed values of the same variable indefinitely approach a fixed value, so that finally they differ from it by as little as desired, the last is called the *limit* of all the others.



Rigorizing The Calculus

- Turning “calculus” into “real analysis” is largely due to Austin-Luis Cauchy and Karl Weierstrass.
- Cauchy thought in terms of limits. For example, he expressed $\lim_{x \rightarrow a} f(x) = L$ as

When the successively attributed values of the same variable indefinitely approach a fixed value, so that finally they differ from it by as little as desired, the last is called the *limit* of all the others.



- Weierstrass: For all $\epsilon > 0$, there exists a $\delta > 0$ such that, if $0 < |x - a| < \delta$, then $|f(x) - L| < \epsilon$.

Infinite Series

Infinite Series

Infinite Series


- So far, only series that had been solved were geometric series.

Infinite Series

- So far, only series that had been solved were geometric series.
- Next one involves *triangular numbers*.


Infinite Series

- So far, only series that had been solved were geometric series.
- Next one involves *triangular numbers*.



Infinite Series

- So far, only series that had been solved were geometric series.
- Next one involves *triangular numbers*.



- Leibniz wanted to find $\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \dots$

Infinite Series

Infinite Series

- Leibniz found

$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \dots$$

Infinite Series

- Leibniz found

$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \dots$$

$$= (2 - 1) + \left(1 - \frac{2}{3}\right) + \left(\frac{2}{3} - \frac{2}{4}\right) + \left(\frac{2}{4} - \frac{2}{5}\right) + \left(\frac{2}{5} - \frac{2}{6}\right) + \dots$$

Infinite Series

- Leibniz found

$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \dots$$

$$= (2 - 1) + \left(1 - \frac{2}{3}\right) + \left(\frac{2}{3} - \frac{2}{4}\right) + \left(\frac{2}{4} - \frac{2}{5}\right) + \left(\frac{2}{5} - \frac{2}{6}\right) + \dots$$

$$= 2 - (1 - 1) - \left(\frac{2}{3} - \frac{2}{3}\right) - \left(\frac{2}{4} - \frac{2}{4}\right) - \left(\frac{2}{5} - \frac{2}{5}\right) - \left(\frac{2}{6} - \frac{2}{6}\right) - \dots$$

Infinite Series

- Leibniz found

$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \dots$$

$$= (2 - 1) + \left(1 - \frac{2}{3}\right) + \left(\frac{2}{3} - \frac{2}{4}\right) + \left(\frac{2}{4} - \frac{2}{5}\right) + \left(\frac{2}{5} - \frac{2}{6}\right) + \dots$$

$$= 2 - (1 - 1) - \left(\frac{2}{3} - \frac{2}{3}\right) - \left(\frac{2}{4} - \frac{2}{4}\right) - \left(\frac{2}{5} - \frac{2}{5}\right) - \left(\frac{2}{6} - \frac{2}{6}\right) - \dots$$

$$= 2.$$

Infinite Series

Infinite Series

- Johann Bernoulli used this result to prove that

$$\sum_{k=1}^{\infty} \frac{1}{k} = \infty .$$

This is called the *harmonic series*. Details in notes.

Infinite Series

- Johann Bernoulli used this result to prove that

$$\sum_{k=1}^{\infty} \frac{1}{k} = \infty .$$

This is called the *harmonic series*. Details in notes.

- This sum diverges suuuuper slowly. In 1968, J.W. Wrench Jr. showed that $\sum_{k=1}^n \frac{1}{k} > 100$ for the first time when

Infinite Series

- Johann Bernoulli used this result to prove that

$$\sum_{k=1}^{\infty} \frac{1}{k} = \infty .$$

This is called the *harmonic series*. Details in notes.

- This sum diverges suuuuper slowly. In 1968, J.W. Wrench Jr. showed that $\sum_{k=1}^n \frac{1}{k} > 100$ for the first time when

$$n = 15,092,688,622,113,788,323,693,563,264,538,101,449,859,497.$$

Infinite Series

Infinite Series

- Next big question: What is. $\sum_{k=1}^{\infty} \frac{1}{k^2}$?

Infinite Series

- Next big question: What is. $\sum_{k=1}^{\infty} \frac{1}{k^2}$?
- Since $\frac{1}{k^2} \leq \frac{2}{k(k+1)}$, the answer is *finite*.

formula for the
kth triangular
number

Infinite Series

- Next big question: What is. $\sum_{k=1}^{\infty} \frac{1}{k^2}$?
- Since $\frac{1}{k^2} \leq \frac{2}{k(k+1)}$, the answer is *finite*.

formula for the
kth triangular
number

Infinite Series

- Next big question: What is $\sum_{k=1}^{\infty} \frac{1}{k^2}$?
- Since $\frac{1}{k^2} \leq \frac{2}{k(k+1)}$, the answer is *finite*.
- Jacob Bernoulli tried but failed to solve it. He challenged the math community to find it, writing:

formula for the
kth triangular
number

Infinite Series

- Next big question: What is $\sum_{k=1}^{\infty} \frac{1}{k^2}$?
- Since $\frac{1}{k^2} \leq \frac{2}{k(k+1)}$, the answer is *finite*.
- Jacob Bernoulli tried but failed to solve it. He challenged the math community to find it, writing:

If anyone finds and communicates to us that which thus far has eluded our efforts, great will be our gratitude.

formula for the
kth triangular
number

Infinite Series

- Next big question: What is. $\sum_{k=1}^{\infty} \frac{1}{k^2}$?
- Since $\frac{1}{k^2} \leq \frac{2}{k(k+1)}$, the answer is *finite*.
- Jacob Bernoulli tried but failed to solve it. He challenged the math community to find it, writing:

If anyone finds and communicates to us that which thus far has eluded our efforts, great will be our gratitude.

- Jacob never solved it. But his younger brother Johann mentored the person who would.

Leonhard Euler



Leonhard Euler

- Born in Basel, Switzerland in 1707.
Student of Johann Bernoulli.



Leonhard Euler



- Born in Basel, Switzerland in 1707.
Student of Johann Bernoulli.
- Made a name for himself when at the age of 24 he proved

$$1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots =$$

Leonhard Euler



- Born in Basel, Switzerland in 1707.
Student of Johann Bernoulli.
- Made a name for himself when at
the age of 24 he proved

$$1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots = \frac{\pi^2}{6}$$

Leonhard Euler



- Born in Basel, Switzerland in 1707.
Student of Johann Bernoulli.

- Made a name for himself when at the age of 24 he proved

$$1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots = \frac{\pi^2}{6}$$

- Euler was amazing at determining how far his techniques would go.

Leonhard Euler



Leonhard Euler

- Here, he found the answer for
$$\sum_{k=1}^{\infty} \frac{1}{k^p}$$
for every even value of p from 2 to 26.



Leonhard Euler



- Here, he found the answer for
$$\sum_{k=1}^{\infty} \frac{1}{k^p}$$
for every even value of p from 2 to 26.

- He finished with

$$\sum_{k=1}^{\infty} \frac{1}{k^{26}} = \frac{1,315,862}{11,094,481,976,030,578,125} \pi^{26}.$$

Leonhard Euler



- Here, he found the answer for
$$\sum_{k=1}^{\infty} \frac{1}{k^p}$$
for every even value of p from 2 to 26.
- He finished with
$$\sum_{k=1}^{\infty} \frac{1}{k^{26}} = \frac{1,315,862}{11,094,481,976,030,578,125} \pi^{26}.$$
- He kick-started many new areas of math, such as graph theory and topology.

Leonhard Euler



Leonhard Euler

- He made major contributions to real analysis, complex analysis, number theory, combinatorics, applied math and physics.



Leonhard Euler



- He made major contributions to real analysis, complex analysis, number theory, combinatorics, applied math and physics.
- He is the productive mathematician in history. He published about 800 pages of math work per year. Publishing his collective works is still ongoing, and is currently 92 volumes long (about 35,000 pages). During the 18th century, among all published math, he was responsible for a *third* of it.

Leonhard Euler



- He made major contributions to real analysis, complex analysis, number theory, combinatorics, applied math and physics.
- He is the productive mathematician in history. He published about 800 pages of math work per year. Publishing his collective works is still ongoing, and is currently 92 volumes long (about 35,000 pages). During the 18th century, among all published math, he was responsible for a *third* of it.
- *When he went blind, he became more productive.*

Leonhard Euler



Leonhard Euler

Another Euler result: Recall the Taylor Series:



Leonhard Euler

Another Euler result: Recall the Taylor Series:

$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$



Leonhard Euler

Another Euler result: Recall the Taylor Series:

$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$



Leonhard Euler

Another Euler result: Recall the Taylor Series:



$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \dots$$

Leonhard Euler



Another Euler result: Recall the Taylor Series:

$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \dots$$

And this works for imaginary numbers too!

Leonhard Euler

Another Euler result: Recall the Taylor Series:



$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \dots$$

And this works for imaginary numbers too!

$$i^2 = -1 \quad , \quad i^3 = -i \quad , \quad i^4 = 1 \quad , \quad i^5 = i$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$e^{ix} = 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$\begin{aligned} e^{ix} &= 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots \\ &= 1 + ix + \frac{i^2 x^2}{2!} + \frac{i^3 x^3}{3!} + \frac{i^4 x^4}{4!} + \frac{i^5 x^5}{5!} + \frac{i^6 x^6}{6!} + \frac{i^7 x^7}{7!} + \dots \end{aligned}$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

$$i^5 = i$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$e^{ix} = 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots$$

$$= 1 + ix + \frac{i^2 x^2}{2!} + \frac{i^3 x^3}{3!} + \frac{i^4 x^4}{4!} + \frac{i^5 x^5}{5!} + \frac{i^6 x^6}{6!} + \frac{i^7 x^7}{7!} + \dots$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

$$i^5 = i$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$e^{ix} = 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots$$

$$= 1 + ix + \frac{i^2 x^2}{2!} + \frac{i^3 x^3}{3!} + \frac{i^4 x^4}{4!} + \frac{i^5 x^5}{5!} + \frac{i^6 x^6}{6!} + \frac{i^7 x^7}{7!} + \dots$$

$$= 1 + ix - \frac{x^2}{2!} - \frac{ix^3}{3!} + \frac{x^4}{4!} + \frac{ix^5}{5!} - \frac{x^6}{6!} - \frac{ix^7}{7!} + \dots$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

$$i^5 = i$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$\begin{aligned} e^{ix} &= 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots \\ &= 1 + ix + \frac{i^2 x^2}{2!} + \frac{i^3 x^3}{3!} + \frac{i^4 x^4}{4!} + \frac{i^5 x^5}{5!} + \frac{i^6 x^6}{6!} + \frac{i^7 x^7}{7!} + \dots \\ &= 1 + ix - \frac{x^2}{2!} - \frac{ix^3}{3!} + \frac{x^4}{4!} + \frac{ix^5}{5!} - \frac{x^6}{6!} - \frac{ix^7}{7!} + \dots \\ &= (1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots) + (ix - \frac{ix^3}{3!} + \frac{ix^5}{5!} - \frac{ix^7}{7!} + \dots) \end{aligned}$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

$$i^5 = i$$

Leonhard Euler



Plugging $x = i\theta$ into the e^x Taylor series:

$$e^{ix} = 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots$$

$$= 1 + ix + \frac{i^2 x^2}{2!} + \frac{i^3 x^3}{3!} + \frac{i^4 x^4}{4!} + \frac{i^5 x^5}{5!} + \frac{i^6 x^6}{6!} + \frac{i^7 x^7}{7!} + \dots$$

$$= 1 + ix - \frac{x^2}{2!} - \frac{ix^3}{3!} + \frac{x^4}{4!} + \frac{ix^5}{5!} - \frac{x^6}{6!} - \frac{ix^7}{7!} + \dots$$

$$= \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots\right) + \left(ix - \frac{ix^3}{3!} + \frac{ix^5}{5!} - \frac{ix^7}{7!} + \dots\right)$$

$$= \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

Recall:

$$\cos x = 1 - \frac{x}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Plugging in $x = \pi$,

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Plugging in $x = \pi$,

$$e^{i\pi} = \cos \pi + i \sin \pi$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Plugging in $x = \pi$,

$$e^{i\pi} = \cos \pi + i \sin \pi$$

$$e^{i\pi} = -1 + i \cdot 0$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Plugging in $x = \pi$,

$$e^{i\pi} = \cos \pi + i \sin \pi$$

$$e^{i\pi} = -1 + i \cdot 0$$

$$e^{i\pi} = -1$$

Leonhard Euler

$$e^{ix} = \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots\right) + i\left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots\right)$$

$$e^{ix} = \cos x + i \sin x$$

Plugging in $x = \pi$,

$$e^{i\pi} = \cos \pi + i \sin \pi$$

$$e^{i\pi} = -1 + i \cdot 0$$

$$e^{i\pi} = -1$$

$$\boxed{e^{i\pi} + 1 = 0}$$

Leonhard Euler

One more gem from Euler:



Leonhard Euler

One more gem from Euler:







- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

Leonhard Euler

One more gem from Euler:






- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:






Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices				
Edges				
Faces				





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4			
Edges				
Faces				





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4			
Edges	6			
Faces				




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4			
Edges	6			
Faces	4			




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8		
Edges	6			
Faces	4			




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8		
Edges	6	12		
Faces	4			





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8		
Edges	6	12		
Faces	4	6		





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	
Edges	6	12		
Faces	4	6		




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	
Edges	6	12	12	
Faces	4	6		





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	
Edges	6	12	12	
Faces	4	6	8	





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	
Faces	4	6	8	





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20
$V - E + F$				




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20
$V - E + F$	2			





Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20
$V - E + F$	2	2		




Leonhard Euler

- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20
$V - E + F$	2	2	2	2

Leonhard Euler


- *Polyhedra* are 3D objects comprised of faces which meet at edges, and edges which meet at vertices. Euler noticed a property in their counts:

				
Vertices	4	8	6	12
Edges	6	12	12	30
Faces	4	6	8	20
$V - E + F$	2	2	2	2

Bernoulli Family

Bernoulli Family

Bernouli Family Tree



Srinivasa Ramanujan

Srinivasa Ramanujan

- Ramanujan was a devout Hindu, with particular devotion to his family deity, the Goddess Namagiri. His family was Brahmin—the top of the caste system—but was poor.



Srinivasa Ramanujan


- Ramanujan was a devout Hindu, with particular devotion to his family deity, the Goddess Namagiri. His family was Brahmin—the top of the caste system—but was poor.
- At age 16, he got a copy of *A Synopsis of Elementary Results in Pure and Applied Mathematics*, which is a book of ~5,000 math facts/theorems, with essentially no proofs or justification.



Srinivasa Ramanujan


Srinivasa Ramanujan

- He neglected his non-math classes and flunked out of college. Instead, he worked on his *notebooks*—in which he wrote his mathematical discoveries.



Srinivasa Ramanujan

- He neglected his non-math classes and flunked out of college. Instead, he worked on his *notebooks*—in which he wrote his mathematical discoveries.
- For 5 years he had lots of time to work and no collaborators or math teachers. They allowed him to blaze his own trail in math.



Srinivasa Ramanujan

Srinivasa Ramanujan

- He was also a college dropout who was living with his parents.

Srinivasa Ramanujan

- He was also a college dropout who was living with his parents.
- His mom's solution: An arranged marriage. Ramanujan married a girl named Janaki, but they wouldn't live together for several years.

Srinivasa Ramanujan

- He was also a college dropout who was living with his parents.
- His mom's solution: An arranged marriage. Ramanujan married a girl named Janaki, but they wouldn't live together for several years.
- Ramanujan traveled to Madras where a patron supported him while he did math, including by publishing his first paper.

Srinivasa Ramanujan

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$3 = \sqrt{9}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$3 = \sqrt{9} = \sqrt{1 + 8}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$3 = \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$3 = \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$3 = \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned}3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\&= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}}\end{aligned}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned}3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\&= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{25}}}\end{aligned}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned} 3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\ &= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{25}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 24}}} \end{aligned}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned} 3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\ &= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{25}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 24}}} \\ &= \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4 \cdot 6}}} \end{aligned}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned} 3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\ &= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{25}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 24}}} \\ &= \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4 \cdot 6}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{36}}}} = \dots, \end{aligned}$$

Srinivasa Ramanujan

- He also published a problem for readers of the Journal of the Indian Mathematical Society. He asked for the value of $\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}}$
- His answer: 3. Reasoning:

$$\begin{aligned} 3 &= \sqrt{9} = \sqrt{1 + 8} = \sqrt{1 + 2 \cdot 4} = \sqrt{1 + 2\sqrt{16}} = \sqrt{1 + 2\sqrt{1 + 15}} \\ &= \sqrt{1 + 2\sqrt{1 + 3 \cdot 5}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{25}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 24}}} \\ &= \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4 \cdot 6}}} = \sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{36}}}} = \dots, \end{aligned}$$

- Because:

$$\sqrt{n^2} = \sqrt{1 + (n^2 - 1)} = \sqrt{1 + (n - 1)(n + 1)} = \sqrt{1 + (n - 1)\sqrt{(n + 1)^2}}$$

Srinivasa Ramanujan

Srinivasa Ramanujan

- Ramanujan needed support that he wasn't able to get from India.

Srinivasa Ramanujan

- Ramanujan needed support that he wasn't able to get from India.
- In 1913, he wrote to Cambridge Professor G. H. Hardy, one of the top mathematicians in Europe.

Srinivasa Ramanujan

- Ramanujan needed support that he wasn't able to get from India.
- In 1913, he wrote to Cambridge Professor G. H. Hardy, one of the top mathematicians in Europe.

Madras, 16th January 1913,
Dear Sir,

I beg to introduce myself to you as a clerk in the Accounts Department of the Port Trust Office at Madras on a salary of only £20 per annum. I am now about 23 years of age. I have no University education but I have undergone the ordinary school course. After leaving school I have been employing the spare time at my disposal to work at Mathematics. I have not trodden through the conventional regular course followed in a University course, but I am striking out a new path for myself. I have made special investigation of divergent series in general and the results I get are termed by the local mathematicians as "startling."

Srinivasa Ramanujan

Srinivasa Ramanujan

- The letter also contained 9 pages of theorems, but no proofs. For example, Ramanujan claimed that if

$$u = \cfrac{x}{1 + \cfrac{x^5}{1 + \cfrac{x^{10}}{1 + \cfrac{x^{15}}{1 + \dots}}}}$$

and

$$v = \cfrac{x^{1/5}}{1 + \cfrac{x}{1 + \cfrac{x^2}{1 + \cfrac{x^3}{1 + \dots}}}},$$

Srinivasa Ramanujan

- The letter also contained 9 pages of theorems, but no proofs. For example, Ramanujan claimed that if

$$u = \cfrac{x}{1 + \cfrac{x^5}{1 + \cfrac{x^{10}}{1 + \cfrac{x^{15}}{1 + \dots}}}}$$

and

$$v = \cfrac{x^{1/5}}{1 + \cfrac{x}{1 + \cfrac{x^2}{1 + \cfrac{x^3}{1 + \dots}}}},$$

then $v^5 = u \frac{1 - 2u + 4u^2 - 3u^3 + u^4}{1 + 3u + 4u^2 + 2u^3 + u^4}.$

Srinivasa Ramanujan

- The letter also contained 9 pages of theorems, but no proofs. For example, Ramanujan claimed that if

$$u = \cfrac{x}{1 + \cfrac{x^5}{1 + \cfrac{x^{10}}{1 + \cfrac{x^{15}}{1 + \dots}}}}$$

and

$$v = \cfrac{x^{1/5}}{1 + \cfrac{x}{1 + \cfrac{x^2}{1 + \cfrac{x^3}{1 + \dots}}}},$$

then $v^5 = u \frac{1 - 2u + 4u^2 - 3u^3 + u^4}{1 + 3u + 4u^2 + 2u^3 + u^4}$.

- Hardy: “They must be true because, if they were not true, no one would have had the imagination to invent them.”

Srinivasa Ramanujan



Srinivasa Ramanujan

- Hardy wanted to bring Ramanujan to Cambridge. Initially Ramanujan resisted for religious reasons, but in 1914 he went.



Srinivasa Ramanujan

- Hardy wanted to bring Ramanujan to Cambridge. Initially Ramanujan resisted for religious reasons, but in 1914 he went.
- They had a very productive few years, making advances on highly composite numbers, the partition function, and much more.



Srinivasa Ramanujan

Srinivasa Ramanujan


- When WWI started, Ramanujan's health deteriorated. Was treated for tuberculosis. Once tried to commit suicide.

Srinivasa Ramanujan

- When WWI started, Ramanujan's health deteriorated. Was treated for tuberculosis. Once tried to commit suicide.
- He published 9 papers in 1915, earning the reputation he desired.

Srinivasa Ramanujan

- When WWI started, Ramanujan's health deteriorated. Was treated for tuberculosis. Once tried to commit suicide.
- He published 9 papers in 1915, earning the reputation he desired.
- He was elected a Fellow of the Royal Society, a tremendous honor.



Srinivasa Ramanujan

Srinivasa Ramanujan

- In 1919, he returned to India. He was widely celebrated for his work and being named an Fellow of the Royal Society.

Srinivasa Ramanujan

- In 1919, he returned to India. He was widely celebrated for his work and being named an Fellow of the Royal Society.
- The following year he died, at the age of 32.

Srinivasa Ramanujan

- In 1919, he returned to India. He was widely celebrated for his work and being named an Fellow of the Royal Society.
- The following year he died, at the age of 32.
- Mathematicians marvel at his work, which feels otherworldly. Example:

Srinivasa Ramanujan

- In 1919, he returned to India. He was widely celebrated for his work and being named an Fellow of the Royal Society.
- The following year he died, at the age of 32.
- Mathematicians marvel at his work, which feels otherworldly. Example:


$$\frac{1}{\pi} = \frac{2\sqrt{2}}{9801} \sum_{k=0}^{\infty} \frac{(4k)!(1103 + 26390k)}{(k!)^4 396^{4k}}$$

Poem: Joined By Math


Who Is The GMOAT?

The Contenders


The Contenders




The Contenders




The Contenders




The Contenders



The Contenders



The Contenders



Infinities

Hilbert's overly-accommodating Infinity Hotel





“No other question has ever moved so profoundly the spirit of mankind; no other idea has so fruitfully stimulated our intellect; yet no other concept stands in greater need of clarification than that of the infinite.”

—David Hilbert

1

2

3

4

5

6

7

...



1

2

3


4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3




4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3


4

5

6

7

...



1

2

3


4

5

6

7

...



Person in room $n \rightarrow$ room $n + 1$



1

2

3


4

5


6

7

...



Person in room $n \rightarrow$ room $n + 1$
 $n \rightarrow n + 1$



1

2

3


4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3


4

5

6

7

...



1

2

3


4

5

6

7

...



1

2

3


4

5

6

7

...



$$n \rightarrow n + 2$$



1

2

3


4

5

6

7

...



$$n \rightarrow n + 2$$



1

2

3


4

5

6

7

...



$$n \rightarrow n + 2$$



1

2

3


4

5

6

7

...



1

2

3



4

5

6

7

...



1

2

3


4

5


6

7

...



Room for
all of us?



1

2

3



4

5

6

7

...



1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Sure!



Room for
all of us?



1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Sure!



Room for
all of us?



...

1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Sure!



Room for
all of us?



1

2

3


4

5



6

7

...



$$n \rightarrow 2n$$



1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Sure!



Room for
all of us?



...

1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Sure!



Room for
all of us?



1

2

3


4

5

6

7

...



$$n \rightarrow 2n$$

Hilbert's
overly-accommodating
Infinity Hotel



Sure!



Room for
all of us?

...

1

2

3


4

5

6

7

...




$$n \rightarrow 2n$$



Room for
all of us?

Moral: ∞ is weird

What does size mean?



*“In mathematics the art
of proposing a question
must be held of higher
value than solving it.”*

—Georg Cantor

What does size mean?

What does size mean?

- Until the mid 1800s, infinity was widely treated carelessly or with suspicion.

What does size mean?

- Until the mid 1800s, infinity was widely treated carelessly or with suspicion.
- Example: Carl Friedrich Gauss in 1831 wrote

What does size mean?

- Until the mid 1800s, infinity was widely treated carelessly or with suspicion.
- Example: Carl Friedrich Gauss in 1831 wrote

I protest above all against the use of infinite quantity as something *completed*, which in mathematics is never allowed. The Infinite is only a manner of speaking.

What does size mean?

- Until the mid 1800s, infinity was widely treated carelessly or with suspicion.
- Example: Carl Friedrich Gauss in 1831 wrote

I protest above all against the use of infinite quantity as something *completed*, which in mathematics is never allowed. The Infinite is only a manner of speaking.

- It was made precise Georg Cantor (1845-1918). Who was a PhD student of Weierstrass.

What does size mean?

What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:


What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:

$$\{1,2,3\} \quad \{a,b,c\}$$

What does size mean?


- What does it mean for two things to have the same size? Cantor's definition:

$$\{1,2,3\} \quad \{a,b,c\}$$


What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:


$$\{ \ ,2,3\} \qquad \{ \ ,b,c\}$$



What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:

$$\{ \ , \ , 3 \} \qquad \{ \ , \ , c \}$$



What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:

$$\{ , , \} \qquad \{ , , \}$$

$$\begin{array}{ccc} 1 & \longleftrightarrow & a \\ 2 & \longleftrightarrow & b \\ 3 & \longleftrightarrow & c \end{array}$$

What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:

$$\{ , , \} \qquad \{ , , \}$$

$$\begin{array}{ccc} 1 & \longleftrightarrow & b \\ 2 & \longleftrightarrow & a \\ 3 & \longleftrightarrow & c \end{array}$$

What does size mean?

- What does it mean for two things to have the same size? Cantor's definition:

$$\{ , , \} \qquad \{ , , \}$$

$$\begin{array}{ccc} 1 & \longleftrightarrow & b \\ 2 & \longleftrightarrow & a \\ 3 & \longleftrightarrow & c \end{array}$$

- Sets are the same size if we can “pair up” the elements

Fact. There are the same number of positive integers as negative integers

Fact. There are the same number of positive integers as negative integers

Proof.

Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.

Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.


$$\{-1, -2, -3, -4, -5, \dots\} \quad \{1, 2, 3, 4, 5, \dots\}$$

Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.


Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.





Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.





Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.





Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.





Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.




Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.



Fact. There are the same number of positive integers as negative integers

Proof. We need to find a way to pair them all up.



Fact. There are the same number of positive integers as positive *even* integers

Fact. There are the same number of positive integers as positive even integers

Proof.

Fact. There are the same number of positive integers as positive *even* integers

Proof. We need to find a way to pair them all up.

Fact. There are the same number of positive integers as positive *even* integers

Proof. We need to find a way to pair them all up.

$$\{1, 2, 3, 4, 5, \dots\}$$

Fact. There are the same number of positive integers as positive *even* integers


Proof. We need to find a way to pair them all up.

$$\{1, 2, 3, 4, 5, \dots\} \quad \{2, 4, 6, 8, 10, \dots\}$$

Fact. There are the same number of positive integers as positive *even* integers

Proof. We need to find a way to pair them all up.


$$\{1, 2, 3, 4, 5, \dots\} \quad \{2, 4, 6, 8, 10, \dots\}$$



Fact. There are the same number of positive integers as positive *even* integers

Proof. We need to find a way to pair them all up.


$$\{ \ , 2, 3, 4, 5, \dots \} \quad \{ \ , 4, 6, 8, 10, \dots \}$$



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.


$$\{ \ , \ , 3, 4, 5, \dots \} \quad \{ \ , \ , 6, 8, 10, \dots \}$$



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.


$$\{ \ , \ , \ , 4, 5, \dots \} \quad \{ \ , \ , \ , 8, 10, \dots \}$$



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.


$$\{ \ , \ , \ , \ , \ , 5, \dots \} \quad \{ \ , \ , \ , \ , \ , 10, \dots \}$$



Fact. There are the same number of positive integers as positive even integers


Proof. We need to find a way to pair them all up.

$$\{ , , , , , \dots \} \quad \{ , , , , , \dots \}$$



Fact. There are the same number of positive integers as positive even integers


Proof. We need to find a way to pair them all up.



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.


$$\{ , , , , , \dots \} \quad \{ , , , , , \dots \}$$



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.

$$\{ , , , , , \dots \} \quad \{ , , , , , \dots \}$$



Fact. There are the same number of positive integers as positive *even* integers

Proof. We need to find a way to pair them all up.

Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.

1

2

3

4

5

6

7

...



Fact. There are the same number of positive integers as positive even integers

Proof. We need to find a way to pair them all up.

1

2

3

4

5

6

7

...

2

4

6


8

10

12

14

...



Different sizes of ∞ ?

Different sizes of ∞ ?

Theorem (Cantor).

There are the same number of positive integers as rational numbers.

Different sizes of ∞ ?

Theorem (Cantor).

There are the same number of positive integers as rational numbers.


Proof. Pair up the n th positive integer with the n th new rational number in this order:

Different sizes of ∞ ?

Theorem (Cantor).

There are the same number of positive integers as rational numbers.

Proof. Pair up the n th positive integer with the n th new rational number in this order:



Different sizes of ∞ ?

- Sets are *different* sizes if we *can't* pair up the elements

Different sizes of ∞ ?

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers than natural numbers.

Different sizes of ∞ ?

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers than natural numbers.


Proof. By contradiction — we assume we can pair them up.

Different sizes of ∞ ?

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers than natural numbers.

Proof. By contradiction — we assume we can pair them up.




Different sizes of ∞ ?

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers than natural numbers.

Proof. By contradiction — we assume we can pair them up.



- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

Proof. Assume for a contradiction that we *can* pair these up.

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.7 4 9 6 8 2 6 5 9 ...
2	0.4 8 6 8 0 3 6 1 3 ...
3	0.9 4 6 3 2 6 4 6 8 ...
4	0.1 3 2 4 5 6 4 6 7 ...
5	0.2 4 5 3 5 7 8 4 5 ...
6	0.3 3 4 5 4 3 8 8 3 ...
7	0.8 4 6 6 4 2 5 3 7 ...
8	0.3 3 5 1 3 7 4 7 9 ...
9	0.0 0 9 8 3 1 4 0 3 ...
:	:

Proof. Assume for a contradiction that we *can* pair these up.

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.7 4 9 6 8 2 6 5 9 ...
2	0.4 8 6 8 0 3 6 1 3 ...
3	0.9 4 6 3 2 6 4 6 8 ...
4	0.1 3 2 4 5 6 4 6 7 ...
5	0.2 4 5 3 5 7 8 4 5 ...
6	0.3 3 4 5 4 3 8 8 3 ...
7	0.8 4 6 6 4 2 5 3 7 ...
8	0.3 3 5 1 3 7 4 7 9 ...
9	0.0 0 9 8 3 1 4 0 3 ...
:	:

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659 ...
2	0.486803613 ...
3	0.946326468 ...
4	0.132456467 ...
5	0.245357845 ...
6	0.334543883 ...
7	0.846642537 ...
8	0.3351374079 ...
9	0.009831403 ...
:	:

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

	0.	
1	0.749682659...	
2	0.486803613...	
3	0.946326468...	
4	0.132456467...	
5	0.245357845...	
6	0.334543883...	
7	0.846642537...	
8	0.3351374079...	
9	0.009831403...	
:	:	

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.7	4	9	6	8	2	6	5	9	...
2	0.4	8	6	8	0	3	6	1	3	...
3	0.9	4	6	3	2	6	4	6	8	...
4	0.1	3	2	4	5	6	4	6	7	...
5	0.2	4	5	3	5	7	8	4	5	...
6	0.3	3	4	5	4	3	8	8	3	...
7	0.8	4	6	6	4	2	5	3	7	...
8	0.3	3	5	1	3	7	4	7	9	...
9	0.0	0	9	8	3	1	4	0	3	...
:						:				
	0.3	1	2							

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.7	4	9	6	8	2	6	5	9	...
2	0.4	8	6	8	0	3	6	1	3	...
3	0.9	4	6	3	2	6	4	6	8	...
4	0.1	3	2	4	5	6	4	6	7	...
5	0.2	4	5	3	5	7	8	4	5	...
6	0.3	3	4	5	4	3	8	8	3	...
7	0.8	4	6	6	4	2	5	3	7	...
8	0.3	3	5	1	3	7	4	7	9	...
9	0.0	0	9	8	3	1	4	0	3	...
:						:				
	0.3	1	2	2						

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.7	4	9	6	8	2	6	5	9	...
2	0.4	8	6	8	0	3	6	1	3	...
3	0.9	4	6	3	2	6	4	6	8	...
4	0.1	3	2	4	5	6	4	6	7	...
5	0.2	4	5	3	5	7	8	4	5	...
6	0.3	3	4	5	4	3	8	8	3	...
7	0.8	4	6	6	4	2	5	3	7	...
8	0.3	3	5	1	3	7	4	7	9	...
9	0.0	0	9	8	3	1	4	0	3	...
:						:				
	0.3	1	2	2	8	2	4	3	8	...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

0.312282438...

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

	0.749682659...
1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.3351374079...
9	0.009831403...
:	:
1,240,449	0.312282438...
:	:

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
⋮	⋮
1,240,449	0.009831403...
⋮	⋮
	0.312282438...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
:	:
1,240,449	0.009831403...
:	:
	0.312282438...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
:	:
1,240,449	0.009831403...
:	:
	0.312282438...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

	1	0.749682659...
2	0.486803613...	
3	0.946326468...	
4	0.132456467...	
5	0.245357845...	
6	0.334543883...	
7	0.846642537...	
8	0.335137479...	
9	0.009831403...	
	⋮	⋮
1,240,449	0.009831403...	
	⋮	⋮

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
⋮	⋮
1,240,449	0.009831403...
⋮	⋮
	0.312282438...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

1,240,449th digit
?

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
⋮	⋮
1,240,449	0.009831403...
⋮	⋮

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

1,240,449th digit

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

	1	0.749682659...	
2	0.486803613...		
3	0.946326468...		
4	0.132456467...		
5	0.245357845...		
6	0.334543883...		
7	0.846642537...		
8	0.335137479...		
9	0.009831403...		
:	:		
1,240,449	0.009831403.....?		
:	:		
	0.312282438.....?		

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

1,240,449th digit

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

	1	0.749682659...	
2	0.486803613...		
3	0.946326468...		
4	0.132456467...		
5	0.245357845...		
6	0.334543883...		
7	0.846642537...		
8	0.335137479...		
9	0.009831403...		
:	:		
1,240,449	0.009831403.....3...		
:	:		
	0.312282438.....?		

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

1,240,449th digit

- Sets are *different* sizes if we *can't* pair up the elements

Theorem (Cantor). There is a greater number of real numbers between 0 and 1 than positive integers.

1	0.749682659...
2	0.486803613...
3	0.946326468...
4	0.132456467...
5	0.245357845...
6	0.334543883...
7	0.846642537...
8	0.335137479...
9	0.009831403...
⋮	⋮
1,240,449	0.009831403...
⋮	⋮
	0.312282438...

Proof. Assume for a contradiction that we *can* pair these up.

Build a new number between 0 and 1. Its n th digit is anything besides 0, 9 or the n th digit of the n th number in this table

 1,240,449th digit

The Power Set

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

$$1 \longleftrightarrow \{1, 3, 27\}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

$$\begin{array}{ccc} 1 & \longleftrightarrow & \{1, 3, 27\} \\ 2 & \longleftrightarrow & \{4\} \end{array}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

$$\begin{array}{ccc} 1 & \longleftrightarrow & \{1, 3, 27\} \\ 2 & \longleftrightarrow & \{4\} \end{array}$$

$$3 \longleftrightarrow \{22, 101\}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

$$1 \longleftrightarrow \{1, 3, 27\}$$

$$2 \longleftrightarrow \{4\}$$

$$3 \longleftrightarrow \{22, 101\}$$

$$4 \longleftrightarrow \{3, 4, 5\}$$

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$


$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$



The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$



Call a number n *selfish* if n is in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$

selfish

$$1 \longleftrightarrow \{1, 3, 27\}$$

$$2 \longleftrightarrow \{4\}$$

:

$$3 \longleftrightarrow \{22, 101\}$$

$$4 \longleftrightarrow \{3, 4, 5\}$$

:

Call a number n *selfish* if n is in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$




Call a number n *selfish* if n is in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$




Call a number n *selfish* if n is in the set its paired with.

Call a number n *unselfish* if n is not in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$




Call a number n *selfish* if n is in the set its paired with.

Call a number n *unselfish* if n is not in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$




Call a number n *selfish* if n is in the set its paired with.

Call a number n *unselfish* if n is not in the set its paired with.

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$



Call a number n *selfish* if n is in the set its paired with.


Call a number n *unselfish* if n is not in the set its paired with.

Let D be the set of all unselfish numbers. Assume n is paired with D . Question: Is n selfish?

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$



Call a number n *selfish* if n is in the set its paired with.

Call a number n *unselfish* if n is not in the set its paired with.


Let D be the set of all unselfish numbers. Assume n is paired with D . Question: Is n selfish?

- If n is selfish then n is in D — contradiction to definition of D !

The Power Set

$$\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$$

$$\text{PowerSet}(\mathbb{N}) = \{\{1, 2, 3\}, \{2\}, \{543, 235\}, \{2, 4, 6, 8, 10, \dots\}, \dots\}$$



Call a number n *selfish* if n is in the set its paired with.

Call a number n *unselfish* if n is not in the set its paired with.

Let D be the set of all unselfish numbers. Assume n is paired with D . Question: Is n selfish?

- If n is selfish then n is in D — contradiction to definition of D !
- If n is unselfish then n is not in D — same contradiction!

The Power Set

Fact. The power set of any set is bigger than that set.


The Power Set

Fact. The power set of any set is bigger than that set.

\mathbb{N}


The Power Set

Fact. The power set of any set is bigger than that set.




The Power Set

Fact. The power set of any set is bigger than that set.




The Power Set

Fact. The power set of any set is bigger than that set.




The Power Set

Fact. The power set of any set is bigger than that set.




The Power Set

Fact. The power set of any set is bigger than that set.




The Power Set

Fact. The power set of any set is bigger than that set.



The Power Set

Fact. The power set of any set is bigger than that set.



The Continuum Hypothesis

The Continuum Hypothesis

- In Chapter 4 we discussed *undecidable statements*, which are statements that (within your axiomatic system) can neither be proven nor disproven.

The Continuum Hypothesis

- In Chapter 4 we discussed *undecidable statements*, which are statements that (within your axiomatic system) can neither be proven nor disproven.

Question: Is there an infinity between the number of natural numbers and the number of real numbers?

The Continuum Hypothesis

- In Chapter 4 we discussed *undecidable statements*, which are statements that (within your axiomatic system) can neither be proven nor disproven.

Question: Is there an infinity between the number of natural numbers and the number of real numbers?

The continuum hypothesis claims that the answer is ‘no’

The Continuum Hypothesis

- In Chapter 4 we discussed *undecidable statements*, which are statements that (within your axiomatic system) can neither be proven nor disproven.

Question: Is there an infinity between the number of natural numbers and the number of real numbers?

The continuum hypothesis claims that the answer is ‘no’


- The continuum hypothesis is undecidable in ZFC. That is, its truth value is independent of the standard axioms of math.

Cantor




Cantor

- Cantor struggled with mental illness. These were worsened due to the harsh criticism he received for his work on infinities.




Cantor

- Cantor struggled with mental illness. These were worsened due to the harsh criticism he received for his work on infinities.
- Poincaré called it a “grave disease,” Kronecker said Cantor was a “corrupter of youth.” Some Christians objected to it as challenging “God’s exclusive claim to supreme infinity.”



Cantor


- Cantor struggled with mental illness. These were worsened due to the harsh criticism he received for his work on infinities.
- Poincaré called it a “grave disease,” Kronecker said Cantor was a “corrupter of youth.” Some Christians objected to it as challenging “God’s exclusive claim to supreme infinity.”
- Today, his legacy is entirely secured. Hilbert:



Cantor

- Cantor struggled with mental illness. These were worsened due to the harsh criticism he received for his work on infinities.
- Poincaré called it a “grave disease,” Kronecker said Cantor was a “corrupter of youth.” Some Christians objected to it as challenging “God’s exclusive claim to supreme infinity.”
- Today, his legacy is entirely secured. Hilbert:

“No one shall expel us from the paradise that Cantor has created.”



Shout-outs!

Shout-outs!

- The person most responsible for bringing the calculus to France was Émilie du Châtelet. Her translation was richly annotated and is still the standard French translation today.

Shout-outs!

- The person most responsible for bringing the calculus to France was Émilie du Châtelet. Her translation was richly annotated and is still the standard French translation today.
- She also included in it the addition of a new law of physics: The law of conservation of total energy.



Shout-outs!

Shout-outs!

- Another group of historically-significant women who applied calculus to great effect was recently portrayed in the book and movie *Hidden Figures*.

Shout-outs!

- Another group of historically-significant women who applied calculus to great effect was recently portrayed in the book and movie *Hidden Figures*.



Katherine G. Johnson-1983-NASA



Dorothy J. Vaughan-NASA



Mary W. Jackson– NASA

Shout-outs!

Shout-outs!

- While discussing Newton and Einstein, should also discuss Carl Friedrich Bass, who discovered the existence of non-Euclidean geometries (which led to tensor calculus and eventually to Einstein's theory of general relativity). Einstein's theories assert that the natural universe is non-Euclidean.



Shout-outs!

- While discussing Newton and Einstein, should also discuss Carl Friedrich Bass, who discovered the existence of non-Euclidean geometries (which led to tensor calculus and eventually to Einstein's theory of general relativity). Einstein's theories assert that the natural universe is non-Euclidean.



Paul Dirac quote: “The mathematician plays a game in which he himself invents the rules while the physicist plays a game in which the rules are provided by nature, but as time goes on it becomes increasingly evident that the rules which the mathematician finds interesting are the same as those which nature has chosen.”


Shout-outs!

Shout-outs!

- An important problem in real analysis is the study of how heat diffuses through a region. Jean-Baptiste Joseph Fourier developed *Fourier analysis* to study this.

Shout-outs!

- An important problem in real analysis is the study of how heat diffuses through a region. Jean-Baptiste Joseph Fourier developed *Fourier analysis* to study this.
- Bernhard Riemann (1826-1866) also made important contributions to Fourier analysis, but he is best remembered in real analysis for being the first to construct the modern, rigorous formulation of the integral, which today is called the *Riemann integral*.



Shout-outs!

Shout-outs!

- A final notable person in the modern study of integration is Henri Lebesgue.



Shout-outs!

- A final notable person in the modern study of integration is Henri Lebesgue.
- In his 1902 PhD dissertation, he used measure theory to develop a theory of integration that is central to modern real analysis and modern probability.



Shout-outs!

- A final notable person in the modern study of integration is Henri Lebesgue.
- In his 1902 PhD dissertation, he used measure theory to develop a theory of integration that is central to modern real analysis and modern probability.
- All these people helped to give a rigorous foundation to one of the biggest accomplishments in history: the creation of calculus.



A People's History of Infinity

Cultural Influence

Cultural Influence

- What mathematicians have chosen to care about is influenced by the culture in which they lived.

Cultural Influence

- What mathematicians have chosen to care about is influenced by the culture in which they lived.
- Cultures have also influenced who did math, and who was excluded.

Cultural Influence

- What mathematicians have chosen to care about is influenced by the culture in which they lived.
- Cultures have also influenced who did math, and who was excluded.
- One of the largest cultural influences is religion. We have already seen in this course examples of religion influencing mathematics.

Cultural Influence

- What mathematicians have chosen to care about is influenced by the culture in which they lived.
- Cultures have also influenced who did math, and who was excluded.
- One of the largest cultural influences is religion. We have already seen in this course examples of religion influencing mathematics.
- Many mathematicians were also deeply religious.

Duality

Duality

- In the West, mathematicians remained uncomfortable with negative numbers until the 18th century!

Duality

- In the West, mathematicians remained uncomfortable with negative numbers until the 18th century!
- In the East, they were comfortable with them far earlier. Example: In the *Nine Chapters*.



Duality

- In the West, mathematicians remained uncomfortable with negative numbers until the 18th century!
- In the East, they were comfortable with them far earlier. Example: In the *Nine Chapters*.
- Perhaps Ancient Chinese religious philosophy, which emphasized duality, explains this. Liu Hui:



Duality



- In the West, mathematicians remained uncomfortable with negative numbers until the 18th century!
- In the East, they were comfortable with them far earlier. Example: In the *Nine Chapters*.
- Perhaps Ancient Chinese religious philosophy, which emphasized duality, explains this. Liu Hui:


I read the Nine Chapters as a boy, and studied it in full detail when I was older. I observed the division between the dual natures of Yin and Yang [the positive and negative aspects] which sum up the fundamentals of mathematics.

Order and Beauty

Order and Beauty


- Religion suggests that there is an order and beauty to the world. This seems to have influenced many mathematicians.

Order and Beauty



- Religion suggests that there is an order and beauty to the world. This seems to have influenced many mathematicians.
- Example: Isaac Newton believed that there *should* be universal laws. Without this belief, he may not have spent so long searching for them.

Order and Beauty



- Religion suggests that there is an order and beauty to the world. This seems to have influenced many mathematicians.
- Example: Isaac Newton believed that there *should* be universal laws. Without this belief, he may not have spent so long searching for them.
 - Srinivasa Ramanujan: “An equation has no meaning unless it expresses a thought of God.”




Infinity

Infinity

- Many more mathematicians who dealt with infinity had religious leanings.


Infinity

- Many more mathematicians who dealt with infinity had religious leanings.
- Georg Cantor believed God communicated mathematical truths to him.



Infinity

- Many more mathematicians who dealt with infinity had religious leanings.
- Georg Cantor believed God communicated mathematical truths to him.



The fear of infinity is a form of myopia that destroys the possibility of seeing the actual infinite, even though it in its highest form has created and sustains us, and in its secondary transfinite forms occurs all around us and even inhabits our minds.

People's History of Infinity

People's History of Infinity

- The study of how culture influences math is called *ethnomathematics*.

People's History of Infinity

- The study of how culture influences math is called *ethnomathematics*.
- Would these people have cared about and discovered all this without being religious? It is impossible to say.

People's History of Infinity

- The study of how culture influences math is called *ethnomathematics*.
- Would these people have cared about and discovered all this without being religious? It is impossible to say.
- But culture influencing math is undeniable.

People's History of Infinity

- The study of how culture influences math is called *ethnomathematics*.
- Would these people have cared about and discovered all this without being religious? It is impossible to say.
- But culture influencing math is undeniable.
- It is still happening today, and it will likely happen forever. It's worth being aware of it.

