

Iteration of linear differential operators

Jake Edmonstone

Matthew Tchouikine

Mentor: Paul Cusson

University of Waterloo, Directed Research Program

2026-07-25

1. The original problem

Does iterating the derivative infinitely many times give a smooth function whenever it converges?

Asked 4 years, 6 months ago Modified 3 years, 6 months ago Viewed 5k times



I am a graduate student and I've been thinking about this fun but frustrating problem for some time. Let $d = \frac{d}{dx}$, and let $f \in C^\infty(\mathbb{R})$ be such that for every real x ,

76

$$g(x) := \lim_{n \rightarrow \infty} d^n f(x)$$



converges. A simple example for such an f would be $ce^x + h(x)$ for any constant c where $h(x)$ converges to 0 everywhere under this iteration (in fact my hunch is that every such f is of this form), eg. $h(x) = e^{x/2}$ or simply a polynomial, of course.



I've been trying to show that g is, in fact, differentiable, and thus is a fixed point of d . Whether this is true would provide many interesting properties from a dynamical systems point of view if one can generalize to arbitrary smooth linear differential operators, although they might be too good to be true.

Perhaps this is a known result? If so I would greatly appreciate a reference. If it has a trivial counterexample I've missed, please let me know. Otherwise, I'll try to come up with some tricky double limit using tricks such as in [this MSE answer](#), to

Any help is kindly appreciated.

asked Jan 5, 2022 at 6:05



Paul Cusson

1.1 The original question

1. The original problem

Let $f \in C^\infty(\mathbb{R})$, and write

$$L_1 := \frac{d}{dx}.$$

Suppose that for every $x \in \mathbb{R}$ the pointwise limit

$$g(x) := \lim_{n \rightarrow \infty} L_1^n f(x)$$

exists.

Question. Must g be smooth? Moreover, can we prove $L_1 g = g$?

A direct answer:



64



I was able to adapt [the accepted answer](#) to [this MathOverflow post](#) to positively answer the question. The point is that one can squeeze more out of Petrov's Baire category argument if one applies it to the "singular set" of the function, rather than to an interval.

The key step is to establish



Theorem 1. Let $f \in C^\infty(\mathbf{R})$ be such that the quantity $M(x) := \sup_{m \geq 0} |f^{(m)}(x)|$ is finite for all x . Then f is the restriction to $\mathbf{R}$ of an entire function (or equivalently, f is real analytic with an infinite radius of convergence).

The convergence hypothesis actually forces f to be a restriction of an entire function.

answered Jan 6, 2022 at 4:06



Terry Tao

1.2 Tao's analyticity theorem

1. The original problem

Theorem 1

Let $f \in C^\infty(\mathbb{R})$, and suppose that for every $x \in \mathbb{R}$,

$$M(x) := \sup_{m \geq 0} |f^{(m)}(x)| < \infty.$$

Then f is the restriction to $\mathbb{R}$ of an entire function. Equivalently, f is real analytic with infinite radius of convergence.

1.3 Why Tao's theorem applies

1. The original problem

For each fixed $x \in \mathbb{R}$, the sequence

$$(f^{(n)}(x))_{n \geq 0}$$

converges, hence is bounded. Therefore

$$M(x) = \sup_{n \geq 0} |f^{(n)}(x)| < \infty,$$

so Tao's theorem applies immediately: f is the restriction of an entire function.

1.4 One bound for every derivative

1. The original problem

Since f is the restriction of an entire function, expand $f^{(k)}$ at 0. Every coefficient is controlled by the *single* number $M(0)$:

$$f^{(k)}(x) = \sum_{n=0}^{\infty} \frac{f^{(n+k)}(0)}{n!} x^n,$$
$$\implies |f^{(k)}(x)| \leq \sum_{n=0}^{\infty} \frac{M(0)}{n!} |x|^n = M(0)e^{|x|}.$$

Note that for all $n \in \mathbb{N}$ and $x \in [a, b]$,

$$|f^{(n+1)}(x)| \leq C_{[a,b]} := \sup_{x \in [a,b]} M(0)e^{|x|} < \infty.$$

1.5 Pass the limit through the integral

1. The original problem

Fix $a < b$. For every n , the Fundamental Theorem of Calculus gives the identity

$$f^{(n)}(b) - f^{(n)}(a) = \int_a^b f^{(n+1)}(x) \, dx$$

For $x \in [a, b]$, we have

$$\lim_{n \rightarrow \infty} f^{(n+1)}(x) = g(x) \quad \text{and} \quad |f^{(n+1)}(x)| \leq C_{[a,b]}.$$

Therefore, by Lebesgue's Dominated Convergence Theorem,

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_a^b f^{(n+1)}(x) \, dx &= \int_a^b \lim_{n \rightarrow \infty} f^{(n+1)}(x) \, dx \\ &= \int_a^b g(x) \, dx. \end{aligned}$$

1.6 The limit is continuous

1. The original problem

Using the locally uniform bound from before, we show that g is continuous,

$$g(b) - g(a) = \int_a^b g(x) \, dx,$$

Therefore,

$$\begin{aligned} |g(b) - g(a)| &\leq \int_a^b |g(x)| \, dx \\ &\leq C_{[a,b]}(b - a). \end{aligned}$$

1.7 The limit satisfies $L_1 g = g$

1. The original problem

Fix b . The integral identity becomes

$$g(x) = g(b) + \int_b^x g(t) \, dt.$$

Since g is continuous, the Fundamental Theorem of Calculus yields

$$g'(x) = g(x).$$

Thus $L_1 g = g$, so g is smooth. Solving the ODE gives

$$g(x) = Ce^x$$

for some $C \in \mathbb{R}$.

2. Extensions

$$2.1 \ L_2 f := af'$$

2.1.1 The result

Let $a \in C^\infty(\mathbb{R})$ be nowhere zero, and define

$$L_2 f := a f'.$$

Suppose that the pointwise limit

$$g(x) := \lim_{n \rightarrow \infty} L_2^n f(x)$$

exists for every $x \in \mathbb{R}$.

We will show that

$$L_2 g = g.$$

2.1.2 Change coordinates

Introduce the new coordinate

$$\Phi : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \int_0^x \frac{1}{a(t)} dt.$$

Since a is nowhere zero, Φ is a diffeomorphism from $\mathbb{R}$ onto an open interval $I := \Phi(\mathbb{R})$. Set

$$F := f \circ \Phi^{-1} : I \rightarrow \mathbb{R}.$$

The chain rule gives

$$F' = (L_2 f) \circ \Phi^{-1},$$

and therefore, for every $n \geq 0$,

$$F^{(n)} = (L_2^n f) \circ \Phi^{-1}.$$

2.1.3 Pull back the ordinary result

$$2.1 \quad L_2 f := a f'$$

Note that

$$\begin{aligned} \lim_{n \rightarrow \infty} F^{(n)}(y) &= \lim_{n \rightarrow \infty} (L_2^n f)(\Phi^{-1}(y)) \\ &= g(\Phi^{-1}(y)) \end{aligned}$$

exists. So, we can define $G(y) := \lim_{n \rightarrow \infty} F^{(n)}(y)$. The ordinary derivative result on the interval I gives

$$G' = G.$$

2.1.4 Pull back the ordinary result

$$2.1 \quad L_2 f := a f'$$

Since $G = g \circ \Phi^{-1}$, then $g = G \circ \Phi$. Since $\Phi' = \frac{1}{a}$ (by FTC), then

$$\begin{aligned} g' &= (G' \circ \Phi) \cdot \Phi' \\ &= (G \circ \Phi) \cdot \frac{1}{a} \\ &= g \cdot \frac{1}{a}. \end{aligned}$$

Hence

$$\begin{aligned} L_2 g &= a g' \\ &= a \left(g \cdot \frac{1}{a} \right) \\ &= g. \end{aligned}$$

$$2.2 \ L_3 f := af' + f$$

2.2.1 The result

$$2.2 \quad L_3 f := af' + f$$

$$2.3 \quad L_4 f := af' + bf$$

2.3.1 The result

$$2.3 \quad L_4 f := a f' + b f$$

$$2.4 \ L_5 f := x f'$$

2.4.1 A simple zero

$$2.4 \quad L_5 f := x f'$$

$$2.5 \quad L_6 f := x^2 f'$$

2.5.1 A second-order zero

Suppose $g(x) := \lim_{n \rightarrow \infty} L_6^n f(x)$ exists pointwise.

Change coordinates. On each half-line, $\Phi(x) := -\frac{1}{x}$ turns L_6 into ordinary differentiation, so

$$g(x) = C_+ e^{-\frac{1}{x}} \quad (x > 0), \quad g(x) = C_- e^{-\frac{1}{x}} \quad (x < 0).$$

At the origin. As $x \rightarrow 0^-$, $e^{-\frac{1}{x}} \rightarrow \infty$, so smoothness forces $C_- = 0$. Also, $(L_6 h)(0) = 0$ for every smooth h , hence $g(0) = 0$.

Conclusion.

$$g(x) = \begin{cases} C e^{-\frac{1}{x}} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

The right-hand term vanishes at 0. Therefore $g \in C^\infty(\mathbb{R})$ and $L_6 g = g$.

3. Future directions

3.1.1 The spaces A , B , and C

3.1 Structure and open problems

For a linear differential operator $L : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R})$, define

$$A := \left\{ h \in C^\infty(\mathbb{R}) \mid \lim_{n \rightarrow \infty} L^n h(x) = 0 \text{ for every } x \in \mathbb{R} \right\}.$$

Transient space: the iterates of h disappear pointwise.

$$B := \left\{ f \in C^\infty(\mathbb{R}) \mid \lim_{n \rightarrow \infty} L^n f(x) \text{ exists for every } x \in \mathbb{R} \right\}.$$

Input space: these are precisely the functions for which the iteration problem makes sense.

$$C := \{ g \in C^\infty(\mathbb{R}) \mid Lg = g \}.$$

Fixed-point space: every iterate of g is already equal to g .

3.1.2 Why $B = A + C$

3.1 Structure and open problems

Assume every pointwise limit of iterates is smooth and fixed by L . If $f \in B$, let

$$g(x) := \lim_{n \rightarrow \infty} L^n f(x), \quad h := f - g.$$

Then $g \in C$, while

$$L^n h = L^n f - g \rightarrow 0,$$

so $h \in A$. Hence $f = h + g \in A + C$.

Conversely, if $f = h + g$ with $h \in A$ and $g \in C$, then

$$L^n f = L^n h + g \rightarrow g.$$

Thus $f \in B$, and therefore

$$B = A + C.$$

In other words: **every convergent orbit is a transient part plus its limiting fixed point.**

3.1.3 Beyond first-order ODEs

3.1 Structure and open problems

The guiding question is:

$$L^n f \rightarrow g \text{ pointwise} \implies g \in C^\infty \text{ and } Lg = g?$$

1. Higher-order ODEs. Study

$$Lf(x) = \sum_{k=0}^m a_{k(x)} f^{(k)}(x), \quad m \geq 2.$$

Which conditions on the coefficients recover local derivative bounds when the first-order change of coordinates is unavailable?

2. Partial differential operators. For

$$Lf(x) = \sum_{|\alpha| \leq m} a_{\alpha(x)} \partial^\alpha f(x) \quad \text{on } \mathbb{R}^d,$$

Begin with constant-coefficient or elliptic operators. Can their regularity theory justify passing the limit through L ?

Questions?