Pointwise Limits of Differential-Operator Iterates

Analytic Input and a Singular Counterexample

Jake Edmonstone and Matthew Tchouikine
Mentor: Paul Cusson
University of Waterloo Directed Research Program

Abstract. Let L be a linear differential operator of positive order with real-analytic coefficients on an open interval, allowing zeros of the leading coefficient. For real-analytic input f, we prove that pointwise boundedness of $(L^n f)$ implies local uniform bounds on every derivative of the iterates. If the orbit converges pointwise, its limit g is real analytic, the convergence holds locally uniformly with all derivatives, and $Lg = g$. For one such first-order operator with polynomial coefficients, we construct a smooth, nonanalytic input whose iterates converge to the indicator of the origin.

Contents

1 Introduction 2
1.1 Background and results 2
1.2 Notation 2
2 Elementary estimates 3
2.1 Interpolation 3
2.2 Smooth convergence 4
3 Analytic coefficients and input 5
3.1 Uniform propagation on a complex disk 5
3.2 The main theorem 7
4 A singular counterexample 11
4.1 Construction of the operator 11
4.2 Smoothness of the input 12
4.3 Orbit formula and convergence 13
4.4 Failure of local uniform convergence 14
4.5 The positive-order assumption 15
References 15

1 Introduction

1.1 Background and results

Our previous work (Edmonstone and Tchouikine 2026) studied the following question of Paul Cusson (Tao 2022).

> **Problem** **Pointwise limits of successive derivatives**
>
> Suppose $f \in C^\infty(\mathbb{R})$ and $f^{(n)}(x) \to g(x)$ for every real x. Must g be differentiable, and must $g' = g$?

Tao's analyticity theorem (Tao 2022) gives an affirmative answer: pointwise boundedness of all derivatives forces an entire extension of f and local uniform derivative bounds. These bounds justify passing to the limit in $f^{(n+1)} = Df^{(n)}$.

The first-order theorem (Edmonstone and Tchouikine 2026, Section 2) treats $Lh = ah' + bh$, where $a, b \in C^\infty(\mathbb{R})$ and a is nowhere zero. A change of variable and an integrating factor turn repeated application of L into repeated ordinary differentiation. For $f \in C^\infty(\mathbb{R})$, pointwise convergence of $L^n f$ therefore implies that the limit is smooth and satisfies $Lg = g$.

In Theorem 3.2 we consider operators of arbitrary positive order with analytic coefficients and input, allowing zeros of the leading coefficient. Pointwise boundedness of the orbit gives local uniform bounds on every derivative; pointwise convergence then gives a real-analytic limit satisfying $Lg = g$. The proof obtains bounds on one real interval, propagates them along complex paths avoiding the zeros, and uses the maximum-modulus principle to extend the bounds across those zeros.

In Theorem 4.2, a first-order operator with polynomial coefficients and a single zero of its leading coefficient has a smooth input whose orbit converges to a discontinuous limit. This shows that the analytic-input assumption in Theorem 3.2 cannot be replaced by smoothness. It also shows that the nowhere-zero hypothesis in the smooth first-order theorem cannot be omitted.

1.2 Notation

Throughout, $I \subseteq \mathbb{R}$ is a nonempty open interval. Inputs and coefficients on I are real valued. Holomorphic extensions are denoted by the same symbols as the original functions. Write $\mathbb{N}_0 = \{0, 1, 2, \ldots\}$, $D = \frac{\mathrm{d}}{\mathrm{d}x}$, and $h^{(0)} = h$. For an operator

$$Lh = \sum_{j=0}^{m} a_j h^{(j)},$$

write $u_n = L^n f$ for $n \in \mathbb{N}_0$, with $u_0 = f$. Thus $Lu_n = u_{n+1}$.

For a continuous function h on a nonempty compact set K, put $\|h\|_K = \sup_{x\in K}|h(x)|$. We say that v_n converges to g **locally uniformly with all derivatives** if g is smooth and

$$\left\|v_n^{(q)} - g^{(q)}\right\|_K \longrightarrow 0 \quad \text{for every } q \in \mathbb{N}_0 \text{ and every compact interval } K \subseteq I.$$

We write $v_n \to g$ in $C^\infty_{\mathrm{loc}}(I)$ for this convergence. In the complex plane, $B(z,R) = \{w \in \mathbb{C} \mid |w - z| < R\}$ denotes an open disk, and $\overline{B(z,R)}$ its closure.

2 Elementary estimates

2.1 Interpolation

Lemma 2.1 **Scaled interpolation**

Fix an integer $m \geq 1$. There is a constant C_m, depending only on m, such that for every compact interval $J = [\alpha, \beta]$ with $\alpha < \beta$, every real function h of class C^m on a neighborhood of J, and $0 < \delta \leq \frac{\beta-\alpha}{2m}$,

$$\left\|h^{(j)}\right\|_J \leq C_m\delta^{-j}\|h\|_J + C_m\delta^{m-j}\left\|h^{(m)}\right\|_J \quad (0 \leq j < m).$$

Proof: Fix $x \in J$. Choose $\varepsilon = 1$ if $x \leq \frac{\alpha+\beta}{2}$, and $\varepsilon = -1$ otherwise. Then all points $x + \varepsilon\ell\delta$, $1 \leq \ell \leq m$, lie in J. Taylor's theorem gives

$$h(x + \varepsilon\ell\delta) = \sum_{j=0}^{m-1} \frac{(\varepsilon\ell)^j}{j!}\delta^j h^{(j)}(x) + R_\ell,$$

where

$$|R_\ell| \leq \frac{(m\delta)^m}{m!}\left\|h^{(m)}\right\|_J.$$

These are m linear equations for the unknowns $\delta^j h^{(j)}(x)$, $0 \leq j < m$. Their coefficient matrix has entries $V_{\ell,j} = \frac{(\varepsilon\ell)^j}{j!}$.

If $Vv = 0$, the polynomial $p(t) = \sum_{j=0}^{m-1} \frac{v_j}{j!}t^j$ vanishes at the m distinct points $\varepsilon, 2\varepsilon, \ldots, m\varepsilon$. Since its degree is at most $m-1$, it is zero. Thus V is invertible.

Let A_m be the maximum sum of absolute values of entries in a row of V^{-1}, over both choices of ε. Multiplication by V^{-1} gives

$$\left|h^{(j)}(x)\right| \le A_m \delta^{-j}\left(\|h\|_J + \frac{(m\delta)^m}{m!}\left\|h^{(m)}\right\|_J\right).$$

Take the supremum over $x \in J$ and set $C_m = A_m \max\{1, \frac{m^m}{m!}\}$. □

We will also use the following first-derivative estimate. Let J be a compact interval and let h be twice continuously differentiable on a neighborhood of J. If $t > 0$ and $[x, x+t] \subseteq J$, Taylor's theorem gives

$$|h'(x)| \le \frac{2}{t}\|h\|_J + \frac{t}{2}\|h''\|_J.$$

Indeed, subtract $h(x)$ from the formula for $h(x+t)$ and divide by t.

2.2 Smooth convergence

Lemma 2.2 **Pointwise convergence and local derivative bounds**

Suppose $v_n \in C^\infty(I)$ and $v_n(x) \to g(x)$ for every $x \in I$. Assume that for every compact interval $J \subseteq I$ and every $j \ge 0$ there is a finite constant $B_{J,j}$ such that

$$\left\|v_n^{(j)}\right\|_J \le B_{J,j} \quad \text{for every } n.$$

Then g is smooth and $v_n \to g$ locally uniformly with all derivatives.

Proof: **First, the functions converge uniformly on compact intervals.** Fix a compact interval $K \subseteq I$ and let $B = B_{K,1}$. The mean value theorem gives

$$|v_n(x) - v_n(y)| \le B|x - y| \quad (x, y \in K).$$

Given $\varepsilon > 0$, choose $\delta > 0$ with $2B\delta < \frac{\varepsilon}{2}$. Cover K by finitely many intervals of radius δ centered at points $x_1, \dots, x_q$ of K. At these finitely many points, pointwise convergence gives one N such that

$$|v_n(x_i) - v_m(x_i)| < \frac{\varepsilon}{2} \quad (n, m \ge N, 1 \le i \le q).$$

For any $x \in K$, choose x_i within δ of x. The triangle inequality gives

$$|v_n(x) - v_m(x)| < 2B\delta + \frac{\varepsilon}{2} < \varepsilon.$$

Thus (v_n) is uniformly Cauchy on K. Letting m tend to infinity shows uniform convergence to its pointwise limit g. In particular, g is continuous.

Next, the derivatives are uniformly Cauchy as well. Suppose that $\left(v_n^{(j)}\right)$ is uniformly Cauchy on every compact interval. Choose $K = [c, d]$ and a larger $J = [a, b] \subseteq I$ with $a < c \leq d < b$. Apply the preceding first-derivative estimate to $h = v_n^{(j)} - v_m^{(j)}$. For $0 < t < b - d$ it gives

$$\left\| v_n^{(j+1)} - v_m^{(j+1)} \right\|_K \leq \frac{2}{t} \left\| v_n^{(j)} - v_m^{(j)} \right\|_J + tB_{J,j+2}.$$

Given $\varepsilon > 0$, first fix t small enough that the last term is less than $\frac{\varepsilon}{2}$. The induction hypothesis then makes the first term less than $\frac{\varepsilon}{2}$ for all sufficiently large n, m. This proves the induction step. Starting at $j = 0$, all derivative sequences converge uniformly on compact intervals. Denote their continuous limits by g_j, with $g_0 = g$.

Finally, identify these limits as derivatives. For $x, y \in I$, the fundamental theorem of calculus gives

$$v_n^{(j)}(y) - v_n^{(j)}(x) = \int_x^y v_n^{(j+1)}(s)\,\mathrm{d}s.$$

Uniform convergence on the interval between x and y permits passage to the limit, so

$$g_j(y) - g_j(x) = \int_x^y g_{j+1}(s)\,\mathrm{d}s.$$

Since g_{j+1} is continuous, this identity implies $g_j{}' = g_{j+1}$. Induction gives $g_j = g^{(j)}$ for every j. □

3 Analytic coefficients and input

3.1 Uniform propagation on a complex disk

Away from a zero of the leading coefficient, the relation $Lu_n = u_{n+1}$ expresses $u_n^{(m)}$ in terms of u_{n+1} and the lower derivatives of u_n. The following lemma propagates a bound for these functions from the center of a disk to a smaller disk. The smaller radius depends only on the coefficients and the original radius, so it can be used uniformly along a fixed path.

Lemma 3.1 **Uniform propagation**

Fix an integer $m \geq 1$, a radius $R > 0$, and $z_* \in \mathbb{C}$. Suppose that $c, b_0, \ldots, b_{m-1}$ and the functions $v_{n,j}$, $n \in \mathbb{N}_0$, $0 \leq j < m$, are holomorphic on a common neighborhood of $\overline{B(z_*, R)}$ and satisfy

$$v_{n,j}{}' = v_{n,j+1} \quad (0 \leq j < m-1),$$
$$v_{n,m-1}{}' = c v_{n+1,0} + \sum_{j=0}^{m-1} b_j v_{n,j}.$$

Put $E = \overline{B(z_*, R)}$. Choose

$$C_0 \geq \max\left\{1, \|c\|_E + \sum_{j=0}^{m-1} \|b_j\|_E\right\}, \quad A = \max\left\{\frac{2}{R}, 2C_0\right\}.$$

If a finite constant $M \geq 0$ satisfies $|v_{n,j}(z_*)| \leq M$ for all n, j, then

$$|v_{n,j}(z)| \leq 2M \quad \text{whenever } |z - z_*| \leq \frac{1}{2A},$$

for every $n \in \mathbb{N}_0$ and $0 \leq j < m$.

Proof: Expand at z_*:

$$v_{n,j}(z) = \sum_{k=0}^{\infty} d_{n,j,k}(z - z_*)^k,$$
$$c(z) = \sum_{\ell=0}^{\infty} c_\ell (z - z_*)^\ell, \quad b_j(z) = \sum_{\ell=0}^{\infty} b_{j,\ell}(z - z_*)^\ell.$$

Cauchy's estimates give

$$|c_\ell| + \sum_{j=0}^{m-1} |b_{j,\ell}| \leq C_0 R^{-\ell}.$$

Set $T_k = \sup_{n \geq 0, 0 \leq j < m} |d_{n,j,k}|$, initially allowing the value ∞. The hypothesis gives $T_0 \leq M$. Comparing coefficients in the differential relations, we obtain

$$(k+1) d_{n,j,k+1} = d_{n,j+1,k} \quad (j < m-1),$$
$$(k+1) d_{n,m-1,k+1} = \sum_{\ell=0}^{k} \left(c_\ell d_{n+1,0,k-\ell} + \sum_{j=0}^{m-1} b_{j,\ell} d_{n,j,k-\ell} \right).$$

Consequently, whenever $T_0, \dots, T_k$ are finite,

$$(k+1)T_{k+1} \le C_0 \sum_{\ell=0}^{k} R^{-\ell} T_{k-\ell}.$$

For the first set of relations, the $\ell = 0$ term suffices, since $C_0 \ge 1$. For the last relation, use the coefficient bound above. This also proves inductively that every T_k is finite.

We claim that $T_k \le MA^k$ for all $k \ge 0$. It holds at $k = 0$. Assuming it through degree k and using $RA \ge 2$, we get

$$\begin{aligned}(k+1)T_{k+1} &\le C_0 M A^k \sum_{\ell=0}^{k} (RA)^{-\ell} \\ &\le 2C_0 M A^k \le M A^{k+1}.\end{aligned}$$

Since $k + 1 \ge 1$, this proves the induction step. The Taylor series of every $v_{n,j}$ represents it on $B(z_*, R)$, and $\frac{1}{2A} \le \frac{R}{4}$. Hence, for $|z - z_*| \le \frac{1}{2A}$,

$$\left|v_{n,j}(z)\right| \le \sum_{k=0}^{\infty} MA^k |z - z_*|^k \le 2M.$$

□

3.2 The main theorem

Theorem 3.2 **Analytic input with zeros of the leading coefficient**

Let $m \ge 1$ be an integer, let $a_0, \dots, a_m$ be real analytic on I, and suppose a_m is not identically zero. Define

$$Lh = \sum_{j=0}^{m} a_j h^{(j)}.$$

Let f be real analytic on I, and put $u_n = L^n f$.

If $\left(u_n(x)\right)_{n\ge0}$ is bounded for every $x \in I$, then for every compact interval $K \subseteq I$ and every $q \in \mathbb{N}_0$,

$$\sup_{n\ge0} \left\|u_n^{(q)}\right\|_K < \infty.$$

If $u_n(x) \to g(x)$ for every $x \in I$, then g is real analytic,

$$Lg = g, \quad u_n \longrightarrow g \text{ in } C^\infty_{\mathrm{loc}}(I).$$

Proof: **Bounds on one real interval.** First assume pointwise boundedness. Since a_m is not identically zero, there is a compact interval $J_0 = [\alpha_0, \beta_0] \subseteq I$, with $\alpha_0 < \beta_0$, on which a_m does not vanish. For each positive integer N, set

$$E_N = \{x \in J_0 \mid |u_n(x)| \leq N \text{ for every } n \geq 0\}.$$

These sets are closed and cover J_0. Baire's theorem gives an E_N with nonempty relative interior. Choose a compact interval $J = [\alpha, \beta] \subseteq E_N$ with $\alpha < \beta$. Thus $\|u_n\|_J \leq N$ for all n.

On J, set $c = \frac{1}{a_m}$ and $b_j = -\frac{a_j}{a_m}$ for $0 \leq j < m$. The orbit relation becomes

$$u_n^{(m)} = cu_{n+1} + \sum_{j=0}^{m-1} b_j u_n^{(j)}.$$

Write $H_{n,j} = \left\|u_n^{(j)}\right\|_J$. Applying Lemma 2.1 to each u_n, for $0 < \delta \leq \frac{\beta-\alpha}{2m}$, yields

$$H_{n,m} \leq \|c\|_J N + C_m N \sum_{j=0}^{m-1} \|b_j\|_J \delta^{-j}$$

$$+C_m H_{n,m} \sum_{j=0}^{m-1} \|b_j\|_J \delta^{m-j}.$$

All powers $m - j$ in the last sum are positive. Choose a fixed $\delta > 0$ small enough that the coefficient of $H_{n,m}$ on the right is at most $\frac{1}{2}$. Each $H_{n,m}$ is finite, so moving that term to the left gives

$$H_{n,m} \leq 2\|c\|_J N + 2C_m N \sum_{j=0}^{m-1} \|b_j\|_J \delta^{-j}.$$

This bound is independent of n. Interpolation now bounds every $H_{n,j}$ for $j < m$, uniformly in n. In particular, at a chosen $x_* \in (\alpha, \beta)$,

$$\sup_{n \geq 0, 0 \leq j < m} \left|u_n^{(j)}(x_*)\right| \leq M_*$$

for some finite M_*.

A common holomorphic domain. The functions $f, a_0, \ldots, a_m$ have holomorphic extensions on a common disk centered at each point of I. Choose these disks with real diameters contained in I, and let Ω be their union. If two disks intersect, their intersection contains a real interval. The extensions agree on that interval, hence on the

intersection by the identity theorem. They therefore define holomorphic functions on Ω. The set Ω is connected, since each disk meets I.

Differentiation and multiplication preserve holomorphy on Ω, so all iterates $u_n = L^n f$ are holomorphic on this same domain. The zero set

$$Z = \{z \in \Omega \mid a_m(z) = 0\}$$

has no accumulation point in Ω, because a_m is holomorphic and not identically zero. It is also closed in Ω. The functions $c = \frac{1}{a_m}$ and $b_j = -\frac{a_j}{a_m}$ are holomorphic on $\Omega \setminus Z$.

Propagation along paths. The domain $\Omega \setminus Z$ is path connected. Indeed, two points can be joined by a polygonal path in Ω. Its compact image meets only finitely many zeros, and small detours around those zeros give a path in $\Omega \setminus Z$.

Fix $z \in \Omega \setminus Z$ and a continuous path from x_* to z in this domain. Its image Γ is compact. Choose $R > 0$ small enough that the compact set

$$E = \bigcup_{w \in \Gamma} \overline{B(w, R)}$$

is contained in $\Omega \setminus Z$. The coefficients $c, b_0, \ldots, b_{m-1}$ are bounded on E. Set

$$C_0 = \max\left\{1, \|c\|_E + \sum_{j=0}^{m-1} \|b_j\|_E\right\}, \quad A = \max\left\{\frac{2}{R}, 2C_0\right\}.$$

These choices satisfy Lemma 3.1 on every disk of radius R centered on the path.

Apply that lemma with $v_{n,j} = u_n^{(j)}$, $0 \le j < m$. Uniform continuity of the path allows a finite subdivision whose consecutive centers have distance less than $\frac{1}{2A}$. Starting with the bound M_* at x_*, each application gives a bound on the next disk at most twice the preceding bound. After finitely many steps, all the u_n are uniformly bounded on a neighborhood of z. Thus the family (u_n) is locally bounded on $\Omega \setminus Z$.

Bounds across a zero. Fix $z_0 \in Z$. Choose $\rho > 0$ such that $\overline{B(z_0, \rho)} \subseteq \Omega$ and this disk contains no other zero. Its boundary circle is a compact subset of $\Omega \setminus Z$. A finite cover by neighborhoods with uniform orbit bounds gives a constant C such that

$$|u_n(z)| \le C \quad (n \ge 0, |z - z_0| = \rho).$$

Every u_n is holomorphic on a neighborhood of the closed disk, including at z_0. The maximum-modulus principle gives

$$|u_n(z)| \le C \quad (n \ge 0, |z - z_0| \le \rho).$$

Consequently, (u_n) is locally bounded on all of Ω.

Bounds on derivatives. Let $K \subseteq I$ be a compact interval. Choose $r > 0$ so that its closed complex r-neighborhood is contained in Ω. Local boundedness and a finite cover give a constant C_K bounding every u_n on this compact neighborhood. Cauchy's estimates on disks of radius r centered at $x \in K$ give

$$\left\|u_n^{(q)}\right\|_K \leq C_K q! r^{-q} \quad (n, q \in \mathbb{N}_0).$$

This proves the first assertion.

Convergence and analyticity of the limit. Suppose now that $u_n(x) \to g(x)$ for every $x \in I$. Each pointwise orbit is bounded, so the preceding estimates apply. By Lemma 2.2, g is smooth and $u_n \to g$ locally uniformly with all derivatives. Hence we can pass to the limit in

$$\sum_{j=0}^{m} a_j u_n^{(j)} = u_{n+1}$$

to obtain $Lg = g$ throughout I, including at zeros of a_m.

Fix $x_0 \in I$ and choose $r > 0$ such that $[x_0 - r, x_0 + r] \subseteq I$ and $\overline{B(x_0, 2r)} \subseteq \Omega$. Let C bound all the u_n on this closed disk. For every real x with $|x - x_0| \leq r$, Cauchy's estimates on $B(x, r)$ and convergence of derivatives give

$$\left|g^{(q)}(x)\right| \leq C q! r^{-q} \quad (q \geq 0).$$

Taylor's theorem therefore bounds the remainder at x_0 by

$$\left| g(x) - \sum_{q=0}^{N} \frac{g^{(q)}(x_0)}{q!} (x - x_0)^q \right| \leq C \left(\frac{|x - x_0|}{r} \right)^{N+1}.$$

For $|x - x_0| < r$ this tends to zero as N tends to infinity. Thus g is real analytic at every $x_0 \in I$. □

For analytic input, the theorem applies to D^r for every integer $r \geq 1$, and to xD and $x^2 D$, including on intervals containing zero. It gives the corresponding fixed-point equations $g^{(r)} = g$, $xg' = g$, and $x^2 g' = g$. Our previous results (Edmonstone and Tchouikine 2026, Section 3) allow smooth input for xD and $x^2 D$.

4 A singular counterexample

4.1 Construction of the operator

We construct the operator by reversing the change of variable and integrating factor from the first-order theorem (Edmonstone and Tchouikine 2026, Section 2). The construction starts with a sequence satisfying $H_n{}' = H_{n+1}$.

For $t > 0$ and $n \in \mathbb{N}_0$, define

$$H_n(t) := \int_0^1 s^n e^{st} \, \mathrm{d}s.$$

Differentiating under the integral gives $H_n{}' = H_{n+1}$. This is justified on each compact t-interval by continuity and uniform boundedness of the integrand and its t-derivative on the rectangle. Also,

$$H_0(t) = \frac{e^t - 1}{t}.$$

Set $p(t) = te^{-t}$, so that $p(t)H_0(t) = 1 - e^{-t}$, and put $t(x) = \frac{1}{x^2}$ for $x \neq 0$. Then $p(t(x))H_0(t(x)) = 1 - e^{-\frac{1}{x^2}}$, which extends smoothly to the origin with value one by Lemma 4.1.

For $x \neq 0$,

$$\frac{\mathrm{d}t}{\mathrm{d}x} = -\frac{2}{x^3}, \quad -\frac{x^3}{2}\frac{\mathrm{d}t}{\mathrm{d}x} = 1.$$

Thus the operator $-\frac{x^3}{2}D$ becomes differentiation with respect to t. Multiplication by p introduces an extra term, which we cancel using $\frac{p'(t)}{p(t)} = \frac{1}{t} - 1$. This leads to

$$Lh := -\frac{x^3}{2}h' + (1 - x^2)h.$$

Both coefficients are polynomials, and the leading coefficient has exactly one zero, of order three, at the origin.

For $H \in C^\infty((0, \infty))$, define $w_H(x) = p(t(x))H(t(x))$ on $\mathbb{R} \setminus \{0\}$. Writing $t = t(x)$, the chain rule gives

$$\begin{aligned}(Lw_H)(x) &= \frac{\mathrm{d}}{\mathrm{d}t}[p(t)H(t)] + \left(1 - \frac{1}{t}\right)p(t)H(t) \\ &= p(t)H'(t) + \left[p'(t) + \left(1 - \frac{1}{t}\right)p(t)\right]H(t) \\ &= p(t)H'(t).\end{aligned}$$

In particular, $Lw_{H_n} = w_{H_{n+1}}$ on each half-line.

4.2 Smoothness of the input

Lemma 4.1 **The flat exponential**

Define

$$\eta(x) := \begin{cases} e^{-\frac{1}{x^2}} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}.$$

Then $\eta \in C^\infty(\mathbb{R})$ and $\eta^{(j)}(0) = 0$ for every $j \geq 0$.

Proof: First, for every integer $N \geq 0$,

$$|x|^{-N} e^{-\frac{1}{x^2}} \longrightarrow 0 \quad \text{as } x \longrightarrow 0.$$

To verify this, put $y = \frac{1}{x^2}$ and choose an integer $q > \frac{N}{2}$. Since $e^y \geq \frac{y^q}{q!}$ for $y > 0$, we have $0 \leq y^{\frac{N}{2}} e^{-y} \leq q! y^{\frac{N}{2} - q} \to 0$.

Off zero, repeated differentiation gives $\eta^{(j)}(x) = P_j\left(\frac{1}{x}\right) e^{-\frac{1}{x^2}}$ for polynomials P_j. Indeed, starting with $P_0 = 1$, the chain and product rules give

$$P_{j+1}(y) = -y^2 P_j'(y) + 2y^3 P_j(y).$$

Each of these expressions extends continuously by zero at the origin, by the preceding decay estimate. Its difference quotient at zero is $\frac{P_j\left(\frac{1}{x}\right)}{x} e^{-\frac{1}{x^2}}$, which also tends to zero. Inductively, each continuous extension is differentiable at zero with derivative zero, and its derivative is the next continuous extension. This proves smoothness and all the stated derivative values. □

Set $f = 1 - \eta$. Then $f \in C^\infty(\mathbb{R})$, $f(0) = 1$, and $f^{(j)}(0) = 0$ for $j \geq 1$. Its Taylor series at zero is the constant one. Since $f(x) < 1$ for every $x \neq 0$, it cannot agree with that Taylor series on any neighborhood of zero. Thus f is not real analytic at zero.

4.3 Orbit formula and convergence

Theorem 4.2 **A discontinuous limit with polynomial coefficients**

On $\mathbb{R}$, define

$$Lh = -\frac{x^3}{2}h' + (1-x^2)h, \quad f = 1-\eta,$$

where η is the function in Lemma 4.1. Write $u_n = L^n f$. Then every u_n is smooth, and for every $n \in \mathbb{N}_0$,

$$u_n(x) = \begin{cases} \frac{1}{x^2}\int_0^1 s^n e^{-\frac{1-s}{x^2}}\,\mathrm{d}s & \text{if } x \neq 0 \\ 1 & \text{if } x = 0. \end{cases}$$

Moreover, $0 \leq u_{n+1} \leq u_n \leq 1$ on $\mathbb{R}$, and the sequence converges pointwise to the discontinuous function

$$g(x) = \begin{cases} 1 & \text{if } x = 0 \\ 0 & \text{if } x \neq 0. \end{cases}$$

Proof: Smoothness of f follows from Lemma 4.1. Since differentiation and multiplication by polynomial coefficients preserve smoothness, induction shows that every u_n is smooth on all of $\mathbb{R}$.

On $\mathbb{R} \setminus \{0\}$, we have $f = w_{H_0}$ and $Lw_{H_n} = w_{H_{n+1}}$. Induction therefore gives, with $t = \frac{1}{x^2}$,

$$u_n(x) = p(t)H_n(t) = t\int_0^1 s^n e^{-(1-s)t}\,\mathrm{d}s.$$

Substituting $t = \frac{1}{x^2}$ gives the asserted formula.

At the origin, every smooth h satisfies

$$(Lh)(0) = h(0).$$

Since $f(0) = 1$, induction gives $u_n(0) = 1$ for all n.

For a fixed $x \neq 0$, the integral formula implies

$$0 \leq u_n(x) \leq \frac{1}{x^2}\int_0^1 s^n\,\mathrm{d}s = \frac{1}{x^2(n+1)}.$$

The right side tends to zero as n tends to infinity. Together with the value at zero, this proves the pointwise limit.

Since $0 \le s^{n+1} \le s^n$ on $[0,1]$, the same formula shows $0 \le u_{n+1}(x) \le u_n(x)$ off zero; the inequalities also hold at zero. Finally, $u_0 = f \le 1$. Hence the orbit is nonnegative, decreasing, and bounded by one everywhere. □

4.4 Failure of local uniform convergence

The convergence is uniform on every compact set avoiding zero: if $|x| \ge \delta > 0$, the bound in the proof is at most $\frac{1}{\delta^2(n+1)}$. On any interval containing zero in its interior, however, convergence cannot be uniform. In fact, each u_n is continuous and equals one at zero, while g is zero at every nearby nonzero point. Consequently

$$\sup_{0<|x|\le\varepsilon} |u_n(x) - g(x)| = 1 \quad (\varepsilon > 0).$$

There is also an explicit failure of uniform derivative bounds. Integrating the formula for u_n by parts in s, for $n \ge 1$, gives

$$u_n(x) = 1 - nx^2 u_{n-1}(x) \quad (x \neq 0).$$

Indeed, the differential of $e^{-(1-s)t}$ is $te^{-(1-s)t}\,\mathrm{d}s$; the boundary term is one and the remaining integral is $\frac{n}{t}u_{n-1}$. By continuity the recurrence holds at zero as well. Differentiating it twice at zero therefore gives

$$u_n''(0) = -2nu_{n-1}(0) = -2n.$$

Consequently, the local bound on second derivatives required in Lemma 2.2 fails at the origin.

For this same operator, Theorem 3.2 guarantees an analytic fixed-point limit whenever an analytic input has a pointwise convergent orbit. The failure here occurs for the smooth input $f = 1 - \eta$, whose nonanalyticity at zero was established above.

4.5 The positive-order assumption

Positive order is also necessary. On $I = (-1, 1)$, let

$$Lh = (1 - x^2)h, \quad f = 1.$$

Both the coefficient and the input are real analytic, but

$$L^n f(x) = (1 - x^2)^n \longrightarrow \begin{cases} 1 & \text{if } x = 0 \\ 0 & \text{if } x \neq 0 \end{cases}.$$

Indeed, $0 < 1 - x^2 < 1$ whenever $0 < |x| < 1$, whereas the value at zero is always one. Thus Theorem 3.2 does not extend to operators of order zero.

References

Edmonstone, Jake, and Matthew Tchouikine. 2026. *Iteration of Linear Differential Operators*. Directed Research Program report. University of Waterloo.

Tao, Terence. 2022. "Answer to: Does Iterating the Derivative Infinitely Many Times Give a Smooth Function Whenever It Converges?." January 6. https://mathoverflow.net/a/413247.