

Calculus 3, Final Formulas

December 18, 2018

Vector Operations/Identities

$$v \cdot w = \|v\| \|w\| \cos \theta$$

$$v \cdot v = \|v\|^2$$

$$v \perp w \Leftrightarrow v \cdot w = 0$$

$$\text{Unit vector: } e_u = \frac{u}{\|u\|}$$

$$\text{Projection of } u \text{ along } v: u_{\parallel v} = \left(\frac{u \cdot v}{v \cdot v} \right) v = \left(\frac{u \cdot v}{\|v\|^2} \right) v = \left(\frac{u \cdot v}{\|v\|} \right) e_v$$

$$\text{Component of } u \text{ along } v: \frac{u \cdot v}{\|v\|} = \|u\| \cos \theta$$

$$u \text{ perpendicular to } v: u_{\perp v} = u - u_{\parallel v}$$

$$\text{Decomposition of vector } u \text{ with respect to } v \text{ (derived from above): } u = u_{\perp v} + u_{\parallel v}$$

Cross product of v and w :

$$v \times w = \begin{pmatrix} i & j & k \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{pmatrix} = \begin{pmatrix} v_2 & v_3 \\ w_2 & w_3 \end{pmatrix} i - \begin{pmatrix} v_1 & v_3 \\ w_1 & w_3 \end{pmatrix} j + \begin{pmatrix} v_1 & v_2 \\ w_1 & w_2 \end{pmatrix} k$$

$$(v \times w) \perp v \text{ and } (v \times w) \perp w$$

$$\|v \times w\| = \|v\| \|w\| \sin \theta$$

$$\text{Area of parallelogram spanned by } v \text{ and } w = \|v \times w\|$$

Lines and Planes in Space

Equation of a line:

A line through $p_0 = (x_0, y_0, z_0)$ in the direction of $v = \langle a, b, c \rangle$:

$$r(t) = r_0 + t v = \langle x_0, y_0, z_0 \rangle + t \langle a, b, c \rangle$$

Parametric equations for line (derived from above):

$$x(t) = x_0 + at$$

$$y(t) = y_0 + bt$$

$$z(t) = z_0 + ct$$

Equation of a plane:

Vector form:

$$n \cdot \langle x, y, z \rangle = d$$

Scalar form:

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

$$\text{or } ax + by + cz = d \text{ where } d = ax_0 + by_0 + cz_0.$$

Arc length:

$$s(t) = \int_0^t \|r'(t)\| dt$$

Speed:

$$\frac{d}{dt} s(t) = \|r'(t)\|$$

Tangent plane:

For $z = f(x, y)$:

tangent plane at (x_0, y_0) : $f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$

Gradients, Directional Derivatives

Chain Rule (Alternate Method and Multivariate Method):

Given $y = f(x)$, $x = g(t)$, then $y = f(x) = f(g(t))$ and $\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}$. The following is then true:

Given $z = f(x, y)$, where $x = g(t)$, $y = h(t)$, $z = f(x, y) = f(g(t), h(t))$

and $\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$.

Gradient: Given $f(x, y)$, the gradient of f is $\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$.

Directional Derivative:

In the direction of unit vector $\vec{u} = \langle a, b \rangle$: $D_{\vec{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b$.

In three variables: $D_{\vec{u}}f(x, y, z) = f_x(x, y, z)a + f_y(x, y, z)b + f_z(x, y, z)c$.

or in a simpler form:

Given $f(x, y)$, the directional derivative of f at $p = (a, b)$ in the direction of unit vector \vec{u} is

$$D_{\vec{u}}f(p) = D_{\vec{u}}f(a, b) = \nabla f(a, b) \cdot \vec{u}$$

Rate of change of a function f in the direction of $\nabla f(p)$:
 $\|\nabla f(p)\|$.

Rate of change of a function f in the direction of a unit vector \vec{u} making an angle θ with $\nabla f(p)$:

$$\nabla f(p) \cdot \vec{u} = \|\nabla f(p)\| \|\vec{u}\| \cos \theta$$

(This comes from the following identity): $\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta$.

Optimization

Critical points

A point $p = (a, b)$ in the domain of f is a *critical point* if:

$f_x(a, b) = 0$ or $f_x(a, b)$ does not exist, and

$f_y(a, b) = 0$ or $f_y(a, b)$ does not exist.

Solving the system of $f_x = 0$, $f_y = 0$ will find the critical point (if it exists).

Second Derivative Test

The second derivative test finds local max., min., and saddle points.

A critical point of $f(x, y)$ is needed, as is the discriminant of $f(x, y)$ which is:

$$D(a, b) = \begin{bmatrix} f_{xx}(a, b) & f_{xy}(a, b) \\ f_{yx}(a, b) & f_{yy}(a, b) \end{bmatrix}$$

which yields

$$D(a, b) = f_{xx}(a, b)f_{yy}(a, b) - f_{xy}^2(a, b)$$

Then, the second derivative test's rules are:

$p = (a, b)$ is a *critical point* of $f(x, y)$.

1. If $D > 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minimum.
2. If $D > 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maximum.
3. If $D < 0$, then f has a saddle point at (a, b) .
4. If $D = 0$, then the test is inconclusive.

Global Extrema

Let $f(x, y)$ be defined over a closed domain D . Then, f 's extreme values occur at either critical points in

the interior of D , or at points on the boundary of D .

First, find and examine critical points. Then, evaluate f at the boundaries of D . Compare these points to find f_{max} and f_{min} .

Lagrange Multipliers

The Lagrange condition is: $\nabla f = \lambda \nabla g$.

Then, solve for λ in terms of x and y : $\lambda = m x$, $\lambda = n y$. (Taking m and n to represent some expressions).

Then, let these two λ expressions equal each other and solve for x and y : $m x = n y$.

This new x and y is the crit. point. Sub. the newly found x and y into constraint g to find max. and min. of f .