

# 4: Expectation

Spring 2023

Matthew Blackwell

Gov 2002 (Harvard)

# Where are we? Where are we going?

- We've defined random variables and their distributions.

# Where are we? Where are we going?

- We've defined random variables and their distributions.
- Distributions give full information about the probabilities of an r.v.

# Where are we? Where are we going?

- We've defined random variables and their distributions.
- Distributions give full information about the probabilities of an r.v.
- Today: begin to summarize distributions with a few numbers.

# 1/ Definition of Expectation

# How can we summarize distributions?

- Probability distributions describe the uncertainty about r.v.s.

# How can we summarize distributions?

- Probability distributions describe the uncertainty about r.v.s.
- Can we summarize probability distributions?


# How can we summarize distributions?

- Probability distributions describe the uncertainty about r.v.s.
- Can we summarize probability distributions?
- **Question:** What is the difference between these two p.m.f.s? How might we summarize this difference?



# How can we summarize distributions?

- Probability distributions describe the uncertainty about r.v.s.
- Can we summarize probability distributions?
- **Question:** What is the difference between these two p.m.f.s? How might we summarize this difference?



# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.
2. **Spread:** how spread out the distribution is around the center.

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.
2. **Spread:** how spread out the distribution is around the center.
  - We'll focus on the variance/standard deviation.

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.
2. **Spread:** how spread out the distribution is around the center.
  - We'll focus on the variance/standard deviation.
  - These are **population parameters** so we don't get to observe them.

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.
2. **Spread:** how spread out the distribution is around the center.
  - We'll focus on the variance/standard deviation.
  - These are **population parameters** so we don't get to observe them.
    - We won't get to observe them...

# Goals for summarizing

1. **Central tendency:** where the center of the distribution is.
  - We'll focus on the mean/expectation.
2. **Spread:** how spread out the distribution is around the center.
  - We'll focus on the variance/standard deviation.
  - These are **population parameters** so we don't get to observe them.
    - We won't get to observe them...
    - but we'll use our sample to learn about them



# Two ways to calculate averages

- Calculate the average of:  $\{1, 1, 1, 3, 4, 4, 5, 5\}$

# Two ways to calculate averages

- Calculate the average of:  $\{1, 1, 1, 3, 4, 4, 5, 5\}$

$$\frac{1 + 1 + 1 + 3 + 4 + 4 + 5 + 5}{8} = 3$$

# Two ways to calculate averages

- Calculate the average of:  $\{1, 1, 1, 3, 4, 4, 5, 5\}$

$$\frac{1 + 1 + 1 + 3 + 4 + 4 + 5 + 5}{8} = 3$$

- Alternative way to calculate average based on **frequency weights**:

# Two ways to calculate averages

- Calculate the average of:  $\{1, 1, 1, 3, 4, 4, 5, 5\}$

$$\frac{1 + 1 + 1 + 3 + 4 + 4 + 5 + 5}{8} = 3$$

- Alternative way to calculate average based on **frequency weights**:

$$1 \times \frac{3}{8} + 3 \times \frac{1}{8} + 4 \times \frac{2}{8} + 5 \times \frac{2}{8} = 3$$

# Two ways to calculate averages

- Calculate the average of: {1, 1, 1, 3, 4, 4, 5, 5}

$$\frac{1 + 1 + 1 + 3 + 4 + 4 + 5 + 5}{8} = 3$$

- Alternative way to calculate average based on **frequency weights**:

$$1 \times \frac{3}{8} + 3 \times \frac{1}{8} + 4 \times \frac{2}{8} + 5 \times \frac{2}{8} = 3$$

- Each value times how often that value occurs in the data.

# Two ways to calculate averages

- Calculate the average of:  $\{1, 1, 1, 3, 4, 4, 5, 5\}$

$$\frac{1 + 1 + 1 + 3 + 4 + 4 + 5 + 5}{8} = 3$$

- Alternative way to calculate average based on **frequency weights**:

$$1 \times \frac{3}{8} + 3 \times \frac{1}{8} + 4 \times \frac{2}{8} + 5 \times \frac{2}{8} = 3$$

- Each value times how often that value occurs in the data.
- We'll use this intuition to create an average/mean for r.v.s.

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.



# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.
  - $E[X]$  is a constant!

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.
  - $E[X]$  is a constant!
- Example:  $X \sim \text{Bern}(p)$ , then  $\mathbb{E}[X] =$

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.
  - $E[X]$  is a constant!
- Example:  $X \sim \text{Bern}(p)$ , then  $\mathbb{E}[X] = 1p + 0(1 - p) = p$ .

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.
  - $E[X]$  is a constant!
- Example:  $X \sim \text{Bern}(p)$ , then  $\mathbb{E}[X] = 1p + 0(1 - p) = p$ .
- If  $X$  and  $Y$  have the same distribution, then  $\mathbb{E}[X] = \mathbb{E}[Y]$ .

# Expectation

## Definition

The **expected value** (or **expectation** or **mean**) of a discrete r.v.  $X$  with possible values,  $x_1, x_2, \dots$  is

$$\mathbb{E}[X] = \sum_{j=1}^{\infty} x_j \mathbb{P}(X = x_j)$$

- Weighted average of the values of the r.v. weighted by the probability of each value occurring.
  - $E[X]$  is a constant!
- Example:  $X \sim \text{Bern}(p)$ , then  $\mathbb{E}[X] = 1p + 0(1 - p) = p$ .
- If  $X$  and  $Y$  have the same distribution, then  $\mathbb{E}[X] = \mathbb{E}[Y]$ .
  - Converse isn't true!

## Example - number of treated units

- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

## Example - number of treated units

- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

- Calculate the expectation of  $X$ :

## Example - number of treated units

- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

- Calculate the expectation of  $X$ :

$$\mathbb{E}[X] = \sum_{j=1}^k x_j \mathbb{P}(X = x_j)$$



## Example - number of treated units

- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

- Calculate the expectation of  $X$ :

$$\begin{aligned}\mathbb{E}[X] &= \sum_{j=1}^k x_j \mathbb{P}(X = x_j) \\ &= 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) + 2 \cdot \mathbb{P}(X = 2) + 3 \cdot \mathbb{P}(X = 3)\end{aligned}$$

# Example - number of treated units

- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$	$x p_X(x)$
0	1/8	0
1	3/8	3/8
2	3/8	6/8
3	1/8	3/8

- Calculate the expectation of  $X$ :

$$\begin{aligned}\mathbb{E}[X] &= \sum_{j=1}^k x_j \mathbb{P}(X = x_j) \\ &= 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) + 2 \cdot \mathbb{P}(X = 2) + 3 \cdot \mathbb{P}(X = 3) \\ &= 0 \cdot \frac{1}{8} + 1 \cdot \frac{3}{8} + 2 \cdot \frac{3}{8} + 3 \cdot \frac{1}{8}\end{aligned}$$

## Example - number of treated units



- Randomized experiment with 3 units.  $X$  is number of treated units.

$x$	$p_X(x)$	$x p_X(x)$
0	1/8	0
1	3/8	3/8
2	3/8	6/8
3	1/8	3/8

- Calculate the expectation of  $X$ :

$$\begin{aligned}\mathbb{E}[X] &= \sum_{j=1}^k x_j \mathbb{P}(X = x_j) \\ &= 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) + 2 \cdot \mathbb{P}(X = 2) + 3 \cdot \mathbb{P}(X = 3) \\ &= 0 \cdot \frac{1}{8} + 1 \cdot \frac{3}{8} + 2 \cdot \frac{3}{8} + 3 \cdot \frac{1}{8} = \frac{12}{8} = 1.5\end{aligned}$$

# Expectation as balancing point



## **2/** Linearity of Expectations

# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s

# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s
- Possible for **linear** functions because expectation is **linear**:

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[aX] = a\mathbb{E}[X] \quad \text{if } a \text{ is a constant}$$

# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s
- Possible for **linear** functions because expectation is **linear**:

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[aX] = a\mathbb{E}[X] \quad \text{if } a \text{ is a constant}$$

- True even if  $X$  and  $Y$  are dependent!



# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s
- Possible for **linear** functions because expectation is **linear**:

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[aX] = a\mathbb{E}[X] \quad \text{if } a \text{ is a constant}$$

- True even if  $X$  and  $Y$  are dependent!
- But this isn't always true for nonlinear functions:

# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s
- Possible for **linear** functions because expectation is **linear**:

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[aX] = a\mathbb{E}[X] \quad \text{if } a \text{ is a constant}$$

- True even if  $X$  and  $Y$  are dependent!
- But this isn't always true for nonlinear functions:
  - $\mathbb{E}[g(X)] \neq g(\mathbb{E}[X])$  unless  $g(\cdot)$  is a linear function.

# Properties of the expected value

- Often want to derive expectation of **transformations** of other r.v.s
- Possible for **linear** functions because expectation is **linear**:

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[aX] = a\mathbb{E}[X] \quad \text{if } a \text{ is a constant}$$

- True even if  $X$  and  $Y$  are dependent!
- But this isn't always true for nonlinear functions:
  - $\mathbb{E}[g(X)] \neq g(\mathbb{E}[X])$  unless  $g(\cdot)$  is a linear function.
  - $\mathbb{E}[XY] \neq \mathbb{E}[X]\mathbb{E}[Y]$  unless  $X$  and  $Y$  are independent.

# Expectation of a binomial

- Let  $X \sim \text{Bin}(n, p)$ , what's  $\mathbb{E}[X]$ ? Could just plug in formula:

$$\mathbb{E}[X] = \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} = ??$$

# Expectation of a binomial

- Let  $X \sim \text{Bin}(n, p)$ , what's  $\mathbb{E}[X]$ ? Could just plug in formula:

$$\mathbb{E}[X] = \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} = ??$$

- Use the story of the binomial as a sum of  $n$  Bernoulli  $X_i \sim \text{Bern}(p)$

$$X = X_1 + \dots + X_n$$

# Expectation of a binomial

- Let  $X \sim \text{Bin}(n, p)$ , what's  $\mathbb{E}[X]$ ? Could just plug in formula:

$$\mathbb{E}[X] = \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} = ??$$

- Use the story of the binomial as a sum of  $n$  Bernoulli  $X_i \sim \text{Bern}(p)$

$$X = X_1 + \dots + X_n$$

- Use linearity:

$$\mathbb{E}[X] = \mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n] = np$$

# Expectation of the sample mean

- Let  $X_1, \dots, X_n$  be identically distributed with  $\mathbb{E}[X_i] = \mu$ .

# Expectation of the sample mean

- Let  $X_1, \dots, X_n$  be identically distributed with  $\mathbb{E}[X_i] = \mu$ .
- Define the **sample mean** to be  $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$ .



# Expectation of the sample mean

- Let  $X_1, \dots, X_n$  be identically distributed with  $\mathbb{E}[X_i] = \mu$ .
- Define the **sample mean** to be  $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$ .
  - $\bar{X}$  is a r.v.!

# Expectation of the sample mean

- Let  $X_1, \dots, X_n$  be identically distributed with  $\mathbb{E}[X_i] = \mu$ .
- Define the **sample mean** to be  $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$ .
  - $\bar{X}$  is a r.v.!
- We can find the expectation of the sample mean using linearity:

$$\mathbb{E}[\bar{X}_n] = \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \frac{1}{n} n\mu = \mu$$

# Expectation of the sample mean

- Let  $X_1, \dots, X_n$  be identically distributed with  $\mathbb{E}[X_i] = \mu$ .
- Define the **sample mean** to be  $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$ .
  - $\bar{X}$  is a r.v.!
- We can find the expectation of the sample mean using linearity:

$$\mathbb{E}[\bar{X}_n] = \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \frac{1}{n} n\mu = \mu$$

- Intuition: on average, the sample mean is equal to the population mean.

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.
- But it must be between the highest and lowest possible value of an r.v.

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.
- But it must be between the highest and lowest possible value of an r.v.
  - If  $\mathbb{P}(X \geq c) = 1$ , then  $\mathbb{E}[X] \geq c$ .

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.
- But it must be between the highest and lowest possible value of an r.v.
  - If  $\mathbb{P}(X \geq c) = 1$ , then  $\mathbb{E}[X] \geq c$ .
  - If  $\mathbb{P}(X \leq c) = 1$ , then  $\mathbb{E}[X] \leq c$ .



# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.
- But it must be between the highest and lowest possible value of an r.v.
  - If  $\mathbb{P}(X \geq c) = 1$ , then  $\mathbb{E}[X] \geq c$ .
  - If  $\mathbb{P}(X \leq c) = 1$ , then  $\mathbb{E}[X] \leq c$ .
- Useful application of linearity: expectation is **monotone**.

# Monotonicity of expectations

- Expectations don't have to be in the support of the data.
  - $X \sim \text{Bern}(p)$  has  $E[X] = p$  which isn't 0 or 1.
- But it must be between the highest and lowest possible value of an r.v.
  - If  $\mathbb{P}(X \geq c) = 1$ , then  $\mathbb{E}[X] \geq c$ .
  - If  $\mathbb{P}(X \leq c) = 1$ , then  $\mathbb{E}[X] \leq c$ .
- Useful application of linearity: expectation is **monotone**.
  - If  $X \geq Y$  with probability 1, then  $\mathbb{E}(X) \geq \mathbb{E}(Y)$ .

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

- How much would you be willing to pay to play the game?

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

- How much would you be willing to pay to play the game?
- Let payout be  $Y = 2^X$ , we want  $\mathbb{E}[Y]$ :

$$\mathbb{E}[Y] = \sum_{k=1}^{\infty} 2^k \frac{1}{2^k} = \sum_{k=1}^{\infty} 1 = \infty$$

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

- How much would you be willing to pay to play the game?
- Let payout be  $Y = 2^X$ , we want  $\mathbb{E}[Y]$ :

$$\mathbb{E}[Y] = \sum_{k=1}^{\infty} 2^k \frac{1}{2^k} = \sum_{k=1}^{\infty} 1 = \infty$$

- Two ways to resolve the “paradox”:

# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

- How much would you be willing to pay to play the game?
- Let payout be  $Y = 2^X$ , we want  $\mathbb{E}[Y]$ :

$$\mathbb{E}[Y] = \sum_{k=1}^{\infty} 2^k \frac{1}{2^k} = \sum_{k=1}^{\infty} 1 = \infty$$

- Two ways to resolve the “paradox”:
  - No infinite money: max payout of  $2^{40}$  (around a trillion)  $\rightsquigarrow \mathbb{E}[Y] = 41$



# St. Petersburg Paradox

- Game of chance: stranger pays you  $\$2^X$  where  $X$  is the number of flips with a fair coin until the first heads.
  - Probability of reaching  $X = k$  is:

$$\mathbb{P}(X = k) = \mathbb{P}(T_1 \cap T_2 \cap \dots \cap T_{k-1} \cap H_k) = \mathbb{P}(T_1)\mathbb{P}(T_2) \dots \mathbb{P}(T_{k-1})\mathbb{P}(H_k) = \frac{1}{2^k}$$

- How much would you be willing to pay to play the game?
- Let payout be  $Y = 2^X$ , we want  $\mathbb{E}[Y]$ :

$$\mathbb{E}[Y] = \sum_{k=1}^{\infty} 2^k \frac{1}{2^k} = \sum_{k=1}^{\infty} 1 = \infty$$

- Two ways to resolve the “paradox”:
  - No infinite money: max payout of  $2^{40}$  (around a trillion)  $\rightsquigarrow \mathbb{E}[Y] = 41$
  - Risk avoidance/concave utility  $U = Y^{1/2} \rightsquigarrow \mathbb{E}[U(Y)] \approx 2.41$

# Undefined expectations\*

- We saw  $\mathbb{E}[X]$  can be infinite, but it can also be undefined.

# Undefined expectations\*

- We saw  $\mathbb{E}[X]$  can be infinite, but it can also be undefined.
- Example:  $X$  takes  $2^k$  and  $-2^k$  each with prob  $2^{-k-1}$ .

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} 2^k 2^{-k-1} - \sum_{k=1}^{\infty} 2^k 2^{-k-1} = \sum_{k=1}^{\infty} \frac{1}{2} - \sum_{k=1}^{\infty} \frac{1}{2} = \infty - \infty$$

# Undefined expectations\*

- We saw  $\mathbb{E}[X]$  can be infinite, but it can also be undefined.
- Example:  $X$  takes  $2^k$  and  $-2^k$  each with prob  $2^{-k-1}$ .

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} 2^k 2^{-k-1} - \sum_{k=1}^{\infty} 2^k 2^{-k-1} = \sum_{k=1}^{\infty} \frac{1}{2} - \sum_{k=1}^{\infty} \frac{1}{2} = \infty - \infty$$

- Often, both of these are assumed away by assuming  $\mathbb{E}[|X|] < \infty$  which implies  $\mathbb{E}[X]$  exists and is finite.

## **3/** Indicator Variables

# Indicator variables/fundamental bridge

- The probability of an event is equal to the expectation of its indicator:

$$\mathbb{P}(A) = \mathbb{E}[\mathbb{I}(A)]$$

# Indicator variables/fundamental bridge

- The probability of an event is equal to the expectation of its indicator:

$$\mathbb{P}(A) = \mathbb{E}[\mathbb{I}(A)]$$

- **Fundamental bridge** between probability and expectation

# Indicator variables/fundamental bridge

- The probability of an event is equal to the expectation of its indicator:

$$\mathbb{P}(A) = \mathbb{E}[\mathbb{I}(A)]$$

- **Fundamental bridge** between probability and expectation
- Makes it easy to prove probability results like **Bonferroni's inequality**

$$\mathbb{P}(A_1 \cup \dots \cup A_n) \leq \mathbb{P}(A_1) + \dots + \mathbb{P}(A_n)$$



# Indicator variables/fundamental bridge

- The probability of an event is equal to the expectation of its indicator:

$$\mathbb{P}(A) = \mathbb{E}[\mathbb{I}(A)]$$

- **Fundamental bridge** between probability and expectation
- Makes it easy to prove probability results like **Bonferroni's inequality**

$$\mathbb{P}(A_1 \cup \dots \cup A_n) \leq \mathbb{P}(A_1) + \dots + \mathbb{P}(A_n)$$

- Use the fact that  $\mathbb{I}(A_1 \cup \dots \cup A_n) \leq \mathbb{I}(A_1) + \dots + \mathbb{I}(A_n)$  and then take expectations.

# Using indicators to find expectations

- Suppose we are assigning  $n$  units to  $k$  treatments and all possibilities equally likely. What is the expected number of treatment conditions without any units?

# Using indicators to find expectations

- Suppose we are assigning  $n$  units to  $k$  treatments and all possibilities equally likely. What is the expected number of treatment conditions without any units?
- Use indicators!  $I_j = 1$  if  $j$ th condition is empty. So  $I_1 + \dots + I_k$  is the number of empty conditions.

$$\begin{aligned}\mathbb{E}[I_j] &= \mathbb{P}(\text{cond } j \text{ empty}) \\ &= \mathbb{P}(\{\text{unit } 1 \text{ not in cond } j\} \cap \dots \cap \{\text{unit } n \text{ not in cond } j\}) \\ &= \mathbb{P}(\{\text{unit } 1 \text{ not in cond } j\}) \dots \mathbb{P}(\{\text{unit } n \text{ not in cond } j\}) \\ &= \left(1 - \frac{1}{n}\right)^k\end{aligned}$$

# Using indicators to find expectations

- Suppose we are assigning  $n$  units to  $k$  treatments and all possibilities equally likely. What is the expected number of treatment conditions without any units?
- Use indicators!  $I_j = 1$  if  $j$ th condition is empty. So  $I_1 + \dots + I_k$  is the number of empty conditions.

$$\begin{aligned}\mathbb{E}[I_j] &= \mathbb{P}(\text{cond } j \text{ empty}) \\ &= \mathbb{P}(\{\text{unit 1 not in cond } j\} \cap \dots \cap \{\text{unit } n \text{ not in cond } j\}) \\ &= \mathbb{P}(\{\text{unit 1 not in cond } j\}) \dots \mathbb{P}(\{\text{unit } n \text{ not in cond } j\}) \\ &= \left(1 - \frac{1}{n}\right)^k\end{aligned}$$

- Thus, we have  $\mathbb{E} \left[ \sum_j I_j \right] = k(1 - 1/k)^n$ .

## 4/ Variance

# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- Could also use  $\mathbb{E}[|X - \mathbb{E}[X]|]$  but more clunky as a function.

# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- Could also use  $\mathbb{E}[|X - \mathbb{E}[X]|]$  but more clunky as a function.
- Weighted average of the squared distances from the mean.



# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- Could also use  $\mathbb{E}[|X - \mathbb{E}[X]|]$  but more clunky as a function.
- Weighted average of the squared distances from the mean.
  - Larger deviations (+ or -)  $\rightsquigarrow$  higher variance

# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- Could also use  $\mathbb{E}[|X - \mathbb{E}[X]|]$  but more clunky as a function.
- Weighted average of the squared distances from the mean.
  - Larger deviations (+ or -)  $\rightsquigarrow$  higher variance
- The **standard deviation** is the (positive) square root of the variance:

$$SD(X) = \sqrt{\mathbb{V}[X]}$$

# Variance

- The **variance** measures the spread of the distribution:

$$\mathbb{V}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- Could also use  $\mathbb{E}[|X - \mathbb{E}[X]|]$  but more clunky as a function.
- Weighted average of the squared distances from the mean.
  - Larger deviations (+ or -)  $\rightsquigarrow$  higher variance
- The **standard deviation** is the (positive) square root of the variance:

$$SD(X) = \sqrt{\mathbb{V}[X]}$$

- Useful equivalent representation of the variance:

$$\mathbb{V}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$$

- How do we calculate  $\mathbb{E}[X^2]$  since it's nonlinear?

- How do we calculate  $\mathbb{E}[X^2]$  since it's nonlinear?

## Definition

The **Law of the Unconscious Statistician**, or LOTUS, states that if  $g(X)$  is a function of a discrete random variable, then

$$\mathbb{E}[g(X)] = \sum_x g(x) \mathbb{P}(X = x)$$

- How do we calculate  $\mathbb{E}[X^2]$  since it's nonlinear?

## Definition

The **Law of the Unconscious Statistician**, or LOTUS, states that if  $g(X)$  is a function of a discrete random variable, then

$$\mathbb{E}[g(X)] = \sum_x g(x) \mathbb{P}(X = x)$$

- Example:  $\mathbb{E}[X^2]$  where  $X \sim \text{Bin}(n, p)$ .

$$\mathbb{E}[X] = \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k}$$

$$\mathbb{E}[X^2] = \sum_{k=0}^n k^2 \binom{n}{k} p^k (1-p)^{n-k}$$

## Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

## Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$
0	1/8
1	3/8
2	3/8
3	1/8

- Let's go back to the number of treated units to figure out the variance of the number of treated units:



# Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$	$x - \mathbb{E}[X]$
0	1/8	-1.5
1	3/8	-0.5
2	3/8	0.5
3	1/8	1.5

- Let's go back to the number of treated units to figure out the variance of the number of treated units:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 p_X(x_j)$$

# Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$	$x - \mathbb{E}[X]$	$(x - \mathbb{E}[X])^2$
0	1/8	-1.5	2.25
1	3/8	-0.5	0.25
2	3/8	0.5	0.25
3	1/8	1.5	2.25

- Let's go back to the number of treated units to figure out the variance of the number of treated units:

$$\begin{aligned}\mathbb{V}[X] &= \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 p_X(x_j) \\ &= (-1.5)^2 \times \frac{1}{8} + (-0.5)^2 \times \frac{3}{8} + 0.5^2 \times \frac{3}{8} + 1.5^2 \times \frac{1}{8}\end{aligned}$$

# Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$	$x - \mathbb{E}[X]$	$(x - \mathbb{E}[X])^2$
0	1/8	-1.5	2.25
1	3/8	-0.5	0.25
2	3/8	0.5	0.25
3	1/8	1.5	2.25

- Let's go back to the number of treated units to figure out the variance of the number of treated units:

$$\begin{aligned}\mathbb{V}[X] &= \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 p_X(x_j) \\ &= (-1.5)^2 \times \frac{1}{8} + (-0.5)^2 \times \frac{3}{8} + 0.5^2 \times \frac{3}{8} + 1.5^2 \times \frac{1}{8} \\ &= 2.25 \times \frac{1}{8} + 0.25 \times \frac{3}{8} + 0.25 \times \frac{3}{8} + 2.25 \times \frac{1}{8}\end{aligned}$$

# Example - number of treated units

- Use LOTUS to calculate the variance for a discrete r.v.:

$$\mathbb{V}[X] = \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 \mathbb{P}(X = x_j)$$

$x$	$p_X(x)$	$x - \mathbb{E}[X]$	$(x - \mathbb{E}[X])^2$
0	1/8	-1.5	2.25
1	3/8	-0.5	0.25
2	3/8	0.5	0.25
3	1/8	1.5	2.25

- Let's go back to the number of treated units to figure out the variance of the number of treated units:

$$\begin{aligned}\mathbb{V}[X] &= \sum_{j=1}^k (x_j - \mathbb{E}[X])^2 p_X(x_j) \\ &= (-1.5)^2 \times \frac{1}{8} + (-0.5)^2 \times \frac{3}{8} + 0.5^2 \times \frac{3}{8} + 1.5^2 \times \frac{1}{8} \\ &= 2.25 \times \frac{1}{8} + 0.25 \times \frac{3}{8} + 0.25 \times \frac{3}{8} + 2.25 \times \frac{1}{8} = 0.75\end{aligned}$$

# Properties of variances

1.  $\mathbb{V}[X + c] = \mathbb{V}[X]$  for any constant  $c$ .

# Properties of variances

1.  $\mathbb{V}[X + c] = \mathbb{V}[X]$  for any constant  $c$ .
2. If  $a$  is a constant,  $\mathbb{V}[aX] = a^2\mathbb{V}[X]$ .

# Properties of variances

1.  $\mathbb{V}[X + c] = \mathbb{V}[X]$  for any constant  $c$ .
2. If  $a$  is a constant,  $\mathbb{V}[aX] = a^2\mathbb{V}[X]$ .
3. If  $X$  and  $Y$  are **independent**, then  $\mathbb{V}[X + Y] = \mathbb{V}[X] + \mathbb{V}[Y]$ .

# Properties of variances

1.  $\mathbb{V}[X + c] = \mathbb{V}[X]$  for any constant  $c$ .
2. If  $a$  is a constant,  $\mathbb{V}[aX] = a^2\mathbb{V}[X]$ .
3. If  $X$  and  $Y$  are **independent**, then  $\mathbb{V}[X + Y] = \mathbb{V}[X] + \mathbb{V}[Y]$ .
  - But this doesn't hold for dependent r.v.s



# Properties of variances

1.  $\mathbb{V}[X + c] = \mathbb{V}[X]$  for any constant  $c$ .
2. If  $a$  is a constant,  $\mathbb{V}[aX] = a^2\mathbb{V}[X]$ .
3. If  $X$  and  $Y$  are **independent**, then  $\mathbb{V}[X + Y] = \mathbb{V}[X] + \mathbb{V}[Y]$ .
  - But this doesn't hold for dependent r.v.s
4.  $\mathbb{V}[X] \geq 0$  with equality holding only if  $X$  is a constant,  $\mathbb{P}(X = b) = 1$ .

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?
  - Use stories and indicator variables!

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?
  - Use stories and indicator variables!
- $X \sim \text{Bin}(n, p)$  is equivalent to  $X_1 + \dots + X_n$  where  $X_i \sim \text{Bern}(p)$

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?
  - Use stories and indicator variables!
- $X \sim \text{Bin}(n, p)$  is equivalent to  $X_1 + \dots + X_n$  where  $X_i \sim \text{Bern}(p)$
- Variance of a Bernoulli:

$$\mathbb{V}[X_i] = \mathbb{E}[X_i^2] - (\mathbb{E}[X_i])^2 = p - p^2 = p(1 - p)$$

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?
  - Use stories and indicator variables!
- $X \sim \text{Bin}(n, p)$  is equivalent to  $X_1 + \dots + X_n$  where  $X_i \sim \text{Bern}(p)$
- Variance of a Bernoulli:

$$\mathbb{V}[X_i] = \mathbb{E}[X_i^2] - (\mathbb{E}[X_i])^2 = p - p^2 = p(1 - p)$$

- (Used  $X_i^2 = X_i$  for indicator variables)

# Binomial variance

- Clunky to use LOTUS to calculate variances. Other ways?
  - Use stories and indicator variables!
- $X \sim \text{Bin}(n, p)$  is equivalent to  $X_1 + \dots + X_n$  where  $X_i \sim \text{Bern}(p)$
- Variance of a Bernoulli:

$$\mathbb{V}[X_i] = \mathbb{E}[X_i^2] - (\mathbb{E}[X_i])^2 = p - p^2 = p(1 - p)$$

- (Used  $X_i^2 = X_i$  for indicator variables)
- Binomials are the sum of **independent** Bernoulli r.v.s so:

$$\mathbb{V}[X] = \mathbb{V}[X_1 + \dots + X_n] = \mathbb{V}[X_1] + \dots + \mathbb{V}[X_n] = np(1 - p)$$

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$



# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?
- We can apply the rules of variances:

$$\mathbb{V}[\bar{X}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{V}[X_i] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}$$

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?
- We can apply the rules of variances:

$$\mathbb{V}[\bar{X}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{V}[X_i] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}$$

- Note: we needed independence and identically distributed for this.

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?
- We can apply the rules of variances:

$$\mathbb{V}[\bar{X}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{V}[X_i] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}$$

- Note: we needed independence and identically distributed for this.
- $SD(\bar{X}_n) = \sigma/\sqrt{n}$

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?
- We can apply the rules of variances:

$$\mathbb{V}[\bar{X}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{V}[X_i] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}$$

- Note: we needed independence and identically distributed for this.
  - $SD(\bar{X}_n) = \sigma/\sqrt{n}$
- Under i.i.d. sampling we know the expectation and variance of  $\bar{X}_n$  without any other assumptions about the distribution of the  $X_i$ !

# Variance of the sample mean

- Let  $X_1, \dots, X_n$  be i.i.d. with  $\mathbb{E}[X_i] = \mu$  and  $\mathbb{V}[X_i] = \sigma^2$ 
  - Earlier we saw that  $\mathbb{E}[\bar{X}_n] = \mu$ , what about  $\mathbb{V}[\bar{X}_n]$ ?
- We can apply the rules of variances:

$$\mathbb{V}[\bar{X}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \mathbb{V}[X_i] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}$$

- Note: we needed independence and identically distributed for this.
  - $SD(\bar{X}_n) = \sigma/\sqrt{n}$
- Under i.i.d. sampling we know the expectation and variance of  $\bar{X}_n$  without any other assumptions about the distribution of the  $X_i$ !
  - We don't know what distribution it takes though!

## 5/ Inequalities

# Inequalities

- Bounds are very important establishing unknown probabilities.



# Inequalities

- Bounds are very important establishing unknown probabilities.
  - Also very helpful in establishing limit results later on.



# Inequalities

- Bounds are very important establishing unknown probabilities.
  - Also very helpful in establishing limit results later on.
- Remember that  $\mathbb{E}[a + bX] = a + b\mathbb{E}[X]$  is linear, but  $\mathbb{E}[g(X)] \neq g(\mathbb{E}[X])$  for nonlinear functions.

# Inequalities

- Bounds are very important establishing unknown probabilities.
  - Also very helpful in establishing limit results later on.
- Remember that  $\mathbb{E}[a + bX] = a + b\mathbb{E}[X]$  is linear, but  $\mathbb{E}[g(X)] \neq g(\mathbb{E}[X])$  for nonlinear functions.
- Can we relate those? Yes for **convex** and **concave** functions.

# Concave and convex



# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\begin{aligned}\mathbb{E}[g(X)] &\geq g(\mathbb{E}[X]) && \text{if } g \text{ is convex} \\ \mathbb{E}[g(X)] &\leq g(\mathbb{E}[X]) && \text{if } g \text{ is concave}\end{aligned}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.

# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\begin{aligned}\mathbb{E}[g(X)] &\geq g(\mathbb{E}[X]) && \text{if } g \text{ is convex} \\ \mathbb{E}[g(X)] &\leq g(\mathbb{E}[X]) && \text{if } g \text{ is concave}\end{aligned}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.
  - $g(x) = x^2$  is convex, so  $\mathbb{E}[X^2] \geq (\mathbb{E}[X])^2$ .

# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\begin{aligned}\mathbb{E}[g(X)] &\geq g(\mathbb{E}[X]) && \text{if } g \text{ is convex} \\ \mathbb{E}[g(X)] &\leq g(\mathbb{E}[X]) && \text{if } g \text{ is concave}\end{aligned}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.
  - $g(x) = x^2$  is convex, so  $\mathbb{E}[X^2] \geq (\mathbb{E}[X])^2$ .
- Allows us to easily reason about complicated functions:

# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\mathbb{E}[g(X)] \geq g(\mathbb{E}[X]) \quad \text{if } g \text{ is convex}$$

$$\mathbb{E}[g(X)] \leq g(\mathbb{E}[X]) \quad \text{if } g \text{ is concave}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.
  - $g(x) = x^2$  is convex, so  $\mathbb{E}[X^2] \geq (\mathbb{E}[X])^2$ .
- Allows us to easily reason about complicated functions:
  - $\mathbb{E}[|X|] \geq |\mathbb{E}[X]|$



# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\mathbb{E}[g(X)] \geq g(\mathbb{E}[X]) \quad \text{if } g \text{ is convex}$$

$$\mathbb{E}[g(X)] \leq g(\mathbb{E}[X]) \quad \text{if } g \text{ is concave}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.
  - $g(x) = x^2$  is convex, so  $\mathbb{E}[X^2] \geq (\mathbb{E}[X])^2$ .
- Allows us to easily reason about complicated functions:
  - $\mathbb{E}[|X|] \geq |\mathbb{E}[X]|$
  - $\mathbb{E}[1/X] \geq 1/\mathbb{E}[X]$

# Jensen's inequality

## Jensen's inequality

Let  $X$  be a r.v. Then, we have

$$\mathbb{E}[g(X)] \geq g(\mathbb{E}[X]) \quad \text{if } g \text{ is convex}$$

$$\mathbb{E}[g(X)] \leq g(\mathbb{E}[X]) \quad \text{if } g \text{ is concave}$$

with equality only holding if  $g$  is linear.

- Makes proving variance positive simple.
  - $g(x) = x^2$  is convex, so  $\mathbb{E}[X^2] \geq (\mathbb{E}[X])^2$ .
- Allows us to easily reason about complicated functions:
  - $\mathbb{E}[|X|] \geq |\mathbb{E}[X]|$
  - $\mathbb{E}[1/X] \geq 1/\mathbb{E}[X]$
  - $\mathbb{E}[\log(X)] \leq \log(\mathbb{E}[X])$

## **6/** Poisson Distribution

## Definition

An r.v.  $X$  has the **Poisson distribution** with parameter  $\lambda > 0$ , written  $X \sim \text{Pois}(\lambda)$  if the p.m.f. of  $X$  is:

$$\mathbb{P}(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

- One more discrete distribution is very popular, especially for counts.

## Definition

An r.v.  $X$  has the **Poisson distribution** with parameter  $\lambda > 0$ , written  $X \sim \text{Pois}(\lambda)$  if the p.m.f. of  $X$  is:

$$\mathbb{P}(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

- One more discrete distribution is very popular, especially for counts.
  - Number of contributions a candidate for office receives in a day.

## Definition

An r.v.  $X$  has the **Poisson distribution** with parameter  $\lambda > 0$ , written  $X \sim \text{Pois}(\lambda)$  if the p.m.f. of  $X$  is:

$$\mathbb{P}(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

- One more discrete distribution is very popular, especially for counts.
  - Number of contributions a candidate for office receives in a day.
- Key calculus fact that makes this a valid p.m.f.:  $\sum_{k=0}^{\infty} \lambda^k / k! = e^{\lambda}$ .

# Poisson properties

- A Poisson r.v.  $X \sim \text{Pois}(\lambda)$  has an unusual property:

$$\mathbb{E}[X] = \mathbb{V}[X] = \lambda$$

# Poisson properties

- A Poisson r.v.  $X \sim \text{Pois}(\lambda)$  has an unusual property:

$$\mathbb{E}[X] = \mathbb{V}[X] = \lambda$$

- The sum of independent Poisson r.v.s is Poisson:

$$X \sim \text{Pois}(\lambda_1) \quad Y \sim \text{Pois}(\lambda_2) \quad \implies \quad X + Y \sim \text{Pois}(\lambda_1 + \lambda_2)$$



# Poisson properties

- A Poisson r.v.  $X \sim \text{Pois}(\lambda)$  has an unusual property:

$$\mathbb{E}[X] = \mathbb{V}[X] = \lambda$$

- The sum of independent Poisson r.v.s is Poisson:

$$X \sim \text{Pois}(\lambda_1) \quad Y \sim \text{Pois}(\lambda_2) \quad \implies \quad X + Y \sim \text{Pois}(\lambda_1 + \lambda_2)$$

- If  $X \sim \text{Bin}(n, p)$  with  $n$  large and  $p$  small, then  $X$  is approx  $\text{Pois}(np)$ .