

Module 3: Inference for the Average Treatment Effect

Fall 2021

Matthew Blackwell

Gov 2003 (Harvard)

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.
- Neyman: use the difference in means as an **estimator** of the ATE.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.
- Neyman: use the difference in means as an **estimator** of the ATE.
 - No assumptions to fill in the potential outcomes.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.
- Neyman: use the difference in means as an **estimator** of the ATE.
 - No assumptions to fill in the potential outcomes.
 - No exact derivation of the randomization distribution.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.
- Neyman: use the difference in means as an **estimator** of the ATE.
 - No assumptions to fill in the potential outcomes.
 - No exact derivation of the randomization distribution.
 - \rightsquigarrow asymptotic approximations.

Where are we? Where are we going?

- Fisher: use sharp null to fill in science table + permutation tests.
 - No way to estimate or infer about “average” effects, just the sharp null.
- Neyman: use the difference in means as an **estimator** of the ATE.
 - No assumptions to fill in the potential outcomes.
 - No exact derivation of the randomization distribution.
 - \rightsquigarrow asymptotic approximations.
- What’s common: the focus on **randomization** as generating variation in estimators.

Social pressure effect

- Gerber, Green, and Larimer (APSR, 2008)

Dear Registered Voter:

WHAT IF YOUR NEIGHBORS KNEW WHETHER YOU VOTED?

Why do so many people fail to vote? We've been talking about the problem for years, but it only seems to get worse. This year, we're taking a new approach. We're sending this mailing to you and your neighbors to publicize who does and does not vote.

The chart shows the names of some of your neighbors, showing which have voted in the past. After the August 8 election, we intend to mail an updated chart. You and your neighbors will all know who voted and who did not.

DO YOUR CIVIC DUTY — VOTE!

MAPLE DR	Aug 04	Nov 04	Aug 06
9995 JOSEPH JAMES SMITH	Voted	Voted	_____
9995 JENNIFER KAY SMITH		Voted	_____
9997 RICHARD B JACKSON		Voted	_____
9999 KATHY MARIE JACKSON		Voted	_____

Social pressure results

TABLE 2. Effects of Four Mail Treatments on Voter Turnout in the August 2006 Primary Election

	Experimental Group				
	Control	Civic Duty	Hawthorne	Self	Neighbors
Percentage Voting	29.7%	31.5%	32.2%	34.5%	37.8%
N of Individuals	191,243	38,218	38,204	38,218	38,201

- Typical reporting of the Neighbors vs Control effect:

$$\text{estimate} = \frac{1}{n_1} \sum_{i=1}^n D_i Y_i - \frac{1}{n_0} \sum_{i=1}^n (1 - D_i) Y_i \approx 8.1$$

$$\text{standard error} = \sqrt{\frac{\widehat{\sigma}_1^2}{n_1} + \frac{\widehat{\sigma}_0^2}{n_0}} \approx 0.27$$

$$95\% \text{ CI} = [\text{est} - 1.96 \cdot SE, \text{est} + 1.96 \cdot SE] \approx [7.57, 8.63]$$

Social pressure results

TABLE 2. Effects of Four Mail Treatments on Voter Turnout in the August 2006 Primary Election

	Experimental Group				
	Control	Civic Duty	Hawthorne	Self	Neighbors
Percentage Voting	29.7%	31.5%	32.2%	34.5%	37.8%
N of Individuals	191,243	38,218	38,204	38,218	38,201

- Typical reporting of the Neighbors vs Control effect:

$$\text{estimate} = \frac{1}{n_1} \sum_{i=1}^n D_i Y_i - \frac{1}{n_0} \sum_{i=1}^n (1 - D_i) Y_i \approx 8.1$$

$$\text{standard error} = \sqrt{\frac{\widehat{\sigma}_1^2}{n_1} + \frac{\widehat{\sigma}_0^2}{n_0}} \approx 0.27$$

$$95\% \text{ CI} = [\text{est} - 1.96 \cdot SE, \text{est} + 1.96 \cdot SE] \approx [7.57, 8.63]$$

- Can this analysis be justified by randomization?

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:
 - We want to find an estimator that is **unbiased** for the SATE.

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:
 - We want to find an estimator that is **unbiased** for the SATE.
 - But also derive the **sampling variance** of the estimator.

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:
 - We want to find an estimator that is **unbiased** for the SATE.
 - But also derive the **sampling variance** of the estimator.
- Properties of the estimators across repeated samples from:

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:
 - We want to find an estimator that is **unbiased** for the SATE.
 - But also derive the **sampling variance** of the estimator.
- Properties of the estimators across repeated samples from:
 - the randomization distribution.

Estimand of interest

- Common estimand in experiments: **sample average treatment effect**

$$\text{SATE} = \tau_{\text{fs}} = \frac{1}{n} \sum_{i=1}^n [Y_i(1) - Y_i(0)]$$

- Neyman/our goals:
 - We want to find an estimator that is **unbiased** for the SATE.
 - But also derive the **sampling variance** of the estimator.
- Properties of the estimators across repeated samples from:
 - the randomization distribution.
 - the randomization distribution + sampling from the population.

Finite sample results

- Setting: **completely randomized experiment**

Finite sample results

- Setting: **completely randomized experiment**
 - n units, n_1 treated and n_0 control.

Finite sample results

- Setting: **completely randomized experiment**
 - n units, n_1 treated and n_0 control.
- Natural estimator for the SATE, **difference-in-means**:

$$\hat{\tau}_{\text{diff}} = \underbrace{\frac{1}{n_1} \sum_{i=1}^n D_i Y_i}_{\text{mean among treated}} - \underbrace{\frac{1}{n_0} \sum_{i=1}^n (1 - D_i) Y_i}_{\text{mean among control}}$$

Finite sample results

- Setting: **completely randomized experiment**
 - n units, n_1 treated and n_0 control.
- Natural estimator for the SATE, **difference-in-means**:

$$\hat{\tau}_{\text{diff}} = \underbrace{\frac{1}{n_1} \sum_{i=1}^n D_i Y_i}_{\text{mean among treated}} - \underbrace{\frac{1}{n_0} \sum_{i=1}^n (1 - D_i) Y_i}_{\text{mean among control}}$$

- Conditional on the sample, $\hat{\tau}_{\text{diff}}$ only varies because of D_i

Repeated samples/randomizations

randomization 1




Repeated samples/randomizations




Repeated samples/randomizations




Repeated samples/randomizations




Repeated samples/randomizations




Repeated samples/randomizations




Repeated samples/randomizations




Repeated samples/randomizations



Repeated samples/randomizations




Repeated samples/randomizations



randomization
distribution

Repeated samples/randomizations



- **Randomization distribution** = sampling distribution of this estimator.

Finite-sample properties

- How does $\hat{\tau}_{\text{diff}}$ across randomizations?

Finite-sample properties

- How does $\hat{\tau}_{\text{diff}}$ across randomizations?
- Key properties of the randomization distribution we'd like to know:

Finite-sample properties

- How does $\hat{\tau}_{\text{diff}}$ across randomizations?
- Key properties of the randomization distribution we'd like to know:
 - **Unbiasedness:** is mean of the randomization distribution equal to the true SATE?

Finite-sample properties

- How does $\hat{\tau}_{\text{diff}}$ across randomizations?
- Key properties of the randomization distribution we'd like to know:
 - **Unbiasedness:** is mean of the randomization distribution equal to the true SATE?
 - **Sampling variance:** variance of the randomization distribution?

Finite-sample properties

- How does $\hat{\tau}_{\text{diff}}$ across randomizations?
- Key properties of the randomization distribution we'd like to know:
 - **Unbiasedness:** is mean of the randomization distribution equal to the true SATE?
 - **Sampling variance:** variance of the randomization distribution?
- Use these properties to construct confidence intervals, conduct tests.

Unbiasedness

- In a completely randomized experiment, $\hat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] = \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}]$$

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\begin{aligned}\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}] \\ &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i(1) \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i(0) \mid \mathbf{O}]\end{aligned}$$

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\begin{aligned}\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}] \\ &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i(1) \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i(0) \mid \mathbf{O}] \\ &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i \mid \mathbf{O}] Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) \mid \mathbf{O}] Y_i(0)\end{aligned}$$

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\begin{aligned}\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i(1) \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i(0) \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i \mid \mathbf{O}] Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) \mid \mathbf{O}] Y_i(0) \\&= \frac{1}{n_1} \sum_{i=1}^n \left(\frac{n_1}{n} \right) Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \left(\frac{n_0}{n} \right) Y_i(0)\end{aligned}$$

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\begin{aligned}\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i(1) \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i(0) \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i \mid \mathbf{O}] Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) \mid \mathbf{O}] Y_i(0) \\&= \frac{1}{n_1} \sum_{i=1}^n \left(\frac{n_1}{n}\right) Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \left(\frac{n_0}{n}\right) Y_i(0) \\&= \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = \tau_{\text{fs}}\end{aligned}$$

Unbiasedness

- In a completely randomized experiment, $\widehat{\tau}_{\text{diff}}$ is unbiased for τ_{fs}
- Let $\mathbf{O} = \{\mathbf{Y}(1), \mathbf{Y}(0)\}$ be the the potential outcomes.

$$\begin{aligned}\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}] &= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i Y_i(1) \mid \mathbf{O}] - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) Y_i(0) \mid \mathbf{O}] \\&= \frac{1}{n_1} \sum_{i=1}^n \mathbb{E}_D[D_i \mid \mathbf{O}] Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \mathbb{E}_D[(1 - D_i) \mid \mathbf{O}] Y_i(0) \\&= \frac{1}{n_1} \sum_{i=1}^n \left(\frac{n_1}{n}\right) Y_i(1) - \frac{1}{n_0} \sum_{i=1}^n \left(\frac{n_0}{n}\right) Y_i(0) \\&= \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = \tau_{\text{fs}}\end{aligned}$$

- Note: number treated/control doesn't matter for unbiasedness!

Finite-sample sampling variance

- Sampling variance of the difference-in-means estimator is:

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n},$$

Finite-sample sampling variance

- Sampling variance of the difference-in-means estimator is:

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n},$$

- S_0^2 and S_1^2 are the in-sample variances of $Y_i(0)$ and $Y_i(1)$, respectively.

$$S_0^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(0) - \overline{Y}(0))^2 \quad S_1^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(1) - \overline{Y}(1))^2$$

Finite-sample sampling variance

- Sampling variance of the difference-in-means estimator is:

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n},$$

- S_0^2 and S_1^2 are the in-sample variances of $Y_i(0)$ and $Y_i(1)$, respectively.

$$S_0^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(0) - \overline{Y}(0))^2 \quad S_1^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(1) - \overline{Y}(1))^2$$

- Here, $\overline{Y}(d) = (1/n) \sum_{i=1}^n Y_i(d)$.

Finite-sample sampling variance

- Sampling variance of the difference-in-means estimator is:

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n},$$

- S_0^2 and S_1^2 are the in-sample variances of $Y_i(0)$ and $Y_i(1)$, respectively.

$$S_0^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(0) - \overline{Y}(0))^2 \quad S_1^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(1) - \overline{Y}(1))^2$$

- Here, $\overline{Y}(d) = (1/n) \sum_{i=1}^n Y_i(d)$.
- Last term is the in-sample variation of the individual treatment effects:

Finite-sample sampling variance

- Sampling variance of the difference-in-means estimator is:

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n},$$

- S_0^2 and S_1^2 are the in-sample variances of $Y_i(0)$ and $Y_i(1)$, respectively.

$$S_0^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(0) - \overline{Y}(0))^2 \quad S_1^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(1) - \overline{Y}(1))^2$$

- Here, $\overline{Y}(d) = (1/n) \sum_{i=1}^n Y_i(d)$.
- Last term is the in-sample variation of the individual treatment effects:

$$S_{\tau_i}^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i(1) - Y_i(0) - \tau_{\text{fs}})^2$$

- None of these are directly observable!

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0
$Y_i(1)$	10	-10	0
τ_i	0	0	0

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0
$Y_i(1)$	10	-10	0
τ_i	0	0	0

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0
$Y_i(1)$	10	-10	0
τ_i	0	0	0

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	-10	10	0
$Y_i(1)$	10	-10	0
τ_i	20	-20	0

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0
$Y_i(1)$	10	-10	0
τ_i	0	0	0

	$i = 1$	$i = 2$	Avg.
$Y_i(0)$	-10	10	0
$Y_i(1)$	10	-10	0
τ_i	20	-20	0

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.		$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0	$Y_i(0)$	-10	10	0
$Y_i(1)$	10	-10	0	$Y_i(1)$	10	-10	0
τ_i	0	0	0	τ_i	20	-20	0

- Both have $\tau_{\text{fs}} = 0$, first has constant effects.

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.		$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0	$Y_i(0)$	-10	10	0
$Y_i(1)$	10	-10	0	$Y_i(1)$	10	-10	0
τ_i	0	0	0	τ_i	20	-20	0

- Both have $\tau_{\text{fs}} = 0$, first has constant effects.
- In first setup, $\widehat{\tau}_{\text{diff}} = 20$ or $\widehat{\tau}_{\text{diff}} = -20$ depending on the randomization.

Finite-sample sampling variance

$$\mathbb{V}_D(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{S_0^2}{n_0} + \frac{S_1^2}{n_1} - \frac{S_{\tau_i}^2}{n}$$

- If the treatment effects are constant across units, then $S_{\tau_i}^2 = 0$.
 - \rightsquigarrow in-sample variance is largest when treatment effects are constant.
- Intuition looking at two-unit samples:

	$i = 1$	$i = 2$	Avg.		$i = 1$	$i = 2$	Avg.
$Y_i(0)$	10	-10	0	$Y_i(0)$	-10	10	0
$Y_i(1)$	10	-10	0	$Y_i(1)$	10	-10	0
τ_i	0	0	0	τ_i	20	-20	0

- Both have $\tau_{\text{fs}} = 0$, first has constant effects.
- In first setup, $\widehat{\tau}_{\text{diff}} = 20$ or $\widehat{\tau}_{\text{diff}} = -20$ depending on the randomization.
- In second setup, $\widehat{\tau}_{\text{diff}} = 0$ in either randomization.

Estimating the sampling variance

- We can use sample variances within levels of D_i to estimate S_0^2 and S_1^2 :

$$\hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i=1}^n \mathbb{1}\{D_i = d\} (Y_i - \bar{Y}_d)^2$$

Estimating the sampling variance

- We can use sample variances within levels of D_i to estimate S_0^2 and S_1^2 :

$$\hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i=1}^n \mathbb{1}\{D_i = d\} (Y_i - \bar{Y}_d)^2$$

- Here, $\bar{Y}_0 = (1/n_0) \sum_{i=1}^n (1 - D_i) Y_i$ and $\bar{Y}_1 = (1/n_1) \sum_{i=1}^n D_i Y_i$.

Estimating the sampling variance

- We can use sample variances within levels of D_i to estimate S_0^2 and S_1^2 :

$$\hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i=1}^n \mathbb{1}\{D_i = d\} (Y_i - \bar{Y}_d)^2$$

- Here, $\bar{Y}_0 = (1/n_0) \sum_{i=1}^n (1 - D_i) Y_i$ and $\bar{Y}_1 = (1/n_1) \sum_{i=1}^n D_i Y_i$.
- But what about $S_{\tau_i}^2$?

Estimating the sampling variance

- We can use sample variances within levels of D_i to estimate S_0^2 and S_1^2 :

$$\hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i=1}^n \mathbb{1}\{D_i = d\} (Y_i - \bar{Y}_d)^2$$

- Here, $\bar{Y}_0 = (1/n_0) \sum_{i=1}^n (1 - D_i) Y_i$ and $\bar{Y}_1 = (1/n_1) \sum_{i=1}^n D_i Y_i$.
- But what about $S_{\tau_i}^2$?

$$S_{\tau_i}^2 = \frac{1}{n-1} \sum_{i=1}^n \underbrace{(Y_i(1) - Y_i(0))}_{\text{???}} - \tau_{fs})^2$$

Estimating the sampling variance

- We can use sample variances within levels of D_i to estimate S_0^2 and S_1^2 :

$$\hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i=1}^n \mathbb{1}\{D_i = d\} (Y_i - \bar{Y}_d)^2$$

- Here, $\bar{Y}_0 = (1/n_0) \sum_{i=1}^n (1 - D_i) Y_i$ and $\bar{Y}_1 = (1/n_1) \sum_{i=1}^n D_i Y_i$.
- But what about $S_{\tau_i}^2$?

$$S_{\tau_i}^2 = \frac{1}{n-1} \sum_{i=1}^n \underbrace{(Y_i(1) - Y_i(0))}_{???} - \tau_{fs})^2$$

- What to do?

Bounding the sampling variance

- First approach: find the worst possible (largest) variance.

Bounding the sampling variance

- First approach: find the worst possible (largest) variance.
- We can rewrite the variance as:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_{01} \right)$$

Bounding the sampling variance

- First approach: find the worst possible (largest) variance.
- We can rewrite the variance as:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{0}) = \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_{01} \right)$$

- Last term is the **covariance** between potential outcomes:

$$S_{01} = \frac{1}{n-1} \sum_{i=1}^n \{Y_i(1) - \bar{Y}(1)\} \{Y_i(0) - \bar{Y}(0)\}$$

Bounding the sampling variance

- First approach: find the worst possible (largest) variance.
- We can rewrite the variance as:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{O}) = \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_{01} \right)$$

- Last term is the **covariance** between potential outcomes:

$$S_{01} = \frac{1}{n-1} \sum_{i=1}^n \{Y_i(1) - \bar{Y}(1)\} \{Y_i(0) - \bar{Y}(0)\}$$

- We can use the **Cauchy-Schwarz** inequality: $S_{01} \leq S_0 S_1$

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{O}) \leq \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_0 S_1 \right) = \frac{n_0 n_1}{n} \left(\frac{S_0}{n_0} + \frac{S_1}{n_1} \right)^2$$

Bounding the sampling variance

- First approach: find the worst possible (largest) variance.
- We can rewrite the variance as:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{O}) = \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_{01} \right)$$

- Last term is the **covariance** between potential outcomes:

$$S_{01} = \frac{1}{n-1} \sum_{i=1}^n \{Y_i(1) - \bar{Y}(1)\} \{Y_i(0) - \bar{Y}(0)\}$$

- We can use the **Cauchy-Schwarz** inequality: $S_{01} \leq S_0 S_1$

$$\mathbb{V}(\widehat{\tau}_{\text{diff}} \mid \mathbf{O}) \leq \frac{1}{n} \left(\frac{n_1}{n_0} S_0^2 + \frac{n_0}{n_1} S_1^2 + 2S_0 S_1 \right) = \frac{n_0 n_1}{n} \left(\frac{S_0}{n_0} + \frac{S_1}{n_1} \right)^2$$

- Upper bound that is only a function of identified parameters.

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{V} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{V} \mid \mathbf{o}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{0}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{0})$, but that bias is always positive.

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{0}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{0})$, but that bias is always positive.
- Leads to **conservative inferences**:

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{0}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{0})$, but that bias is always positive.
- Leads to **conservative inferences**:
 - Standard errors, $\sqrt{\hat{\mathbb{V}}}$ will be at least as big as they should be.

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{0}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{0})$, but that bias is always positive.
- Leads to **conservative inferences**:
 - Standard errors, $\sqrt{\hat{\mathbb{V}}}$ will be at least as big as they should be.
 - Confidence intervals using $\sqrt{\hat{\mathbb{V}}}$ will be at least wide as they should be.

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{O}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{O})$, but that bias is always positive.
- Leads to **conservative inferences**:
 - Standard errors, $\sqrt{\hat{\mathbb{V}}}$ will be at least as big as they should be.
 - Confidence intervals using $\sqrt{\hat{\mathbb{V}}}$ will be at least wide as they should be.
 - Type I error rates will still be correct, power will be lower.

Conservative variance estimation

- Usual variance estimator is the Neyman (or robust) estimator:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \mathbb{E} [\hat{\mathbb{V}} \mid \mathbf{0}] = \frac{S_1^2}{n_1} + \frac{S_0^2}{n_0}$$

- Notice that $\hat{\mathbb{V}}$ is biased for $\mathbb{V}(\hat{\tau}_{\text{diff}} \mid \mathbf{0})$, but that bias is always positive.
- Leads to **conservative inferences**:
 - Standard errors, $\sqrt{\hat{\mathbb{V}}}$ will be at least as big as they should be.
 - Confidence intervals using $\sqrt{\hat{\mathbb{V}}}$ will be at least wide as they should be.
 - Type I error rates will still be correct, power will be lower.
 - Both will be exactly right if treatment effects are constant.

Inference in the Neyman approach

- If n is large, CLT will imply $\widehat{\tau}_{\text{diff}}$ will be approximately normal.

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

- Testing very similar to standard normal-approximation tests:

$$H_0 : \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = 0 \quad T = \frac{\hat{\tau}_{\text{diff}}}{\sqrt{\hat{V}}} \stackrel{a}{\sim} N(0, 1)$$

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

- Testing very similar to standard normal-approximation tests:

$$H_0 : \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = 0 \quad T = \frac{\hat{\tau}_{\text{diff}}}{\sqrt{\hat{V}}} \stackrel{a}{\sim} N(0, 1)$$

- Contains more situations than the sharp null, but...

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

- Testing very similar to standard normal-approximation tests:

$$H_0 : \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = 0 \quad T = \frac{\hat{\tau}_{\text{diff}}}{\sqrt{\hat{V}}} \stackrel{a}{\sim} N(0, 1)$$

- Contains more situations than the sharp null, but...
 - Fisher tests might not be well-powered against $\tau_{\text{fs}} = 0$ alternatives.

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

- Testing very similar to standard normal-approximation tests:

$$H_0 : \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = 0 \quad T = \frac{\hat{\tau}_{\text{diff}}}{\sqrt{\hat{V}}} \stackrel{a}{\sim} N(0, 1)$$

- Contains more situations than the sharp null, but...
 - Fisher tests might not be well-powered against $\tau_{\text{fs}} = 0$ alternatives.
- Can improve approximations using t -distribution.

Inference in the Neyman approach

- If n is large, CLT will imply $\hat{\tau}_{\text{diff}}$ will be approximately normal.
- Formulate confidence intervals in the usual way:

$$\text{CI}^{95}(\tau_{\text{fs}}) = (\hat{\tau}_{\text{diff}} - 1.96 \cdot \hat{V}^{1/2}, \hat{\tau}_{\text{diff}} + 1.96 \cdot \hat{V}^{1/2})$$

- Testing very similar to standard normal-approximation tests:

$$H_0 : \frac{1}{n} \sum_{i=1}^n Y_i(1) - Y_i(0) = 0 \quad T = \frac{\hat{\tau}_{\text{diff}}}{\sqrt{\hat{V}}} \stackrel{a}{\sim} N(0, 1)$$

- Contains more situations than the sharp null, but...
 - Fisher tests might not be well-powered against $\tau_{\text{fs}} = 0$ alternatives.
- Can improve approximations using t -distribution.
 - Works since \hat{V} will be approximately χ^2_{n-1} in large samples.

Population estimands

- What if we want to make inference to a (super)population?

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.
- Difference-in-means is **unbiased** across repeated samples:

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.
- Difference-in-means is **unbiased** across repeated samples:

$$\mathbb{E}[\widehat{\tau}_{\text{diff}}]$$

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.
- Difference-in-means is **unbiased** across repeated samples:

$$\mathbb{E}[\hat{\tau}_{\text{diff}}] = \underbrace{\mathbb{E}\{\mathbb{E}_D[\hat{\tau}_{\text{diff}} \mid \mathbf{O}]\}}_{\text{iterated expectations}}$$

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.
- Difference-in-means is **unbiased** across repeated samples:

$$\mathbb{E}[\widehat{\tau}_{\text{diff}}] = \underbrace{\mathbb{E}\{\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}]\}}_{\text{iterated expectations}} = \underbrace{\mathbb{E}[\tau_{\text{fs}}]}_{\text{SATE unbiasedness}}$$

Population estimands

- What if we want to make inference to a (super)population?
 - n units are a **simple random sample** from the population.
 - $Y_i(1), Y_i(0)$ are now random variables (induced by sampling)
- New goal: inference for the PATE, $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$.
 - Average of the SATEs across repeated samples: $\text{PATE} = \mathbb{E}[\text{SATE}]$.
- Difference-in-means is **unbiased** across repeated samples:

$$\mathbb{E}[\widehat{\tau}_{\text{diff}}] = \underbrace{\mathbb{E}\{\mathbb{E}_D[\widehat{\tau}_{\text{diff}} \mid \mathbf{O}]\}}_{\text{iterated expectations}} = \underbrace{\mathbb{E}[\tau_{\text{fs}}]}_{\text{SATE unbiasedness}} = \tau$$

Population sampling variance

- What about the sampling variance of $\hat{\tau}_{\text{diff}}$ when estimating the PATE?

Population sampling variance

- What about the sampling variance of $\hat{\tau}_{\text{diff}}$ when estimating the PATE?
 - Variation comes from random sample **and** random assignment.

Population sampling variance

- What about the sampling variance of $\hat{\tau}_{\text{diff}}$ when estimating the PATE?
 - Variation comes from random sample **and** random assignment.
- It turns out that the sampling variance of the estimator is simply:

$$\mathbb{V}(\hat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1} = \frac{\mathbb{V}[Y_i(0)]}{n_0} + \frac{\mathbb{V}[Y_i(1)]}{n_1}$$

Population sampling variance

- What about the sampling variance of $\widehat{\tau}_{\text{diff}}$ when estimating the PATE?
 - Variation comes from random sample **and** random assignment.
- It turns out that the sampling variance of the estimator is simply:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1} = \frac{\mathbb{V}[Y_i(0)]}{n_0} + \frac{\mathbb{V}[Y_i(1)]}{n_1}$$

- Here, σ_0^2 and σ_1^2 are the population-level variances of $Y_i(1)$ and $Y_i(0)$.

Population sampling variance

- What about the sampling variance of $\widehat{\tau}_{\text{diff}}$ when estimating the PATE?
 - Variation comes from random sample **and** random assignment.
- It turns out that the sampling variance of the estimator is simply:

$$\mathbb{V}(\widehat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1} = \frac{\mathbb{V}[Y_i(0)]}{n_0} + \frac{\mathbb{V}[Y_i(1)]}{n_1}$$

- Here, σ_0^2 and σ_1^2 are the population-level variances of $Y_i(1)$ and $Y_i(0)$.
- The variance of τ_i term drops out \rightsquigarrow higher variance for PATE than SATE.

Estimating pop. sampling variance

$$\mathbb{V}(\hat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1},$$

- Notice that the Neyman estimator $\hat{\mathbb{V}}$ is now unbiased for $\mathbb{V}(\hat{\tau}_{\text{diff}})$:

$$\hat{\mathbb{V}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}$$

Estimating pop. sampling variance

$$\mathbb{V}(\widehat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1},$$

- Notice that the Neyman estimator $\widehat{\mathbb{V}}$ is now unbiased for $\mathbb{V}(\widehat{\tau}_{\text{diff}})$:

$$\widehat{\mathbb{V}} = \frac{\widehat{\sigma}_0^2}{n_0} + \frac{\widehat{\sigma}_1^2}{n_1}$$

- Two interpretations of $\widehat{\mathbb{V}}$:

Estimating pop. sampling variance

$$\mathbb{V}(\widehat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1},$$

- Notice that the Neyman estimator $\widehat{\mathbb{V}}$ is now unbiased for $\mathbb{V}(\widehat{\tau}_{\text{diff}})$:

$$\widehat{\mathbb{V}} = \frac{\widehat{\sigma}_0^2}{n_0} + \frac{\widehat{\sigma}_1^2}{n_1}$$

- Two interpretations of $\widehat{\mathbb{V}}$:
 1. Unbiased estimator for sampling variance of the traditional estimator of the PATE

Estimating pop. sampling variance

$$\mathbb{V}(\widehat{\tau}_{\text{diff}}) = \frac{\sigma_0^2}{n_0} + \frac{\sigma_1^2}{n_1},$$

- Notice that the Neyman estimator $\widehat{\mathbb{V}}$ is now unbiased for $\mathbb{V}(\widehat{\tau}_{\text{diff}})$:

$$\widehat{\mathbb{V}} = \frac{\widehat{\sigma}_0^2}{n_0} + \frac{\widehat{\sigma}_1^2}{n_1}$$

- Two interpretations of $\widehat{\mathbb{V}}$:
 1. Unbiased estimator for sampling variance of the traditional estimator of the PATE
 2. Conservative estimator for the sampling variance of the traditional estimator of the SATE