

Module 6(b): Two Stage Least Squares

Fall 2021

Matthew Blackwell

Gov 2003 (Harvard)

1/ Basic two-stage least squares

- **Two stage least squares** (TSLs) is the classical approach to IV.

- **Two stage least squares** (TSLS) is the classical approach to IV.
- Basic idea is to assume two constant effects linear models:

$$Y_i = \alpha + \tau D_i + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \eta_i$$

- **Two stage least squares** (TSLs) is the classical approach to IV.
- Basic idea is to assume two constant effects linear models:

$$Y_i = \alpha + \tau D_i + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \eta_i$$

- Here the treatment D_i is **endogenous** so $\mathbb{E}[\varepsilon_i \mid D_i] \neq 0$

- **Two stage least squares** (TSLs) is the classical approach to IV.
- Basic idea is to assume two constant effects linear models:

$$Y_i = \alpha + \tau D_i + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \eta_i$$

- Here the treatment D_i is **endogenous** so $\mathbb{E}[\varepsilon_i \mid D_i] \neq 0$
- But we have an **instrument** Z_i that is exogenous $\mathbb{E}[\varepsilon_i \mid Z_i] = 0$

- **Two stage least squares** (TSLs) is the classical approach to IV.
- Basic idea is to assume two constant effects linear models:

$$Y_i = \alpha + \tau D_i + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \eta_i$$

- Here the treatment D_i is **endogenous** so $\mathbb{E}[\varepsilon_i \mid D_i] \neq 0$
- But we have an **instrument** Z_i that is exogenous $\mathbb{E}[\varepsilon_i \mid Z_i] = 0$
 - It also is exogenous for treatment, so $\mathbb{E}[\eta_i \mid Z_i] = 0$.

- **Two stage least squares** (TSLs) is the classical approach to IV.
- Basic idea is to assume two constant effects linear models:

$$Y_i = \alpha + \tau D_i + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \eta_i$$

- Here the treatment D_i is **endogenous** so $\mathbb{E}[\varepsilon_i \mid D_i] \neq 0$
- But we have an **instrument** Z_i that is exogenous $\mathbb{E}[\varepsilon_i \mid Z_i] = 0$
 - It also is exogenous for treatment, so $\mathbb{E}[\eta_i \mid Z_i] = 0$.
- This implies the following CEF form for Y_i conditional on Z_i :

$$\mathbb{E}[Y_i \mid Z_i] = \alpha + \tau \mathbb{E}[D_i \mid Z_i] = \alpha + \tau \cdot (\gamma Z_i)$$

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

- Applying this to above CEF we have:

$$\tau = \frac{\text{cov}(Y_i, \gamma Z_i)}{\mathbb{V}[\gamma Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\gamma \mathbb{V}[Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)}$$

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

- Applying this to above CEF we have:

$$\tau = \frac{\text{cov}(Y_i, \gamma Z_i)}{\mathbb{V}[\gamma Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\gamma \mathbb{V}[Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)}$$

- TSLS estimator:

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

- Applying this to above CEF we have:

$$\tau = \frac{\text{cov}(Y_i, \gamma Z_i)}{\mathbb{V}[\gamma Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\gamma \mathbb{V}[Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)}$$

- TSLS estimator:
 - Estimate $\hat{\gamma}$ from regression of treatment D_i on instrument Z_i

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

- Applying this to above CEF we have:

$$\tau = \frac{\text{cov}(Y_i, \gamma Z_i)}{\mathbb{V}[\gamma Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\gamma \mathbb{V}[Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)}$$

- TSLS estimator:
 - Estimate $\hat{\gamma}$ from regression of treatment D_i on instrument Z_i
 - Estimate $\hat{\tau}_{2SLS}$ as the slope of a regression of Y_i on $\hat{\gamma} Z_i$

TSLS estimands

- Under the model, we have the following CEF: $\mathbb{E}[Y_i | Z_i] = \alpha + \tau \cdot (\gamma Z_i)$
 - \rightsquigarrow a regression of Y_i on γZ_i would have τ as the slope.
- If the CEF is linear, we have this simple relationship slopes:

$$\mathbb{E}[D_i | Z_i] = \delta + \gamma Z_i \quad \rightsquigarrow \quad \gamma = \frac{\text{cov}(D_i, Z_i)}{\mathbb{V}[Z_i]}$$

- Applying this to above CEF we have:

$$\tau = \frac{\text{cov}(Y_i, \gamma Z_i)}{\mathbb{V}[\gamma Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\gamma \mathbb{V}[Z_i]} = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)}$$

- TSLS estimator:
 - Estimate $\hat{\gamma}$ from regression of treatment D_i on instrument Z_i
 - Estimate $\hat{\tau}_{2SLS}$ as the slope of a regression of Y_i on $\hat{\gamma} Z_i$
 - Under this model, $\hat{\tau}_{2SLS} \xrightarrow{p} \tau$ (but don't use SEs from second stage)

Binary treatment and instrument

- Under binary treatment/instrument, TSLS estimand is the LATE:

$$\tau = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)} = \frac{\mathbb{E}[Y_i | Z_i = 1] - \mathbb{E}[Y_i | Z_i = 0]}{\mathbb{E}[D_i | Z_i = 1] - \mathbb{E}[D_i | Z_i = 0]} = \frac{\text{ITT}_Y}{\text{ITT}_D} = \tau_{\text{LATE}}$$

Binary treatment and instrument

- Under binary treatment/instrument, TSLS estimand is the LATE:

$$\tau = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)} = \frac{\mathbb{E}[Y_i | Z_i = 1] - \mathbb{E}[Y_i | Z_i = 0]}{\mathbb{E}[D_i | Z_i = 1] - \mathbb{E}[D_i | Z_i = 0]} = \frac{\text{ITT}_Y}{\text{ITT}_D} = \tau_{\text{LATE}}$$

- And the TSLS estimator is the Wald estimator:

$$\hat{\tau}_{\text{TSLS}} = \frac{\widehat{\text{cov}}(Y_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} = \frac{\bar{Y}_1 - \bar{Y}_0}{\bar{D}_1 - \bar{D}_0} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_D} = \hat{\tau}_{\text{iv}}$$

Binary treatment and instrument

- Under binary treatment/instrument, TSLS estimand is the LATE:

$$\tau = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)} = \frac{\mathbb{E}[Y_i | Z_i = 1] - \mathbb{E}[Y_i | Z_i = 0]}{\mathbb{E}[D_i | Z_i = 1] - \mathbb{E}[D_i | Z_i = 0]} = \frac{\text{ITT}_Y}{\text{ITT}_D} = \tau_{\text{LATE}}$$

- And the TSLS estimator is the Wald estimator:

$$\hat{\tau}_{2SLS} = \frac{\widehat{\text{cov}}(Y_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} = \frac{\bar{Y}_1 - \bar{Y}_0}{\bar{D}_1 - \bar{D}_0} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_D} = \hat{\tau}_{iv}$$

- \rightsquigarrow constant effects model not required for TSLS in this setting.

Binary treatment and instrument

- Under binary treatment/instrument, TSLS estimand is the LATE:

$$\tau = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)} = \frac{\mathbb{E}[Y_i | Z_i = 1] - \mathbb{E}[Y_i | Z_i = 0]}{\mathbb{E}[D_i | Z_i = 1] - \mathbb{E}[D_i | Z_i = 0]} = \frac{\text{ITT}_Y}{\text{ITT}_D} = \tau_{\text{LATE}}$$

- And the TSLS estimator is the Wald estimator:

$$\hat{\tau}_{\text{TSLS}} = \frac{\widehat{\text{cov}}(Y_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} = \frac{\bar{Y}_1 - \bar{Y}_0}{\bar{D}_1 - \bar{D}_0} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_D} = \hat{\tau}_{\text{iv}}$$

- \rightsquigarrow constant effects model not required for TSLS in this setting.
- But we need constant effects when we add covariates:

$$Y_i = \alpha + \tau D_i + \mathbf{X}_i' \beta_y + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \mathbf{X}_i' \beta_d + \eta_i$$

Binary treatment and instrument

- Under binary treatment/instrument, TSLS estimand is the LATE:

$$\tau = \frac{\text{cov}(Y_i, Z_i)}{\text{cov}(D_i, Z_i)} = \frac{\mathbb{E}[Y_i | Z_i = 1] - \mathbb{E}[Y_i | Z_i = 0]}{\mathbb{E}[D_i | Z_i = 1] - \mathbb{E}[D_i | Z_i = 0]} = \frac{\text{ITT}_Y}{\text{ITT}_D} = \tau_{\text{LATE}}$$

- And the TSLS estimator is the Wald estimator:

$$\hat{\tau}_{\text{TSLS}} = \frac{\widehat{\text{cov}}(Y_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} = \frac{\bar{Y}_1 - \bar{Y}_0}{\bar{D}_1 - \bar{D}_0} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_D} = \hat{\tau}_{\text{iv}}$$

- \rightsquigarrow constant effects model not required for TSLS in this setting.
- But we need constant effects when we add covariates:

$$Y_i = \alpha + \tau D_i + \mathbf{X}_i' \beta_y + \varepsilon_i$$

$$D_i = \delta + \gamma Z_i + \mathbf{X}_i' \beta_d + \eta_i$$

- Otherwise, τ is an odd weighted function of causal effects and $\tau \neq \tau_{\text{LATE}}$

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$Y_i = \tau D_i + \varepsilon_i \qquad \mathbb{E}[\varepsilon_i \mid D_i] \neq 0$$

$$D_i = 0 \times Z_i + \eta_i \quad \mathbb{E}[\varepsilon_i \mid Z_i] = \mathbb{E}[\eta_i \mid Z_i] = 0$$

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$Y_i = \tau D_i + \varepsilon_i \qquad \mathbb{E}[\varepsilon_i \mid D_i] \neq 0$$

$$D_i = 0 \times Z_i + \eta_i \quad \mathbb{E}[\varepsilon_i \mid Z_i] = \mathbb{E}[\eta_i \mid Z_i] = 0$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$\begin{aligned} Y_i &= \tau D_i + \varepsilon_i & \mathbb{E}[\varepsilon_i \mid D_i] &\neq 0 \\ D_i &= 0 \times Z_i + \eta_i & \mathbb{E}[\varepsilon_i \mid Z_i] = \mathbb{E}[\eta_i \mid Z_i] &= 0 \end{aligned}$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.
- We can write the bias of the Wald estimator as:

$$\widehat{\tau}_{iv} - \tau = \frac{\widehat{\text{cov}}(\tau D_i + \varepsilon_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} - \tau$$

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$\begin{aligned} Y_i &= \tau D_i + \varepsilon_i & \mathbb{E}[\varepsilon_i \mid D_i] &\neq 0 \\ D_i &= 0 \times Z_i + \eta_i & \mathbb{E}[\varepsilon_i \mid Z_i] &= \mathbb{E}[\eta_i \mid Z_i] = 0 \end{aligned}$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.
- We can write the bias of the Wald estimator as:

$$\widehat{\tau}_{iv} - \tau = \frac{\widehat{\text{cov}}(\tau D_i + \varepsilon_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} - \tau = \frac{\frac{1}{n} \sum_{i=1}^n \varepsilon_i Z_i}{\frac{1}{n} \sum_{i=1}^n \eta_i Z_i}$$

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$\begin{aligned} Y_i &= \tau D_i + \varepsilon_i & \mathbb{E}[\varepsilon_i \mid D_i] &\neq 0 \\ D_i &= 0 \times Z_i + \eta_i & \mathbb{E}[\varepsilon_i \mid Z_i] &= \mathbb{E}[\eta_i \mid Z_i] = 0 \end{aligned}$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.
- We can write the bias of the Wald estimator as:

$$\widehat{\tau}_{iv} - \tau = \frac{\widehat{\text{cov}}(\tau D_i + \varepsilon_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} - \tau = \frac{\frac{1}{n} \sum_{i=1}^n \varepsilon_i Z_i}{\frac{1}{n} \sum_{i=1}^n \eta_i Z_i} \xrightarrow{d} \underbrace{\frac{\text{cov}(\varepsilon_i, \eta_i)}{\mathbb{V}[\varepsilon_i]}}_{\text{bias}} + \underbrace{W_i}_{\text{Cauchy r.v.}}$$

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$\begin{aligned} Y_i &= \tau D_i + \varepsilon_i & \mathbb{E}[\varepsilon_i \mid D_i] &\neq 0 \\ D_i &= 0 \times Z_i + \eta_i & \mathbb{E}[\varepsilon_i \mid Z_i] &= \mathbb{E}[\eta_i \mid Z_i] = 0 \end{aligned}$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.
- We can write the bias of the Wald estimator as:

$$\widehat{\tau}_{iv} - \tau = \frac{\widehat{\text{cov}}(\tau D_i + \varepsilon_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} - \tau = \frac{\frac{1}{n} \sum_{i=1}^n \varepsilon_i Z_i}{\frac{1}{n} \sum_{i=1}^n \eta_i Z_i} \xrightarrow{d} \underbrace{\frac{\text{cov}(\varepsilon_i, \eta_i)}{\mathbb{V}[\varepsilon_i]}}_{\text{bias}} + \underbrace{W_i}_{\text{Cauchy r.v.}}$$

- Inconsistent and asymptotically heavy tails (bc of Cauchy)

Weak instruments

- IV is unstable when instrument weakly affects treatment $\text{cov}(D_i, Z_i) \approx 0$.
- **Example** completely irrelevant instrument:

$$\begin{aligned} Y_i &= \tau D_i + \varepsilon_i & \mathbb{E}[\varepsilon_i \mid D_i] &\neq 0 \\ D_i &= 0 \times Z_i + \eta_i & \mathbb{E}[\varepsilon_i \mid Z_i] &= \mathbb{E}[\eta_i \mid Z_i] = 0 \end{aligned}$$

- Note that we only assume mean independence, so $\text{cov}(D_i, Z_i)$ could be nonzero.
- We can write the bias of the Wald estimator as:

$$\widehat{\tau}_{iv} - \tau = \frac{\widehat{\text{cov}}(\tau D_i + \varepsilon_i, Z_i)}{\widehat{\text{cov}}(D_i, Z_i)} - \tau = \frac{\frac{1}{n} \sum_{i=1}^n \varepsilon_i Z_i}{\frac{1}{n} \sum_{i=1}^n \eta_i Z_i} \xrightarrow{d} \underbrace{\frac{\text{cov}(\varepsilon_i, \eta_i)}{\mathbb{V}[\varepsilon_i]}}_{\text{bias}} + \underbrace{W_i}_{\text{Cauchy r.v.}}$$

- Inconsistent and asymptotically heavy tails (bc of Cauchy)
 - When $Z \rightarrow D$ effect is small but non-zero we see similar behavior.

What to do about weak instruments?

- Detecting weak instruments:

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff
- Anderson-Rubin (1949) test (simplified setting, binary Z/D)

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff
- Anderson-Rubin (1949) test (simplified setting, binary Z/D)
 - $H_0 : \tau = \tau_0$ equivalent to $H_0 : \text{ITT}_Y - \text{ITT}_D \cdot \tau_0 = 0$

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff
- Anderson-Rubin (1949) test (simplified setting, binary Z/D)
 - $H_0 : \tau = \tau_0$ equivalent to $H_0 : \text{ITT}_Y - \text{ITT}_D \cdot \tau_0 = 0$
 - Under the null, asymptotically we have

$$g(\tau_0) = \widehat{\text{ITT}}_Y - \widehat{\text{ITT}}_D \tau_0 \sim N(0, \Omega(\tau_0))$$

$$\Omega(\tau_0) = \mathbb{V}[\widehat{\text{ITT}}_Y] + \tau_0^2 \mathbb{V}[\widehat{\text{ITT}}_D] - 2\tau_0 \text{cov}(\widehat{\text{ITT}}_Y, \widehat{\text{ITT}}_D)$$

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff
- Anderson-Rubin (1949) test (simplified setting, binary Z/D)
 - $H_0 : \tau = \tau_0$ equivalent to $H_0 : \text{ITT}_Y - \text{ITT}_D \cdot \tau_0 = 0$
 - Under the null, asymptotically we have

$$g(\tau_0) = \widehat{\text{ITT}}_Y - \widehat{\text{ITT}}_D \tau_0 \sim N(0, \Omega(\tau_0))$$

$$\Omega(\tau_0) = \mathbb{V}[\widehat{\text{ITT}}_Y] + \tau_0^2 \mathbb{V}[\widehat{\text{ITT}}_D] - 2\tau_0 \text{cov}(\widehat{\text{ITT}}_Y, \widehat{\text{ITT}}_D)$$

- AR test statistic: $g(\tau_0)^2 / \Omega(\tau_0) \sim \chi^2$ no matter first-stage effect.

What to do about weak instruments?

- Detecting weak instruments:
 - F-test on instruments (excluded from second stage): $H_0 : \gamma = 0$.
 - Rule of thumb: bias is small when F-stat ≥ 10 (Stock & Yogo, 2005)
 - Correct coverage may require cutoff $F \geq 104.7$ (Lee et al, 2020)
 - The latter is a worst-case, typical data maybe ok with 10 cutoff
- Anderson-Rubin (1949) test (simplified setting, binary Z/D)
 - $H_0 : \tau = \tau_0$ equivalent to $H_0 : \text{ITT}_Y - \text{ITT}_D \cdot \tau_0 = 0$
 - Under the null, asymptotically we have

$$g(\tau_0) = \widehat{\text{ITT}}_Y - \widehat{\text{ITT}}_D \tau_0 \sim N(0, \Omega(\tau_0))$$
$$\Omega(\tau_0) = \mathbb{V}[\widehat{\text{ITT}}_Y] + \tau_0^2 \mathbb{V}[\widehat{\text{ITT}}_D] - 2\tau_0 \text{cov}(\widehat{\text{ITT}}_Y, \widehat{\text{ITT}}_D)$$

- AR test statistic: $g(\tau_0)^2 / \Omega(\tau_0) \sim \chi^2$ no matter first-stage effect.
- Can invert (analytically!) to get confidence intervals

Multi-valued treatments

- Generalization of these ideas:

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata
 - Affected: $(D_i(0), D_i(1)) \in \{(0, 1), (0, 2), (1, 2)\}$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata
 - Affected: $(D_i(0), D_i(1)) \in \{(0, 1), (0, 2), (1, 2)\}$
 - Unaffected: $(D_i(0), D_i(1)) \in \{(0, 0), (1, 1), (2, 2)\}$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata
 - Affected: $(D_i(0), D_i(1)) \in \{(0, 1), (0, 2), (1, 2)\}$
 - Unaffected: $(D_i(0), D_i(1)) \in \{(0, 0), (1, 1), (2, 2)\}$
 - Negatively affected: $(D_i(0), D_i(1)) \in \{(1, 0), (2, 0), (2, 1)\}$

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata
 - Affected: $(D_i(0), D_i(1)) \in \{(0, 1), (0, 2), (1, 2)\}$
 - Unaffected: $(D_i(0), D_i(1)) \in \{(0, 0), (1, 1), (2, 2)\}$
 - Negatively affected: $(D_i(0), D_i(1)) \in \{(1, 0), (2, 0), (2, 1)\}$
 - Last ruled out by monotonicity.

Multi-valued treatments

- Generalization of these ideas:
 - Multi-valued treatment: $D_i \in \{0, 1, \dots, K - 1\}$
 - Binary instrument: $Z_i \in \{0, 1\}$
- Assumptions:
 - Randomization: $[\{Y_i(d, z), \forall d, z\}, D_i(1), D_i(0)] \perp\!\!\!\perp Z_i$
 - Monotonicity: $D_i(1) \geq D_i(0)$ (instrument only increases treatment)
 - Exclusion restriction: $Y_i(1, d) = Y_i(0, d)$ for all $d = 0, 1, \dots, K - 1$
- Can't identify the proportion of all compliance types here.
- Example: $K = 3 \rightsquigarrow 9$ principal strata
 - Affected: $(D_i(0), D_i(1)) \in \{(0, 1), (0, 2), (1, 2)\}$
 - Unaffected: $(D_i(0), D_i(1)) \in \{(0, 0), (1, 1), (2, 2)\}$
 - Negatively affected: $(D_i(0), D_i(1)) \in \{(1, 0), (2, 0), (2, 1)\}$
 - Last ruled out by monotonicity.
 - 5 unknowns and 4 knowns under monotonicity.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.
 - Let $\rho_{jk} = \mathbb{P}(D_i(1) = j, D_i(0) = k)$ be the strata size.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.
 - Let $\rho_{jk} = \mathbb{P}(D_i(1) = j, D_i(0) = k)$ be the strata size.
- We can show that the 2SLS estimator converges to:

$$\hat{\tau}_{2SLS} \xrightarrow{p} \sum_{k=0}^{K-1} \sum_{j=k+1}^{K-1} \omega_{jk} \mathbb{E} \left(\frac{Y_i(1) - Y_i(0)}{j - k} \mid C_i = jk \right)$$
$$\omega_{jk} = \frac{(j - k)\rho_{jk}}{\sum_{s=0}^{K-1} \sum_{t=s+1}^{K-1} (s - t)\rho_{st}}$$

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.
 - Let $\rho_{jk} = \mathbb{P}(D_i(1) = j, D_i(0) = k)$ be the strata size.
- We can show that the 2SLS estimator converges to:

$$\hat{\tau}_{2SLS} \xrightarrow{p} \sum_{k=0}^{K-1} \sum_{j=k+1}^{K-1} \omega_{jk} \mathbb{E} \left(\frac{Y_i(1) - Y_i(0)}{j - k} \mid C_i = jk \right)$$
$$\omega_{jk} = \frac{(j - k)\rho_{jk}}{\sum_{s=0}^{K-1} \sum_{t=s+1}^{K-1} (s - t)\rho_{st}}$$

- Intuition: a weighted average of effects per dose for each affected type.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.
 - Let $\rho_{jk} = \mathbb{P}(D_i(1) = j, D_i(0) = k)$ be the strata size.
- We can show that the 2SLS estimator converges to:

$$\hat{\tau}_{2SLS} \xrightarrow{p} \sum_{k=0}^{K-1} \sum_{j=k+1}^{K-1} \omega_{jk} \mathbb{E} \left(\frac{Y_i(1) - Y_i(0)}{j - k} \mid C_i = jk \right)$$
$$\omega_{jk} = \frac{(j - k)\rho_{jk}}{\sum_{s=0}^{K-1} \sum_{t=s+1}^{K-1} (s - t)\rho_{st}}$$

- Intuition: a weighted average of effects per dose for each affected type.
 - Weights are proportional to size of the strata and how big the effect of the instrument is for that strata.

TSLS with multivalued treatments

- Let $C_i = jk$ be an indicator for compliance type $D_i(1) = j$ and $D_i(0) = k$.
 - People that are moved from k to j by the instrument.
 - Let $\rho_{jk} = \mathbb{P}(D_i(1) = j, D_i(0) = k)$ be the strata size.
- We can show that the 2SLS estimator converges to:

$$\hat{\tau}_{2SLS} \xrightarrow{p} \sum_{k=0}^{K-1} \sum_{j=k+1}^{K-1} \omega_{jk} \mathbb{E} \left(\frac{Y_i(1) - Y_i(0)}{j - k} \mid C_i = jk \right)$$
$$\omega_{jk} = \frac{(j - k)\rho_{jk}}{\sum_{s=0}^{K-1} \sum_{t=s+1}^{K-1} (s - t)\rho_{st}}$$

- Intuition: a weighted average of effects per dose for each affected type.
 - Weights are proportional to size of the strata and how big the effect of the instrument is for that strata.
 - If instrument can only increase by 1 dose, then simplifies to weighted average of principal strata effects.

2/ General two-stage least squares

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

General 2SLS

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.

General 2SLS

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i \mid \mathbf{Z}_i] = 0$.

General 2SLS

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i \mid \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i \mid \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.
 - Rank condition: $\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i']$ and $\mathbb{E}[\mathbf{X}_i \mathbf{Z}_i']$ have full rank.

General 2SLS

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i \mid \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.
 - Rank condition: $\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i']$ and $\mathbb{E}[\mathbf{X}_i \mathbf{Z}_i']$ have full rank.
- Identification:

General 2SLS

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i | \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i | \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.
 - Rank condition: $\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i']$ and $\mathbb{E}[\mathbf{X}_i \mathbf{Z}_i']$ have full rank.
- Identification:
 - $k = \ell$: just-identified.

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i \mid \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i \mid \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.
 - Rank condition: $\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i']$ and $\mathbb{E}[\mathbf{X}_i \mathbf{Z}_i']$ have full rank.
- Identification:
 - $k = \ell$: just-identified.
 - $k < \ell$: over-identified (can test the exclusion restriction, kinda)

- Linear model for each i :

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

- \mathbf{X}_i is $k \times 1$ now includes D_i and any pretreatment covariates.
- Parts of \mathbf{X}_i are endogenous so that $\mathbb{E}[\varepsilon_i | \mathbf{X}_i] \neq 0$
- Instruments \mathbf{Z}_i that is $\ell \times 1$ vector such that $\mathbb{E}[\varepsilon_i | \mathbf{Z}_i] = 0$.
 - \mathbf{Z}_i might include exogenous/pretreatment variables from \mathbf{X}_i as well.
 - Rank condition: $\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i']$ and $\mathbb{E}[\mathbf{X}_i \mathbf{Z}_i']$ have full rank.
- Identification:
 - $k = \ell$: just-identified.
 - $k < \ell$: over-identified (can test the exclusion restriction, kinda)
 - $k > \ell$: unidentified (fails rank condition)

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \boldsymbol{\beta} + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \boldsymbol{\beta} + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta} + \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i]$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \boldsymbol{\beta} + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta} + \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta} + \mathbf{\Pi}' \mathbb{E}[\mathbf{Z}_i \varepsilon_i]$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \boldsymbol{\beta} + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \boldsymbol{\beta} + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta} + \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta} + \mathbf{\Pi}' \mathbb{E}[\mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \boldsymbol{\beta}$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \beta + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta + \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta + \mathbf{\Pi}' \mathbb{E}[\mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta$$

$$\mathbb{E}[\tilde{\mathbf{X}}_i Y_i] = \mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}_i'] \beta$$

Nasty Matrix Algebra

- Projection matrix projects values of \mathbf{X}_i onto \mathbf{Z}_i :

$$\mathbf{\Pi} = (\mathbb{E}[\mathbf{Z}_i \mathbf{Z}_i'])^{-1} \mathbb{E}[\mathbf{Z}_i \mathbf{X}_i'] \quad (\text{projection matrix})$$

$$\tilde{\mathbf{X}}_i = \mathbf{\Pi}' \mathbf{Z}_i \quad (\text{projected values})$$

- To derive the 2SLS estimator, take the fitted values, $\mathbf{\Pi}' \mathbf{Z}_i$ and multiply both sides of the outcome equation by them:

$$Y_i = \mathbf{X}_i' \beta + \varepsilon_i$$

$$\mathbf{\Pi}' \mathbf{Z}_i Y_i = \mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i' \beta + \mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta + \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta + \mathbf{\Pi}' \mathbb{E}[\mathbf{Z}_i \varepsilon_i]$$

$$\mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i Y_i] = \mathbb{E}[\mathbf{\Pi}' \mathbf{Z}_i \mathbf{X}_i'] \beta$$

$$\mathbb{E}[\tilde{\mathbf{X}}_i Y_i] = \mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}_i'] \beta$$

$$\beta = (\mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}_i'])^{-1} \mathbb{E}[\tilde{\mathbf{X}}_i Y_i]$$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$
 - Fitted values of regression of \mathbb{X} on \mathbb{Z} .

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$
 - Fitted values of regression of \mathbb{X} on \mathbb{Z} .
 - Matrix party trick: $\mathbb{X}'\mathbb{Z}/n = (1/n) \sum_i \mathbf{X}_i \mathbf{Z}'_i \xrightarrow{p} \mathbb{E}[\mathbf{X}_i \mathbf{Z}'_i]$.

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$
 - Fitted values of regression of \mathbb{X} on \mathbb{Z} .
 - Matrix party trick: $\mathbb{X}'\mathbb{Z}/n = (1/n) \sum_i \mathbf{X}_i \mathbf{Z}'_i \xrightarrow{P} \mathbb{E}[\mathbf{X}_i \mathbf{Z}'_i]$.
- Take the population formula for the parameters:

$$\beta = (\mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}'_i])^{-1} \mathbb{E}[\tilde{\mathbf{X}}_i Y_i]$$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$
 - Fitted values of regression of \mathbb{X} on \mathbb{Z} .
 - Matrix party trick: $\mathbb{X}'\mathbb{Z}/n = (1/n) \sum_i \mathbf{X}_i \mathbf{Z}'_i \xrightarrow{P} \mathbb{E}[\mathbf{X}_i \mathbf{Z}'_i]$.
- Take the population formula for the parameters:

$$\beta = (\mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}'_i])^{-1} \mathbb{E}[\tilde{\mathbf{X}}_i Y_i]$$

- And plug in the sample values (the n cancels out):

$$\hat{\beta}_{2SLS} = (\widehat{\mathbb{X}}'\widehat{\mathbb{X}})^{-1}\widehat{\mathbb{X}}'\mathbf{y} \xrightarrow{P} \beta$$

How to estimate the parameters

- Collect \mathbf{X}_i into a $n \times k$ matrix $\mathbb{X} = (\mathbf{X}'_1, \dots, \mathbf{X}'_n)$
- Collect \mathbf{Z}_i into a $n \times \ell$ matrix $\mathbb{Z} = (\mathbf{Z}'_1, \dots, \mathbf{Z}'_n)$
- In-sample projection matrix produces fitted values: $\widehat{\mathbb{X}} = \mathbb{Z}(\mathbb{Z}'\mathbb{Z})^{-1}\mathbb{Z}'\mathbb{X}$
 - Fitted values of regression of \mathbb{X} on \mathbb{Z} .
 - Matrix party trick: $\mathbb{X}'\mathbb{Z}/n = (1/n) \sum_i \mathbf{X}_i \mathbf{Z}'_i \xrightarrow{P} \mathbb{E}[\mathbf{X}_i \mathbf{Z}'_i]$.
- Take the population formula for the parameters:

$$\beta = (\mathbb{E}[\tilde{\mathbf{X}}_i \mathbf{X}'_i])^{-1} \mathbb{E}[\tilde{\mathbf{X}}_i Y_i]$$

- And plug in the sample values (the n cancels out):

$$\hat{\beta}_{2SLS} = (\widehat{\mathbb{X}}'\widehat{\mathbb{X}})^{-1}\widehat{\mathbb{X}}'\mathbf{y} \xrightarrow{P} \beta$$

- This is how R/Stata estimates the 2SLS parameters

Asymptotic variance for 2SLS

- We can write the centered, normalized TSLS estimator as:

$$\sqrt{n}(\hat{\beta}_{2SLS} - \beta) = \underbrace{\left(n^{-1} \sum_i \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'\right)^{-1}}_{\xrightarrow{p} (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}} \underbrace{\left(n^{-1/2} \sum_i \hat{\mathbf{X}}_i \varepsilon_i\right)}_{\xrightarrow{d} N(0, \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i \varepsilon_i \hat{\mathbf{X}}_i])}$$

Asymptotic variance for 2SLS

- We can write the centered, normalized TSLS estimator as:

$$\sqrt{n}(\hat{\beta}_{2SLS} - \beta) = \underbrace{\left(n^{-1} \sum_i \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'\right)^{-1}}_{\xrightarrow{p} (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}} \underbrace{\left(n^{-1/2} \sum_i \hat{\mathbf{X}}_i \varepsilon_i\right)}_{\xrightarrow{d} N(0, \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i])}$$

- Thus, we have that $\sqrt{n}(\hat{\beta}_{2SLS} - \beta)$ has asymptotic variance:

$$(\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1} \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i] (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}$$

Asymptotic variance for 2SLS

- We can write the centered, normalized TSLS estimator as:

$$\sqrt{n}(\hat{\beta}_{2SLS} - \beta) = \underbrace{\left(n^{-1} \sum_i \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'\right)^{-1}}_{\xrightarrow{p} (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}} \underbrace{\left(n^{-1/2} \sum_i \hat{\mathbf{X}}_i \varepsilon_i\right)}_{\xrightarrow{d} N(0, \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i])}$$

- Thus, we have that $\sqrt{n}(\hat{\beta}_{2SLS} - \beta)$ has asymptotic variance:

$$(\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1} \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i] (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}$$

- **Robust 2SLS variance estimator** with residuals $\hat{u}_i = Y_i - \mathbf{X}_i' \hat{\beta}$:

$$\widehat{\text{var}}(\hat{\beta}_{2SLS}) = (\widehat{\mathbb{X}}' \widehat{\mathbb{X}})^{-1} \left(\sum_i \hat{u}_i^2 \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i' \right) (\widehat{\mathbb{X}}' \widehat{\mathbb{X}})^{-1}$$

Asymptotic variance for 2SLS

- We can write the centered, normalized TSLS estimator as:

$$\sqrt{n}(\hat{\beta}_{2SLS} - \beta) = \underbrace{\left(n^{-1} \sum_i \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'\right)^{-1}}_{\xrightarrow{p} (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}} \underbrace{\left(n^{-1/2} \sum_i \hat{\mathbf{X}}_i \varepsilon_i\right)}_{\xrightarrow{d} N(0, \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i])}$$

- Thus, we have that $\sqrt{n}(\hat{\beta}_{2SLS} - \beta)$ has asymptotic variance:

$$(\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1} \mathbb{E}[\hat{\mathbf{X}}_i' \varepsilon_i' \varepsilon_i \hat{\mathbf{X}}_i] (\mathbb{E}[\hat{\mathbf{X}}_i \hat{\mathbf{X}}_i'])^{-1}$$

- Robust 2SLS variance estimator** with residuals $\hat{u}_i = Y_i - \mathbf{X}_i' \hat{\beta}$:

$$\widehat{\text{var}}(\hat{\beta}_{2SLS}) = (\widehat{\mathbb{X}}' \widehat{\mathbb{X}})^{-1} \left(\sum_i \hat{u}_i^2 \hat{\mathbf{X}}_i \hat{\mathbf{X}}_i' \right) (\widehat{\mathbb{X}}' \widehat{\mathbb{X}})^{-1}$$

- HC2, clustering, and autocorrelation versions exist