

Empirical Methods for Policy Evaluation

Matteo Bobba

Toulouse School of Economics (TSE)

TSE PhD Program (MRes)

Fall 2026

Part 2: Regression Discontinuity Designs and School Choice Models

① RD methods

- ⇒ Local randomization approach
- ⇒ Continuity-based approach
- ⇒ Extensions

② School choice models

- ⇒ Agarwal-Somaini (ARE, 2020)


③ Teacher sorting and student outcomes

- ⇒ Bobba-Ederer-Leon-Nielson-Nieddu (JPE, forthcoming)

Setup and Notation

- Potential outcomes: $Y_i(1), Y_i(0)$, with $\tau_i = Y_i(1) - Y_i(0)$
- Running variable (score): X_i
- Treatment indicator: $D_i = D_i(X_i) = 1$ if treated, 0 otherwise
- Observed outcome: $Y_i = Y_i(1)D_i + Y_i(0)(1 - D_i)$
- RD exploits a **discontinuity in $P[D_i = 1|X_i]$** at some cutoff c

Graphical Intuition



The local randomization approach

RD as a Randomized Experiment

- Close enough to the cutoff, some units were “lucky”
- Treatment **as if randomly assigned** in window $W_0 \in [c - w, c + w]$ for $w > 0$:
 ⇒ Probability distribution of X_i is uncounfounded


$$P[X_i \leq x | y_i(0), y_i(1)] = P[X_i \leq x], \quad \forall X_i \in W_0$$

- ⇒ Potential outcomes not affected by value of the score


$$Y_i(d, x) = Y_i(d)$$

- Example: $X_i \sim U[0, 1]$, $D_i = \mathbb{I}(X_i \geq c)$, then $P[D_i = 1] = 1 - c$

RD as a Randomized Experiment



(a) Randomized Experiment



(b) RD Design

Window Selection

- Under random assignment, **covariates should be balanced**

$$P[V_i \leq v | D_i = 1] = P[V_i \leq v | D_i = 0]$$

- Can use this idea as a **windows selection criterion**

⇒ Find window in which all covariates are balanced

- Heuristic procedure:

⇒ Choose a test statistic (diff. means, Kolmogorov-Smirnov,...)

⇒ Choose an **initial "small" window** $W_0^{(1)} = [c - w_{(1)}, c + w_{(1)}]$

⇒ Test $H_0 : E[X_i | c + w_{(1)}] = E[X_i | c - w_{(1)}]$

⇒ Pick a new, enlarged $W_0^{(k)}$ and **repeat for $k > 1$** until H_0 is rejected

Estimation and inference


- Once W_0 is found, proceed as in a randomized experiment

$$\hat{\tau} = \bar{Y}_1 - \bar{Y}_0$$

- Covariate-balance criterion may yield windows with few observations
- Inference based on large-sample approximations may not be reliable
- Alternative (and better!) approach: [randomization inference](#)

Example: Incumbency Advantage in U.S. Senate

- Y_i = election outcome at $t + 2$ (= 1 if party wins)
- D_i = election outcome at t (= 1 if party wins)
- X_i = margin of victory at t ($c = 0$)



The Continuity-based approach

Identification

- **Sharp design:** $D_i = \mathbb{I}(X_i \geq c)$
- **Smoothness:** $\mathbb{E}[Y_i(0)|X_i = x]$, $\mathbb{E}[Y_i(1)|X_i = x]$ continuous at $x = c$

$$\tau_{\text{SRD}} = \mathbb{E}[Y_i(1) - Y_i(0)|X_i = c]$$

- Treatment effect is nonparametrically identified **only at the cutoff**
 - $\Rightarrow$ We have zero observations at $X_i = c$, so need to approximate the regression functions
 - $\Rightarrow$ Approximation error may directly affect estimation and inference

Identification

- Naive difference in means:

$$\begin{aligned}\Delta(h) &= \mathbb{E}\{Y_i \mid X_i \in [c, c+h]\} - \mathbb{E}\{Y_i \mid X_i \in [c-h, c]\} \\ &= \mathbb{E}\{Y_i(1) \mid X_i \in [c, c+h]\} - \mathbb{E}\{Y_i(0) \mid X_i \in [c-h, c]\} \\ &= \mathbb{E}\{\tau_i \mid X_i \in [c, c+h]\} + \text{Bias}(h)\end{aligned}$$

where $\text{Bias}(h) = E\{Y_i(0) \mid X_i \in [c, c+h]\} - E\{Y_i(0) \mid X_i \in [c-h, c]\}$

- If $\mathbb{E}[Y_i(d) \mid X_i = x]$ is **continuous at $x = c$** for $d = 0, 1$, then:

$$\lim_{h \downarrow 0} \Delta(h) = \mathbb{E}[\tau_i \mid X_i = c]$$

Estimation

- Global Parametric

- ⇒ Estimate a polynomial on full sample
- ⇒ Sensitive to misspecification
- ⇒ Erratic behavior at boundary points

- Local parametric

- ⇒ Estimate a polynomial within an ad-hoc bandwidth
- ⇒ Sensitive to misspecification and bandwidth choice

- Local non-parametric

- ⇒ Data-driven bandwidth selection
- ⇒ Accounts for misspecification when performing inference

Global Parametric Approach

- Parametric assumption on conditional expectations, e.g.


$$\mathbb{E}[Y_i(d)|X_i] = \alpha_d + \beta_d(X_i - c)$$

- This implies


$$Y_i = \alpha_0 + (\alpha_1 - \alpha_0)D_i + \beta_0(X_i - c) + (\beta_1 - \beta_0)(X_i - c)D_i + u_i$$

- Easily estimated by OLS on full sample
- Coefficient $\alpha_1 - \alpha_0$ recovers the treatment effect at the cutoff


Global Parametric Approach: $p = 1$



Global Parametric Approach: $p = 2$



Global Parametric Approach: $p = 3$



Local Linear (Polynomial) Regression

- Suppose $c = 0$ (otherwise, use $X_i - c$)
- Choose **some bandwidth** $h > 0$ and estimate by OLS:


$$(\hat{\alpha}^+, \hat{\beta}^+) = \underset{(\alpha, \beta)}{\operatorname{argmin}} \sum_{i=1}^n (Y_i - \alpha - \beta X_i)^2 \mathbb{I}(0 \leq X_i \leq h)$$

$$(\hat{\alpha}^-, \hat{\beta}^-) = \underset{(\alpha, \beta)}{\operatorname{argmin}} \sum_{i=1}^n (Y_i - \alpha - \beta X_i)^2 \mathbb{I}(-h \leq X_i < 0)$$

- Estimated **treatment effect at the cutoff**

$$\hat{\tau} = \hat{\alpha}^+ - \hat{\alpha}^-$$

Local Linear Regression: Graphical Intuition



Local Linear Regression: OLS Estimands

- By standard OLS algebra (symmetrically for $\hat{\beta}^-, \hat{\alpha}^-$):

$$\hat{\beta}^+ = \frac{\sum_{i=1}^n Y_i (X_i - \bar{X}_h) \mathbb{I}(0 \leq X_i \leq h)}{\sum_{i=1}^n X_i (X_i - \bar{X}_h) \mathbb{I}(0 \leq X_i \leq h)}$$

$$\hat{\alpha}^+ = \bar{Y}_h - \bar{X}_h \hat{\beta}^+$$

- $\bar{X}_h, \bar{Y}_h$ are **local means**

$$\bar{X}_h = \frac{\sum_{i=1}^n X_i \mathbb{I}(0 \leq X_i \leq h)}{\sum_{i=1}^n \mathbb{I}(0 \leq X_i \leq h)}$$

$$\bar{Y}_h = \frac{\sum_{i=1}^n Y_i \mathbb{I}(0 \leq X_i \leq h)}{\sum_{i=1}^n \mathbb{I}(0 \leq X_i \leq h)}$$

Local Linear Regression: Bias

- If $h = 0$ the estimator would be unbiased, but with $h = 0$ there are no obs. within the bandwidth!

$$\mathbb{E}[\hat{\alpha}^+ | \mathbf{X}] = \mu_1(0) + h^2 \mathcal{B}_+ + o_p(h^2)$$

- Smaller h implies small bias but fewer observations: more variance
- $\mathcal{B}_+$ is a constant that depends on:
 - $\Rightarrow$ The curvature of $\mu_1(x)$
 - $\Rightarrow$ The kernel function (i.e. the choice of how we weight observations within the bandwidth)
 - $\Rightarrow$ The order of polynomial, p

Variance

- Similarly, it can be shown that:

$$\mathbb{V}[\hat{\alpha}^+|\mathbf{X}] = \frac{\mathcal{V}_+}{nh} + o_p(h)$$

- Decreasing the variance requires $nh \rightarrow \infty$ (effective sample size)

- $\mathcal{V}_+$ is a constant that depends on:

- $\Rightarrow \mathbb{V}[Y_i(1)|X_i = 0]$

- $\Rightarrow$ The density of the score variable at the cutoff

- $\Rightarrow$ The kernel function

- $\Rightarrow$ The order of polynomial, p

MSE and Bandwidth Selection

- Mean-squared error (MSE):

$$\begin{aligned}\text{MSE}(\hat{\tau}) &= \text{Bias}(\hat{\tau})^2 + V[\hat{\tau}] \\ &= h^4 \mathcal{B}^2 + \frac{\mathcal{V}}{nh}\end{aligned}$$

- Trade-off in bandwidth selection: bias vs variance
- MSE-optimal bandwidth:

$$\begin{aligned}h_{\text{MSE}}^* &= \underset{h}{\operatorname{argmin}} \text{MSE}(\hat{\tau}) \\ &= \left(\frac{\mathcal{V}}{4\mathcal{B}^2} \right)^{1/5} n^{-1/5}\end{aligned}$$

Inference

- Under regularity and rate restrictions on $h \rightarrow 0$

$$\sqrt{nh}(\hat{\tau} - \tau - h^2\mathcal{B}) \rightarrow_{\mathcal{D}} \mathcal{N}(0, \Omega)$$

- Where $\Omega \propto \frac{\nu_+ + \nu_-}{f}$ can be estimated using residuals $\epsilon_{+,i} = Y_i - \hat{\mu}_+(X_i)$
- Two main approaches to handle the leading bias term $\mathcal{B}$:
 - $\Rightarrow$ **Undersmoothing**: use a “smaller” bandwidth
 - $\Rightarrow$ **Bias correction**

Undersmoothing

- Choose a small enough bandwidth and proceed as if $\mathcal{B} \approx 0$

$$\sqrt{nh}(\hat{\tau} - \tau - h^2\mathcal{B}) = \sqrt{nh}(\hat{\tau} - \tau) + o_p(1) \rightarrow_{\mathcal{D}} \mathcal{N}(0, \Omega)$$

$\Rightarrow$ But recall that $h_{\text{MSE}}^* \propto n^{-1/5}$ so this approximation will have a bias

Robust Bias Correction

- Consider the following de-biasing approach:

$$\sqrt{nh}(\hat{\tau} - \tau) = \sqrt{nh}(\hat{\tau} - \mathbb{E}[\hat{\tau}|\mathbf{X}]) + \sqrt{nh}\mathcal{B}_n$$

⇒ Bias correction:

$$\sqrt{nh}(\hat{\tau} - \tau - \mathcal{B}_n) = \sqrt{nh}(\hat{\tau} - \mathbb{E}[\hat{\tau}|\mathbf{X}]) \rightarrow_{\mathcal{D}} \mathcal{N}(0, \Omega)$$

- But the **bias is unknown**, use $\mathcal{B}_p \propto \mu_+^{(p+1)} - \mu_-^{(p+1)}$ with pilot bandwidth b_n

$$\sqrt{nh}(\hat{\tau} - \tau - \hat{\mathcal{B}}_n) \rightarrow_{\mathcal{D}} \mathcal{N}(0, \Omega + \Sigma)$$

- Σ accounts for the estimation of the bias

Density discontinuity tests


- RDDs can be invalid if individuals manipulate X_i
 - Manipulation can imply sorting on one side of the cutoff
- ⇒ Test whether the density of X_i is continuous around c

Continuity in covariates / placebo outcomes


- Some variables should reveal no treatment effect:
 - ⇒ Outcomes not targeted by treatment (**placebo outcomes**)
 - ⇒ Exogenous or **predetermined covariates**
 - Estimate an RD effect on these variables
- ⇒ Finding a **non-zero effect suggests an invalid RDD**
- Existence of other (unobserved) treatments at the cutoff
 - Selection

Example: Impact of Head Start on Child Mortality

- Federal program that provides health and social services for children aged 5-9
 - ⇒ HS assistance for 300 counties based on poverty index ($X_i \geq 59.19$)
 - ⇒ Y_i = county-level mortality rates per 100,000



(a) Scatter Plot, Raw Data, $N^- = 2,455$, $N^+ = 290$



(b) RD Plot, ES, and MV, $J_- = 37$, $J_+ = 38$

Impact of Head Start on Child Mortality: Falsification

⇒ Non-parametric test for continuity of the PDF of X_i near the cutoff

	Density tests				
	h_-	h_+	N_W^-	N_W^+	p -value
Method					
Unrestricted, 2-h	10.151	9.213	351	221	0.788
Unrestricted, 1-h	9.213	9.213	316	221	0.607
Restricted (1-h)	13.544	13.544	482	255	0.655

Notes: (i) Cutoff is $\bar{r} = 59.1984$ and $W = [\bar{r} - h, \bar{r} + h]$ denotes the symmetric window around the cutoff used for each choice of bandwidth; (ii) Density test p -values are computed using Gaussian distributional approximation to bias-corrected local-linear polynomial estimator with triangular kernel and robust standard errors; (iii) column "Method" reports unrestricted inference with two distinct estimated bandwidths ("U, 2- h "), unrestricted inference with one common estimated bandwidth ("U, 1- h "), and restricted inference with one common estimated bandwidth ("R, 1- h "). See Cattaneo, Jansson, and Ma (2016a, 2016b) for methodological and implementation details.

Impact of Head Start on Child Mortality: RD Estimates

• Local parametric RD

$$\Rightarrow \hat{\tau}_{\{p=4, \text{full sample}\}} = -3.065, p\text{-value} = 0.005$$

$$\Rightarrow \hat{\tau}_{\{p=1, h=18\}} = -1.198, p\text{-value} = 0.071$$

$$\Rightarrow \hat{\tau}_{\{p=1, h=9\}} = -1.895, p\text{-value} = 0.055$$

• Local non-parametric RD

$$\Rightarrow \hat{\tau}_{\{p=0, \hat{h}_{MSE}=3.24\}} = -2.114, \text{robust } p\text{-value} = 0.037$$

$$\Rightarrow \hat{\tau}_{\{p=1, \hat{h}_{MSE}=6.81\}} = -2.409, \text{robust } p\text{-value} = 0.042$$

Extensions

Sharp Vs. Fuzzy RD

- Sharp RD: **score perfectly determines treatment** status
 - ⇒ All units scoring above the cutoff receive the treatment
 - ⇒ $D_i = \mathbb{I}(X_i \geq c)$
 - ⇒ Probability of treatment jumps from 0 to 1 at c
- Fuzzy RD: **imperfect compliance** wrt treatment assignment
 - ⇒ Some units below c may be treated or vice versa
 - ⇒ Jump in probability at c may be < 1 (but > 0)

Intention-to-treat (ITT) Parameter

- Sharp RD design on the treatment assignment variable

$$\tau_{\text{ITT}} = \mathbb{E}[Y_i(1) - Y_i(0) | X_i = c]$$

- Under usual continuity assumption

$$\tau_{\text{ITT}} = \mathbb{E}[\underbrace{(Y_i(1) - Y_i(0))}_{\tau_i} (\underbrace{D_{1i} - D_{0i}}_{\substack{= 1 \text{ for compliers} \\ = -1 \text{ for defiers} \\ = 0 \text{ for always/never takers}}}) | X_i = c]$$

⇒ ITT can be ≈ 0 even if τ is large

The Monotonicity Assumption

- Rule out the presence of **defiers**

$$P[\text{defier} | X_i = c] = 0$$

- This assumption is called **monotonicity**, since it implies that:

$$D_{1i} \geq D_{0i}, \quad \forall i$$

$\Rightarrow X_i \geq c$ does not decrease the probability of treatment

First Stage

- Effect of treatment assignment on treatment status

$$\tau_{FS} = \mathbb{E}[D_{1i} - D_{0i} | X_i = c]$$

- Under monotonicity

$$\tau_{FS} = P[D_{1i} > D_{0i} | X_i = c] = P[\text{complier} | X_i = c]$$

⇒ First stage identifies the proportion of compliers at the cutoff

Recovering the ATE on Compliers (LATE)

- Instrument D_i with $\mathbb{I}(X_i \geq c)$

$$\tau_{\text{LATE}} = \mathbb{E}[Y_i(1) - Y_i(0) | X_i = c, D_{1i} > D_{0i}] = \frac{\mathbb{E}[Y_i(1) - Y_i(0) | X_i = c]}{\mathbb{E}[D_{1i} - D_{0i} | X_i = c]}$$

⇒ Fuzzy RD parameter is “doubly local”

- At the cutoff
- On the subpopulation of compliers

Estimation in Fuzzy Designs

- ITT and FS are sharp RD estimators
- The FRD parameter can be estimated using [two-stage least squares](#)
- Can adapt all previous tools/packages to this case
 - ⇒ Data driven bandwidth selection
 - ⇒ Local polynomial estimation
 - ⇒ Robust bias-corrected inference

Multicutoff and Multiscore RD

1 Multiple cutoffs:

- ⇒ Cutoffs change across regions, time periods, etc
- ⇒ All units receive the same treatment when they exceed their cutoff

2 Cumulative cutoffs:

- ⇒ Treatment is multivalued
- ⇒ Different dosage of treatment depending on value of X_i

3 Multiple scores:

- ⇒ Treatment assigned based on multiple running variables

RD with Multiple Cutoffs

- Common empirical approach: **pooling**

⇒ Discrete cutoffs: $\mathcal{C} = \{c_0, c_1, \dots, c_J\}$

⇒ Re-centered running variable: $\tilde{X}_i = X_i - C_i$

⇒ Pooled estimand:

$$\tau^p = \lim_{x \downarrow 0} \mathbb{E}[Y_i | \tilde{X}_i = x] - \lim_{x \uparrow 0} \mathbb{E}[Y_i | \tilde{X}_i = x]$$

Identification under the Pooling Approach

- If the CEFs and $f_{X|C}(x|c)$ are continuous at the cutoffs, then

$$WATE = \sum_{c \in \mathcal{C}} \mathbb{E}[Y_i(1, c) - Y_i(0, c) | X_i = c, C_i = c] \omega(c)$$

⇒ Where the **weights reflect the densities at each cutoff**

$$\omega(c) = \frac{f_{X|C}(c)P[C_i = c]}{\sum_{c \in \mathcal{C}} f_{X|C}(c)P[C_i = c]}$$

Exploiting Multiple Cutoffs

- Two features of the pooling approach:
 - Combines TEs for different populations
 - Discards variation that can identify parameters of interest

⇒ Potential CEFs:

$$\mu_d(x, c) = \mathbb{E}[Y_i(d) | X_i = x, C_i = c], \quad d \in \{0, 1\}$$

⇒ CATE

$$\tau(x, c) = \mathbb{E}[Y_i(1) - Y_i(0) | X_i = x, C_i = c] = \mu_1(x, c) - \mu_0(x, c)$$

RD with Cumulative Cutoffs

- Multivalued treatment $D_i \in \{d_1, d_2, \dots, c_J\}$

⇒ Effect of switching to one dosage to the next one

$$\tau_j = \mathbb{E}[Y_i(d_j) - Y_i(d_{j-1}) | X = c_j]$$

- Under continuity assumption

$$\tau_j = \lim_{x \downarrow c_j} \mathbb{E}[Y_i | X_i = x] - \lim_{x \uparrow c_j} \mathbb{E}[Y_i | X_i = x]$$

RD with Multiple Scores (Boundary RD)

- Bivariate score: $\mathbf{X}_i = (X_{1i}, X_{2i})$
- Treat/control regions $\mathcal{A}_1, \mathcal{A}_0$ and **boundary** $\mathcal{B} = \text{bd}(\mathcal{A}_1) \cap \text{bd}(\mathcal{A}_0)$
- **Multidimensional RD** parameter:

$$\tau(\mathbf{b}) = \mathbb{E}[Y_i(1) - Y_i(0) | \mathbf{X}_i = \mathbf{b}], \quad \mathbf{b} \in \mathcal{B}$$

⇒ ATE at each point in the boundary $\mathcal{B}$

RD with Multiple Scores (Boundary RD)

1 Univariate distance to boundary

- ⇒ Define distance measure, e.g. $d(\mathbf{X}_i, \mathbf{b}) = \sqrt{(X_{1i} - b_1)^2 + (X_{2i} - b_2)^2}$
- ⇒ Run RD on (unidimensional) normalized running variable $\tilde{d}_i$
- ⇒ Misspecification bias when $\mathcal{B}$ is non-smooth (e.g. near a kink, next example)

2 Bivariate location relative to boundary


- ⇒ Additional (mild) regularity condition on $\mathcal{B}$ is needed
- ⇒ Estimator is well behaved at any point of $\mathcal{B}$
- ⇒ Identification, estimation, and inference are standard (rd2d package for R)

Teacher sorting and student outcomes (Bobba et al, 2024)


Teacher Compensation and Structural Inequality: Evidence from Centralized Teacher School Choice in Peru

- ① Descriptives on nation-wide allocation of public teachers in Peru
 - ⇒ Document **large urban-rural gaps** in teacher quality and student test scores
- ② RD-based evidence of teacher **wage bonuses in rural locations**
 - ⇒ Teacher competency $\uparrow$ by 0.39σ + student test scores $\uparrow$ by $0.23\text{--}0.32\sigma$
- ③ Model of teacher school choice/value added to study aggregate policy effects
 - ⇒ Framework to **design cost-effective wage policy for equity/efficiency** objectives



Inequality of Education Inputs and Output




a) Teacher Competency by Urban/Rural




b) Student Achievement by Urban/Rural




The Rural Wage Bonus Policy





Density Tests Around Extremely Rural Cutoff



a. Population (2016)



b. Population (2018)



Sharp RD Along Population Cutoff

- We rely on **pop-based assignment** rule for rural schools with $\text{dist} > 30\text{min}$

⇒ Weighted average increase in wages of 11%


- Given continuity of potential outcomes around the population cutoff

$$y_{ijt} = \gamma_0 + \gamma_1 \mathbf{1}(\text{pop}_{jt} < \text{pop}_c) + g(\text{pop}_{jt}, \text{pop}_c) + \delta_t + u_{ijt}$$


- $g(\cdot)$: flexible polynomial on population of the locality of school j
- δ_t : indicator for year of assignment
- u_{ijt} : error term, clustered at the school-year level

⇒ Estimate γ_1 **non-parametrically within MSE-optimal** bandwidths

Rural Bonus and Teacher Choices over Job Postings



a) Vacancy Filled



b) Competency Score

(1)

(2)

(3)

Rural Bonus and Student Achievement


Panel A: Dependent Variable is Spanish Test (z-score)

	(1) Vacancy	(2) No vacancy	(3) All
High Bonus	0.395 (0.152)	-0.004 (0.127)	0.232 (0.088)
Bandwidth	107.818	148.920	105.822
Schools	264	451	832
Observations	4635	6773	16681


Panel B: Dependent Variable is Math Test (z-score)

	(1) Vacancy	(2) No vacancy	(3) All
High Bonus	0.579 (0.193)	0.067 (0.143)	0.317 (0.105)
Bandwidth	85.848	155.174	95.638
Schools	220	470	764
Observations	3939	7039	15363


RD Robustness



a. Vacancy filled



b. Teacher scores (std)



What is the Rationale for the Model?

- The RD evidence is limited to the local effect of the rural wage bonus
- ⇒ What is the **effect of the policy** on urban-rural gaps in education outcomes?
- ⇒ Can we characterize **more effective teacher-school allocations**?
- ⇒ Can we achieve those with **alternative wage schedules**?

An Empirical Matching Model of Teachers and Schools

- ① A discrete choice framework recovers **teachers' preferences**
 - ⇒ Flexibly parametrize pref. distributions over wages and non-wage amenities
- ② A matching equilibrium that maps preferences into **assignment outcomes**
- ③ A value-added model that maps teacher sorting into **student achievement**
 - ⇒ Teacher VA varies with students' prior achievement and demographics
 - ⇒ Teacher VA parameters correlate with teachers' preferences

Teacher Preferences

- Teacher i utility from school j (off-plat. $j = J + 1$ + outside option $j = 0$):

$$\begin{aligned} u_{ijt} &= \alpha_i w_{jt} + x'_i \Theta_1 z_{jt} + x'_i \Theta_2 d_{ijt} + \epsilon_{ijt}, \\ u_{iJ+1t} &= \gamma_{0t} + x'_i \gamma_1, \\ u_{i0t} &= \beta_{0t} + \beta_i. \end{aligned}$$

- Serial dictatorship $\Rightarrow$ discrete choice model with observed choice sets

$$\mu_w^*(i, t) = \arg \max_{j \in \Omega(s_{it})} U_{ijt}$$

Teacher Value Added

- Student l **potential outcome** when matched with teacher i :

$$Y_{lij} = + \underbrace{c_{jt}\beta}_{\text{school/classroom effect}} + \underbrace{z_{lt}\bar{\delta}}_{\text{student ability}} + \underbrace{z_{lt}(\delta_i - \bar{\delta})}_{\text{teacher ATE + match effects}} + \nu_{lij}$$

- We allow **value-added to correlate with preferences** $\theta_i = (\log \alpha_i, \beta_i)$

$$(\theta_i, \delta_i) | x_{it} \sim \mathcal{N} \left[\begin{pmatrix} x'_{1it} \gamma^\theta \\ x'_{2it} \gamma^\delta \end{pmatrix}, \begin{pmatrix} \Sigma_{\theta, \theta} & \Sigma_{\theta, \delta} \\ \Sigma_{\delta, \theta} & \Sigma_{\delta, \delta} \end{pmatrix} \right]$$

$\Rightarrow \Sigma_{\theta, \theta}$: how teachers value **non-wage benefits/outside option** against wages

$\Rightarrow \Sigma_{\delta, \theta}$: **selection on teacher quality** resulting from changes in wages


Identification

- We identify the achievement prod. function using **teacher-classroom data**
 - Estimate teacher effectiveness as fixed effects δ_i
 - Use variation in observables x_{2it} to recover $(\gamma^\delta, \Sigma_{\delta,\delta})$
- We identify choice parameters using **panel data on choices/choice sets**
 - Repeated choice data help identify the distribution of random coefficients θ_i
 - **Wages vary only with observables** $\Rightarrow$ residual variation is RD effect
- We identify $\Sigma_{\theta,\delta}$ by linking assignments with **teacher-classroom data**
 - Conditional on knowing $\Sigma_{\delta,\delta}$ we can recover $\Sigma_{\theta,\delta}$

Monthly Willingness to Pay for Non-Wage Characteristics

	Mean		10% Quantile		90% Quantile	
	Soles (1)	% Wage (2)	Soles (3)	% Wage (4)	Soles (5)	% Wage (6)
<i>Amenities, Infrastructure and Remoteness</i>						
Amenity/Infrastructures	200	10	30	2	440	22
Closer to Home by 1km						
$0 \leq \text{Distance} < 20$	200	10	33	2	443	22
$20 \leq \text{Distance} < 100$	113	6	23	1	243	12
$\text{Distance} \geq 100$	20	1	3	0	43	2
<i>Ethnolinguistic Proximity</i>						
Same Language: Spanish	2,777	139	393	20	6,180	309
Same Language: Quechua	986	49	303	15	1,929	96
Same Language: Aymara	3,264	163	656	33	6,976	349
<i>Teaching Conditions</i>						
No Border	406	20	-97	-5	1,122	56
No Multigrade	962	48	147	7	2,121	106
No Single Teacher	1,758	88	100	6	4,102	206

Rural vs. Urban Non-Pecuniary Utility Differences




Standard Deviation of TVA Coefficients


	Math (1)	Spanish (2)
ATE	0.465 (0.006)	0.408 (0.006)
Lagged Score	0.145 (0.005)	0.150 (0.005)
Lagged Score ²	0.049 (0.004)	0.061 (0.003)
Female	0.098 (0.010)	0.083 (0.013)
Quechua - Aymara	0.040 (0.030)	0.067 (0.019)
Age	0.115 (0.007)	0.110 (0.008)

- ⇒ One SD increase in TVA ⇒ ↑ in test scores by 0.44-0.50 SD
- ⇒ Significant **match effects** on lagged measures of student achievement
- ⇒ 12-18% of variance in TVA explained by **teachers' comparative advantage**

Model Fit



a. Sorting



b. Value Added

Counterfactual 1: Aggregate Effects of the Rural Bonus

- Predict teachers' choices over schools with and without rural bonus
 - Simulate U_{ijt} from estimated parameters and a random draw of ϵ_{ijt} and θ_i
- Compute the stable matching eq. using the teacher-proposing DA algorithm
- Predict the distribution of teacher value-added without and with rural bonus
 - Use the mean of the posterior distribution of δ_i (see Appendix D.3)

Counterfactual 1: Aggregate Effects of the Rural Bonus

	Status Quo (1)	No Rural Bonus (2)	Policy Effect (3)
<i>Panel A: Total Value Added</i>			
Urban-Rural Gap	0.077	0.164	-0.087
Urban	0.024	0.059	-0.036
Rural	-0.053	-0.105	0.052
<i>Moderately Rural</i>	-0.033	-0.055	0.022
<i>Rural</i>	-0.111	-0.049	-0.063
<i>Extremely Rural</i>	0.067	-0.099	0.166
<i>Panel B: Match Effects</i>			
Urban	-0.007	0.002	-0.009
Rural	0.008	0.001	0.007

⇒ Small average effects on TVA, mostly concentrated in very remote schools

⇒ Rural bonus does not induce sorting based on comparative advantages


Counterfactual 2: Gains from Teachers' Reallocation

- We consider the following the linear program:

$$\max_{\mu} \sum_{i=1}^{n_w} \sum_{j=1}^{n_m} \pi_j \bar{z}'_j \hat{\delta}_i \mathbb{I}\{\mu(i) = j\}$$

- $\bar{z}'_j \hat{\delta}_i$ is the **predicted (shrunk) average TVA** for teacher i in school j
- Solution $\mu^*(\pi_j)$ depends on **weight put on students in school j**

Counterfactual 2: Gains from Teachers Reallocation



Counterfactual 3: Optimal Wage Policy

- Policymaker can set priorities and wages in each school such that:

$$\min_w \sum_j w_j, \text{ s.t. } \begin{cases} \max_{i \in \mu(j)} z'_{lt} \delta_i \geq c_j, \forall j \in \mathcal{S} & \text{(C1)} \\ \mu \text{ is stable given } w \text{ and using } z'_{lt} \delta_i \text{ as priorities} & \text{(C2)} \end{cases}$$

- For a fixed wage, schools strictly rank teachers according to $z'_{lt} \delta_i$
- Otherwise, the allocation with the lower wage is always strictly preferred
- A **stable set of contracts** always exists in this counterfactual economy
 - Each school $j \in \mathcal{S}$ bids upward until (C1) is satisfied
 - Outcome is (μ, w) that satisfies (C1)-(C2) while **minimizing total wage bill**

Counterfactual 3: Optimal Wage Policy

	Status Quo (1)	Teacher Value Added Threshold				
		$c = -0.4$ (2)	$c = -0.3$ (3)	$c = -0.2$ (4)	$c = -0.1$ (5)	$c = 0$ (6)
<i>Panel A: Teacher Value Added</i>						
Urban	0.055	0.036	0.035	0.019	-0.009	-0.058
Rural	-0.048	0.015	0.076	0.133	0.197	0.258
<i>Moderately Rural</i>	0.025	0.007	0.058	0.040	0.127	0.203
<i>Rural</i>	-0.154	-0.060	0.034	0.094	0.117	0.199
<i>Extremely Rural</i>	-0.022	0.080	0.131	0.225	0.296	0.357
<i>Panel B: Match Effects</i>						
Urban	0.019	0.017	0.018	0.018	0.013	0.022
Rural	0.040	0.063	0.111	0.137	0.180	0.191
<i>Moderately Rural</i>	0.008	0.002	0.031	0.022	0.065	0.089
<i>Rural</i>	0.039	0.085	0.141	0.107	0.154	0.161
<i>Extremely Rural</i>	0.070	0.106	0.168	0.218	0.247	0.300
<i>Panel C: Monthly Total Cost (in Soles)</i>						
% Base Wage	0.111	0.086	0.140	0.234	0.379	0.621
Mean Bonus per School	223	171	279	467	759	1,242
SD Bonus per School	220	407	576	839	1,184	1,698

RDD $\Leftrightarrow$ Economic Model

- RD evidence is local in nature and may neglect indirect/equilibrium effects
 - The model is a tool to extrapolate the policy effect away from the RD cutoff
 - $\Rightarrow$ Quantify aggregate effect of the policy reform
 - $\Rightarrow$ Characterize gains from alternative teacher assignments
 - $\Rightarrow$ Design optimal teacher compensation schemes
- $\Rightarrow$ RD variation is embedded in the model, enhancing its credibility