

On the Origin of Stochastic, Low-Frequency Photometric Variability in Massive Stars

MATTEO CANTIello,^{1,2} DANIEL LECOANET,^{3,4,2} ADAM S. JERMYN,¹ AND LUCA GRASSITELLI⁵

¹*Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA*

²*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

³*Department of Engineering Sciences & Applied Mathematics, Northwestern University, Evanston, IL 60201, USA*

⁴*CIERA, Northwestern University, Evanston, IL 60201, USA*

⁵*Argelander-Institut für Astronomie der Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany*

Submitted to ApJ

ABSTRACT

High-precision photometric observations have revealed ubiquitous stochastic low-frequency photometric variability in early type stars. It has been suggested that this variability arises due to either subsurface convection or internal gravity waves launched by the convective core. Here we show that relevant properties of convection in subsurface convective layers correlate very well with the timescale and amplitude of stochastic low-frequency photometric variability, as well as with the amplitude of macroturbulence. We suggest that low-frequency, stochastic photometric variability and surface turbulence in massive stars are caused by the the presence of subsurface convection. We show that an explanation for the observed surface photometric variability and macroturbulence relying on convective core driven internal gravity waves encounters a number of difficulties and seems unlikely to be able to explain the observed trends.

Keywords: convection, stars: magnetic field, stars: variability

1. INTRODUCTION

Massive stars largely drive the dynamical and chemical evolution of gas in galaxies (e.g. Hopkins et al. 2014). They accomplish this via their stellar winds, eruptions, and explosive deaths, ultimately producing neutron stars and black holes (Langer 2012). These compact remnants can merge and generate the gravitational waves observed by LIGO/Virgo. The evolutionary trajectory starting with a massive star burning hydrogen in its core and ending with a compact remnant is understood only qualitatively. We still do not know how to map initial properties of the star, like mass, rotation rate, and metallicity, to e.g. the final mass and spin of the compact remnant it leaves behind. The picture is further complicated by the fact that the majority of massive stars are found in multiple systems (Sana et al. 2012), with a large fraction expected to interact their companions (de Mink et al. 2014).

The path towards a detailed understanding of massive stars begins with a quantitative study of their internal structure on the main sequence. In recent years asteroseismology has opened a new window on these challenging astrophysical environments, with high precision photometry from space delivering many new exciting results (e.g., MOST, CoRoT, BRITE, Kepler/K2 and TESS, see Bowman 2020). The latest discovery is the detection of a new ubiquitous phenomenon in massive stars: stochastic low-frequency photometric variability (SLF variability; Bowman et al. 2019a,b; Pedersen et al. 2019; Bowman et al. 2020). This joins a number of other surface and wind phenomena that are routinely observed in early-type stars, including surface velocity fluctuations (Macroturbulence; Simón-Díaz & Herrero 2014), line profile variability (Fullerton et al. 1996), and discrete absorption components in UV spectra (Howarth & Prinja 1989; Cranmer & Owocki 1996; Fullerton et al. 1997; Kaper et al. 1997). Surface magnetism and bright spots are harder to observe but could still be common in these stars (e.g. Ramaramananantsoa et al. 2014).

The origin of this SLF variability is currently debated. It could be caused by sub-surface convec-

tion zones (Cantiello et al. 2009; Blomme et al. 2011; Lecoanet et al. 2019) or by internal gravity waves (IGWs) launched by turbulent core convection (Edelmann et al. 2019; Ratnasingam et al. 2020)¹. Regardless of its origin, this photometric signal likely carries important information about stellar structure, complementing asteroseismic studies that use well-identified oscillation modes (e.g. Aerts 2019; Aerts et al. 2019; Bowman 2020).

Recently the use of high resolution ground-based spectroscopy for the targets observed by K2 and TESS (Bowman et al. 2020) has allowed precise determination of stellar parameters, including mass, luminosity and macroturbulence. The latter is particularly important if, as seems likely, the mechanism exciting surface turbulent velocities is the same as that which produces the observed SLF variability (Grassitelli et al. 2016; Bowman et al. 2020).

Here we combine spectroscopic and photometric data to compare the observed properties of the stochastic photometric variability and macroturbulence with predictions from 1D stellar models. We make simple predictions for the amplitude and frequency of the variability that subsurface convection induces at the stellar surface and examine how these vary with stellar temperature and luminosity. We find that the predicted trends coming from a subsurface convection zone driven by the iron opacity peak (FeCZ) at $\approx 150kK$ match the observations well.

We next show that one way to differentiate between the two proposed mechanisms is by examining macroturbulence and SLF in massive stars with surface magnetic fields, since magnetic effects have a larger impact on core IGWs than on the FeCZ. Macroturbulence is observed in stars with fairly strong magnetic fields, sufficient to suppress IGWs from the core, favoring a model based on subsurface convection. Furthermore, the only stars with no observed macroturbulence are ones where the magnetic field is strong enough to shut off the FeCZ, and so far as we know all stars with such strong magnetic fields lack macroturbulence, consistent with a subsurface origin of surface perturbations (Jermyn & Cantiello 2020).

Based on these considerations we suggest that subsurface convection represents a possible unifying mechanism causing SLF variability, surface turbulence, and magnetic spots in massive stars.

¹ Classical heat-mechanism pulsations could be responsible for spectroscopic and photometric variability in specific parts of the HRD, but they can hardly justify the apparent ubiquity of macroturbulence and SLF in massive stars (Godart et al. 2017; Simón-Díaz et al. 2017a).




Figure 1. Evolution of the normalized radial location of the FeCZ in a $20M_{\odot}$ model during the main sequence (first 4.7 Myr of evolution). The HeCZ is also visible very close to the surface. Locations of $P(r)/P(R_*) = e^n$ for $n = [1, 3, 5, 7]$ are also shown and labelled as nH_P .

2. METHODS

We calculated stellar evolution models using the Modules for Experiments in Stellar Astrophysics (MESA Paxton et al. 2011, 2013, 2015, 2018, 2019) software instrument. Details on the microphysics inputs to this software instrument are given in Appendix A. We focused on the structure and convective properties of our models in order to calculate typical frequencies and convective fluxes. Convection is calculated in the framework of the Mixing Length Theory (Böhm-Vitense 1958, MLT hereafter), and we adopted $\alpha_{MLT} = 1.6$. While the properties of efficient convection zones (e.g. stellar cores) are insensitive to the choice of this parameter, those of inefficient convective regions close to the stellar surface *do* depend on α_{MLT} (Jiang et al. 2015; Cantiello & Braithwaite 2019). In Section 5 we discuss how uncertainties in the choice of the α_{MLT} parameter affect our results.

Since we are interested in the excitation of surface phenomena we focus on the upper part of convective zones. Following Cantiello et al. (2009) we define the **average of a generic quantity q** as

$$\bar{q} \equiv \frac{1}{\alpha_{MLT}H_P} \int_{R_c - \alpha_{MLT}H_P}^{R_c} q(r) dr, \quad (1)$$

where H_P is the pressure scale height calculated at the upper boundary (R_c) of the convective zone of interest. We tested a variety of different average prescriptions and found that our results do not depend much on the specific choice of prescription.

Using equation (1) we extracted the average Mach number $\bar{M}_c$, convective velocity $\bar{v}_c$ and density $\bar{\rho}_c$ in the convective core and in the subsurface convection zones of our models. The most important subsurface convection zone for the massive stars we focus on is the iron convection zone (FeCZ, see Fig. 1), although in the low luminosity regime He-driven convection zones could play a role as well (Cantiello et al. 2009; Cantiello & Braithwaite 2019).

We compare relevant properties of our theoretical models with the observed characteristic frequency ν_{char} and amplitude α_0 of SLF variability. These quantities are derived by fitting the stochastically-excited, low-frequency power excess in a power density spectrum using a Lorentzian function (e.g. Bowman et al. 2019b):

$$\alpha(\nu) = \frac{\alpha_0}{1 + \left(\frac{\nu}{\nu_{\text{char}}}\right)^\gamma} + P_W. \quad (2)$$

This shows that α_0 represents the amplitude at zero frequency. ν_{char} is defined as in eqn. 3 with τ the typical timescale of the SLF variability. Finally, γ is the gradient of the linear part of the profile and P_W is a white noise term. **Due to its stochastic, low-frequency manifestation in the power density spectrum, in the literature the SLF variability is also referred to as “red noise” (e.g. Blomme et al. 2011).**

3. RESULTS

We want to test a possible correlation between the properties of the FeCZ and observed photospheric phenomena, in particular SLF variability and turbulent velocities at the stellar surface (macroturbulence). We proceed by calculating quantities that measure the amplitude of perturbations in the FeCZ. We then check if some of these properties correlate with the amplitude of observed photospheric phenomena, including turbulent velocity fluctuations and SLF variability.

We define the characteristic frequency as

$$\nu_{\text{char}} \equiv \frac{1}{2\pi\tau}, \quad (3)$$

where τ is a characteristic timescale. For comparing with observations we set τ to be the average convective turnover time

$$\tau_c = \alpha_{\text{MLT}} \bar{H}_P / \bar{v}_c, \quad (4)$$

calculated either in the FeCZ or in the convective core.

We calculated the convective flux $F_c = \bar{\rho}_c \bar{v}_c^3$, where $\bar{\rho}_c$ and $\bar{v}_c$ are the average density and convective velocity calculated according to eqn. 1. We did this as a function of both mass and evolutionary history for stars ranging

from $5 - 120 M_\odot$. Here we present results for an initial metallicity of $Z=0.02$, but in Appendix C we report results for model grids with $Z = 0.006$ and 0.002 as well.

3.1. FeCZ and Macroturbulence

We show the average convective velocity in the FeCZ in Fig. 2, left panel. Velocities of the order $10 \dots 100 \text{ km s}^{-1}$ are found across the upper spectroscopic H-R diagram ($\mathcal{L} \equiv T_{\text{eff}}^4/g$; Langer & Kudritzki 2014), with a trend of increasing v_c for higher luminosities. Note that, contrary to core convection, the FeCZ is just mildly subsonic, with Mach numbers ranging from 0.01 to 0.3, though these are uncertain by a factor of ≈ 8 due to a dependence on the uncertain α_{MLT} (Cantiello & Braithwaite 2019).

In the right panel of the same figure we also show the maximum of the ratio of turbulent pressure to total pressure ($P_{\text{turb}} \propto v_c^2$), which, in agreement with the results of Grassitelli et al. (2015b), shows a strong correlation with the spectroscopically-derived macroturbulent velocities in massive stars.

The two quantities in Fig. 2 measure the strength of the inefficient convection, and are related to different possible mechanisms for how motions in the FeCZ could lead to surface velocities. If one assumes convective elements conserve their inertia as they reach layers stable against convection (e.g., because they are thermally diffusive, see Cantiello et al. 2009), then the surface velocities should be proportional to the convective velocity; see the left panel of Fig. 2. Alternatively, one can assume IGWs are excited with pressure perturbations $\delta p \sim P_{\text{turb}} \propto v_c^2$, the turbulent pressure of the convection (Press 1981). Using the polarization relations of adiabatic IGWs (?), a wave with pressure perturbation δp has an associated horizontal velocity $u_h \sim (\delta p / \rho_0) k_h / \omega$, where k_h is the horizontal wavenumber of the wave and ω is its frequency. The dominant waves will have $k_h \sim 1/H_P$ and $\omega \sim 1/\tau_c$. So IGWs would have surface velocities which also scale like $u_h \sim v_c$ at the surface. In the right panel of Fig. 2 we also report the ratio of turbulent pressure to total pressure in the FeCZ. According to Grassitelli et al. (2015b), this quantity traces the stochastic Lagrangian pressure perturbation associated with the convective motions, which is responsible for local deviation from hydrostatic equilibrium and the excitation of high-order pulsations with frequencies close to the spectrum of the fluctuations.

We confirm that convective velocities and the ratio of turbulent pressure to total pressure in the FeCZ correlate very well with the amplitude of macroturbulence.




Figure 2. Left: Average convective velocities in the FeCZ as a function of the location of stellar models in the spectroscopic H-R Diagram ($\mathcal{L} \equiv T_{\text{eff}}^4/g$). We also show observed stars with detected macroturbulent velocity as grey dots. The area of the dot is proportional to the observed macroturbulence. **The FeCZ is absent in models with $\log \mathcal{L}/\mathcal{L}_{\odot} \lesssim 2.5$.** Right: Same as left, but showing predictions for the maximum of the ratio between turbulent pressure and total pressure in any subsurface convection zone. The FeCZ largely dominates, except for stars at low luminosities where turbulent pressure is provided by a helium convection zone (HeCZ).

3.2. FeCZ and Stochastic, Low-Frequency Variability

If the SLF variability is caused by the FeCZ, a natural choice of proxy for the typical timescale of this variability is the convective turnover timescale. We find typical values of the convective turnover timescale to be about $\sim 0.1 \dots 2$ d in the FeCZ, with a tendency for shorter values in models with high effective temperature and surface luminosity (Cantiello et al. 2009). We computed ν_{char} using eqn. 3 and plot this alongside observed characteristic frequencies² of SLF variability on the spectroscopic H-R diagram, see left panel in Fig. 3.

We see good agreement, with our models reproducing both the typical values of the observed ν_{char} and the trend with $\log \mathcal{L}$ and $\log T_{\text{eff}}$. While the characteristic frequencies found by Bowman et al. (2020) seem to be larger by a factor of ≈ 3 , our predictions for the turnover timescale are affected by uncertainty in the convective velocities arising from the MLT treatment ($v_c \propto \alpha_{\text{MLT}}^3$, so $\nu_{\text{char}} \propto \alpha_{\text{MLT}}^2$), **as well as our limited knowledge of the frequency spectrum generated by turbulent convection.** Since α_{MLT} is uncertain by a factor of 2 or so, our estimates of the characteristic frequency

in the FeCZ are uncertain by a factor of ≈ 4 and so are consistent with the observations.

We also compare the amplitude of SLF variability with the relative convective flux in the FeCZ (right panel in Fig. 3). Values of F_c/F_* tend to increase with increasing $\log \mathcal{L}$ and decreasing $\log T_{\text{eff}}$, a trend that is also found for the values of observed SLF variability.

Overall the turnover timescale and relative flux in the FeCZ correlate very well with the timescale and amplitude of the observed SLF variability.

3.3. Core Convection and Stochastic, Low-Frequency Variability

Early-type stars possess convective cores during their main sequence. Turbulent convection in these convective cores excites internal gravity waves (Goldreich & Kumar 1990; Rogers et al. 2013; Lecoanet & Quataert 2013; Shiode et al. 2013; Edelmann et al. 2019; Horst et al. 2020), that can propagate through the stellar envelope and reach the stellar surface (e.g. Ratnasingam et al. 2019; Lecoanet et al. 2019; Ratnasingam et al. 2020). **Some groups have argued that such waves could be responsible for both the observed macroturbulence and SLF variability in early-type stars** (Aerts et al. 2009; Simón-Díaz et al. 2017b; Bowman et al. 2019b,a).

² These were inferred via eqn. (2).




Figure 3. Left panel: Characteristic frequency ν_{char} in the FeCZ as a function of the location of stellar models in the spectroscopic H-R Diagram (black contour lines). Evolutionary tracks are shown as grey solid lines. We also show the observed stars with detected stochastic photometric variability as grey dots. The area of the dot is proportional to the observed ν_{char} from eqn. (2). Right panel: Ratio of FeCZ convective flux to the total stellar flux in the spectroscopic H-R Diagram ($\mathcal{L} \equiv T_{\text{eff}}^4/g$). We also show the observed stars with detected stochastic, low-frequency photometric variability as grey dots. The area of the dot is proportional to α_0 .

Shiode et al. (2013) estimated that IGWs propagating from the core should lead to surface brightness fluctuations with typical amplitudes $\approx 10^{-2} - 10^2 \mu\text{mag}$, at odds with the relatively large amplitudes $\approx 10 - 10^4 \mu\text{mag}$ observed by e.g. Bowman et al. (2020). Lecoanet et al. (2019) also argue that the low-frequency, stochastic variability is unlikely to be caused by linearly propagating waves from the core. Their predicted wave transfer function shows little power at very low frequencies due to radiative damping, as well as standing modes at frequencies above 0.5 d^{-1} . None of these features are present in the observed spectra.

Recent numerical simulations of wave generation by convection in a $3M_{\odot}$ star (Edelmann et al. 2019; Horst et al. 2020) produce wave fluctuation spectra which are qualitatively similar to those observed. However, those simulations artificially boost the stellar luminosity by factors ranging from 10^3 to 10^7 . This makes it difficult to compare the amplitudes of the waves in those simulations to, e.g., Shiode et al. (2013), which predicts the surface variability in stars with realistic luminosities.

Although the detailed physics of wave generation by convection is uncertain, we can still analyze the properties of core convective of our models. If the surface variability is due to core convection, one would expect

the characteristic frequency and amplitude of the SLF variability to correlate with the properties of the core convection. The flux of IGWs excited by turbulent convection at the core boundary is of order $F_{\text{IGW}} = F_* \bar{M}_c$ (Goldreich & Kumar 1990), where we evaluate the average Mach number $\bar{M}_c$ using eqn. 1 and the local adiabatic sound speed. We expect the luminosity fluctuations to be proportional to the relative surface radial displacement ξ_r/R produced by these waves at the stellar surface (e.g. Dziembowski 1977; Aerts et al. 2010). It can be shown that $\xi_r/R \propto \sqrt{F_{\text{IGW}}/F_0}$ (see Appendix B). This quantity is presented in Fig. 4 along with the observed amplitudes of SFL variability.

One important caveat is that this estimate neglects the important role of radiative damping, which is essential in shaping both the amplitude and the shape of the spectrum of waves at the surface (e.g. Lecoanet et al. 2019). Despite neglecting the important effect of radiative damping, the trend for the relative radial displacement at the surface do show some interesting correlations with the observed trend in SLF variability, though they generally proceed the wrong way, with increasing amplitude towards higher luminosities where we expect IGW to show the smallest effects (left panel in Fig. 4).




Figure 4. Left panel: Characteristic frequency ν_{char} in the convective core as function of the location of stellar models in the spectroscopic H-R diagram. We also show the observed stars with detected photometric variability as grey dots. The area of the dot is proportional to ν_{char} . Right panel: Square root of the ratio between the flux of gravity waves launched by the convective core and $F_0 = 1/2\rho(r)r^3\omega^2\sqrt{N(r)^2 - \omega^2}$, evaluated at the stellar surface. In the absence of damping, this quantity is expected to be proportional to the relative radial displacement (Appendix B) and hence to the relative luminosity fluctuations at the surface. The IGW flux is calculated multiplying the core convective flux by the average convective Mach number in the top pressure scale height of the convective region. We also show the observed stars with detected stochastic, low-frequency photometric variability as grey dots. The area of the dot is proportional to α_0 .

Next, we focus on the characteristic timescale of waves excited by core convection. We expect the maximum of the IGW flux to be launched at frequencies close to $\nu_{\text{char}} = (2\pi\tau_c)^{-1}$, with τ_c defined in eqn. 4. Fig. 4 shows that the predicted values of ν_{char} are in the range $0.02 \dots 0.008 \text{ d}^{-1}$ ($0.2 \dots 0.08 \mu\text{Hz}$), in agreement with the results of Shiode et al. (2013). These values are about 2 orders of magnitude smaller than the typical characteristic frequencies of SLF variability observed by (Bowman et al. 2019b,a).

As mentioned earlier, a significant caveat in correlating core and surface quantities is that wave propagation through the stellar envelope affects the spectrum, changing the frequency of maximum power of waves reaching the surface (Lecoanet et al. 2019). Radiative damping efficiently suppresses low-frequency IGWs (because these waves have high radial wavenumbers), and the peak of the spectrum observed at the surface is expected to move to higher frequencies. The amplitude of this effect depends on the envelope properties, which are a function of $\log T_{\text{eff}}$ and $\log L$. Nevertheless we will assume that trends of ν_{char} , as function of $\log T_{\text{eff}}$ and $\log L$, are still set by the core convective properties.

Under this assumption, Fig. 4 shows that ν_{char} should increase with both $\log T_{\text{eff}}$ and $\log L$, so that the characteristic frequencies of peak IGWs should increase as stars evolve on the main sequence. This is exactly the opposite of what is observed: the characteristic frequencies of SLF variability is largest for stars early on on the main sequence, and seem to decrease as stars evolve. Therefore either radiative damping is able to revert this trend or else the observed variability is unlikely to be caused by IGWs launched by the core.

4. MACROTURBULENCE IN MAGNETIC STARS

The hypothesis that subsurface convection is responsible for surface turbulence is corroborated by the match between trends in the FeCZ properties (i.e. convective velocities and turbulent pressure) and the observed micro and macroturbulence (Cantiello et al. 2009; Grasielli et al. 2015b). One important test to this hypothesis is provided by the disappearance of macroturbulence in stars with surface magnetic fields above a critical strength (Sundqvist et al. 2013), closely corresponding to the critical field needed to stabilize the FeCZ (MacDonald & Petit 2019; Jermyn & Cantiello 2020). Note that in OB stars, the FeCZ is deeper and more vigorous

than the H and He convection zones, so a magnetic field stabilizing the FeCZ will necessarily also stabilize the other subsurface convection zones.

We can analogously define a critical magnetic field strength B_{crit} which suffices to reflect IGWs before they reach the photosphere. Using the dispersion relation for IGWs in a magnetic medium, the radial component of this field is (Fuller et al. 2015)

$$B_{r,\text{crit}} = \frac{\omega}{2k_r} \sqrt{4\pi\rho}, \quad (5)$$

where $\omega = \nu/2\pi$ is the angular frequency of the waves, ρ is the density, and k_r is the radial wave-number. For IGWs the radial wave-number is related to the spherical harmonic degree ℓ by

$$k_r \approx \frac{\sqrt{\ell(\ell+1)}}{r} \left(\frac{N}{\omega} \right), \quad (6)$$

where N is the Brünt-Väisälä frequency. So

$$B_{r,\text{crit}} \approx \frac{\omega^2 r}{N} \sqrt{\frac{4\pi\rho}{2\ell(\ell+1)}} \quad (7)$$

$$\approx \frac{\omega^2 r}{\ell N} \sqrt{4\pi\rho}. \quad (8)$$

Note that because this decreases with increasing ℓ , all waves of a given frequency are reflected if the $\ell = 1$ waves are reflected.

We computed $B_{r,\text{crit}}$ for several values of ℓ and ν for a main-sequence model of a $30M_\odot$ star as a function of radius, shown in Fig. 5. For frequencies comparable to those of core convection the critical magnetic field strength is of order 10^{-2} G to 10^{-1} G. Strong macroturbulence is observed in similar-mass O-type stars with magnetic fields up to 2.5 kG (e.g. HD 191612, Sundqvist et al. 2013), so macroturbulence in those stars is unlikely to be due to IGWs coming from their cores if those waves have similar frequencies to that of core convection.

For frequencies comparable to the observed ν_{char} , on the order of 3 d^{-1} , the critical magnetic field is much larger, on the order of 300 G to 1 kG. This makes an explanation of macroturbulence based on IGWs marginally inconsistent with observations showing strong macroturbulence up to field strengths of 2.5 kG. However, because equation (7) is a strong function of ω , and hence of ν , it is possible that these strongly magnetized stars just have larger ν_{char} . For instance if HD 191612 has $\nu_{\text{char}} \approx 5 \text{ d}^{-1}$ then it would have $B_{r,\text{crit}} \approx 3 \text{ kG}$ for the $\ell = 1$ mode.

A further prediction of this calculation is that, if IGWs are the cause of macroturbulence in these stars, we should expect the strength of macroturbulence to decline with increasing magnetic field strength as more

and more modes are reflected before they reach the surface. Sundqvist et al. (2013) find no such trend, and the only strong correlation they see of macroturbulence with magnetic field strength is that it vanishes when the magnetic field exceeds the FeCZ shutoff strength. This again points against an explanation of macroturbulence based on core-generated IGWs.




Figure 5. The critical radial magnetic field strength needed to reflect internal gravity waves of different ℓ and ν is shown as a function of fractional depth for a $30M_\odot$ stellar model at an age of 4 Myr. The frequency $\nu_{\text{char,MLT}}$ is that given by equation (3) but evaluated for the core convection zone instead of the subsurface FeCZ. The frequency $\nu_{\text{char,Obs}}$ is a typical value of 3 d^{-1} for the observations.

5. DISCUSSION

The presence of a subsurface convection zone can result in variability of photospheric properties via a number of processes, including wave excitation (Cantiello et al. 2009; Grassitelli et al. 2015b) and magnetic buoyancy (Cantiello & Braithwaite 2011). Multi dimensional simulations including radiation have been performed in a restricted range of the parameter space, and show that the full turbulent manifestation of these convective regions extends up to the stellar surface (Jiang et al. 2015, 2017, 2018). In their calculations of OB stars envelopes including the stellar photosphere, Jiang et al. (2015) observe turbulent velocities reaching the isothermal sound speed ($\approx 50 \text{ km s}^{-1}$) at the stellar surface, demonstrating that velocity fields of amplitude comparable to the observed macroturbulence are naturally explained by the presence of the FeCZ.

Therefore it could be that the observed SLF variability and macroturbulence simply represent the direct man-

ifestation of turbulent, radiation-dominated convection at the stellar surface. The simplified MLT treatment in our one dimensional calculations is unable to capture the complex phenomenology of these layers, but the fact that it can reproduce both the timescales and the trends in amplitude of the observed SLF variability is compelling. It calls for extending the radiation hydrodynamics simulations to cover the parameter space of the TESS observations, in order to unravel the precise mechanism connecting subsurface convection zones to the observed surface variability.

If the FeCZ is responsible for surface macro-turbulence and SLF variability, then these phenomena should be affected by the stellar metallicity. On the other hand we do not expect a significant metallicity dependence in the context of a core-convection origin. The results of (Bowman et al. 2019b) show that SLF variability is also observed in low-metallicity, LMC stars. The presence of the FeCZ depends on the luminosity and metallicity of the star, and in 1D stellar evolution calculations the FeCZ occurs above a luminosity of $L \approx 10^{3.2} L_{\odot}$ for $Z=0.02$. This corresponds to a zero age main sequence star of about $7M_{\odot}$ (see e.g. Fig. 2). **The LMC has a metallicity about half solar, and in models at $Z=0.008$ the FeCZ only appears at $L \approx 10^{3.9} L_{\odot}$, corresponding to a zero age main sequence star of about $11M_{\odot}$ (Cantiello et al. 2009).** In our models with metallicity $Z=0.006$ (See Appendix C) the FeCZ appears at $\log \mathcal{L}/L_{\odot} \approx 3.2$.

Note that these limits could move downward in the presence of atomic diffusion and radiative acceleration (Richer et al. 2000), or due to an upward revision of the uncertain values of Fe opacity (e.g. Bailey et al. 2015). We notice that the TESS observations of LMC stars reported by Bowman et al. (2019a) reveal a trend of lower ν_{char} compared to the galactic sample (See their Fig.4). Interestingly, this trend is reproduced by the characteristic frequency of convection in the FeCZ in our models (compare Fig. 3 with Fig. 7). It will be interesting to see if the TESS observations can probe the transition region between stars with and without a FeCZ, and determine a possible change in surface properties. At the same time, X-shooter within the Ulysses program could more firmly establish the metallicity-dependence of the macroturbulent line-broadening (Penny & Gies 2009).

It is important to note that some low-luminosity, main sequence A stars in Bowman et al. (2019b) show SLF variability. For these stars models do not predict the

presence of the FeCZ. However, these stars still show (sub)surface convection zones triggered by ionization of H and He (Cantiello & Braithwaite 2019). The characteristic frequencies of convection in these regions are also in the range $\sim$ tens of μHz , although the amplitude of the velocity fluctuations they can induce is much smaller than for the FeCZ, so it is not clear if they could be linked to observed SLF variability and macroturbulence. On the other hand, since the relative perturbation induced by subsurface convection is weakest in the regime of A and late-B type (Cantiello & Braithwaite 2019; Jermyn & Cantiello 2020), these stars are the best targets for detecting core-generated IGWs. Depending on the amplitude of core generated IGWs, the impact of subsurface convection could well be subdominant in these objects, allowing for a detection. Shiode et al. (2013) predict that in this regime, the peak rms surface brightness fluctuations induced by core-generated IGWs should be in the range $\approx 0.01\text{--}100\mu\text{mag}$ (See their Fig. 6).

5.1. Stochastic, Low-Frequency Variability in Evolved Stars

SLF variability with similar properties to the massive main sequence stars discussed by (Bowman et al. 2019b, 2020) was recently observed in evolved massive stars. Naze et al. (2021) detected SLF variability in both luminous blue variables (LBV) and Wolf-Rayet (WR) stars, and Dorn-Wallenstein et al. (2020) found the same photometric signature in yellow supergiants (YSG). Compared to OB stars, LBV, WR and YSG correspond to later stages of evolution. In particular, WR stars and YSG are likely burning helium in their cores.

The internal structure of OB, LBV, WR and YSG stars changes dramatically, and this can affect substantially the generation and propagation of internal gravity waves. The amount of radiative damping is expected to change due to the large differences in envelope temperature and densities. We recall here that the radiative damping rate γ_{rad} for a traveling g-mode is given by

$$\gamma_{\text{rad}}(\omega, \ell, r) = K_{\text{rad}}(r) k_r^2, \\ K_{\text{rad}}(r) = \frac{16 \sigma T(r)^3}{3 \rho(r)^2 \kappa(r) c_p(r)}.$$

where k_r is the radial wavenumber (eqn. 6) and κ is the opacity. Compact WR stars have surface temperatures that can exceed $\approx 10^5\text{K}$, while OB stars and LBVs have effective temperatures $\approx 10^{4.4\text{--}4.7}\text{K}$, with LBVs found at the cooler end of this range. The surface of extended, low-density, YSGs is cooler than 10^4K . With different types of core convection (H-burning vs He-burning) and radiative damping rates, it would be surprising if core-

generated IGWs in e.g. OB and WR stars showed up at the surface with similar properties. On the other hand, the FeCZ is present below/at the surface in both OB and WR stars, and with similar velocities and convective turnover times. This seems to strengthen the main thesis of this work, adding support to a (sub)surface origin of the observed SLF variability.

The discussion is more complicated for LBVs and YSGs, since at lower temperatures other convective regions can become prominent (driven by H and He recombination, see e.g. Jiang et al. 2018). Dorn-Wallenstein et al. (2020) disfavor a near-surface origin for the observed SLF variability in YSG, on the ground that the observed timescale do not follow the predicted scaling for granulation (Kallinger et al. 2014). However, we believe that the scaling of Kallinger et al. (2014) is not applicable in the regime explored by Dorn-Wallenstein et al. (2020). This scaling was derived for a sample of red giants and solar-like stars, which all have temperatures well-below the temperature for the recombination of hydrogen, and it is not directly applicable to earlier spectral subtypes.

Interestingly, the characteristic timescales of 0.1-1 days found by Dorn-Wallenstein et al. (2020) at $\log T_{\text{eff}} \approx 4$ are consistent with the convective turnover timescale in the FeCZ (e.g. Fig. 3). Moreover, in their lower temperature sample ($\log T_{\text{eff}} < 3.75$) the rapid increase in amplitudes and characteristic frequencies of the variability is in agreement with the development of near-surface convection induced by the large opacities associated with the recombination of hydrogen (Grassitelli et al. 2015a). We suggest that surface and near-surface convection could be indeed responsible for the variability observed by Dorn-Wallenstein et al. (2020), and we plan to systematically study the properties of (sub)surface convection in these evolved, cool stars in future work.

5.2. Towards a unified model for surface phenomena in massive stars

The presence of subsurface convection can simultaneously account for a large variety of puzzling phenomena that appear ubiquitous at the surface of early-type stars.

- **Microturbulence and Macroturbulence** can be accounted for by velocity fields excited by the underlying FeCZ subsurface convection zone (Cantiello et al. 2009; Grassitelli et al. 2015b; Jiang et al. 2015). The only stars with no

macroturbulence appear to have magnetic fields strong enough to shut off convection in the FeCZ, while stars with slightly lower surface magnetic fields show normal values of macroturbulence (Sundqvist et al. 2013).

- **Bright spots** in early-type stars have been observed (e.g. Ramaramanantsoa et al. 2014), and can be explained with the presence of magnetic spots rising from subsurface convective layers (Cantiello & Braithwaite 2011, 2019).
- **Discrete absorption components (DACs)** in UV spectra (e.g. Howarth & Prinja 1989; Cranmer & Owocki 1996; Fullerton et al. 1997; Kaper et al. 1997) can then be caused by the aforementioned bright spots (Cantiello & Braithwaite 2011).
- **Line profile variability** is another ubiquitous phenomena in hot stars (Fullerton et al. 1996), and can be explained by surface velocity and density perturbations seeded by subsurface convection (e.g. Cantiello et al. 2009; Jiang et al. 2015).
- **Wind clumping** can also be seeded by these surface density and velocity perturbations, which are amplified by the development of instabilities in the stellar wind (Owocki et al. 1988, but see also Sundqvist & Owocki (2013)).
- **SLF variability** can also be caused by the presence of subsurface convection zones, as discussed in this work.

An economical hypothesis emerges: the presence of subsurface convection, and in particular of the FeCZ, could be the common underlying physical cause for the appearance of turbulence, magnetic spots, SLF variability, as well spectroscopic variability in early-type stars.

6. CONCLUSIONS

We used one-dimensional stellar evolution calculations to study the predicted trends in the properties of subsurface convection in the spectroscopic H-R diagram. We found that the trends of relative convective flux and convective turnover timescale in the FeCZ of our models match very well the trends in timescale and amplitude of stochastic, low-frequency photometric variability in OB stars observed by TESS and K2.

Similar to previous works, we show that the observed trends in stellar macroturbulence are also well reproduced assuming the FeCZ is its driver. This connection is also supported by the observations of strongly magnetized early-type stars, showing no macroturbulence only for magnetic fields above the critical value required to

shut-off turbulent convection in the FeCZ. We find that IGWs coming from the stellar core would be reflected or damped for values of the magnetic field well below this critical value. The fact that stars with strong but subcritical magnetic fields show typical values of macroturbulence points against a convective core origin of this surface perturbation.

Overall the observations support a picture in which subsurface convection, and in particular the FeCZ, is responsible for the ubiquitous low-frequency, stochastic photometric variability and macroturbulence detected in OB stars. These surface manifestations join a number of phenomena observed in early-type stars and attributed to the presence of subsurface convection, including the observations of (magnetic) bright spots as

well as wind and spectroscopic variability. Radiation (magneto)hydrodynamics simulations of the outer envelope regions of early-type stars are required to understand the details of how subsurface convection zones cause the observed surface perturbations.

ACKNOWLEDGMENTS

The Center for Computational Astrophysics at the Flatiron Institute is supported by the Simons Foundation. This research was supported in part by the National Science Foundation under Grant No. NSF PHY-1748958 and by the Gordon and Betty Moore Foundation through Grant GBMF7392. DL is supported by a Lyman Spitzer Jr. Fellowship.

APPENDIX

A. SOFTWARE DETAILS

Calculations were done with MESA version 11701. The MESA EOS is a blend of the OPAL Rogers & Nayfonov (2002), SCVH Saumon et al. (1995), PTEH Pols et al. (1995), HELM Timmes & Swesty (2000), and PC Potekhin & Chabrier (2010) EOSes. Radiative opacities are primarily from OPAL (Iglesias & Rogers 1993, 1996), with low-temperature data from Ferguson et al. (2005) and the high-temperature, Compton-scattering dominated regime by Buchler & Yueh (1976). Electron conduction opacities are from Cassisi et al. (2007). Nuclear reaction rates are a combination of rates from NACRE (Angulo et al. 1999), JINA REACLIB (Cyburt et al. 2010), plus additional tabulated weak reaction rates Fuller et al. (1985); Oda et al. (1994); Langanke & Martínez-Pinedo (2000). Screening is included via the prescription of Chugunov et al. (2007). Thermal neutrino loss rates are from Itoh et al. (1996). The inlists, processing scripts, and model output are available at [Zenodo.org](https://zenodo.org).

B. RADIAL DISPLACEMENT FROM IGWS

Let's consider traveling internal gravity waves in the WKB approximation. The wave flux is $F_{\text{IGW}} = \frac{1}{2}\rho|u|^2u_{g,r}$ with the group velocity $u_{g,r} = \partial\omega/\partial\kappa$. Since $u_{g,r} = (\omega/N)^2\sqrt{N^2 - \omega^2}/k_h$, $|u|^2 = (N/\omega)^2u_r^2$, and $u_r = \omega\xi_r$ we obtain:

$$F_{\text{IGW}} = \frac{1}{2}\rho\xi_r^2\omega^2\frac{\sqrt{N^2 - \omega^2}}{k_h}. \quad (\text{B1})$$

Using $k_h = \sqrt{\ell(\ell+1)}/r$ we can then write the relative radial displacement as:

$$\frac{\xi_r}{r} = \left[\frac{F_{\text{IGW}}\sqrt{\ell(\ell+1)}}{1/2\rho r^3\omega^2\sqrt{N^2 - \omega^2}} \right]^{1/2} \propto \sqrt{\frac{F_{\text{IGW}}}{F_0}} \quad (\text{B2})$$

C. GRIDS AT DIFFERENT METALLICITIES

Here we present results for the critical and surface magnetic field at metallicities of $Z=0.006$ and $Z=0.002$, representing early-type stars in the Large Magellanic Cloud (LMC, Fig. 6 and 7) and Small Magellanic Cloud (SMC, Fig. 8 and 9) respectively (e.g. Yusof et al. 2013). The initial Helium content of the grids is $Y=0.2559$ (LMC) and $Y=0.2508$ (SMC). The metallicity is initialized scaling the standard solar composition of Grevesse & Sauval (1998).

REFERENCES

- Aerts, C. 2019, arXiv e-prints, arXiv:1912.12300.
<https://arxiv.org/abs/1912.12300>
- Aerts, C., Christensen-Dalsgaard, J., & Kurtz, D. W. 2010, Asteroseismology




Figure 6. Left: Average convective velocities in the FeCZ as function of the location of stellar models in the spectroscopic H-R Diagram for models with $Z=0.006$ (LMC). The area of the dot is proportional to the observed macroturbulence. **The FeCZ is absent in models with $\log L/L_\odot \lesssim 3.2$.** Right: Same as left, but showing predictions for the maximum of the ratio between turbulent pressure and total pressure in any subsurface convection zone. The FeCZ largely dominates, except for stars at low luminosity where turbulent pressure is provided by a helium convection zone (HeCZ).

- Aerts, C., Mathis, S., & Rogers, T. M. 2019, *ARA&A*, 57, 35, doi: [10.1146/annurev-astro-091918-104359](https://doi.org/10.1146/annurev-astro-091918-104359)
- Aerts, C., Puls, J., Godart, M., & Dupret, M. A. 2009, *A&A*, 508, 409, doi: [10.1051/0004-6361/200810471](https://doi.org/10.1051/0004-6361/200810471)
- Angulo, C., Arnould, M., Rayet, M., et al. 1999, *Nuclear Physics A*, 656, 3, doi: [10.1016/S0375-9474\(99\)00030-5](https://doi.org/10.1016/S0375-9474(99)00030-5)
- Bailey, J. E., Nagayama, T., Loisel, G. P., et al. 2015, *Nature*, 517, 56, doi: [10.1038/nature14048](https://doi.org/10.1038/nature14048)
- Blomme, R., Mahy, L., Catala, C., et al. 2011, *A&A*, 533, A4, doi: [10.1051/0004-6361/201116949](https://doi.org/10.1051/0004-6361/201116949)
- Böhm-Vitense, E. 1958, *ZA*, 46, 108
- Bowman, D. M. 2020, arXiv e-prints, arXiv:2008.11162, <https://arxiv.org/abs/2008.11162>
- Bowman, D. M., Burssens, S., Simón-Díaz, S., et al. 2020, *A&A*, 640, A36, doi: [10.1051/0004-6361/202038224](https://doi.org/10.1051/0004-6361/202038224)
- Bowman, D. M., Burssens, S., Pedersen, M. G., et al. 2019a, *Nature Astronomy*, 3, 760, doi: [10.1038/s41550-019-0768-1](https://doi.org/10.1038/s41550-019-0768-1)
- Bowman, D. M., Aerts, C., Johnston, C., et al. 2019b, *A&A*, 621, A135, doi: [10.1051/0004-6361/201833662](https://doi.org/10.1051/0004-6361/201833662)
- Buchler, J. R., & Yueh, W. R. 1976, *ApJ*, 210, 440, doi: [10.1086/154847](https://doi.org/10.1086/154847)
- Cantiello, M., & Braithwaite, J. 2011, *A&A*, 534, A140, doi: [10.1051/0004-6361/201117512](https://doi.org/10.1051/0004-6361/201117512)
- . 2019, *ApJ*, 883, 106, doi: [10.3847/1538-4357/ab3924](https://doi.org/10.3847/1538-4357/ab3924)
- Cantiello, M., Langer, N., Brott, I., et al. 2009, *A&A*, 499, 279, doi: [10.1051/0004-6361/200911643](https://doi.org/10.1051/0004-6361/200911643)
- Cassisi, S., Potekhin, A. Y., Pietrinferni, A., Catelan, M., & Salaris, M. 2007, *ApJ*, 661, 1094, doi: [10.1086/516819](https://doi.org/10.1086/516819)
- Chugunov, A. I., Dewitt, H. E., & Yakovlev, D. G. 2007, *PhRvD*, 76, 025028, doi: [10.1103/PhysRevD.76.025028](https://doi.org/10.1103/PhysRevD.76.025028)
- Cranmer, S. R., & Owocki, S. P. 1996, *ApJ*, 462, 469, doi: [10.1086/177166](https://doi.org/10.1086/177166)
- Cyburt, R. H., Amthor, A. M., Ferguson, R., et al. 2010, *ApJS*, 189, 240, doi: [10.1088/0067-0049/189/1/240](https://doi.org/10.1088/0067-0049/189/1/240)
- de Mink, S. E., Sana, H., Langer, N., Izzard, R. G., & Schneider, F. R. N. 2014, *ApJ*, 782, 7, doi: [10.1088/0004-637X/782/1/7](https://doi.org/10.1088/0004-637X/782/1/7)
- Dorn-Wallenstein, T. Z., Levesque, E. M., Neugent, K. F., et al. 2020, *ApJ*, 902, 24, doi: [10.3847/1538-4357/abb318](https://doi.org/10.3847/1538-4357/abb318)
- Dziembowski, W. 1977, *AcA*, 27, 203
- Edelmann, P. V. F., Ratnasingam, R. P., Pedersen, M. G., et al. 2019, *ApJ*, 876, 4, doi: [10.3847/1538-4357/ab12df](https://doi.org/10.3847/1538-4357/ab12df)
- Ferguson, J. W., Alexander, D. R., Allard, F., et al. 2005, *ApJ*, 623, 585, doi: [10.1086/428642](https://doi.org/10.1086/428642)
- Fuller, G. M., Fowler, W. A., & Newman, M. J. 1985, *ApJ*, 293, 1, doi: [10.1086/163208](https://doi.org/10.1086/163208)




Figure 7. Left panel: Ratio of FeCZ convective flux to the total stellar flux in the spectroscopic H-R Diagram for models with $Z=0.006$ (LMC). Right panel: Characteristic frequency ν_{char} in the FeCZ as a function of the location of stellar models in the spectroscopic H-R Diagram (black contour lines). Evolutionary tracks are shown as grey solid lines.

- Fuller, J., Cantiello, M., Stello, D., Garcia, R. A., & Bildsten, L. 2015, *Science*, 350, 423, doi: [10.1126/science.aac6933](https://doi.org/10.1126/science.aac6933)
- Fullerton, A. W., Gies, D. R., & Bolton, C. T. 1996, *ApJS*, 103, 475, doi: [10.1086/192285](https://doi.org/10.1086/192285)
- Fullerton, A. W., Massa, D. L., Prinja, R. K., Owocki, S. P., & Cranmer, S. R. 1997, *A&A*, 327, 699
- Godart, M., Simón-Díaz, S., Herrero, A., et al. 2017, *A&A*, 597, A23, doi: [10.1051/0004-6361/201628856](https://doi.org/10.1051/0004-6361/201628856)
- Goldreich, P., & Kumar, P. 1990, *ApJ*, 363, 694, doi: [10.1086/169376](https://doi.org/10.1086/169376)
- Grassitelli, L., Fossati, L., Langer, N., et al. 2015a, *A&A*, 584, L2, doi: [10.1051/0004-6361/201527289](https://doi.org/10.1051/0004-6361/201527289)
- . 2016, *A&A*, 593, A14, doi: [10.1051/0004-6361/201628912](https://doi.org/10.1051/0004-6361/201628912)
- Grassitelli, L., Fossati, L., Simón-Díaz, S., et al. 2015b, *ApJL*, 808, L31, doi: [10.1088/2041-8205/808/1/L31](https://doi.org/10.1088/2041-8205/808/1/L31)
- Grevesse, N., & Sauval, A. J. 1998, *SSRv*, 85, 161, doi: [10.1023/A:1005161325181](https://doi.org/10.1023/A:1005161325181)
- Hopkins, P. F., Kereš, D., Oñorbe, J., et al. 2014, *MNRAS*, 445, 581, doi: [10.1093/mnras/stu1738](https://doi.org/10.1093/mnras/stu1738)
- Horst, L., Edelmann, P. V. F., Andrásy, R., et al. 2020, *A&A*, 641, A18, doi: [10.1051/0004-6361/202037531](https://doi.org/10.1051/0004-6361/202037531)
- Howarth, I. D., & Prinja, R. K. 1989, *ApJS*, 69, 527, doi: [10.1086/191321](https://doi.org/10.1086/191321)
- Iglesias, C. A., & Rogers, F. J. 1993, *ApJ*, 412, 752, doi: [10.1086/172958](https://doi.org/10.1086/172958)
- . 1996, *ApJ*, 464, 943, doi: [10.1086/177381](https://doi.org/10.1086/177381)
- Itoh, N., Hayashi, H., Nishikawa, A., & Kohyama, Y. 1996, *ApJS*, 102, 411, doi: [10.1086/192264](https://doi.org/10.1086/192264)
- Jermyn, A. S., & Cantiello, M. 2020, *ApJ*, 900, 113, doi: [10.3847/1538-4357/ab9e70](https://doi.org/10.3847/1538-4357/ab9e70)
- Jiang, Y.-F., Cantiello, M., Bildsten, L., Quataert, E., & Blaes, O. 2015, *ApJ*, 813, 74, doi: [10.1088/0004-637X/813/1/74](https://doi.org/10.1088/0004-637X/813/1/74)
- . 2017, *ApJ*, 843, 68, doi: [10.3847/1538-4357/aa77b0](https://doi.org/10.3847/1538-4357/aa77b0)
- Jiang, Y.-F., Cantiello, M., Bildsten, L., et al. 2018, *Nature*, 561, 498, doi: [10.1038/s41586-018-0525-0](https://doi.org/10.1038/s41586-018-0525-0)
- Kallinger, T., De Ridder, J., Hekker, S., et al. 2014, *A&A*, 570, A41, doi: [10.1051/0004-6361/201424313](https://doi.org/10.1051/0004-6361/201424313)
- Kaper, L., Henrichs, H. F., Fullerton, A. W., et al. 1997, *A&A*, 327, 281
- Langanke, K., & Martínez-Pinedo, G. 2000, *Nuclear Physics A*, 673, 481, doi: [10.1016/S0375-9474\(00\)00131-7](https://doi.org/10.1016/S0375-9474(00)00131-7)
- Langer, N. 2012, *ARA&A*, 50, 107, doi: [10.1146/annurev-astro-081811-125534](https://doi.org/10.1146/annurev-astro-081811-125534)
- Langer, N., & Kudritzki, R. P. 2014, *A&A*, 564, A52, doi: [10.1051/0004-6361/201423374](https://doi.org/10.1051/0004-6361/201423374)
- Lecoanet, D., & Quataert, E. 2013, *MNRAS*, 430, 2363, doi: [10.1093/mnras/stt055](https://doi.org/10.1093/mnras/stt055)




Figure 8. Left: Average convective velocities in the FeCZ as function of the location of stellar models in the spectroscopic H-R Diagram for models with $Z=0.002$ (SMC). The area of the dot is proportional to the observed macroturbulence. **The FeCZ is absent in models with $\log L/L_\odot \lesssim 3.2$.** Right: Same as left, but showing predictions for the maximum of the ratio between turbulent pressure and total pressure in any subsurface convection zone. The FeCZ largely dominates, except for stars at low luminosity where turbulent pressure is provided by a helium convection zone.

Lecoanet, D., Cantiello, M., Quataert, E., et al. 2019, ApJL, 886, L15, doi: [10.3847/2041-8213/ab5446](https://doi.org/10.3847/2041-8213/ab5446)

MacDonald, J., & Petit, V. 2019, MNRAS, 487, 3904, doi: [10.1093/mnras/stz1545](https://doi.org/10.1093/mnras/stz1545)

Naze, Y., Rauw, G., & Gosset, E. 2021, arXiv e-prints, arXiv:2101.05521. <https://arxiv.org/abs/2101.05521>

Oda, T., Hino, M., Muto, K., Takahara, M., & Sato, K. 1994, Atomic Data and Nuclear Data Tables, 56, 231, doi: [10.1006/adnd.1994.1007](https://doi.org/10.1006/adnd.1994.1007)

Owocki, S. P., Castor, J. I., & Rybicki, G. B. 1988, ApJ, 335, 914, doi: [10.1086/166977](https://doi.org/10.1086/166977)

Paxton, B., Bildsten, L., Dotter, A., et al. 2011, ApJS, 192, 3, doi: [10.1088/0067-0049/192/1/3](https://doi.org/10.1088/0067-0049/192/1/3)

Paxton, B., Cantiello, M., Arras, P., et al. 2013, ApJS, 208, 4, doi: [10.1088/0067-0049/208/1/4](https://doi.org/10.1088/0067-0049/208/1/4)

Paxton, B., Marchant, P., Schwab, J., et al. 2015, ApJS, 220, 15, doi: [10.1088/0067-0049/220/1/15](https://doi.org/10.1088/0067-0049/220/1/15)

Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, ApJS, 234, 34, doi: [10.3847/1538-4365/aaa5a8](https://doi.org/10.3847/1538-4365/aaa5a8)

Paxton, B., Smolec, R., Schwab, J., et al. 2019, ApJS, 243, 10, doi: [10.3847/1538-4365/ab2241](https://doi.org/10.3847/1538-4365/ab2241)

Pedersen, M. G., Chowdhury, S., Johnston, C., et al. 2019, ApJL, 872, L9, doi: [10.3847/2041-8213/ab01e1](https://doi.org/10.3847/2041-8213/ab01e1)

Penny, L. R., & Gies, D. R. 2009, ApJ, 700, 844, doi: [10.1088/0004-637X/700/1/844](https://doi.org/10.1088/0004-637X/700/1/844)

Pols, O. R., Tout, C. A., Eggleton, P. P., & Han, Z. 1995, MNRAS, 274, 964, doi: [10.1093/mnras/274.3.964](https://doi.org/10.1093/mnras/274.3.964)

Potekhin, A. Y., & Chabrier, G. 2010, Contributions to Plasma Physics, 50, 82, doi: [10.1002/ctpp.201010017](https://doi.org/10.1002/ctpp.201010017)

Press, W. H. 1981, ApJ, 245, 286, doi: [10.1086/158809](https://doi.org/10.1086/158809)

Ramiamananantsoa, T., Moffat, A. F. J., Chené, A.-N., et al. 2014, MNRAS, 441, 910, doi: [10.1093/mnras/stu619](https://doi.org/10.1093/mnras/stu619)

Ratnasingam, R. P., Edelmann, P. V. F., & Rogers, T. M. 2019, MNRAS, 482, 5500, doi: [10.1093/mnras/sty3086](https://doi.org/10.1093/mnras/sty3086)

—. 2020, MNRAS, doi: [10.1093/mnras/staa2296](https://doi.org/10.1093/mnras/staa2296)

Richer, J., Michaud, G., & Turcotte, S. 2000, ApJ, 529, 338, doi: [10.1086/308274](https://doi.org/10.1086/308274)

Rogers, F. J., & Nayfonov, A. 2002, ApJ, 576, 1064, doi: [10.1086/341894](https://doi.org/10.1086/341894)

Rogers, T. M., Lin, D. N. C., McElwaine, J. N., & Lau, H. H. B. 2013, ApJ, 772, 21, doi: [10.1088/0004-637X/772/1/21](https://doi.org/10.1088/0004-637X/772/1/21)

Sana, H., de Mink, S. E., de Koter, A., et al. 2012, Science, 337, 444, doi: [10.1126/science.1223344](https://doi.org/10.1126/science.1223344)

Saumon, D., Chabrier, G., & van Horn, H. M. 1995, ApJS, 99, 713, doi: [10.1086/192204](https://doi.org/10.1086/192204)




Figure 9. Left panel: Ratio of FeCZ convective flux to the total stellar flux in the spectroscopic H-R Diagram for models with $Z=0.002$ (SMC). Right panel: Characteristic frequency ν_{char} in the FeCZ as a function of the location of stellar models in the spectroscopic H-R Diagram (black contour lines). Evolutionary tracks are shown as grey solid lines.

Shiode, J. H., Quataert, E., Cantiello, M., & Bildsten, L. 2013, *MNRAS*, 430, 1736, doi: [10.1093/mnras/sts719](https://doi.org/10.1093/mnras/sts719)
 Simón-Díaz, S., Godart, M., Castro, N., et al. 2017a, *A&A*, 597, A22, doi: [10.1051/0004-6361/201628541](https://doi.org/10.1051/0004-6361/201628541)
 —. 2017b, *A&A*, 597, A22, doi: [10.1051/0004-6361/201628541](https://doi.org/10.1051/0004-6361/201628541)
 Simón-Díaz, S., & Herrero, A. 2014, *A&A*, 562, A135, doi: [10.1051/0004-6361/201322758](https://doi.org/10.1051/0004-6361/201322758)
 Sundqvist, J. O., & Owocki, S. P. 2013, *MNRAS*, 428, 1837, doi: [10.1093/mnras/sts165](https://doi.org/10.1093/mnras/sts165)

Sundqvist, J. O., Petit, V., Owocki, S. P., et al. 2013, *Monthly Notices of the Royal Astronomical Society*, 433, 2497, doi: [10.1093/mnras/stt921](https://doi.org/10.1093/mnras/stt921)
 Sutherland, B. R. 2010, *Internal Gravity Waves* (Cambridge University Press), doi: [10.1017/CBO9780511780318](https://doi.org/10.1017/CBO9780511780318)
 Timmes, F. X., & Swesty, F. D. 2000, *ApJS*, 126, 501, doi: [10.1086/313304](https://doi.org/10.1086/313304)
 Yusof, N., Hirschi, R., Meynet, G., et al. 2013, *MNRAS*, 433, 1114, doi: [10.1093/mnras/stt794](https://doi.org/10.1093/mnras/stt794)