

EECS 127/227AT Optimization Models in Engineering

Spring 2020

Discussion 8

1. Complementary slackness

Consider the problem:

$$\begin{aligned} p^* &= \min_{x \in \mathbb{R}} x^2 \\ \text{s.t. } &x \geq 1, x \leq 2. \end{aligned}$$

- (a) Does Slater's condition hold? Is the problem convex? Does strong duality hold?

Solution: We have a strictly feasible point $x = 1.5$ that lies in the relative interior of the domain of the objective function; thus, Slater's condition holds. The objective function x^2 is convex and the inequality constraints are affine and thus convex, so the problem is convex. Since Slater's condition holds and the problem is convex, strong duality holds.

- (b) Find the Lagrangian $\mathcal{L}(x, \lambda_1, \lambda_2)$.

Solution: $\mathcal{L}(x, \lambda_1, \lambda_2) = x^2 + \lambda_1(-x + 1) + \lambda_2(x - 2)$.

- (c) Solve for $x^*, \lambda_1^*, \lambda_2^*$ that satisfy KKT conditions.

Solution: We have:

From stationarity,

$$\begin{aligned} \nabla_x \mathcal{L}(x, \lambda_1, \lambda_2) &= 0 \\ \implies 2x - \lambda_1 + \lambda_2 &= 0. \end{aligned} \tag{1}$$

From primal feasibility,

$$\begin{aligned} x &\geq 1 \\ x &\leq 2. \end{aligned}$$

From dual feasibility,

$$\begin{aligned} \lambda_1 &\geq 0 \\ \lambda_2 &\geq 0. \end{aligned}$$

Finally from complementary slackness,

$$\begin{aligned} \lambda_1(-x + 1) &= 0 \\ \lambda_2(x - 2) &= 0. \end{aligned}$$

First observe that we cannot have $\lambda_1 \neq 0$ and $\lambda_2 \neq 0$ since in this case complementary slackness would not have any feasible solutions for x . Next assume that $\lambda_1 = 0, \lambda_2 \neq 0$. Then from complementary slackness, $x = 2$. Substituting this in equation 1, we get $\lambda_2 = -4$ which violates dual feasibility. Next assume that $\lambda_1 \neq 0, \lambda_2 = 0$. Then from 1 we have $x = 0$ which violates primal feasibility. Finally assume that $\lambda_1 \neq 0, \lambda_2 \neq 0$. From complementary slackness we have $x = 1$ and from 1 we have $\lambda_1 = 2$ which satisfies dual feasibility.

Thus $x^* = 1, \lambda_1^* = 2, \lambda_2^* = 0$ satisfy KKT conditions.

- (d) Can you spot a connection between the values of λ_1^*, λ_2^* in relation to whether the corresponding inequality constraints are strict or not at the optimal x^* ?

Solution: We have $\lambda_1 \neq 0$ and the corresponding inequality $x \geq 1$ is satisfied with equality (and hence is not strict) at $x^* = 1$.

We have $\lambda_2 = 0$ and the corresponding inequality is strict at $x^* = 1$. The non-zero λ_1 tells us that if we relax the constraint $x \geq 1$ (for example, to $x \geq 0.9$) we can reduce the objective function further.

- (e) Find the dual function $g(\lambda_1, \lambda_2)$ so that the dual problem is given by,

$$d^* = \max_{\lambda_1, \lambda_2 \in \mathbb{R}^+} g(\lambda_1, \lambda_2). \quad (2)$$

Solution:

$$g(\lambda_1, \lambda_2) = \inf_x \mathcal{L}(x, \lambda_1, \lambda_2).$$

Note that \mathcal{L} is convex with respect to x , thus setting the gradient with respect to x to 0 we obtain, $x = \frac{\lambda_1 - \lambda_2}{2}$. Thus,

$$g(\lambda_1, \lambda_2) = -\frac{(\lambda_2 - \lambda_1)^2}{4} + \lambda_1 - 2\lambda_2.$$

- (f) Solve the dual problem in (2) for d^* .

Solution: Let us first try setting gradient with respect to λ_1 and λ_2 to 0. This gives us,

$$\begin{aligned} \frac{\lambda_2 - \lambda_1}{2} + 1 &= 0 \\ -\frac{\lambda_2 - \lambda_1}{2} - 2 &= 0. \end{aligned}$$

This has no solution. We can see that a quadratic objective function could be unbounded even if it was convex. To get meaningful solutions we must check for optimal values at the boundaries. Checking at boundary $\lambda_1 = 0$.

$$g(0, \lambda_2) = -\frac{\lambda_2^2}{4} - 2\lambda_2.$$

This is a concave function so taking gradient with respect to λ_2 and setting it to zero we obtain,

$$\begin{aligned} -\frac{\lambda_2}{2} - 2 &= 0 \\ \implies \lambda_2 &= -4. \end{aligned}$$

This is not feasible so we must check value at $\lambda_2 = 0$. We have $g(0, 0) = 0$. Finally let us check at the other boundary $\lambda_2 = 0$.

$$g(\lambda_1, 0) = -\frac{\lambda_1^2}{4} + \lambda_1.$$

Again this is a concave function so taking gradient with respect to λ_1 and setting it to zero we obtain,

$$\begin{aligned} -\frac{\lambda_1}{2} + 1 &= 0 \\ \implies \lambda_1 &= 2. \end{aligned}$$

We have $g(2, 0) = -1 + 2 = 1$. Thus $d^* = 1$.

2. [Optional] Simple constrained optimization problem with duality

Consider the optimization problem

$$\begin{aligned} \min_{x_1, x_2 \in \mathbb{R}} f(x_1, x_2) \\ \text{subject to } 2x_1 + x_2 \geq 1 \\ x_1 + 3x_2 \geq 1 \\ x_1 \geq 0, x_2 \geq 0 \end{aligned}$$

- (a) Express the Lagrangian of the problem $\mathcal{L}(x_1, x_2, \lambda_1, \lambda_2, \lambda_3, \lambda_4)$

Solution:

$$\mathcal{L}(x_1, x_2, \lambda_1, \lambda_2, \lambda_3, \lambda_4) = f(x_1, x_2) + \lambda_1(-2x_1 - x_2 + 1) + \lambda_2(1 - x_1 - 3x_2) - \lambda_3x_1 - \lambda_4x_2$$

Solve the following problems analytically and give the minimizing x_1^*, x_2^* : *Hint:* Use duality if the problem is hard to solve. Use the graphs in Figure 1 to "dualize" only some constraints:

- (b) $f(x_1, x_2) = x_1 + x_2$

Solution: From the drawing, it seems that at the optimal point $x_1 \geq 0$ and $x_2 \geq 0$ are not active. So let's assume that $x_1^* > 0$ and $x_2^* > 0$ so $\lambda_3 = 0$ and $\lambda_4 = 0$ by Complementary Slackness.

So we can solve the following problem

$$\begin{aligned} \min_{\vec{x}, \vec{x} \geq 0} \max_{\lambda_1, \lambda_2 \geq 0} f(x_1, x_2) + \lambda_1(-2x_1 - x_2 + 1) + \lambda_2(1 - x_1 - 3x_2) \\ = \max_{\lambda_1, \lambda_2 \geq 0} \min_{\vec{x}, \vec{x} \geq 0} f(x_1, x_2) + \lambda_1(-2x_1 - x_2 + 1) + \lambda_2(1 - x_1 - 3x_2) \quad (\text{Strong duality}) \\ \implies \nabla_{x_1, x_2} \mathcal{L}(x_1^*, x_2^*, \lambda_1, \lambda_2) = 0 \quad (\text{Stationarity}) \\ \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2\lambda_1 + \lambda_2 \\ \lambda_1 + 3\lambda_2 \end{pmatrix} \end{aligned}$$

This gives $\lambda_1 = \frac{2}{5} > 0$ and $\lambda_2 = \frac{1}{5} > 0$.

And $(-2x_1^* - x_2^* + 1) = 0$ and $(1 - x_1^* - 3x_2^*) = 0$.

So $x_1^* = \frac{2}{5}$ and $x_2^* = \frac{1}{5}$, which satisfies the assumption condition $x_1^* > 0$ and $x_2^* > 0$.

The optimal point is $x_1^* = \frac{2}{5}$ and $x_2^* = \frac{1}{5}$ and $f(x_1^*, x_2^*) = \frac{3}{5}$.

- (c) $f(x_1, x_2) = -x_1 - x_2$

Solution: Here, $t \cdot (1, 1)$ is a feasible point when $t \geq 1$. And $f(t \cdot (1, 1)) = -2t$ which is unbounded below. So the problem is unbounded below.

- (d) $f(x_1, x_2) = x_1$

Solution: Under the constraints $x_1 \geq 0$, we know that $f(x_1, x_2) \geq 0$.

Also $f(x_1, x_2) = 0$ implies that $x_1 = 0$ and the constraints becomes equivalent to $x_2 \geq 1$.

Therefore the set of solutions is $S = \{\vec{x}, x_1 = 0 \text{ and } x_2 \geq 1\}$ and for all $\vec{x} \in S$, $f(\vec{x}) = 0$.




Figure 1: Heatmap of 2(b), 2(c), 2(d), 2(e) and 2(f): $\bar{x}^* = (\frac{2}{5}, \frac{1}{5})$. In red is the unfeasible points, then the level sets are shown with colors; blue points are points (x_1, x_2) with the lowest value $f(x_1, x_2)$, red points are the ones with highest value.

(e) $f(x_1, x_2) = \max\{x_1, x_2\}$

Solution: In order to make the Lagrangian of the problem differentiable with respect to \vec{x} , we can use a slack variable. The problem is equivalent to:

$$\begin{aligned} & \min_{x_1, x_2, t} t \\ & \text{subject to } t \geq x_1, t \geq x_2 \\ & \quad 2x_1 + x_2 \geq 1 \\ & \quad x_1 + 3x_2 \geq 1 \\ & \quad x_1 \geq 0, x_2 \geq 0 \end{aligned}$$

We have the augmented Lagrangian:

$$\begin{aligned} \mathcal{L}(t, x_1, x_2, \lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6) = & t + \lambda_1(-2x_1 - x_2 + 1) + \lambda_2(1 - x_1 - 3x_2) - \lambda_3x_1 - \lambda_4x_2 \\ & + \lambda_5(x_1 - t) + \lambda_6(x_2 - t) \end{aligned}$$

Using the drawing (see figure 1), we see that several constraints are not active at the optimum and we can make the assumption that $\lambda_2 = 0$, $\lambda_3 = 0$ and $\lambda_4 = 0$ by complementary slackness. So we can solve the following problem (with $\mathcal{X} = \{\vec{x} \mid x_1 + 3x_2 \geq 1, x_1 \geq 0 \text{ and } x_2 \geq 0\}$)

$$\begin{aligned} & \min_{(\vec{x}, t), \vec{x} \in \mathcal{X}} \max_{\lambda_1, \lambda_5, \lambda_6} t + \lambda_1(-2x_1 - x_2 + 1) + \lambda_5(x_1 - t) + \lambda_6(x_2 - t) \\ = & \max_{\lambda_1, \lambda_5, \lambda_6} \min_{(\vec{x}, t), \vec{x} \in \mathcal{X}} t + \lambda_1(-2x_1 - x_2 + 1) + \lambda_5(x_1 - t) + \lambda_6(x_2 - t) \quad (\text{Strong duality}) \\ \implies & \nabla_{x_1, x_2, t} \mathcal{L}(x_1^*, x_2^*, t^*, \lambda_1, \lambda_5, \lambda_6) = 0 \quad (\text{Stationarity}) \\ & \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2\lambda_1 - \lambda_5 \\ \lambda_1 - \lambda_6 \\ \lambda_5 + \lambda_6 \end{pmatrix} \end{aligned}$$

This gives that $\lambda_1 = \frac{1}{3}$, $\lambda_5 = \frac{1}{3}$ and $\lambda_6 = \frac{1}{3}$.
Complementary slackness gives that

$$t^* = x_1^* = x_2^* = \frac{1}{3}$$

Here, $x_1^* + 3x_2^* > 1$, $x_1^* > 0$ and $x_2^* > 0$, so the assumptions are valid.

(f) $f(x_1, x_2) = x_1^2 + 9x_2^2$

Solution: Using the drawing (see figure 1), we can make the assumptions that $\lambda_1 = 0$, $\lambda_3 = 0$ and $\lambda_4 = 0$.

So we can solve the following problem (with $\mathcal{X} = \{\vec{x} \mid 2x_1 + x_2 \geq 1, x_1 \geq 0 \text{ and } x_2 \geq 0\}$)

$$\begin{aligned} & \min_{\vec{x}, \vec{x} \geq 0} \max_{\lambda_2} f(x_1, x_2) + \lambda_2(1 - x_1 - 3x_2) \\ = & \max_{\lambda_2} \min_{\vec{x}, \vec{x} \in \mathcal{X}} f(x_1, x_2) + \lambda_2(1 - x_1 - 3x_2) \quad (\text{Strong duality}) \\ \implies & \nabla_{x_1, x_2} \mathcal{L}(x_1^*, x_2^*, \lambda_2) = 0 \quad (\text{Stationarity}) \\ & \begin{pmatrix} 2x_1^* \\ 18x_2^* \end{pmatrix} = \begin{pmatrix} \lambda_1 \\ 3\lambda_2 \end{pmatrix} \end{aligned}$$

So $\lambda_2 = 2x_1^* = 6x_2^*$. $(1 - x_1 - 3x_2) = 0$ gives that $x_1^* = \frac{1}{2}$, $x_2^* = \frac{1}{6}$ and $\lambda_2 = 1$.
We have that $2x_1^* + x_2^* > 1$, $x_1^* > 0$ and $x_2^* > 0$, so the assumptions are valid.