

Machine Learning

Uniform Convergence

Fabio Vandin

November 10th, 2023

Finite Classes are Agnostic PAC Learnable

We prove that finite sets of hypotheses are agnostic PAC learnable under some restriction for the loss.

Proposition

Let \mathcal{H} be a finite hypothesis class, let Z be a domain, and let $\ell : \mathcal{H} \times Z \rightarrow [0, 1]$ be a loss function. Then:

- \mathcal{H} enjoys the uniform convergence property with sample complexity

$$m_{\mathcal{H}}^{UC}(\varepsilon, \delta) \leq \left\lceil \frac{\log(2|\mathcal{H}|/\delta)}{2\varepsilon^2} \right\rceil$$

- \mathcal{H} is agnostically PAC learnable using the ERM algorithm with sample complexity

$$m_{\mathcal{H}}(\varepsilon, \delta) \leq m_{\mathcal{H}}^{UC}(\varepsilon/2, \delta) \leq \left\lceil \frac{2 \log(2|\mathcal{H}|/\delta)}{\varepsilon^2} \right\rceil$$

Idea of the proof:

- ① prove that uniform convergence holds for a finite hypothesis class
- ② use previous result on uniform convergence and PAC learnability

Useful tool: Hoeffding's Inequality

Hoeffding's Inequality

Let $\theta_1, \dots, \theta_m$ be a sequence of i.i.d. random variables and assume that for all i , $\mathbb{E}[\theta_i] = \mu$ and $\mathbb{P}[a \leq \theta_i \leq b] = 1$. Then, for any $\varepsilon > 0$

$$\mathbb{P}\left[\left|\frac{1}{m} \sum_{i=1}^m \theta_i - \mu\right| > \varepsilon\right] \leq 2e^{-\frac{2m\varepsilon^2}{(b-a)^2}}$$

average of the observations \approx expectation

Proof (see also the book) [Corollary 1.6]

Fix $\varepsilon, \delta \in (0,1)$. We need a sample size m such that, for any \mathcal{H} , with probability $\geq 1-\delta$ (over the choice of $S = (z_1, z_2, \dots, z_m)$, $z_i = (\vec{x}_i, y_i)$), we have:
 for all $h \in \mathcal{H}$: $|L_S(h) - L_{\mathcal{O}}(h)| \leq \varepsilon$.

That is: $\mathbb{P}^m(\{S: \forall h \in \mathcal{H}, |L_S(h) - L_{\mathcal{O}}(h)| \leq \varepsilon\}) \geq 1-\delta$,
 where $S = (z_1, z_2, \dots, z_m)$, $z_i = (\vec{x}_i, y_i)$, i.i.d. from \mathcal{O} . Equivalently, we need to show:

$$\mathbb{P}^m(\{S: \exists h \in \mathcal{H}, |L_S(h) - L_{\mathcal{O}}(h)| > \varepsilon\}) < \delta$$

(*)

We have: $\{S: \exists h \in \mathcal{H}, |L_S(h) - L_{\mathcal{O}}(h)| > \varepsilon\} = \bigcup_{h \in \mathcal{H}} \{S: |L_S(h) - L_{\mathcal{O}}(h)| > \varepsilon\}$
 Then: $(*) \leq \sum_{h \in \mathcal{H}} \mathbb{P}^m(\{S: |L_S(h) - L_{\mathcal{O}}(h)| > \varepsilon\})$ (**)

Now we want to bound each term in $L_{\emptyset}(h)$

Recall: $L_{\emptyset}(h) = \mathbb{E}_{z \sim \emptyset} [l(h, z)]$

$$L_s(h) = \frac{1}{m} \sum_{i=1}^m l(h, z_i)$$

Important: each z_i is sampled i.i.d. from \emptyset

$$\mathbb{E}[l(h, z_i)] = \mathbb{E}_{z \sim \emptyset}[l(h, z)] = L_{\emptyset}(h)$$

Therefore: $\mathbb{E}[L_s(h)] = \mathbb{E}\left[\frac{1}{m} \sum_{i=1}^m l(h, z_i)\right]$

by def. of $L_s(h)$

by linearity
of expectation $\rightarrow = \frac{1}{m} \sum_{i=1}^m \underbrace{\mathbb{E}[l(h, z_i)]}_{L_{\emptyset}(h)}$

$$= \frac{1}{m} \cdot m \cdot L_{\emptyset}(h) = L_{\emptyset}(h)$$

Let θ_i be the r.v. given by $l(h, z_i)$ *i-th element of S*

Since h is fixed and z_i is sampled i.i.d. from \mathcal{D}

$\Rightarrow \theta_1, \theta_2, \dots, \theta_m$ are i.i.d. r.v.

Note that: $L_S(h) = \frac{1}{m} \sum_{i=1}^m \theta_i$. Let's define $\mu = L_\Theta(h)$

Given assumption that $l: \mathcal{H} \times \mathcal{Z} \rightarrow [0, 1]$

we have $\theta_i \in [0, 1]$, $\forall i = 1, \dots, m$.

We can apply Hoeffding's inequality with $a_i = 0$, $b_i = 1 \quad \forall i = 1, \dots, m$

$$\mathcal{D}\left(\{S : |L_S(h) - L_\Theta(h)| > \varepsilon\}\right) = \Pr\left[\left| \left(\frac{1}{m} \sum_{i=1}^m \theta_i \right) - \mu \right| > \varepsilon \right]$$

$$\text{by Hoeffding's ineq. } \rightarrow \leq 2 \cdot e^{-2m \cdot \varepsilon^2}$$

Combining the inequality above with $(\star\star)$

$$\mathcal{D}\left(\{S : \exists h \in \mathcal{H}, |L_S(h) - L_\Theta(h)| > \varepsilon\}\right) \leq \sum_{h \in \mathcal{H}} 2 e^{-2m \varepsilon^2}$$

$$= 2|\mathcal{H}| e^{-2m\epsilon^2}$$

By choosing $m \geq \lg \left(\frac{2|\mathcal{H}|}{\delta} \right) \cdot \frac{1}{2\epsilon^2}$ then

$$\begin{aligned} ① (\exists S: \exists h \in \mathcal{H}, |\mathcal{L}_S(h) - \mathcal{L}_0(h)| > \epsilon) &\leq \\ &\leq 2|\mathcal{H}| e^{-2\epsilon^2 \lg \left(\frac{2|\mathcal{H}|}{\delta} \right) \cdot \frac{1}{2\epsilon^2}} \\ &= 2|\mathcal{H}| e^{-\lg \left(\frac{2|\mathcal{H}|}{\delta} \right)} \\ &= 2|\mathcal{H}| \cdot \frac{\delta}{2|\mathcal{H}|} = \delta \end{aligned}$$

□

for example: $m = \lceil \lg \left(\frac{2|\mathcal{H}|}{\delta} \right) \frac{1}{2\epsilon^2} \rceil$

Bibliography

[UML] Chapter 4