

# Maths refreshing course

## HSLU, Semester 1

Matteo Frongillo

September 2, 2024

### Contents

<b>I</b>	<b>Lesson 1</b>	<b>2</b>
<b>1</b>	<b>Algebraic definitions</b>	<b>2</b>
<b>2</b>	<b>Prime numbers</b>	<b>2</b>
<b>3</b>	<b>Positive powers</b>	<b>2</b>
3.1	Property 1 . . . . .	2
3.2	Property 2 . . . . .	2
3.3	Property 3 . . . . .	2
<b>4</b>	<b>Fractions</b>	<b>3</b>
4.1	Property 1 . . . . .	3
4.2	Property 2 . . . . .	3
4.3	Property 3 . . . . .	3
<b>5</b>	<b>Negative powers</b>	<b>3</b>
5.1	Definition . . . . .	3
5.2	Property 4 . . . . .	3
5.3	Property 5 . . . . .	3
<b>6</b>	<b>Fractions and percentages (and back)</b>	<b>4</b>
<b>II</b>	<b>Lesson 2</b>	<b>5</b>

## Part I

# Lesson 1

## 1 Algebraic definitions

- $\mathbb{N} :=$  Natural numbers
- $\mathbb{Z} :=$  Integer numbers
- $\mathbb{Q} :=$  Rational numbers
- $\mathbb{R} :=$  Real numbers

We have that:

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

## 2 Prime numbers

A prime number is a natural number which can be divided only by itself or 1.

$$n \in \mathbb{N}, n \neq \{0, 1\}$$

## 3 Positive powers

Let  $a \in \mathbb{R}, n \in \mathbb{R}^*$  and  $a \in \mathbb{R}$

$$3^1 := 3$$

$$3^2 := 3 \cdot 3$$

$$3^{23} := 3 \cdot 3 \cdot \dots \cdot 3 \text{ (23 times)}$$

### 3.1 Property 1

Let  $a, b \in \mathbb{R}, n, m \in \mathbb{N}$ , then

$$a^n \cdot a^m = a^{n+m}$$

### 3.2 Property 2

Let  $a, b \in \mathbb{R}, n \in \mathbb{N}$ , then

$$(a \cdot b)^n = a^n \cdot b^n$$

Notation: The power  $a^n$ ,  $a$  is the base and  $n$  is the exponent.

### 3.3 Property 3

Let  $a \in \mathbb{R}, m, n \in \mathbb{N}^*$ , then

$$(a^n)^m = a^{n \cdot m}, \text{ which is } \neq a^{(n^m)}$$

## 4 Fractions

Notation 1:  $a \cdot b = a \times b = ab$ ;  $\frac{a}{b} = a \div b = a : b$

Notation 2:  $a$  is called numerator,  $b$  is called denominator.

Notation 3:  $\frac{a}{b}$ ,  $a, b \in \mathbb{R}$ ,  $b \neq 0$

### 4.1 Property 1

Let  $a, b, c, d \in \mathbb{R}$ ,  $a, b \neq 0$

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d}$$

### 4.2 Property 2

Let  $a, b, c, d \in \mathbb{R}$ ,  $a, b \neq 0$

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}$$

### 4.3 Property 3

Let  $a, b, c, d \in \mathbb{R}$ ,  $a, b \neq 0$

$$\frac{a}{b} \pm \frac{c}{d} = \frac{a \cdot d \pm c \cdot b}{b \cdot d}$$

## 5 Negative powers

### 5.1 Definition

$$\forall a \in \mathbb{R}, a \neq 0; \quad a^{-1} := \frac{1}{a}$$

### 5.2 Property 4

Let  $\forall n \in \mathbb{N}$ ,  $\forall a \in \mathbb{R}$

$$a^{-n} = \left(\frac{1}{a}\right)^n$$

This property implies that  $\forall z \in \mathbb{Z}$ ,  $\forall a \in \mathbb{R}$ ,  $a \neq 0$   
We can compute  $a^z$

### 5.3 Property 5

Let  $\forall a \in \mathbb{R}$ ,  $a \neq 0$ ,  $\forall n, m \in \mathbb{Z}$ , then

$$\frac{a^n}{a^m} = a^{n-m}$$

Consequences:

1. Properties 1, 2 and 3 also hold for integral exponents:

$$\forall a \in \mathbb{R}, \forall n, m \in \mathbb{Z} \Rightarrow a^n \cdot a^m = a^{n+m}$$

$$\forall b \in \mathbb{R}, (a \cdot b)^n = a^n \cdot b^n$$

$$(a^n)^m = a^{n \cdot m}$$

2.  $\forall a \in \mathbb{R}^*, a^0 = a^{1-1} = \frac{a^1}{a^1} = 1 \Rightarrow a^0 = 1$

## 6 Fractions and percentages (and back)

$$\alpha \in \mathbb{R}, n\% \text{ of } \alpha \iff \frac{n}{100} \cdot \alpha$$

Part II

Lesson 2