

# Mathematics 1A

## HSLU, Semester 1

Matteo Frongillo

October 14, 2024

## Contents

<b>I</b>	<b>Logic</b>	<b>4</b>
<b>1</b>	<b>Propositional logic</b>	<b>4</b>
1.1	Logical connectives . . . . .	4
1.1.1	Logical conjunction $\wedge$ . . . . .	4
1.1.2	Logical disjunction $\vee$ . . . . .	4
1.1.3	Logical negation $\neg$ . . . . .	4
1.1.4	Implication $\implies$ . . . . .	4
1.1.5	Inference $\Leftarrow$ . . . . .	4
1.1.6	If and only if $\Leftrightarrow$ . . . . .	5
<b>II</b>	<b>Set Theory</b>	<b>5</b>
<b>2</b>	<b>The set theory</b>	<b>5</b>
2.1	Logical symbols . . . . .	5
2.1.1	Definition . . . . .	5
2.1.2	Equal . . . . .	5
2.1.3	Belongs to . . . . .	5
2.1.4	Does not belong to . . . . .	5
2.1.5	Inclusion and contains . . . . .	5
2.1.6	For all/any . . . . .	5
2.2	Numerical sets . . . . .	6
2.2.1	Inclusion of sets . . . . .	6
<b>3</b>	<b>Union <math>\cup</math> and Intersection <math>\cap</math></b>	<b>6</b>
3.1	Universe symbol . . . . .	6
3.2	Venn diagram . . . . .	6
3.2.1	Union $A \cup B$ . . . . .	6
3.2.2	Intersection $A \cap B$ . . . . .	7
3.2.3	Complement $\bar{A}$ . . . . .	7
3.2.4	Difference between sets $\setminus$ . . . . .	8
3.2.5	Symmetrical difference $\Delta$ . . . . .	8
3.2.6	Disjoined sets (Empty sets) $\emptyset$ . . . . .	9
<b>III</b>	<b>Algebra</b>	<b>10</b>
<b>4</b>	<b>Intervals in the real line</b>	<b>10</b>
4.1	Examples . . . . .	10
4.1.1	Interval sets . . . . .	10
4.1.2	Graphical examples . . . . .	10
<b>5</b>	<b>The extended line</b>	<b>10</b>
5.1	Properties . . . . .	10

5.2	Operation in the extended line . . . . .	10
5.2.1	Additions . . . . .	11
5.2.2	Multiplications . . . . .	11
<b>6</b>	<b>Intervals including <math>\pm\infty</math></b>	<b>11</b>
6.1	Examples . . . . .	11
6.1.1	Interval sets . . . . .	11
6.1.2	Graphical examples . . . . .	12
<b>7</b>	<b>The absolute value function</b>	<b>12</b>
7.1	Graph of absolute value functions . . . . .	12
7.2	Properties . . . . .	12
7.3	Triangular inequalities . . . . .	13
<b>8</b>	<b>Concept of functions</b>	<b>13</b>
8.1	Image (Range) . . . . .	13
<b>9</b>	<b>Linear function</b>	<b>14</b>
9.1	Cartesian diagram . . . . .	14
9.2	Straight line . . . . .	14
9.3	Slope-intercept equation . . . . .	14
9.3.1	Slope . . . . .	14
9.3.2	Drawing . . . . .	14
9.4	Vertical lines . . . . .	15
<b>10</b>	<b>Equation of a line</b>	<b>15</b>
10.1	General equation in a cartesian diagram . . . . .	15
<b>11</b>	<b>Increasing and decreasing functions</b>	<b>16</b>
11.1	Increasing functions . . . . .	16
11.2	Decreasing functions . . . . .	16
<b>12</b>	<b>Inverse function</b>	<b>16</b>
12.1	Facts about inverse functions . . . . .	16
<b>13</b>	<b>Expressions and factorization</b>	<b>17</b>
13.1	Expressions, terms and factors . . . . .	17
13.1.1	Expressions . . . . .	17
13.2	Terms . . . . .	17
13.2.1	Factors . . . . .	17
13.2.2	Common factor . . . . .	17
13.3	Notable products . . . . .	17
<b>14</b>	<b>Polynomial function</b>	<b>18</b>
<b>15</b>	<b>Classification of polynomials</b>	<b>18</b>
<b>16</b>	<b>Symmetrical functions</b>	<b>18</b>
16.1	$n$ odd . . . . .	18
16.1.1	Graph examples . . . . .	18
16.2	$n$ even . . . . .	19
16.2.1	Graph examples . . . . .	19
16.3	General case . . . . .	19
16.4	Symmetry of a polynomial . . . . .	19
<b>17</b>	<b>Intersection with axis</b>	<b>20</b>
17.1	Vertical intersection . . . . .	20
17.2	Zeros of a function . . . . .	20
17.3	Graph example . . . . .	20
<b>18</b>	<b>Dominant elements in a function approaching <math>\pm\infty</math></b>	<b>21</b>
18.1	Order of dominance . . . . .	21
18.1.1	Approaching to $+\infty$ . . . . .	21

18.1.2	Approaching to $-\infty$ . . . . .	21
18.1.3	Dominance in rational functions . . . . .	21
<b>19</b>	<b>Exponential and logarithm functions</b>	<b>22</b>
19.1	Exponentials . . . . .	22
19.1.1	General equation . . . . .	22
19.1.2	Euler's number . . . . .	22
19.2	Logarithms . . . . .	22
19.2.1	Natural logarithm . . . . .	22
19.2.2	Logarithms with arbitrary bases . . . . .	22
19.2.3	Common logarithm . . . . .	22
19.3	Exponential growth . . . . .	23
<b>20</b>	<b>Composite functions</b>	<b>23</b>
20.1	Examples . . . . .	23
<b>IV</b>	<b>Trigonometry</b>	<b>24</b>
<b>21</b>	<b>Trigonometry</b>	<b>24</b>
21.1	Conversion table of degrees and radians . . . . .	24
21.2	Trigonometric functions in the unit circle . . . . .	24
21.2.1	Property 1 – Domain and range . . . . .	25
21.2.2	Property 2 – Trigonometric identity . . . . .	25
21.3	Tangent . . . . .	25
21.4	Domain of trigonometric functions . . . . .	25
21.5	Inverse trigonometric functions . . . . .	26
21.5.1	Arccosine . . . . .	26
21.5.2	Arcsine . . . . .	26
21.5.3	Arctan . . . . .	26
21.6	Harmonic oscillation . . . . .	27
<b>V</b>	<b>Calculus I</b>	<b>28</b>
<b>22</b>	<b>Limits</b>	<b>28</b>
22.1	Concept of limit of a real function . . . . .	28
22.1.1	Definition . . . . .	28
22.1.2	Graphic interpretation . . . . .	28
22.2	Limit Value at a finite point . . . . .	28
22.3	One-sided limits . . . . .	29
22.3.1	Graph example . . . . .	29
22.4	Continuity of a function . . . . .	29
22.4.1	Continuity in short . . . . .	30
<b>23</b>	<b>Derivatives</b>	<b>31</b>
23.1	Definition of derivate . . . . .	31
23.1.1	Existence of the derivative . . . . .	31
23.2	Geometric meaning of the derivative . . . . .	31
23.3	Equation of the tangent line . . . . .	32
23.4	Bernoulli – de l'Hôpital Theorem . . . . .	32
23.4.1	The 7 indeterminate forms . . . . .	32
23.4.2	Statement of the theorem . . . . .	32

## Part I

# Logic

## 1 Propositional logic

Propositional logic is a branch of mathematics that deals with propositions and logical operations.

### 1.1 Logical connectives

A	B	$\neg B$	$A \wedge B$	$A \vee B$	$A \implies B$	$A \Leftrightarrow B$
T	T	F	T	T	T	T
T	F	T	F	T	F	F
F	T	F	F	T	T	F
F	F	T	F	F	T	T

#### 1.1.1 Logical conjunction $\wedge$

Given two statements  $P$  and  $Q$ ,  $P \wedge Q$  is true if both  $P$  and  $Q$  are true.

Let  $P = (x > 0)$  and  $Q = (y > 0)$ , then:

$$P \wedge Q = (x > 0 \wedge y > 0)$$

#### 1.1.2 Logical disjunction $\vee$

Given two statements  $P$  and  $Q$ ,  $P \vee Q$  is true if at least one of  $P$  or  $Q$  is true.

Let  $P = (x = 0)$  and  $Q = (y \neq 0)$ , then:

$$P \vee Q = (x = 0 \vee y \neq 0)$$

#### 1.1.3 Logical negation $\neg$

The negation of a statement  $P$ , denoted as  $\neg P$ , is true if  $P$  is false, and false if  $P$  is true.

Let  $P = (x \geq 5)$ , then:

$$\neg P = (x < 5)$$

#### 1.1.4 Implication $\implies$

The symbol  $\implies$  indicates that if statement  $P$  is true, then statement  $Q$  must also be true (i.e.,  $P$  implies  $Q$ ).

Warning: It does not require that  $Q$  implies  $P$ .

$$P = (x = 1) \implies Q = (x \in \mathbb{N})$$

#### 1.1.5 Inference $\Leftarrow$

The symbol  $\Leftarrow$  means that a conclusion or result implies the truth of an earlier statement.

If  $Q$  is true, then  $P$  must be true.

$$Q = (x > 0) \Leftarrow P = (x \in \mathbb{R}^+)$$

### 1.1.6 If and only if $\Leftrightarrow$

The symbol  $\Leftrightarrow$  indicates that two statements  $P$  and  $Q$  are logically equivalent, meaning  $P$  is true if and only if  $Q$  is true.

$$P = (x \in \mathbb{N}, x \neq 0) \Leftrightarrow Q = (x \in \mathbb{N}^*)$$

## Part II

# Set Theory

## 2 The set theory

### 2.1 Logical symbols

#### 2.1.1 Definition

Braces and the definition symbol “:=” are used to define a set giving all its elements:

$$A := \{a, b, c, d, e\}$$

#### 2.1.2 Equal

In this case, the equal symbol means that the set  $A$  is equal to the set  $B$ :

$$A = B$$

#### 2.1.3 Belongs to

The symbols  $\in$  and  $\ni$  describe an element which is part of the set:

$$a \in A \Leftrightarrow A \ni a$$

#### 2.1.4 Does not belong to

The symbols  $\notin$  mean that an element does not belong to the set:

$$f \notin A$$

#### 2.1.5 Inclusion and contains

The symbols  $\subset$  and  $\supset$  mean that a set has another set included in its set:

$$\mathbb{N} \subset \mathbb{Z} \Leftrightarrow \mathbb{Z} \supset \mathbb{N}$$

#### 2.1.6 For all/any

The symbol  $\forall$  means that we are considering any type of element:

$$\forall x \in \mathbb{R}, x > 0$$

In this case, we've defined a new set.

## 2.2 Numerical sets

- $\mathbb{N} :=$  Natural numbers (including 0);
- $\mathbb{Z} :=$  Integer numbers;
- $\mathbb{Q} :=$  Rational numbers;
- $\mathbb{R} :=$  Real numbers  $:= \mathbb{Q} \cup \{\text{irrational numbers}\}$ .

Notation: The “\*” symbol means that the set does not include 0.

### 2.2.1 Inclusion of sets

$$\boxed{\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}}$$

$$B := \{\pi, 1, -1, 0\};$$

$$C := \{\pi, 1\};$$

$$D := \{\pi\}.$$

Then we write some examples:  $\pi \in B$ ,  $D \subset B$ ,  $C \subset B$ ,  $B \not\subset C$ ,  $0 \in B$ ,  $0 \notin C$ .

## 3 Union $\cup$ and Intersection $\cap$

### 3.1 Universe symbol


The symbol  $\bigcup :=$  Universe describes a big set which contains all sets involved in our discussions (not always).

### 3.2 Venn diagram

#### 3.2.1 Union $A \cup B$

If  $A$  and  $B$  are sets, then their union is:


$$\boxed{A \cup B = \{\forall x \in \bigcup \mid x \in A \vee x \in B\}}$$



### 3.2.2 Intersection $A \cap B$

If  $A$  and  $B$  are sets, then their intersection is:


$$A \cap B = \{\forall x \in \mathcal{U} \mid x \in A \wedge x \in B\}$$



### 3.2.3 Complement $\bar{A}$

If  $A$  is a set, its complement is:


$$\bar{A} = \{\forall x \in \mathcal{U} \mid x \notin A\}$$



### 3.2.4 Difference between sets $\setminus$

If  $A$  and  $B$  are sets, then their difference is:


$$A \setminus B = \{\forall x \in \bigcup \mid x \in A, x \notin B\}$$



### 3.2.5 Symmetrical difference $\triangle$

If  $A$  and  $B$  are sets, then their symmetrical difference is:

$$A \triangle B = (A \setminus B) \cup (B \setminus A)$$






### 3.2.6 Disjoined sets (Empty sets) $\emptyset$

$\emptyset$  := the set containing zero elements:

$$A \cap B = \emptyset$$



## Part III

# Algebra

## 4 Intervals in the real line

Intervals describe what happens between two or more elements.

### 4.1 Examples

#### 4.1.1 Interval sets


We have 4 cases:

- $(a, b) = \{\forall x \in \mathbb{R} \mid a < x < b\};$
- $[a, b) = \{\forall x \in \mathbb{R} \mid a \leq x < b\};$
- $(a, b] = \{\forall x \in \mathbb{R} \mid a < x \leq b\};$
- $[a, b] = \{\forall x \in \mathbb{R} \mid a \leq x \leq b\}.$

Notation:  $a$  and  $b$  are often called the “end points” of the interval;

#### 4.1.2 Graphical examples

$\forall x \in \mathbb{R}, x \in [a, b]$




## 5 The extended line

In the real line  $\mathbb{R}$  we add  $\pm\infty$ .

**Real line:**  $(-\infty, +\infty) = \mathbb{R}$

**Extended real line:**  $[-\infty, +\infty] = \overline{\mathbb{R}}$



Remark:  $\pm\infty \notin \mathbb{R}$

### 5.1 Properties

$$\boxed{\forall x \in \mathbb{R} \mid \infty > x \mid -\infty < 0}$$

### 5.2 Operation in the extended line

If  $a, b \in \mathbb{R}$ , then  $a + b$ ,  $a - b$ ,  $a \cdot b$ ,  $\frac{a}{b}$  (with  $b \neq 0$ ) stay the same

### 5.2.1 Additions

Let  $\forall a \in \mathbb{R}$ :

- $a + \infty := \infty$ ;
- $a - \infty := -\infty$ ;
- $+\infty + \infty := +\infty$ ;
- $-\infty - \infty := -\infty$ ;
- $+\infty - \infty := \text{undefined}$ .

### 5.2.2 Multiplications

Let  $\forall a \in \mathbb{R}$ :

- $+\infty \cdot +\infty := +\infty$ ;
- $-\infty \cdot +\infty := -\infty$ ;
- $-\infty \cdot (-\infty) := \infty$ ;
- $a \cdot \infty := \begin{cases} a > 0 & +\infty \\ a < 0 & -\infty \\ a = 0 & \text{undefined} \end{cases}$
- $a \cdot (-\infty) := \begin{cases} a > 0 & -\infty \\ a < 0 & +\infty \\ a = 0 & \text{undefined} \end{cases}$
- $\frac{a}{+\infty} = \frac{a}{-\infty} := 0$ ;
- $\frac{+\infty}{a} := \begin{cases} a > 0 & +\infty \\ a < 0 & -\infty \\ a = 0 & +\infty \end{cases}$
- $\frac{-\infty}{a} := \begin{cases} a > 0 & -\infty \\ a < 0 & +\infty \\ a = 0 & -\infty \end{cases}$
- $\frac{\infty}{\infty} := \text{undefined}$ .

## 6 Intervals including $\pm\infty$

Intervals describe what happens between two or more elements, including  $\pm\infty$ .

### 6.1 Examples


#### 6.1.1 Interval sets

Let  $a \in \mathbb{R}$ , then:

- $(-\infty, a) = \{\forall x \in \mathbb{R} \mid x < a\}$ ;
- $(a, +\infty) = \{\forall x \in \mathbb{R} \mid x > a\}$ ;
- $(-\infty, a] = \{\forall x \in \mathbb{R} \mid x \leq a\}$ ;
- $[a, +\infty] = \{\forall x \in \mathbb{R} \mid x \geq a\}$ ;
- $(-\infty, +\infty) = \mathbb{R}$ ;
- $[-\infty, +\infty] = \overline{\mathbb{R}}$ .

### 6.1.2 Graphical examples

$\forall x \in \mathbb{R}, x \in [a, b] \cup ]c, +\infty[$



Notation: The union of two or more intervals where  $x \in \mathbb{R}$  is denoted by the symbol  $\cup$ .

## 7 The absolute value function


The absolute value is an operator that returns the positive value of a number, regardless of its original sign.

Let  $x \in \mathbb{R}$ , then:

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } -x < 0 \end{cases}$$

### 7.1 Graph of absolute value functions

Let's plot the function  $y = |x|$ :



### 7.2 Properties

Let  $a, b \in \mathbb{R}$ , then:

- $|a \cdot b| = |a| \cdot |b|$ ;
- $\left| \frac{a}{b} \right| = \frac{|a|}{|b|}$  for  $b \neq 0$ ;
- $|a \pm b| \neq |a| \pm |b|$ .

### 7.3 Triangular inequalities

Let  $a, b \in \mathbb{R}$ , then:


$$\begin{array}{l} |a| + |b| \geq |a+b| \\ |a| - |b| \leq |a-b| \end{array}$$

## 8 Concept of functions

Let's take any two sets  $A \{a, b, c, d, e, f, g\}$  and  $B \{a_1, b_1, c_1, d_1, e_1, f_1, g_1\}$ .

$$\begin{array}{l} f : A \Rightarrow B \\ a \mapsto f(a) \end{array}$$

A function is a relation between the sets  $A$  and  $B$ , according to which we associate to each element of  $A$  one and only one element of  $B$ :



Notation:  $f(a) = b_1, f(b) = a_1, f(c) = c_1, f(d) = d_1, \dots$

Each point in set  $A$  is associated with one element of  $B$ . However, it is possible for more than two elements of  $A$  to point to the same element of  $B$ .

The set  $A$  is called *domain* of  $f$ . The set  $B$  is called the *codomain* of  $f$ .

### 8.1 Image (Range)


Let  $f : X \Rightarrow Y$  be a function. The image of  $f$  is defined as:

$$\text{Im}(f) = \{y \in Y \mid y = f(x), x \in X\}$$

Easily, the image is the set containing all the elements of the set  $B$  associated with the elements of the set  $A$ .

## 9 Linear function

### 9.1 Cartesian diagram



### 9.2 Straight line

Let A and B be any two distinct points, then there is one and only one line passing through A and B.

### 9.3 Slope-intercept equation

Let  $m, q \in \mathbb{R}$ , then

$$y = mx + q$$

- $m$ : slope;
- $q$ : vertical intercept.

#### 9.3.1 Slope

The slope of a line can be calculated with the equation


$$m = \frac{y_B - y_A}{x_B - x_A} = \frac{\Delta y}{\Delta x} = \tan(\theta)$$

We have three different slope outcomes:

- $m > 0$ , the line is increasing;
- $m = 0$ , the line is stable;
- $m < 0$ , the line is decreasing.

Warning: This works only if  $x_B \neq x_A$ .

#### 9.3.2 Drawing



## 9.4 Vertical lines

The more the value of  $m$  increases, the closer the line will get to the vertical, without ever reaching it.

Let  $c \in \mathbb{R}$ , then  $x = c$ .

Vertical lines cannot be written as a function.

## 10 Equation of a line


Let  $m, x_A, y_A \in \mathbb{R}$  and  $A(x_A, y_A)$ , then

$$y - y_A = m(x - x_A)$$

e.g.: Find the line with  $m = -1$  and  $A(2, -1)$ .

$$y - 1 = -1(x + 2) \implies y = -x + 1$$

Points:  $A(2, -1)$ ;  $B(0, 1)$



### 10.1 General equation in a cartesian diagram

$$ax + by + c = 0$$

Remark:

- All the lines can be described with this kind of equation;
- When  $b = 0$ ,  $a \neq 0$ , then  $ax = -c \implies x = \frac{-c}{a} \in \mathbb{R}$ ;
- When  $b \neq 0$ , then  $y = -\frac{a}{b}x - \frac{c}{b}$ , where  $m = -\frac{a}{b}$  and  $q = -\frac{c}{b}$ .

## 11 Increasing and decreasing functions

Let  $f : [a, b] \rightarrow \mathbb{R}$

Notation: if you replace  $[a, b]$  with  $\mathbb{R}$ , you obtain the definition in the whole  $\mathbb{R}$ .

### 11.1 Increasing functions

- $f$  is increasing if  $\forall x_1, x_2 \in [a, b] \mid x_2 > x_1$ , then  $f(x_2) \geq f(x_1)$ ;
- $f$  is strictly increasing if  $\forall x_1, x_2 \in [a, b] \mid x_2 > x_1$ , then  $f(x_2) > f(x_1)$ .

### 11.2 Decreasing functions

- $f$  is decreasing if  $\forall x_1, x_2 \in [a, b] \mid x_2 > x_1$ , then  $f(x_2) \leq f(x_1)$ ;
- $f$  is strictly decreasing if  $\forall x_1, x_2 \in [a, b] \mid x_2 > x_1$ , then  $f(x_2) < f(x_1)$ .

## 12 Inverse function

Let's take any two sets  $A$  and  $B$ .

A function  $f : A \Rightarrow B$  is invertible if there exists another function  $f^{-1} : B \Rightarrow A$ , called the inverse function, such that:

$$\boxed{\begin{array}{l} \forall x \in A, f^{-1}(f(x)) = x \\ \forall y \in B, f(f^{-1}(y)) = y \end{array}}$$

Warning: A function is invertible if and only if it is bijective.

### 12.1 Facts about inverse functions

1)

Let  $f : D \Rightarrow \mathbb{R}$

$f$  is invertible in  $D$  when:

- $f$  is strictly increasing;
- $f$  is strictly decreasing.

2)

Let  $f : D \Rightarrow \mathbb{R}$

$f$  is invertible when  $f^{-1} : \text{Im}(f) \Rightarrow D$ .



## 13 Expressions and factorization

### 13.1 Expressions, terms and factors

#### 13.1.1 Expressions

An expression is any formula containing numbers, variables, operations, and brackets.

$$y = ax^2 + bx \cdot c$$

#### 13.2 Terms

A term is any part of the expression separated by “+” or “-”.

$$y = \underbrace{ax^2}_{\text{term}} + \underbrace{bx \cdot c}_{\text{term}}$$

##### 13.2.1 Factors

Each term can be split into a product of factors.

$$x \cdot y \cdot (a - b) \cdot 24 = x \cdot y \cdot (a - b) \cdot 2 \cdot 2 \cdot 2 \cdot 3$$

Notice: the process of splitting a term into several factors is called “factorization”.

The goal of a factorization is to factorize an expression as much as possible.

##### 13.2.2 Common factor

Any expression made of terms is composed of several factors.

$$x^2 + x^3 + x = x(x + x^2 + 1), \forall x \in \mathbb{R}$$

### 13.3 Notable products

- $(a + b)^2 = a^2 + 2ab + b^2$  (square of a binomial);
- $(a - b)^2 = a^2 - 2ab + b^2$  (square of a binomial);
- $(a - b)(a + b) = a^2 - b^2$  (difference of squares);
- $(a + b)(a^2 - ab + b^2) = a^3 + b^3$  (sum of cubes);
- $(a - b)(a^2 + ab + b^2) = a^3 - b^3$  (difference of cubes).

Remark: notable products are useful to factorize expressions when we don’t know a common factor.

## 14 Polynomial function

Let  $n \in \mathbb{N}^*$ , then a polynomial is the sum or difference of n-monomials.

## 15 Classification of polynomials

Polynomials can be classified using two criteria:

1. the number of **terms**;
2. the **degree** of the polynomial.

Number of Terms	Name	Example	Degree
One	Monomial	$ax^2$	1
Two	Binomial	$ax^2 - bx$	2
Three	Trinomial	$ax^2 - bx + c$	3
Four or more	Polynomial	$a_nx^n - a_1x^{n-1} + a_2x^{n-2} \dots a_0$	n-degree

Remark: The degree of a polynomial is the largest exponent of its monomials.



## 16 Symmetrical functions

Let  $y = kx^n$ , then we plot:

### 16.1 $n$ odd

$$f(-x) = -f(x), \quad \forall x \in \mathbb{R}$$


#### 16.1.1 Graph examples




## 16.2 $n$ even

$$f(-x) = f(x), \quad \forall x \in \mathbb{R}$$

### 16.2.1 Graph examples



$k > 0$   
Concave up



$k < 0$   
Concave down

Definition:

- a function  $y = f(x)$  is called **odd** if it is symmetric with respect to the origin;
- a function  $y = f(x)$  is called **even** if it is symmetric with respect to the y-axis.

## 16.3 General case

Let  $y = p(x)$ , where  $p(x)$  is any polynomial with real coefficients:

$$p(x) = a_n \cdot x^n + a_{n-1} \cdot x^{n-1} + a_{n-2} \cdot x^{n-2} + \dots + a_2 \cdot x^2 + a_1 \cdot x^1 + a_0$$

where:

- $n \in \mathbb{N}$ ;
- $n = \deg(p(x))$ ;
- $a_n =$  leading coefficient.

$$p(x) = \sum_{i=0}^n a_i \cdot x^i$$

## 16.4 Symmetry of a polynomial

Let  $y = p(x)$  be a polynomial function, then:

1)

$y = p(x)$  is odd iff all the degrees of all the terms of  $p(x)$  are odd;

2)

$y = p(x)$  is even iff all the degrees of all the terms of  $p(x)$  are even;

3)

$y = p(x)$  has mixed degrees,  $p(x)$  is neither odd nor even.

## 17 Intersection with axis

### 17.1 Vertical intersection

Let  $y = f(x)$  be any function, then we solve for  $y$ :


$$\begin{cases} x = 0 \\ y = f(0) \end{cases}$$

### 17.2 Zeros of a function

Let  $y = f(x)$  be any function, then we solve for  $x$ :

$$\begin{cases} y = 0 \\ 0 = f(x) \end{cases}$$

### 17.3 Graph example



## 18 Dominant elements in a function approaching $\pm\infty$

As  $x$  approaches  $\pm\infty$ , the term with the highest degree in a polynomial function dominates the behavior of the function.

$$p(x) \text{ has, as a dominant, the element } a_n \text{ with the highest degree } x^n$$

### 18.1 Order of dominance

#### 18.1.1 Approaching to $+\infty$

Let  $n \in \mathbb{N}$ ,  $m \in \mathbb{N}$ ,  $2 < n < m$ , then:

$$\ln(x) < x < x^n < x^m < n^x < m^x < x^x$$

In these cases, we always have  $x \Rightarrow +\infty \Rightarrow p(x) \Rightarrow +\infty$

#### 18.1.2 Approaching to $-\infty$

Let  $\lambda > 2$  and odd,  $k > 2$  and even.

$$\begin{array}{l} x^\lambda < -x^2 < x^1 < 0 \\ -x^k < -x^2 < x^1 < 0 \end{array}$$

Functions like  $x^\lambda$  (with  $\lambda$  odd) and  $-x^k$  (with  $k$  even) both approach  $-\infty$ , but at different rates.

#### 18.1.3 Dominance in rational functions

When the dominant element is at the numerator:

$$\lim_{x \rightarrow \infty} \frac{x^n}{x^{n-1}} = \infty$$

When the dominant element is at the denominator:

$$\lim_{x \rightarrow \infty} \frac{x^{n-1}}{x^n} = 0$$

When we have the same degree either in the numerator and in the denominator:

$$\lim_{x \rightarrow \infty} \frac{ax^n}{bx^n} = \frac{a}{b}$$

Definition: **horizontal asymptote** appears when  $x$  approaches to  $\infty$ , which implies that  $y$  approaches to a number  $A$  different from  $\pm\infty$

## 19 Exponential and logarithm functions

The relationship between exponentials and logarithms is based on the following formula:

$$a^{\log_a(x)} = x \iff \log_a(a^x) = x$$

### 19.1 Exponentials

#### 19.1.1 General equation

Let  $\alpha \in \mathbb{R}_+^*$ ,  $x \in \mathbb{R}$ , and  $a > 1$ , then:

$$y = \alpha \cdot a^x$$

#### 19.1.2 Euler's number

Euler's number is defined by the limit:

$$e = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.718 \dots$$

Alternatively, it can be expressed as:

$$e = \sum_{n=0}^{\infty} \frac{1}{n!}$$

### 19.2 Logarithms

#### 19.2.1 Natural logarithm

The inverse function of the Euler's exponential function:

$$f(x) = e^x \iff h(x) = \ln(x)$$

Remark: the domain of  $\ln(x)$  is  $D_n : \forall x \in \mathbb{R}_+^*$

#### 19.2.2 Logarithms with arbitrary bases

The inverse function of any arbitrary exponential function:

$$f(x) = n^x \iff h(x) = \log_n(x)$$

Alternatively, it can be expressed as:

$$\log_a(x) = \frac{\ln(x)}{\ln(a)}$$

#### 19.2.3 Common logarithm

The common logarithm uses base 10:

$$\log_{10}(x) = \frac{\ln(x)}{\ln(10)}$$

### 19.3 Exponential growth

$$N(t) = N_0 \cdot e^{kt}$$

## 20 Composite functions

Let  $y = f(x)$  and  $z = g(y)$  be two functions, then:

$$z = g(f(x))$$

### 20.1 Examples

1)

Let  $f(x) = x^2 + 4x$  and  $g(y) = y^2 + \cos(y)$  be two functions, then:

$$g(f(x)) = (x^2 + 4x)^2 + \cos(x^2 + 4x)$$

2)

Let  $f(x) = x^3$ ,  $h(x) = \arctan(x)$  and  $g(x) = \ln(x)$  be functions, then:

$$g(h(f(x))) = \ln(\arctan(x^3))$$

## Part IV

# Trigonometry

## 21 Trigonometry

### 21.1 Conversion table of degrees and radians

Angles (in Degrees)	0°	30°	45°	60°	90°	180°	270°	360°
Angles (in Radians)	0 <sup>c</sup>	$\pi/6^c$	$\pi/4^c$	$\pi/3^c$	$\pi/2^c$	$\pi^c$	$3\pi/2^c$	$2\pi^c$
$\sin(\theta)$	0	1/2	$\sqrt{2}/2$	$\sqrt{3}/2$	1	0	-1	0
$\cos(\theta)$	1	$\sqrt{3}/2$	$\sqrt{2}/2$	1/2	0	-1	0	1
$\tan(\theta)$	0	$\sqrt{3}/3$	1	$\sqrt{3}$	$\infty$	0	$\infty$	0


Remark:

$$\cos(2\pi + \theta) = \cos(\theta) \quad | \quad \sin(2\pi + \theta) = \sin(\theta)$$

Remark: Let  $\forall k \in \mathbb{Z}$ ,  $\forall \theta \in \mathbb{R}$ , then:

$\cos(\theta + 2\pi k) = \cos(\theta)$

### 21.2 Trigonometric functions in the unit circle



Remark: the circle has center in the origin  $O$ , radius = 1 and function  $x^2 + y^2 = 1$



Trigonometric functions can be extended to angles beyond  $0$  and  $90^\circ$  using the unit circle. For an angle  $\theta$  in the unit circle:

$$\sin \theta := y \quad | \quad \cos \theta := x \quad | \quad \tan \theta := \frac{y}{x}$$

### 21.2.1 Property 1 – Domain and range

Because we are inside a circle of radius 1:

- $-1 \leq \cos(\theta) \leq 1$ ;
- $-1 \leq \sin(\theta) \leq 1$ .

### 21.2.2 Property 2 – Trigonometric identity

Because we have a  $90^\circ$  angle, we can use Pythagoras:

$$\overrightarrow{OH}^2 + \overrightarrow{PH}^2 = \overrightarrow{OP}^2$$

Let  $\forall \theta \in \mathbb{R}$ , then we can compute the following trigonometric identity:

$$\sin^2(\theta) + \cos^2(\theta) = 1$$

## 21.3 Tangent

A tangent of an angle is exactly the slope of a line:

$$m = \frac{\Delta y}{\Delta x} = \tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$$

Remark: the tangent is not defined when the angle is  $\frac{\pi}{2}$  or  $\frac{3\pi}{2}$ , that is when we have a vertical line.


## 21.4 Domain of trigonometric functions

$$\begin{aligned} y &= \cos(x), & x &\in \mathbb{R} \\ y &= \sin(x), & x &\in \mathbb{R} \\ y &= \tan(x), & x &\in \mathbb{R} \setminus \left\{ \frac{\pi}{2} + k\pi \mid k \in \mathbb{Z} \right\} \end{aligned}$$

## 21.5 Inverse trigonometric functions


Warning: in order to be invertible, a function should be either always strictly increasing or always strictly decreasing.

### 21.5.1 Arccosine




### 21.5.2 Arcsine

The domain of the arcsine is  $\forall x \in [-1, 1]$  and the range is  $\forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$



### 21.5.3 Arctan


The domain is  $\forall x \in \mathbb{R}$  and the range is  $\forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$



## 21.6 Harmonic oscillation

Let  $A, B > 0$ , then the function is oscillating harmonically with  $t$  around  $D$ :

$$y = D + A \cdot \sin(Bt + \varphi)$$



## Part V

# Calculus I

## 22 Limits


### 22.1 Concept of limit of a real function

#### 22.1.1 Definition

Let  $f : \mathcal{D} \rightarrow \mathbb{R}$  be a function and  $c$  a point, the limit  $L = \lim_{x \rightarrow c} f(x)$  with  $x$  tending to  $c$  exists only if in a given  $\epsilon > 0$  arbitrarily small, exists another  $\delta > 0$  such that:

$$0 < |x - c| < \delta \implies |f(x) - L| < \epsilon$$

#### 22.1.2 Graphic interpretation



### 22.2 Limit Value at a finite point

The notion of the “limit of  $f(x)$  as  $x$  approaches a (finite) point  $a \in \mathbb{R}$ ” is only meaningful if the point  $a$  can be approximated by points from the domain of definition of  $f$ . We can precisely formulate this concept with the notion of an “accumulation point”.

#### Definition

Given a set  $A \subset \mathbb{R}$  and a real number  $a \in \mathbb{R}$ , the real number  $a$  is called an *accumulation point* of the set  $A$  if every open interval of the form  $(a - \delta, a + \delta)$  with  $\delta > 0$  contains infinitely many points of  $A$ .

In the above definition, it is not required that  $a$  lies in  $A$ . Often, we will consider functions whose domains are unions of intervals of the form:

$$(b, a) \cup (a, c)$$

For example, consider the function defined by  $f(x) = \frac{1}{x}$ , defined on  $(-\infty, 0) \cup (0, \infty)$ . The point 0 is an accumulation point of the domain of definition of  $\frac{1}{x}$ .

#### Definition

Given a real function  $f$ , an accumulation point  $x_0$  of  $\mathcal{D}_f$ , and  $L \in \mathbb{R} = \mathbb{R} \cup \{\pm\infty\}$ , we say that the function  $f$  has the limit  $L$  as  $x \rightarrow x_0$  if  $f(x)$  gets arbitrarily close to  $L$ , provided  $x$  is sufficiently close to (but never equal to)  $x_0$ .

## 22.3 One-sided limits

Often, one considers limits where  $x$  approaches  $x_0$  from only one direction, either from the right or from the left. In these cases, we refer to a right-sided or left-sided limit and use the following notations:

$$\lim_{x \rightarrow x_0^+} f(x) \quad \text{or} \quad \lim_{x \rightarrow x_0, x > x_0} f(x) \quad \text{or} \quad \lim_{x \rightarrow x_0} f(x)$$


for a right-sided limit, and:

$$\lim_{x \rightarrow x_0^-} f(x) \quad \text{or} \quad \lim_{x \rightarrow x_0, x < x_0} f(x) \quad \text{or} \quad \lim_{x \rightarrow x_0} f(x)$$

for a left-sided limit.

$\lim_{x \rightarrow a}$  can indicate a limit as  $x$  approaches an arbitrary point (e.g.,  $a = x_0$  for  $x_0 \in \mathbb{R}$ ), as well as a one-sided limit ( $a = x_0^+$  or  $a = x_0^-$  for  $x_0 \in \mathbb{R}$ ), or a limit at infinity ( $a = \pm\infty$ ).

### 22.3.1 Graph example



## 22.4 Continuity of a function


**Definition** *Continuity of a real function*

Given a real function  $f : D \rightarrow C$ , the function is continuous at the point  $x = c$  where  $c \in D$  if:

$$\lim_{x \rightarrow c} f(x) = f(c)$$

and therefore, if the limit exists and is equal to the value of the function at that point.

In other words, the function is continuous at the point if the limit exists (from both the left and right, coinciding) and the value of the function at that point is equivalent to the value of the limit.



#### 22.4.1 Continuity in short

- A function is said to be continuous at a point if the limit at that point exists and is equal to the value of the function at that point;
- A function is said to be continuous on a subinterval of the domain if it is continuous at all points in that subinterval;
- A function is said to be continuous if it is continuous at all points in its interval.

## 23 Derivatives

### 23.1 Definition of derivate

The derivate of a real function  $f(x)$  is defined as:

$$f'(x) = \lim_{\Delta x \rightarrow 0^{\pm}} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

if the limit exists.

Definition:  $f'(a)$ , if it exists, is calld derivative of  $f(x)$  at  $x = a$ .


It corresponds to the slope of the tangent line at  $x = a$  to the function  $y = f(x)$

#### 23.1.1 Existence of the derivative

The derivative exists if and only if:

$$\lim_{\Delta x \rightarrow 0^+} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0^-} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

### 23.2 Geometric meaning of the derivative





The secant of a function  $f(x)$  between a point  $A$  and  $B$  is given by:

$$\frac{\Delta y}{\Delta x} = \frac{f(B) - f(A)}{B - A}$$

The closer we bring  $A$  and  $B$ , the smaller  $\Delta x$  becomes. As  $\Delta x$  decreases, the slope of the secant becomes more representative of the rate of change of  $f$  in the interval  $[A; B]$ .

When the  $\Delta x$  of the slope becomes infinitesimally small, we obtain the exact slope at a point (instantaneous). This slope is represented by the tangent line:

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$



The derivative of a function  $f(x)$  is therefore another function,  $f'(x)$ , which represents the rate of change of  $f(x)$  at every point. In other words,  $f'(x)$  represents the slope of the tangent at each  $x$  of  $f(x)$ . This is precisely represented by the definition of the derivative, which is the slope  $\frac{\Delta y}{\Delta x}$  calculated with the limit of  $\Delta x \rightarrow 0$ .

### 23.3 Equation of the tangent line

When  $f'(x)$  is defined,  $p = (x, f(x))$ , and  $p \in$  tangent line, then:

$$y - f(x) = f'(x) \cdot (x - a)$$

### 23.4 Bernoulli – de l'Hôpital Theorem

Bernoulli – de l'Hôpital theorem is applicable only if the function results in an indeterminate form.

#### 23.4.1 The 7 indeterminate forms

The seven indeterminate forms are

$$\frac{0}{0}, \quad \frac{\infty}{\infty}, \quad 0 \cdot \infty, \quad \infty - \infty, \quad 0^0, \quad \infty^0, \quad 1^\infty$$

#### 23.4.2 Statement of the theorem

Let us consider two real functions  $f(x)$  and  $g(x)$  that are differentiable in a neighborhood of  $x_0 \in \mathbb{R}$  (not necessarily at  $x_0$ ). Without loss of generality, if we have a limit:

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)}$$

where  $\frac{f(x_0)}{g(x_0)}$  results in an indeterminate form, then:

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow x_0} \frac{f'(x)}{g'(x)}$$

if the limit exists.