

Maths refreshing course

HSLU, Semester 1

Matteo Frongillo

September 4, 2024

Contents

I	Lesson 1	3
1	Numerical sets	3
2	Prime numbers	3
3	Positive powers	3
3.1	Property 1	3
3.2	Property 2	3
3.3	Property 3	3
4	Fractions	4
4.1	Property 1	4
4.2	Property 2	4
4.3	Property 3	4
5	Negative powers	4
5.1	Definition	4
5.2	Property 4	4
5.3	Property 5	4
6	Fractions and percentages (and back)	5
II	Lesson 2	6
7	Symbols	6
8	Brackets	6
9	Latin notations	6
10	The real line	6
10.1	Exercises	6
11	Properties of real numbers	7
11.1	Property 1 - Closure under “+” and “.”	7
11.2	Property 2 - Commutativity	7
11.3	Property 3 - Associative	7
11.4	Property 4 - Distributive	7
11.5	Property 5 - Identity	7
11.6	Property 6 - Inverses and opposites	7
12	The order of operations	7
13	Signed numbers	8

14 Absolute value	8
14.1 Property	8
15 Lesson 3	9

Part I

Lesson 1

1 Numerical sets

- $\mathbb{N} :=$ Natural numbers (including 0)
- $\mathbb{Z} :=$ Integer numbers
- $\mathbb{Q} :=$ Rational numbers
- $\mathbb{R} :=$ Real numbers

Notation: The “*” symbol means that the set does not include 0.

We have that:

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

2 Prime numbers

A prime number is a number $n \in \mathbb{N} \setminus \{0, 1\}$ such that, for every divisor $d \in \mathbb{N}$, if $d \mid n$, then $d = 1$ or $d = n$.

$$n \in \mathbb{N} \setminus \{0, 1\} \text{ is prime} \iff \forall d \in \mathbb{N}, (d \mid n) \Rightarrow (d = 1 \text{ or } d = n)$$

3 Positive powers

Let $a \in \mathbb{R}, n \in \mathbb{N}^*$ and $a \in \mathbb{R}$, then

$$a^1 := a \quad | \quad a^n = \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

3.1 Property 1

Let $a, b \in \mathbb{R}, n, m \in \mathbb{N}$, then

$$a^n \cdot a^m = a^{n+m}$$

3.2 Property 2

Let $a, b \in \mathbb{R}, n \in \mathbb{N}$, then

$$(a \cdot b)^n = a^n \cdot b^n$$

Notation: The power a^n , a is the base and n is the exponent.

3.3 Property 3

Let $a \in \mathbb{R}, m, n \in \mathbb{N}^*$, then

$$(a^n)^m = a^{n \cdot m}, \text{ which is } \neq a^{(n^m)}$$

4 Fractions

Notation 1: $a \cdot b = a \times b = ab$ | $\frac{a}{b} = a \div b = a : b$

Notation 2: “ a ” is called numerator, “ b ” is called denominator.

Notation 3: $\frac{a}{b}$, $a, b \in \mathbb{R}$, $b \neq 0$

4.1 Property 1

Let $a, b \in \mathbb{R}^*$ and $c, d \in \mathbb{R}$, then

$$\boxed{\frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d}}$$

4.2 Property 2

Let $a, b \in \mathbb{R}^*$ and $c, d \in \mathbb{R}$, then

$$\boxed{\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}}$$

4.3 Property 3

Let $a, b \in \mathbb{R}^*$ and $c, d \in \mathbb{R}$, then

$$\boxed{\frac{a}{b} \pm \frac{c}{d} = \frac{a \cdot d \pm c \cdot b}{b \cdot d}}$$

5 Negative powers

5.1 Definition

$$\boxed{\forall a \in \mathbb{R}^*; \quad a^{-1} := \frac{1}{a}}$$

5.2 Property 4

Let $\forall n \in \mathbb{N}$, $\forall a \in \mathbb{R}$, then

$$\boxed{a^{-n} = \left(\frac{1}{a}\right)^n}$$

This property implies that $\forall z \in \mathbb{Z}$, $\forall a \in \mathbb{R}$, $z \neq 0$
We can compute a^z

5.3 Property 5

Let $\forall a \in \mathbb{R}$, $a \neq 0$, $\forall n, m \in \mathbb{Z}$, then

$$\boxed{\frac{a^n}{a^m} = a^{n-m}}$$

Consequences:

1. Properties 1, 2 and 3 also hold for integer exponents:

- $\forall a \in \mathbb{R}, \forall n, m \in \mathbb{Z} \Rightarrow a^n \cdot a^m = a^{n+m}$
- $\forall b \in \mathbb{R}, (a \cdot b)^n = a^n \cdot b^n$
- $(a^n)^m = a^{n \cdot m}$

2. $\forall a \in \mathbb{R}^*, a^0 = a^{1-1} = \frac{a^1}{a^1} = 1 \Rightarrow a^0 = 1$

6 Fractions and percentages (and back)

$\alpha \in \mathbb{R}, n\% \text{ of } \alpha \iff \frac{n}{100} \cdot \alpha$

Part II

Lesson 2

7 Symbols

Let $a, b \in \mathbb{R}$, then

- $a = b \rightarrow$ equality;
- $a \neq b \rightarrow$ inequality (a is not equal to b);
- $a < b \rightarrow$ less than (a is strictly less than b);
- $a \leq b \rightarrow$ less than or equal to (a is less than or equal to b);
- $a > b \rightarrow$ greater than (a is strictly greater than b);
- $a \geq b \rightarrow$ greater than or equal to (a is greater than or equal to b).

Example: $x \in \mathbb{R}$, $x \geq 2 \rightarrow 2 \leq x < \infty$


8 Brackets

- () Parenthesis (round brackets)
- [] Square brackets
- { } Braces

9 Latin notations


- e.g. = for example;
- i.e. = that is / that implies;
- Q.E.D. (\square)= quod erat demonstrandum (we finally prove it).

10 The real line




10.1 Exercises

1) $\forall a, b, x \in \mathbb{R}$, $a \leq x \leq b$



2) $\forall x \in \mathbb{R}$, $x \in]-2, -1] \cup]\frac{3}{2}, +\infty[$



Notation: The union of two or more intervals where $x \in \mathbb{R}$ is denoted by the symbol \cup .

11 Properties of real numbers

11.1 Property 1 - Closure under “+” and “.”

$$\forall x, y \in \mathbb{R}$$

$$x + y \in \mathbb{R}$$

$$x \cdot y \in \mathbb{R}$$

Remark: for $\forall x \in \mathbb{Z}$, closure does not hold for division.

11.2 Property 2 - Commutativity

$$\forall x, y \in \mathbb{R}$$

$$x + y = y + x$$

$$x \cdot y = y \cdot x$$

Remark: commutativity does not hold for divisions and subtractions.

11.3 Property 3 - Associative

$$\forall x, y, z \in \mathbb{R}$$

$$x + (y + z) = (x + y) + z$$

$$x \cdot (y \cdot z) = (x \cdot y) \cdot z$$

Remark: associativity does not hold for divisions and subtractions.

11.4 Property 4 - Distributive

$$\forall x, y, z \in \mathbb{R}$$

$$x(y \pm z) = xy \pm xz$$

11.5 Property 5 - Identity

$$\forall x \in \mathbb{R}$$

a) $0 + x = x$

b) $1 \cdot x = x$

Remark: $\forall x \in \mathbb{R}$, $x \cdot 0 = 0$ is not an identity property.

11.6 Property 6 - Inverses and opposites

$$\forall x \in \mathbb{R}$$

a) $x + (-x) = 0$ (additive inverse)

b) when $x \neq 0$, $x \cdot \frac{1}{x} = 1$ (multiplicative inverse or opposite)

Remark 1: $\forall x \in \mathbb{N}$ does not exist either inverse nor opposite.

Remark 2: $\forall x \in \mathbb{Z}$ has inverses, but not opposites.

12 The order of operations

1. Perform all operations inside grouping symbols beginning with the innermost set:
() inside brackets operations;
2. Perform all exponential operations as you come to them, moving left-to-right:
 x^a ;
3. Perform all multiplications and divisions as you come to them, moving left-to-right:
“.” and “÷”;
4. Perform all additions and subtractions as you come to them, moving left-to-right:
“+” and “-”;
5. When the level of priority is the same (e.g. multiplications and divisions) solve them as you come to them.

13 Signed numbers

A number is denoted as positive if it is directly preceded by a + sign or no sign at all.

A number is denoted as negative if it is directly preceded by a - sign.

$\forall x \in \mathbb{R}$

$$-(-x) = x \qquad +(-x) = -x \qquad +(+x) = x \qquad -(+x) = -x$$

14 Absolute value

Let $x \in \mathbb{R}$, then

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

14.1 Property

$\forall x \in \mathbb{R}$

$$|x| > 0 \quad \text{if } x \neq 0$$

$$|x| = 0 \quad \text{if } x = 0$$

