

The Singular Value Decomposition

Matteo Paz, Dylan Rupel

Apr 26th, 2024

First, we need a matrix. Let $A \in \text{Mat}_n(\mathbb{C})$ be Hermetian. Therefore, we can find an *orthonormal basis of eigenvectors*. There is a small issue with this, what if we want this for a non-square matrix?

Goal: Be able to do the same thing if A is not square.

Assume that $\mathbf{w}_1, \dots, \mathbf{w}_n$ is an orthonormal basis for \mathbb{C}^n , and A not necessarily hermitian. What properties must there be of \mathbf{w}_i if $A\mathbf{w}_1, \dots, A\mathbf{w}_n$ is still an orthogonal set?

$$\langle A\mathbf{w}_i, A\mathbf{w}_j \rangle = 0$$

$$\langle A^H A\mathbf{w}_i, \mathbf{w}_j \rangle = 0$$

This would work if \mathbf{w}_i was an eigenvector for $A^H A$. Let $B = A^H A$. Notice that B is hermetian.

$$B^H = (A^H A)^H = \overline{(A^T A)}^T = (A^T \overline{A})^T = A^H A = B$$

Then we would just want an orthonormal set of eigenvectors of B , which since B is hermetian, does exist. Thus, set those original $\mathbf{w}_1, \dots, \mathbf{w}_n$ to the orthonormal set of eigenvectors for B , with eigenvalues $\lambda_1, \dots, \lambda_n$ respectively. This would imply that $A\mathbf{w}_1, \dots, A\mathbf{w}_n$ is again orthogonal. But would it be orthonormal?

$$|A\mathbf{w}_i| = \sqrt{\langle A\mathbf{w}_i, A\mathbf{w}_i \rangle}$$

$$|A\mathbf{w}_i| = \sqrt{\langle A^H A\mathbf{w}_i, \mathbf{w}_i \rangle}$$

$$|A\mathbf{w}_i| = \sqrt{\langle B\mathbf{w}_i, \mathbf{w}_i \rangle}$$

$$= \sqrt{\lambda_i \langle \mathbf{w}_i, \mathbf{w}_i \rangle}$$

$$= \sqrt{\lambda_i}$$

Thus the magnitude of each basis vector after application of A is the square root of its eigenvalue under $A^H A$. These magnitudes are the **singular values**. We denote $\sigma_i = \sqrt{\lambda_i}$.

Define $\mathbf{v}_i = \frac{A\mathbf{w}_i}{\sigma_i}$ and let $\mathbf{v}_{n+1}, \dots, \mathbf{v}_m$ be an orthonormal basis for $\text{Nul}(A^T)$, which is perpendicular to the image of A . Set

$$U = [\mathbf{w}_1 \mid \dots \mid \mathbf{w}_n]$$

$$V = [\mathbf{v}_1 | \cdots | \mathbf{v}_m]$$

Now, notice that $AU = [A\mathbf{w}_1 \cdots A\mathbf{w}_n] = [\sigma_1 \mathbf{v}_1 \cdots \sigma_n \mathbf{v}_n] = V \begin{bmatrix} \text{diag}(\sigma_1, \dots, \sigma_n) \\ \mathbf{0} \end{bmatrix}$

Let $S = \begin{bmatrix} \text{diag}(\sigma_1, \dots, \sigma_n) \\ \mathbf{0} \end{bmatrix}$. Then we have $AU = VS$.

$$A = VSU^H \tag{1}$$

Now, we have a factorization where:

1. V is an orthonormal basis for \mathbb{C}^m , where the first n columns are the orthonormal results of multiplication by $B = A^H A$.
2. S is a modified diagonal matrix of the singular values of A
3. U is the orthonormal eigenbasis generated by $B = A^H A$

This is the singular value decomposition.