





1a.


```
% q1a
% m:n integer divisor found, I1=4
% 1:1
% m:n integer divisor found, I1=5
% 1:1
% m:n integer divisor found, I1=6
% 1:1
% m:n(1:2), I1=7
% m:n integer divisor found, I1=9
% 2:1
% m:n integer divisor found, I1=10
% 2:1
% m:n integer divisor found, I1=13
% 2:1
% m:n(2:3), I1=14
% m:n integer divisor found, I1=18
% 3:1
```

$I_1=4, I_2=4$ 

$I_1=7, I_2=4$ 

$I_1=9, I_2=4$ 

$I_1=14, I_2=4$ 

$I_1=18, I_2=4$ 

```

parts = [0 0 0];
i=4;
saveDir=uigetdir;
while(true)
    pulsei=[i;4];

    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(pulsei);

    cell1spikes = extractSpikes(spiketimes,1,10);
    cell2spikes = extractSpikes(spiketimes,2,10);

    ratio = length(cell1spikes)/length(cell2spikes);
    if(mod(length(cell1spikes),length(cell2spikes))==0)
        disp(['m:n integer divisor found, I1=',num2str(i)]);
        disp([num2str(ratio),' :1']);
        if(ratio==3)
            parts(1,1)=1;
        end
    end
    if(ratio>1 && ratio<2 && ~parts(1,2))
        disp(['m:n(1:2), I1=',num2str(i)]);
        parts(1,2)=1;
    end
    if(ratio>2 && ratio<3 && ~parts(1,3))
        disp(['m:n(2:3), I1=',num2str(i)]);
        parts(1,3)=1;
    end

    saveas(1,fullfile(saveDir,
['q1_I1=',num2str(i),'_I2=4'],'.png'),'png');
    if(all(parts))
        break;
    end
    i=i+1;
end

```

1b. 1:1 firing stable when $I_1=4,5,6$

1c.

```
% >> q1c  
% 1:1 Firing: gsyn=2,I1=4,I2=4  
% 1:1 Firing: gsyn=2,I1=5,I2=4
```

```
% >> q1c  
% 1:1 Firing: gsyn=5,I1=7,I2=4  
% 1:1 Firing: gsyn=5,I1=8,I2=4
```

```
% >> q1c  
% 1:1 Firing: gsyn=8,I1=8,I2=4  
% 1:1 Firing: gsyn=8,I1=9,I2=4
```

```
% set gsyn=2,5,8
i=4;
while(true)
    pulsei=[i;4];


    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(pulsei);

    cell1spikes = extractSpikes(spiketimes,1,10);
    cell2spikes = extractSpikes(spiketimes,2,10);

    ratio = length(cell1spikes)/length(cell2spikes);
    if(ratio==1)
        disp(['1:1 Firing: gsyn=8,I1=',num2str(i),' ,I2=4' ]);
    end
    if(ratio>1)
        break;
    end
    i=i+1;
end
```

1d.

```
% >> q1d
% 1:1 Firing: gsyn=1,I1=4,I2=4
% 1:1 Firing: gsyn=2,I1=4,I2=4
% 1:1 Firing: gsyn=2,I1=5,I2=4
% 1:1 Firing: gsyn=3,I1=4,I2=4
% 1:1 Firing: gsyn=3,I1=5,I2=4
% 1:1 Firing: gsyn=3,I1=6,I2=4
% 1:1 Firing: gsyn=4,I1=5,I2=4
% 1:1 Firing: gsyn=4,I1=6,I2=4
% 1:1 Firing: gsyn=4,I1=7,I2=4
% 1:1 Firing: gsyn=5,I1=7,I2=4
% 1:1 Firing: gsyn=5,I1=8,I2=4
% 1:1 Firing: gsyn=6,I1=8,I2=4
% 1:1 Firing: gsyn=7,I1=9,I2=4
% 1:1 Firing: gsyn=8,I1=8,I2=4
% 1:1 Firing: gsyn=8,I1=9,I2=4
```



```
data=[0 4]; %stated in qld
for gsyn=1:8
    i=4;
    while(true)
        pulsei=[i;4];
        %temporarily modified to accept gsyn

[t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(pulsei,gsyn);


        cell1spikes = extractSpikes(spiketimes,1,10);
        cell2spikes = extractSpikes(spiketimes,2,10);


        ratio = length(cell1spikes)/length(cell2spikes);
        if(ratio==1)
            disp(['1:1 Firing:
gsyn=',num2str(gsyn),',I1=',num2str(i),',I2=4']);
            data = [data;[gsyn i]];
        end
        if(ratio>1)
            break;
        end
        i=i+1;
    end
end
figure;
plot(data(:,2),data(:,1),'o');
ylabel('gsyn values');
xlabel('I1 values');
title('qld');
```


2a.

```
% >> q2a
% anti-synchrony @ taus:1
% anti-synchrony @ taus:2
% anti-synchrony @ taus:3
% anti-synchrony @ taus:4
% anti-synchrony @ taus:6
% anti-synchrony @ taus:10
% anti-synchrony @ taus:13
% anti-synchrony @ taus:15
% anti-synchrony @ taus:17
% anti-synchrony @ taus:19
% anti-synchrony @ taus:21
% anti-synchrony @ taus:23
% anti-synchrony @ taus:24
% anti-synchrony @ taus:26
% anti-synchrony @ taus:28
% anti-synchrony @ taus:30
% anti-synchrony @ taus:31
% anti-synchrony @ taus:33
% anti-synchrony @ taus:34
% anti-synchrony @ taus:37
% anti-synchrony @ taus:38
% anti-synchrony @ taus:41
% anti-synchrony @ taus:42
% synchrony @ taus:45
% synchrony @ taus:46
% synchrony @ taus:47
% synchrony @ taus:48
% synchrony @ taus:49
% synchrony @ taus:50
% suppression @ taus:53
% suppression @ taus:54
% suppression @ taus:55
% suppression @ taus:56
% suppression @ taus:57
% suppression @ taus:58
% suppression @ taus:59
% suppression @ taus:60
```

$\tau = 1$



$\tau = 45$ 

$\tau = 53$ 

```
saveDir = uigetdir;
for taus=1:60
    %modified to accept taus
    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(taus);


    %after 100ms from start (3rd input)
    cell1spikes = extractSpikes(spiketimes,1,100);
    cell2spikes = extractSpikes(spiketimes,2,100);


    %spikes must be within 10ms of eachother for "synchrony"
    if(length(cell1spikes)==length(cell2spikes))
        if(max(abs(cell1spikes-cell2spikes))<10)
            disp(['synchrony @ taus:',num2str(taus)]);
        end
        if(min(abs(cell1spikes-cell2spikes))>25)
            disp(['anti-synchrony @ taus:',num2str(taus)]);
        end
    end
    if(abs(length(cell1spikes)-length(cell2spikes))>10)
        disp(['suppresion @ taus:',num2str(taus)]);
    end
    saveas(1,fullfile(saveDir,
    ['q2a_taus=',num2str(taus),'.png']),'png');
end
```

2b. As τ increases, frequency decreases. Frequency is constant when $g_{\text{syn}}=0$.

```
% >> q2b
% gsyn=-8,taus=45,freq=13.615
% gsyn=-8,taus=46,freq=13.4995
% gsyn=-8,taus=47,freq=13.3759
% gsyn=-8,taus=48,freq=13.2545
% gsyn=-8,taus=49,freq=13.134
% gsyn=-8,taus=50,freq=13.0202
```

```
% >> q2b
% gsyn=0,taus=45,freq=17.9616
% gsyn=0,taus=46,freq=17.9616
% gsyn=0,taus=47,freq=17.9616
% gsyn=0,taus=48,freq=17.9616
% gsyn=0,taus=49,freq=17.9616
% gsyn=0,taus=50,freq=17.9616
```






```
%values from 2a
figure;
for taus=45:50
    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(taus);
    freq=1000/mean(isis1);
    disp(['gsyn=0,taus=',num2str(taus),',freq=',num2str(freq)]);
    plot(freq,taus,'o');
    hold on;
end
title('q2b');
xlabel('freq (Hz)');
ylabel('taus');
```

2c. As synaptic strength become less negative, frequency increases.

```
% >> q2c
% synchrony @ taus=47,gsyn=-8,freq=13.3759
% synchrony @ taus=47,gsyn=-7.9,freq=13.411
% synchrony @ taus=47,gsyn=-7.8,freq=13.4471
% synchrony @ taus=47,gsyn=-7.7,freq=13.4806
% synchrony @ taus=47,gsyn=-7.6,freq=13.517
% synchrony @ taus=47,gsyn=-7.5,freq=13.5413
```





```
figure;
for gsyn=-8:0.1:-7
    %modified to accept gsyn
    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(gsyn);


    %after 100ms from start (3rd input)
    cell1spikes = extractSpikes(spiketimes,1,100);
    cell2spikes = extractSpikes(spiketimes,2,100);
    freq=1000/mean(isis1);


    if(length(cell1spikes)==length(cell2spikes))
        if(max(abs(cell1spikes-cell2spikes))<10)
            disp(['synchrony @
taus=47,gsyn=',num2str(gsyn),',freq=',...
                num2str(freq)]);
            plot(freq,gsyn,'o');
            hold on;
        end
    end
end
title('q2c');
xlabel('freq');
ylabel('gsyn');
```

2d. No connectivity ($g_{\text{syn}}=0$) results in a constant firing frequency. In general, as τ_{aus} increases (with constant g_{syn}) firing frequency increases. With a high τ_{aus} ($=15$) and constant g_{syn} , firing frequency bounces around, centered around $\sim 18\text{Hz}$.

2b.	$\tau_{\text{aus}} \downarrow$	freq \uparrow
2c.	$g_{\text{syn}} \uparrow$	freq \uparrow
2d.	$\tau_{\text{aus}} \uparrow$	freq \uparrow







```
gsyn=3;
figure;
for taus=5:20
    %modified to accept gsyn and taus

    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(gsyn,taus)
    ;
        freq=1000/mean(isis1);
        plot(freq,taus,'o');
        hold on;
end
title('gsyn=3');
xlabel('freq (Hz)');
ylabel('taus');

taus=15;
figure;
for gsyn=1:8

    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(gsyn,taus)
    ;
        freq=1000/mean(isis1);
        plot(freq,gsyn,'o');
        hold on;
end
title('taus=15');
xlabel('freq (Hz)');
ylabel('gsyn');

gsyn=0;
figure;
for taus=5:20
    %modified to accept gsyn and taus

    [t_tot,v_tot,u_tot,spiketimes,isis1,isis2,s_tot]=ILIF2cells(gsyn,taus)
    ;
        freq=1000/mean(isis1);
        plot(freq,taus,'o');
        hold on;
end
title('gsyn=0');
xlabel('freq (Hz)');
ylabel('taus');
```