

RxConnect SIG Codes: Component Codes

Link codes together with a semi-colon. Combination and medication-specific codes are on page 2.

TABLETS & CAPSULES		INHALER / SPRAY		FREQUENCY		AS NEEDED (PRN)	
1T	Take 1 tablet	1P	Take 1 puff	QD	Every day	PRN	As needed
2T	Take 2 tablets	2P	Take 2 puffs	BID	Twice a day	AN	As needed
12T	Take 1 to 2 tablets	I1P	Inhale 1 puff	TID	3 times a day	PPA	As needed for pain
S1T	Take 1/2 to 1 tablet	I2P	Inhale 2 puffs	QID	4 times a day	PSP	As needed for severe pain
2THS	Take 2 tablets at bedtime	I3P	Inhale 3 puffs	XD	Times a day	PPF	As needed for pain and fever
C1T	Chew 1 tablet	I12P	Inhale 1 to 2 puffs	UTB	Up to 2 times daily	PMS	As needed for muscle spasm
D1T	Dissolve 1 tablet	SP1	1 spray	UTT	Up to 3 times daily	PCR	As needed for cramps
1C	Take 1 capsule	SP2	2 sprays	QOD	Every other day	PHA	As needed for headache
2C	Take 2 capsules	ROUTE POBy mouth ENEach nostril IENInto each nostril OUInto both eyes ODInto right eye OSInto left eye OAInto affected eye AUEach ear ADRight ear ASLeft ear ALLeft ear AAAffected ear VAGVaginally DIRECTIONS UDAs directed TUDTake as directed UADUse as directed AUDApply as directed DURATION F1DFor 1 day F7DFor 7 days F1WFor 1 week FXDFor 10 days UFUntil finished		TIW	Three times a week	PAN	As needed for anxiety
LIQUIDS				WK1	Once per week	PSL	As needed for sleep
1CC	Take 1 CC			Q3	Every 3 hours	PNA	As needed for nausea
15CC	Take 15 CC			Q4	Every 4 hours	PNV	As needed for nausea/vomiting
1ML	Take 1 ML			Q6	Every 6 hours	PDI	As needed for diarrhea
TVF	Take 1 ML daily			Q8	Every 8 hours	PDZ	As needed for dizziness
TV2	1 ML daily to age 2			Q12	Every 12 hours	PCG	As needed for cough
1DRD	Take 1 dropperful daily			Q46	Every 4-6 hours	PCN	As needed for congestion
SAS	Swish and swallow			Q812	Every 8 to 12 hours	PWH	As needed for wheeze
DROPS (EYE / EAR) 1DInstill 1 drop 2DInstill 2 drops I12DInstill 1 to 2 drops G1DGive 1 drop G12DGive 1 to 2 drops G4DGive 4 drops INJECTION IJInject SQSubcutaneously UNSUnits INDICATIONS BPFor blood pressure CGFor congestion CSTFor constipation RDFor rectal discomfort SOBFor shortness of breath WZFor wheezing				828	At 8AM, 2PM, and 8PM	PAL	As needed for allergy
		848	At 8AM, 4PM, and 8PM	PIT	As needed for itching		
		TOPICAL / APPLY					
		AM	In the morning	APDY	Apply daily		
		PM	In the evening	ABID	Apply twice a day		
		APM	And in the evening	ATID	Apply 3 times a day		
		HS	At bedtime	AQID	Apply 4 times a day		
		TIMING & FOOD		APHS	Apply at bedtime		
		AC	Before meals	A1P	Apply 1 patch		
		PC	After meals	A1PD	Apply 1 patch daily		
ACHS	Before meals and at bedtime	AAA	Apply to affected area				
PCHS	After meals and at bedtime	ATR	Apply to rash				
AB	Before breakfast	AAE	Apply to affected eye				
BBS	Before breakfast and supper	ATO	Apply topically				
ASU	With supper	ASP	Apply sparingly				
ES	On empty stomach	APLOUD	Apply locally as directed				
WF	With food	APV	Apply vaginally				
WM	With meals	1A	Insert 1 applicatorful				
WFM	With food and milk	1SP	Insert 1 suppository				
WW	With water						
WOJ	With orange juice						
WBL	With breakfast and lunch						
WBS	With breakfast and supper						

RxConnect SIG Codes: Combination & Medication-Specific

COMBINATION CODES (a complete sig in one entry)			
12TBID	Take 1 to 2 tablets 2 times a day	2PFQ6	2 puffs every 6 hours
12T34D	Take 1 to 2 tablets 3 or 4 times a day	IS1GTID	Instill 1 drop 3 times a day
12TQ34	Take 1 to 2 tablets every 3 to 4 hours	IS1GOUQID	Instill 1 drop in each eye 4 times a day
12TQ8	Take 1 to 2 tablets every 8 hours	1GOUBID	Put 1 drop in each eye twice a day
.5TQAM	Take 1/2 tablet every morning	1GADBID	Put 1 drop in right ear twice a day
.5TQ4PP	Take 1/2 tablet every 4 hours as needed for pain	1GAUBID	Put 1 drop in each ear twice a day
.5TABAL	Take 1/2 tablet before breakfast and lunch	APTIDAA	Apply 3 times a day to affected area
.5TACHS	Take 1/2 tablet before meals and at bedtime	APOUQID	Apply in each eye 4 times a day
1CBIDIF	Take 1 capsule twice a day for infection	AP3-4XDPIT	Apply 3 to 4 times a day as needed for itching
1CBIDNR	Take 1 capsule twice a day for nerves	1SPVHS	Insert 1 suppository vaginally at bedtime
2PFTID	2 puffs 3 times a day	1AVHS	Insert 1 applicatorful vaginally at bedtime
2PQID	2 puffs 4 times a day		

MEDICATION-SPECIFIC CODES	
NIN	NALOXONE - Administer 1 spray into one nostril. Call 911. Repeat after 2-3 minutes if no or minimal response.
NIM	Inject 1 ML intramuscularly as directed. Call 911. Repeat after 2 to 3 minutes if no or minimal response.
EPI	Inject 1 auto injector intramuscularly into thigh as needed for severe allergic reaction.
ADVAIR	Inhale 1 dose by mouth twice daily. Rinse mouth after use.
BC21	Take 1 tablet daily for the first 21 days of each 28 day cycle.
BC28	Take 1 tablet daily for 28 days.
ESTRO	Take 1 tablet daily for days 1 thru 25 of each month.
FLC	To be administered by pharmacist for immunization.
MEDROL	Take 6 tablets on day 1 as directed on package and decrease by 1 tab each day for a total of 6 days.
MIACALCIN	Instill 1 spray into one nostril once daily. Alternate nostrils daily.
MONO	Insert 1 applicatorful vaginally at bedtime for 7 nights.
NITRO	Place 1 tablet under tongue if needed for chest pain.
NUVARING	Insert 1 ring vaginally as directed. Remove after 3 weeks & wait 7 days before inserting a new ring.
PEN	Take 1 tablet 4 times a day for 10 days.
PERIDEX	Swish 15 ML for 30 seconds, then spit. Use twice daily morning and evening after brushing teeth.
SPIRIVA	Inhale 1 capsule via handihaler once daily at the same time every day.
ZPAK	Take 2 tablets by mouth today, then take 1 tablet daily for 4 days.