

RxConnect SIG Codes: Component Codes

Link codes together with a semi-colon. Combination and medication-specific codes are on page 2.

TABLETS & CAPSULES		ROUTE		FREQUENCY		AS NEEDED (PRN)	
1T	Take 1 tablet	PO	By mouth	QD	Every day	PRN	As needed
2T	Take 2 tablets	EN	Each nostril	BID	Twice a day	AN	As needed
12T	Take 1 to 2 tablets	IEN	Into each nostril	TID	3 times a day	PPA	As needed for pain
S1T	Take 1/2 to 1 tablet	OU	Into both eyes	QID	4 times a day	PSP	As needed for severe pain
2THS	Take 2 tablets at bedtime	OD	Into right eye	XD	Times a day	PPF	As needed for pain and fever
C1T	Chew 1 tablet	OS	Into left eye	UTB	Up to 2 times daily	PMS	As needed for muscle spasm
D1T	Dissolve 1 tablet	OA	Into affected eye	UTT	Up to 3 times daily	PCR	As needed for cramps
1C	Take 1 capsule	AU	Each ear	QOD	Every other day	PHA	As needed for headache
2C	Take 2 capsules	AD	Right ear	TIW	Three times a week	PAN	As needed for anxiety
LIQUIDS		AS	Left ear	WK1	Once per week	PSL	As needed for sleep
1CC	Take 1 CC	AL	Left ear	Q3	Every 3 hours	PNA	As needed for nausea
15CC	Take 15 CC	AA	Affected ear	Q4	Every 4 hours	PNV	As needed for nausea/vomiting
1ML	Take 1 ML	VAG	Vaginally	Q6	Every 6 hours	PDI	As needed for diarrhea
TVF	Take 1 ML daily	TOPICAL / APPLY		Q8	Every 8 hours	PDZ	As needed for dizziness
TV2	1 ML daily to age 2	APDY	Apply daily	Q12	Every 12 hours	PCG	As needed for cough
1DRD	Take 1 dropperful daily	ABID	Apply twice a day	Q46	Every 4-6 hours	PCN	As needed for congestion
SAS	Swish and swallow	ATID	Apply 3 times a day	Q812	Every 8 to 12 hours	PWH	As needed for wheeze
DROPS (EYE / EAR)		AQID	Apply 4 times a day	828	At 8AM, 2PM, and 8PM	PAL	As needed for allergy
1D	Instill 1 drop	APHS	Apply at bedtime	848	At 8AM, 4PM, and 8PM	PIT	As needed for itching
2D	Instill 2 drops	A1P	Apply 1 patch	AM	In the morning	INDICATIONS	
I12D	Instill 1 to 2 drops	A1PD	Apply 1 patch daily	PM	In the evening	BD	For breathing difficulty
G1D	Give 1 drop	AAA	Apply to affected area	APM	And in the evening	BP	For blood pressure
G12D	Give 1 to 2 drops	ATR	Apply to rash	HS	At bedtime	CG	For congestion
G4D	Give 4 drops	AAE	Apply to affected eye	TIMING & FOOD		CST	For constipation
INJECTION		ATO	Apply topically	AC	Before meals	DC	For diarrhea and cramps
IJ	Inject	ASP	Apply sparingly	PC	After meals	DI	For diarrhea
IM	Intramuscular	APLOUD	Apply locally as directed	ACHS	Before meals and at bedtime	DN	For diarrhea and nausea
SQ	Subcutaneously	APV	Apply vaginally	PCHS	After meals and at bedtime	DZ	For dizziness
UNS	Units	1A	Insert 1 applicatorful	AB	Before breakfast	FA	For arthritis
INHALER / SPRAY		1SP	Insert 1 suppository	BBS	Before breakfast and supper	FB	For bronchitis
1P	Take 1 puff	DIRECTIONS		ASU	With supper	FC	For control
2P	Take 2 puffs	UD	As directed	ES	On empty stomach	FI	For infection
I1P	Inhale 1 puff	TUD	Take as directed	WF	With food	FM	For migraine
I2P	Inhale 2 puffs	UAD	Use as directed	WM	With meals	FP	For pain
I3P	Inhale 3 puffs	AUD	Apply as directed	WFM	With food and milk	FW	For wheeze
I12P	Inhale 1 to 2 puffs	DURATION		WW	With water	NV	For nausea and vomiting
SP1	1 spray	F1D	For 1 day	WOJ	With orange juice	RD	For rectal discomfort
SP2	2 sprays	F7D	For 7 days	WBL	With breakfast and lunch	SOB	For shortness of breath
		F1W	For 1 week	WBS	With breakfast and supper	TR	For tremor
		FXD	For 10 days	WD	With dinner	WE	For wheeze
		UF	Until finished			WZ	For wheezing

CODE COMPONENTS (combine to build a code)							
B	Back	C	Capsule	D	Daily	F	For fever
J	Joints	N	For nausea	O	Eye	S	1/2
U	Use	W	Water	BL	Blood	FH	Forehead
FT	Foot	ST	Stomach				

RxConnect SIG Codes: Combination & Medication-Specific

COMBINATION CODES (a complete sig in one entry)			
12TBID	Take 1 to 2 tablets 2 times a day	2PFQ6	2 puffs every 6 hours
12T34D	Take 1 to 2 tablets 3 or 4 times a day	IS1GTID	Instill 1 drop 3 times a day
12TQ34	Take 1 to 2 tablets every 3 to 4 hours	IS1GOUQID	Instill 1 drop in each eye 4 times a day
12TQ8	Take 1 to 2 tablets every 8 hours	1GOUBID	Put 1 drop in each eye twice a day
.5TQAM	Take 1/2 tablet every morning	1GADBID	Put 1 drop in right ear twice a day
.5TQ4PP	Take 1/2 tablet every 4 hours as needed for pain	1GAUBID	Put 1 drop in each ear twice a day
.5TABAL	Take 1/2 tablet before breakfast and lunch	APTIDAA	Apply 3 times a day to affected area
.5TACHS	Take 1/2 tablet before meals and at bedtime	APOUQID	Apply in each eye 4 times a day
1CBIDIF	Take 1 capsule twice a day for infection	AP3-4XDPIT	Apply 3 to 4 times a day as needed for itching
1CBIDNR	Take 1 capsule twice a day for nerves	1SPVHS	Insert 1 suppository vaginally at bedtime
2PFTID	2 puffs 3 times a day	1AVHS	Insert 1 applicatorful vaginally at bedtime
2PQID	2 puffs 4 times a day		

MEDICATION-SPECIFIC CODES	
NIN	NALOXONE - Administer 1 spray into one nostril. Call 911. Repeat after 2-3 minutes if no or minimal response.
NIM	Inject 1 ML intramuscularly as directed. Call 911. Repeat after 2 to 3 minutes if no or minimal response.
EPI	Inject 1 auto injector intramuscularly into thigh as needed for severe allergic reaction.
ADVAIR	Inhale 1 dose by mouth twice daily. Rinse mouth after use.
BC21	Take 1 tablet daily for the first 21 days of each 28 day cycle.
BC28	Take 1 tablet daily for 28 days.
ESTRO	Take 1 tablet daily for days 1 thru 25 of each month.
FLC	To be administered by pharmacist for immunization.
MEDROL	Take 6 tablets on day 1 as directed on package and decrease by 1 tab each day for a total of 6 days.
MIACALCIN	Instill 1 spray into one nostril once daily. Alternate nostrils daily.
MONO	Insert 1 applicatorful vaginally at bedtime for 7 nights.
NITRO	Place 1 tablet under tongue if needed for chest pain.
NUVARING	Insert 1 ring vaginally as directed. Remove after 3 weeks & wait 7 days before inserting a new ring.
PEN	Take 1 tablet 4 times a day for 10 days.
PERIDEX	Swish 15 ML for 30 seconds, then spit. Use twice daily morning and evening after brushing teeth.
SPIRIVA	Inhale 1 capsule via handihaler once daily at the same time every day.
ZPAK	Take 2 tablets by mouth today, then take 1 tablet daily for 4 days.