


# Ehrhart Polynomials



Matthias Beck


San Francisco State University

<https://matthbeck.github.io/>

VIII Encuentro Colombiano  
De Combinatoria

“Science is what we understand well enough to explain to a computer, art is all the rest.”

Donald Knuth



# Themes



Discrete-geometric  
polynomials

Computation  
(complexity)

Generating  
functions

Combinatorial  
structures

Polyhedra



# A Sample Problem: Birkhoff–von Neumann Polytope

This site is supported by donations to [The OEIS Foundation](#).

## THE ON-LINE ENCYCLOPEDIA OF INTEGER SEQUENCES®

founded in 1964 by N. J. A. Sloane

[Hints](#)  
(Greetings from [The On-Line Encyclopedia of Integer Sequences!](#))

A037302 Normalized volume of Birkhoff polytope of  $n \times n$  doubly-stochastic square matrices. If the volume is  $v(n)$ , then  $a(n) = ((n-1)^2)! * v(n) / n^{(n-1)}$ .

1, 1, 3, 352, 4718075, 14666561365176, 17832560768358341943028,  
12816077964079346687829905128694016, 7658969897501574748537755050756794492337074203099,  
5091038988117504946842559205930853037841762820367901333706255223000 ([list](#); [graph](#); [refs](#); [listen](#); [history](#);  
[text](#); [internal format](#))

OFFSET 1,3

COMMENTS The Birkhoff polytope is an  $(n-1)^2$ -dimensional polytope in  $n^2$ -dimensional space; its vertices are the  $n!$  permutation matrices.  
Is  $a(n)$  divisible by  $n^2$  for all  $n \geq 4$ ? - [Dean Hickerson](#), Nov 27 2002

$$B_n = \left\{ \begin{pmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \cdots & x_{nn} \end{pmatrix} \in \mathbb{R}_{\geq 0}^{n^2} : \begin{array}{l} \sum_j x_{jk} = 1 \text{ for all } 1 \leq k \leq n \\ \sum_k x_{jk} = 1 \text{ for all } 1 \leq j \leq n \end{array} \right\}$$

# Discrete Volumes



**Rational polyhedron**  $\mathcal{P} \subset \mathbb{R}^d$  – solution set of a system of linear equalities & inequalities with integer coefficients

**Goal:** understand  $\mathcal{P} \cap \mathbb{Z}^d \dots$

► (list)  $\sum_{\mathbf{m} \in \mathcal{P} \cap \mathbb{Z}^d} z_1^{m_1} z_2^{m_2} \dots z_d^{m_d}$

► (count)  $|\mathcal{P} \cap \mathbb{Z}^d|$

► (volume)  $\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^d} \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right|$



# Discrete Volumes



**Rational polyhedron**  $\mathcal{P} \subset \mathbb{R}^d$  – solution set of a system of linear equalities & inequalities with integer coefficients

**Goal:** understand  $\mathcal{P} \cap \mathbb{Z}^d \dots$

► (list)  $\sum_{\mathbf{m} \in \mathcal{P} \cap \mathbb{Z}^d} z_1^{m_1} z_2^{m_2} \dots z_d^{m_d}$

► (count)  $|\mathcal{P} \cap \mathbb{Z}^d|$

► (volume)  $\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^d} \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right|$



**Ehrhart function**  $L_{\mathcal{P}}(t) := \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right| = |t\mathcal{P} \cap \mathbb{Z}^d|$  for  $t \in \mathbb{Z}_{>0}$

# Some Motivation

- ▶ Linear systems are *everywhere*, and so polyhedra are everywhere.

## Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).



## Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.

## Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.


## Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.
- ▶ Polytopes are basic geometric objects, yet even for these basic objects volume computation is **hard** and there remain many open problems.

## Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.
- ▶ Polytopes are basic geometric objects, yet even for these basic objects volume computation is **hard** and there remain many open problems.
- ▶ Also, polytopes are **cool**.

# Today's Menu: Get Your Hands Dirty




# The Unit Cube

Lattice polytope  $\mathcal{P} \subset \mathbb{R}^d$  – convex hull of finitely points in  $\mathbb{Z}^d$

For  $t \in \mathbb{Z}_{>0}$  let  $L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$

The unit cube in  $\mathbb{R}^d$  is  $\mathcal{P} = [0, 1]^d = \{\mathbf{x} \in \mathbb{R}^d : 0 \leq x_j \leq 1\}$




$$\longrightarrow L_{\mathcal{P}}(t) = (t+1)^d$$

# The Unit Cube

Lattice polytope  $\mathcal{P} \subset \mathbb{R}^d$  – convex hull of finitely points in  $\mathbb{Z}^d$

For  $t \in \mathbb{Z}_{>0}$  let  $L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$

The unit cube in  $\mathbb{R}^d$  is  $\mathcal{P} = [0, 1]^d = \{\mathbf{x} \in \mathbb{R}^d : 0 \leq x_j \leq 1\}$




$$\longrightarrow L_{\mathcal{P}}(t) = (t+1)^d$$

$$L_{\mathcal{P}^\circ}(t) = (t-1)^d$$

# The Standard Simplex

The **standard simplex**  $\Delta \in \mathbb{R}^d$  is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{x \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq 1\}$$





# The Standard Simplex

The **standard simplex**  $\Delta \in \mathbb{R}^d$  is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq 1 \}$$

$$\begin{aligned} L_{\Delta}(t) &= \# \{ (x_1, x_2, \dots, x_d) \in \mathbb{Z}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq t \} \\ &= \# \left\{ (x_1, x_2, \dots, x_d, x_{d+1}) \in \mathbb{Z}_{\geq 0}^{d+1} : x_1 + x_2 + \cdots + x_{d+1} = t \right\} \\ &= \binom{d+t}{d} \end{aligned}$$

# The Standard Simplex

The **standard simplex**  $\Delta \in \mathbb{R}^d$  is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \dots + x_d \leq 1\}$$


$$\begin{aligned} L_{\Delta}(t) &= \# \{(x_1, x_2, \dots, x_d) \in \mathbb{Z}_{\geq 0}^d : x_1 + x_2 + \dots + x_d \leq t\} \\ &= \# \{(x_1, x_2, \dots, x_d, x_{d+1}) \in \mathbb{Z}_{\geq 0}^{d+1} : x_1 + x_2 + \dots + x_{d+1} = t\} \\ &= \binom{d+t}{d} \end{aligned}$$

$$L_{\Delta^{\circ}}(t) = \binom{t-1}{d}$$

# The Cross Polytope

The **cross-polytope**  $\diamond \in \mathbb{R}^d$  is

$$\diamond = \{ \mathbf{x} \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1 \}$$




# The Cross Polytope

The **cross-polytope**  $\diamond \in \mathbb{R}^d$  is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute  $L_\diamond(t)$  for  $d = 3 \dots$

► Triangulation




# The Cross Polytope

The **cross-polytope**  $\diamond \in \mathbb{R}^d$  is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute  $L_\diamond(t)$  for  $d = 3 \dots$

- ▶ Triangulation
- ▶ Disjoint triangulation




# The Cross Polytope

The **cross-polytope**  $\diamond \in \mathbb{R}^d$  is

$$\diamond = \{ \mathbf{x} \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1 \}$$

Let's compute  $L_\diamond(t)$  for  $d = 3 \dots$




- ▶ Triangulation
- ▶ Disjoint triangulation
- ▶ Interpolation

# The Cross Polytope

The **cross-polytope**  $\diamond \in \mathbb{R}^d$  is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute  $L_\diamond(t)$  for  $d = 3 \dots$



- ▶ Triangulation
- ▶ Disjoint triangulation
- ▶ Interpolation
- ▶ Generating function

# Ehrhart Polynomials



EH  
1959

**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $\dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

Equivalently,  $\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$  is rational:

$$\text{Ehr}_{\mathcal{P}}(z) = \frac{h(z)}{(1 - z)^{\dim \mathcal{P} + 1}}$$

where the **Ehrhart h-vector**  $h(z)$  satisfies  $h(0) = 1$  and  $h(1) = (\dim \mathcal{P})! \text{vol}(\mathcal{P})$ .



# Ehrhart Polynomials



EH  
1959

**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $\dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

Equivalently,  $\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$  is rational:

$$\text{Ehr}_{\mathcal{P}}(z) = \frac{h(z)}{(1 - z)^{\dim \mathcal{P} + 1}}$$

where the **Ehrhart h-vector**  $h(z)$  satisfies  $h(0) = 1$  and  $h(1) = (\dim \mathcal{P})! \text{vol}(\mathcal{P})$ .

**Seeming dichotomy:**  $\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^{\dim \mathcal{P}}} L_{\mathcal{P}}(t)$  can be computed discretely via a finite amount of data.

# Ehrhart Polynomials



**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $d := \dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Equivalent descriptions of an Ehrhart polynomial:

- ▶  $L_{\mathcal{P}}(t) = c_d t^d + c_{d-1} t^{d-1} + \cdots + c_0$
- ▶ via roots of  $L_{\mathcal{P}}(t)$
- ▶  $\text{Ehr}_{\mathcal{P}}(z) \longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$

# Ehrhart Polynomials



**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $d := \dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Equivalent descriptions of an Ehrhart polynomial:

- ▶  $L_{\mathcal{P}}(t) = c_d t^d + c_{d-1} t^{d-1} + \cdots + c_0$
- ▶ via roots of  $L_{\mathcal{P}}(t)$
- ▶  $\text{Ehr}_{\mathcal{P}}(z) \longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$

**Open Problem** Classify Ehrhart polynomials.

# Ehrhart Polynomials



**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $d := \dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$$

**Theorem** (Macdonald 1971)  $(-1)^d L_{\mathcal{P}}(-t)$  enumerates the **interior** lattice points in  $t\mathcal{P}$ . Equivalently,

$$L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

# Ehrhart Polynomials



**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $d := \dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

**Theorem** (Stanley 1980)  $h_0, h_1, \dots, h_d$  are nonnegative integers.

# Ehrhart Polynomials



**Theorem** (Ehrhart 1962) For any lattice polytope  $\mathcal{P}$ ,  $L_{\mathcal{P}}(t)$  is a polynomial in  $t$  of degree  $d := \dim \mathcal{P}$  with leading coefficient  $\text{vol } \mathcal{P}$  and constant term 1.


$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

**Theorem** (Stanley 1980)  $h_0, h_1, \dots, h_d$  are nonnegative integers.

**Corollary** If  $h_{d+1-k} > 0$  then  $k\mathcal{P}^\circ$  contains an integer point.

# Birkhoff–von Neumann Revisited



For more about roots of  
(Ehrhart) polynomials,  
see Braun (2008) and  
Pfeifle (2010).