


Ehrhart Polynomials

Day I: Appetizers



Matthias Beck


San Francisco State University

<https://matthbeck.github.io/>

VIII Encuentro Colombiano
De Combinatoria

“Science is what we understand well enough to explain to a computer, art is all the rest.”

Donald Knuth



A Sample Problem: Birkhoff–von Neumann Polytope

This site is supported by donations to [The OEIS Foundation](#).

THE ON-LINE ENCYCLOPEDIA OF INTEGER SEQUENCES®

founded in 1964 by N. J. A. Sloane

[Hints](#)
(Greetings from [The On-Line Encyclopedia of Integer Sequences!](#))

A037302 Normalized volume of Birkhoff polytope of $n \times n$ doubly-stochastic square matrices. If the volume is $v(n)$, then $a(n) = ((n-1)^2)! * v(n) / n^{(n-1)}$.

1, 1, 3, 352, 4718075, 14666561365176, 17832560768358341943028,
12816077964079346687829905128694016, 7658969897501574748537755050756794492337074203099,
5091038988117504946842559205930853037841762820367901333706255223000 ([list](#); [graph](#); [refs](#); [listen](#); [history](#);
[text](#); [internal format](#))

OFFSET 1,3

COMMENTS The Birkhoff polytope is an $(n-1)^2$ -dimensional polytope in n^2 -dimensional space; its vertices are the $n!$ permutation matrices.
Is $a(n)$ divisible by n^2 for all $n \geq 4$? - [Dean Hickerson](#), Nov 27 2002

$$B_n = \left\{ \begin{pmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \cdots & x_{nn} \end{pmatrix} \in \mathbb{R}_{\geq 0}^{n^2} : \begin{array}{l} \sum_j x_{jk} = 1 \text{ for all } 1 \leq k \leq n \\ \sum_k x_{jk} = 1 \text{ for all } 1 \leq j \leq n \end{array} \right\}$$

Discrete Volumes



Rational polyhedron $\mathcal{P} \subset \mathbb{R}^d$ – solution set of a system of linear equalities & inequalities with integer coefficients

Goal: understand $\mathcal{P} \cap \mathbb{Z}^d \dots$

► (list)
$$\sum_{\mathbf{m} \in \mathcal{P} \cap \mathbb{Z}^d} z_1^{m_1} z_2^{m_2} \dots z_d^{m_d}$$

► (count) $|\mathcal{P} \cap \mathbb{Z}^d|$

► (volume)
$$\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^d} \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right|$$



Discrete Volumes



Rational polyhedron $\mathcal{P} \subset \mathbb{R}^d$ – solution set of a system of linear equalities & inequalities with integer coefficients

Goal: understand $\mathcal{P} \cap \mathbb{Z}^d \dots$

► (list) $\sum_{\mathbf{m} \in \mathcal{P} \cap \mathbb{Z}^d} z_1^{m_1} z_2^{m_2} \dots z_d^{m_d}$

► (count) $|\mathcal{P} \cap \mathbb{Z}^d|$

► (volume) $\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^d} \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right|$



Ehrhart function $L_{\mathcal{P}}(t) := \left| \mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right| = |t\mathcal{P} \cap \mathbb{Z}^d|$ for $t \in \mathbb{Z}_{>0}$

Some Motivation

- ▶ Linear systems are *everywhere*, and so polyhedra are everywhere.

Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).

Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.

Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.


Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.
- ▶ Polytopes are basic geometric objects, yet even for these basic objects volume computation is **hard** and there remain many open problems.

Some Motivation

- ▶ Linear systems are **everywhere**, and so polyhedra are everywhere.
- ▶ In applications, the **volume** of the polytope represented by a linear system measures some fundamental data of this system (“average”).
- ▶ Many **discrete problems** in various areas are linear problems, thus they ask for the discrete volume of a polytope in disguise.
- ▶ Much discrete geometry can be modeled using **polynomials** and, conversely, many combinatorial polynomials can be modeled geometrically.
- ▶ Polytopes are basic geometric objects, yet even for these basic objects volume computation is **hard** and there remain many open problems.
- ▶ Also, polytopes are **cool**.

Today's Menu: Get Our Hands Dirty




The Unit Cube

Lattice polytope $\mathcal{P} \subset \mathbb{R}^d$ – convex hull of finitely points in \mathbb{Z}^d

For $t \in \mathbb{Z}_{>0}$ let $L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$

The unit cube in \mathbb{R}^d is $\mathcal{P} = [0, 1]^d = \{\mathbf{x} \in \mathbb{R}^d : 0 \leq x_j \leq 1\}$




$$\longrightarrow L_{\mathcal{P}}(t) = (t+1)^d$$

The Unit Cube

Lattice polytope $\mathcal{P} \subset \mathbb{R}^d$ – convex hull of finitely points in \mathbb{Z}^d

For $t \in \mathbb{Z}_{>0}$ let $L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$

The unit cube in \mathbb{R}^d is $\mathcal{P} = [0, 1]^d = \{\mathbf{x} \in \mathbb{R}^d : 0 \leq x_j \leq 1\}$




$$\longrightarrow L_{\mathcal{P}}(t) = (t+1)^d$$

$$L_{\mathcal{P}^\circ}(t) = (t-1)^d$$

The Standard Simplex

The **standard simplex** $\Delta \in \mathbb{R}^d$ is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{x \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq 1\}$$



The Standard Simplex

The **standard simplex** $\Delta \in \mathbb{R}^d$ is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq 1 \}$$

$$\begin{aligned} L_{\Delta}(t) &= \# \{ (x_1, x_2, \dots, x_d) \in \mathbb{Z}_{\geq 0}^d : x_1 + x_2 + \cdots + x_d \leq t \} \\ &= \# \left\{ (x_1, x_2, \dots, x_d, x_{d+1}) \in \mathbb{Z}_{\geq 0}^{d+1} : x_1 + x_2 + \cdots + x_{d+1} = t \right\} \\ &= \binom{d+t}{d} \end{aligned}$$

The Standard Simplex

The **standard simplex** $\Delta \in \mathbb{R}^d$ is the convex hull of the unit vectors and the origin; alternatively,

$$\Delta = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}_{\geq 0}^d : x_1 + x_2 + \dots + x_d \leq 1\}$$


$$\begin{aligned} L_{\Delta}(t) &= \# \{(x_1, x_2, \dots, x_d) \in \mathbb{Z}_{\geq 0}^d : x_1 + x_2 + \dots + x_d \leq t\} \\ &= \# \{(x_1, x_2, \dots, x_d, x_{d+1}) \in \mathbb{Z}_{\geq 0}^{d+1} : x_1 + x_2 + \dots + x_{d+1} = t\} \\ &= \binom{d+t}{d} \end{aligned}$$

$$L_{\Delta^{\circ}}(t) = \binom{t-1}{d}$$

The Cross Polytope

The **cross-polytope** $\diamond \in \mathbb{R}^d$ is

$$\diamond = \{ \mathbf{x} \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1 \}$$




The Cross Polytope

The **cross-polytope** $\diamond \in \mathbb{R}^d$ is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute $L_\diamond(t)$ for $d = 3 \dots$

► Triangulation




The Cross Polytope

The **cross-polytope** $\diamond \in \mathbb{R}^d$ is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute $L_\diamond(t)$ for $d = 3 \dots$

- ▶ Triangulation
- ▶ Disjoint triangulation




The Cross Polytope

The **cross-polytope** $\diamond \in \mathbb{R}^d$ is

$$\diamond = \{ \mathbf{x} \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1 \}$$

Let's compute $L_\diamond(t)$ for $d = 3 \dots$




- ▶ Triangulation
- ▶ Disjoint triangulation
- ▶ Interpolation

The Cross Polytope

The **cross-polytope** $\diamond \in \mathbb{R}^d$ is

$$\diamond = \{x \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\}$$

Let's compute $L_\diamond(t)$ for $d = 3 \dots$



- ▶ Triangulation
- ▶ Disjoint triangulation
- ▶ Interpolation
- ▶ Generating function

Ehrhart Polynomials



EH
1959

Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $\dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

Equivalently, $\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$ is rational:

$$\text{Ehr}_{\mathcal{P}}(z) = \frac{h(z)}{(1 - z)^{\dim \mathcal{P} + 1}}$$

where the **Ehrhart h-vector** $h(z)$ satisfies $h(0) = 1$ and $h(1) = (\dim \mathcal{P})! \text{vol}(\mathcal{P})$.

Ehrhart Polynomials



EH
1959

Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $\dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

Equivalently, $\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$ is rational:

$$\text{Ehr}_{\mathcal{P}}(z) = \frac{h(z)}{(1 - z)^{\dim \mathcal{P} + 1}}$$

where the **Ehrhart h-vector** $h(z)$ satisfies $h(0) = 1$ and $h(1) = (\dim \mathcal{P})! \text{vol}(\mathcal{P})$.

Seeming dichotomy: $\text{vol}(\mathcal{P}) = \lim_{t \rightarrow \infty} \frac{1}{t^{\dim \mathcal{P}}} L_{\mathcal{P}}(t)$ can be computed discretely via a finite amount of data.

Ehrhart Polynomials



Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $d := \dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Equivalent descriptions of an Ehrhart polynomial:

- ▶ $L_{\mathcal{P}}(t) = c_d t^d + c_{d-1} t^{d-1} + \cdots + c_0$
- ▶ via roots of $L_{\mathcal{P}}(t)$
- ▶ $\text{Ehr}_{\mathcal{P}}(z) \longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$

Ehrhart Polynomials



Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $d := \dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Equivalent descriptions of an Ehrhart polynomial:

- ▶ $L_{\mathcal{P}}(t) = c_d t^d + c_{d-1} t^{d-1} + \cdots + c_0$
- ▶ via roots of $L_{\mathcal{P}}(t)$
- ▶ $\text{Ehr}_{\mathcal{P}}(z) \longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$

Open Problem Classify Ehrhart polynomials.

Ehrhart Polynomials



Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $d := \dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}}(t) = h_0 \binom{t+d}{d} + h_1 \binom{t+d-1}{d} + \cdots + h_d \binom{t}{d}$$

Theorem (Macdonald 1971) $(-1)^d L_{\mathcal{P}}(-t)$ enumerates the **interior** lattice points in $t\mathcal{P}$. Equivalently,

$$L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

Ehrhart Polynomials



Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $d := \dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

Theorem (Stanley 1980) h_0, h_1, \dots, h_d are nonnegative integers.

Ehrhart Polynomials



Theorem (Ehrhart 1962) For any lattice polytope \mathcal{P} , $L_{\mathcal{P}}(t)$ is a polynomial in t of degree $d := \dim \mathcal{P}$ with leading coefficient $\text{vol } \mathcal{P}$ and constant term 1.


$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

$$\longrightarrow L_{\mathcal{P}^\circ}(t) = h_d \binom{t+d-1}{d} + h_{d-1} \binom{t+d-2}{d} + \cdots + h_0 \binom{t-1}{d}$$

Theorem (Stanley 1980) h_0, h_1, \dots, h_d are nonnegative integers.

Corollary If $h_{d+1-k} > 0$ then $k\mathcal{P}^\circ$ contains an integer point.

Birkhoff–von Neumann Revisited



For more about roots of
(Ehrhart) polynomials,
see Braun (2008) and
Pfeifle (2010).