

Grid Graphs, Gorenstein Polytopes, and Domino Stackings

Matthias Beck (San Francisco State)

math.sfsu.edu/beck

Joint with Christian Haase (FU Berlin) & Steven Sam (MIT)

[arXiv:0711.4151](https://arxiv.org/abs/0711.4151)

“The hardest thing being with a mathematician is that they always have problems.”

Tendai Chitewere

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d , i.e., the vertices of \mathcal{P} are in \mathbb{Z}^n

$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$ (discrete volume of P)

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d , i.e., the vertices of \mathcal{P} are in \mathbb{Z}^n

$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$ (discrete volume of P)

Ehrhart's Theorem (1962) $L_{\mathcal{P}}(t)$ is a polynomial in $t \in \mathbb{N}$.

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d , i.e., the vertices of \mathcal{P} are in \mathbb{Z}^n

$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$ (discrete volume of P)

Ehrhart's Theorem (1962) $L_{\mathcal{P}}(t)$ is a polynomial in $t \in \mathbb{N}$. Equivalently,

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}},$$

where $h(z)$ is a polynomial, the **Ehrhart h-vector** of \mathcal{P} .

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d , i.e., the vertices of \mathcal{P} are in \mathbb{Z}^n

$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$ (discrete volume of P)

Ehrhart's Theorem (1962) $L_{\mathcal{P}}(t)$ is a polynomial in $t \in \mathbb{N}$. Equivalently,

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}},$$

where $h(z)$ is a polynomial, the **Ehrhart h-vector** of \mathcal{P} .

(Serious) Open Problem Classify Ehrhart polynomials/h-vectors.

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d , i.e., the vertices of \mathcal{P} are in \mathbb{Z}^n

$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$ (discrete volume of P)

Ehrhart's Theorem (1962) $L_{\mathcal{P}}(t)$ is a polynomial in $t \in \mathbb{N}$. Equivalently,

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}},$$

where $h(z)$ is a polynomial, the **Ehrhart h-vector** of \mathcal{P} .

(Serious) Open Problem Classify Ehrhart polynomials/h-vectors.

(Easier) Open Problem Construct and study special classes of lattice polytopes.

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Some sample problems

► Find \mathcal{P} for which the Ehrhart h-vector $h(z)$ is **palindromic**.

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Some sample problems

- ▶ Find \mathcal{P} for which the Ehrhart h-vector $h(z)$ is **palindromic**.
- ▶ For which \mathcal{P} is the Ehrhart h-vector $h(z)$ **unimodal**, i.e.,
 $h_0 \leq \cdots \leq h_{j-1} \leq h_j \geq h_{j+1} \geq \cdots \geq h_d$?

Ehrhart Polynomials

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

Some sample problems

- ▶ Find \mathcal{P} for which the Ehrhart h-vector $h(z)$ is **palindromic**.
- ▶ For which \mathcal{P} is the Ehrhart h-vector $h(z)$ **unimodal**, i.e.,
 $h_0 \leq \cdots \leq h_{j-1} \leq h_j \geq h_{j+1} \geq \cdots \geq h_d$?
- ▶ Study Ehrhart h-vectors of special classes, e.g., simplicial polytopes.

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

\mathcal{P} is **Gorenstein** if there is a $k \in \mathbb{N}$ such that $L_{\mathcal{P}^\circ}(t) = L_{\mathcal{P}}(t - k)$ for all $t \geq k$ and $L_{\mathcal{P}^\circ}(t) = 0$ for $0 < t < k$. We call k the **index** of \mathcal{P} .

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

\mathcal{P} is **Gorenstein** if there is a $k \in \mathbb{N}$ such that $L_{\mathcal{P}^\circ}(t) = L_{\mathcal{P}}(t - k)$ for all $t \geq k$ and $L_{\mathcal{P}^\circ}(t) = 0$ for $0 < t < k$. We call k the **index** of \mathcal{P} .

Examples

► the unit cube $\square = [0, 1]^d$ with $L_{\square}(t) = (t + 1)^d$ and $L_{\square^\circ}(t) = (t - 1)^d$

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

\mathcal{P} is **Gorenstein** if there is a $k \in \mathbb{N}$ such that $L_{\mathcal{P}^\circ}(t) = L_{\mathcal{P}}(t - k)$ for all $t \geq k$ and $L_{\mathcal{P}^\circ}(t) = 0$ for $0 < t < k$. We call k the **index** of \mathcal{P} .

Examples

- ▶ the unit cube $\square = [0, 1]^d$ with $L_{\square}(t) = (t + 1)^d$ and $L_{\square^\circ}(t) = (t - 1)^d$
- ▶ the standard simplex $\Delta = \text{conv}\{0, \mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d\}$ with $L_{\Delta}(t) = \binom{t+d}{d}$ and $L_{\Delta^\circ}(t) = \binom{t-1}{d}$

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

\mathcal{P} is **Gorenstein** if there is a $k \in \mathbb{N}$ such that $L_{\mathcal{P}^\circ}(t) = L_{\mathcal{P}}(t - k)$ for all $t \geq k$ and $L_{\mathcal{P}^\circ}(t) = 0$ for $0 < t < k$. We call k the **index** of \mathcal{P} .

Examples

- ▶ the unit cube $\square = [0, 1]^d$ with $L_{\square}(t) = (t + 1)^d$ and $L_{\square^\circ}(t) = (t - 1)^d$
- ▶ the standard simplex $\Delta = \text{conv}\{0, \mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d\}$ with $L_{\Delta}(t) = \binom{t+d}{d}$ and $L_{\Delta^\circ}(t) = \binom{t-1}{d}$
- ▶ the Birkhoff polytope

$$\left\{ \begin{pmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \cdots & x_{nn} \end{pmatrix} \in \mathbb{R}_{\geq 0}^{n^2} : \begin{array}{l} \sum_j x_{jk} = 1 \text{ for all } 1 \leq k \leq n \\ \sum_k x_{jk} = 1 \text{ for all } 1 \leq j \leq n \end{array} \right\}$$

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

\mathcal{P} is **Gorenstein** if and only if the Ehrhart h-vector $h(z)$ of \mathcal{P} is **palindromic** (this is a nice exercise if one knows the Ehrhart–Macdonald reciprocity theorem).

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

\mathcal{P} is **Gorenstein** if and only if the Ehrhart h-vector $h(z)$ of \mathcal{P} is **palindromic** (this is a nice exercise if one knows the Ehrhart–Macdonald reciprocity theorem).

Remark The Gorenstein property has an extension to Cohen–Macaulay algebras.

Gorenstein Polytopes

$\mathcal{P} \subset \mathbb{R}^n$ – lattice polytope of dimension d

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^n)$$

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t = \frac{h(z)}{(1-z)^{d+1}}$$

\mathcal{P} is **Gorenstein** if and only if the Ehrhart h-vector $h(z)$ of \mathcal{P} is **palindromic** (this is a nice exercise if one knows the Ehrhart–Macdonald reciprocity theorem).

Remark The Gorenstein property has an extension to Cohen–Macaulay algebras.

Goal Construct classes of Gorenstein polytopes.

Suggested Tools

- ▶ LattE macchiato (<http://www.math.ucdavis.edu/~latte/>)
- ▶ barvinok (<http://freshmeat.net/projects/barvinok/>)
- ▶ ehrhart
(<http://icps.u-strasbg.fr/Ehrhart/program/program.html>)

Suggested Tools

- ▶ LattE macchiato (<http://www.math.ucdavis.edu/~latte/>)
- ▶ barvinok (<http://freshmeat.net/projects/barvinok/>)
- ▶ ehrhart
(<http://icps.u-strasbg.fr/Ehrhart/program/program.html>)
- ▶ Normaliz
(<ftp://ftp.mathematik.uni-osnabrueck.de/pub/osm/kommalg/software/>)
- ▶ 4ti2 (www.4ti2.de)

Suggested Tools

- ▶ LattE macchiato (<http://www.math.ucdavis.edu/~latte/>)
- ▶ barvinok (<http://freshmeat.net/projects/barvinok/>)
- ▶ ehrhart
(<http://icps.u-strasbg.fr/Ehrhart/program/program.html>)
- ▶ Normaliz
(<ftp://ftp.mathematik.uni-osnabrueck.de/pub/osm/kommalg/software/>)
- ▶ 4ti2 (www.4ti2.de)
- ▶ polymake (<http://www.math.tu-berlin.de/polymake/>)

Perfect Matchings and Magic Labellings of Graphs

A **perfect matching** of a graph G is a subset $M \subseteq E(G)$ such that every vertex of G is incident with exactly one edge of M .

Perfect Matchings and Magic Labellings of Graphs

A **perfect matching** of a graph G is a subset $M \subseteq E(G)$ such that every vertex of G is incident with exactly one edge of M .

More generally, a **magic labelling** (with sum t) is a function $E(G) \rightarrow \mathbb{Z}_{\geq 0}$ such that for each vertex v , the sum of the labels of the edges incident to v equals t .

Perfect Matchings and Magic Labellings of Graphs

A **perfect matching** of a graph G is a subset $M \subseteq E(G)$ such that every vertex of G is incident with exactly one edge of M .

More generally, a **magic labelling** (with sum t) is a function $E(G) \rightarrow \mathbb{Z}_{\geq 0}$ such that for each vertex v , the sum of the labels of the edges incident to v equals t .

Thus perfect matchings are magic labellings of sum $t = 1$.

Perfect Matchings and Magic Labellings of Graphs

A **perfect matching** of a graph G is a subset $M \subseteq E(G)$ such that every vertex of G is incident with exactly one edge of M .


More generally, a **magic labelling** (with sum t) is a function $E(G) \rightarrow \mathbb{Z}_{\geq 0}$ such that for each vertex v , the sum of the labels of the edges incident to v equals t .

Thus perfect matchings are magic labellings of sum $t = 1$.

The **perfect matching polytope** associated to a graph G is the convex hull in $\mathbb{R}^{E(G)}$ of the incidence vectors of all perfect matchings of G .


Perfect Matchings of Grid Graphs

The $m \times n$ grid graph $\mathcal{G}(m, n)$ has vertex set $\{(i, j) \in \mathbb{Z}^2 : 0 \leq i < n, 0 \leq j < m\}$ and (i, j) and (i', j') are adjacent if $|i - i'| + |j - j'| = 1$.



Perfect Matchings of Grid Graphs


The $m \times n$ grid graph $\mathcal{G}(m, n)$ has vertex set $\{(i, j) \in \mathbb{Z}^2 : 0 \leq i < n, 0 \leq j < m\}$ and (i, j) and (i', j') are adjacent if $|i - i'| + |j - j'| = 1$.



$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t
 $\mathcal{P}(m, n)$ — perfect matching polytope of $\mathcal{G}(m, n)$



Perfect Matchings of Grid Graphs

The $m \times n$ grid graph $\mathcal{G}(m, n)$ has vertex set $\{(i, j) \in \mathbb{Z}^2 : 0 \leq i < n, 0 \leq j < m\}$ and (i, j) and (i', j') are adjacent if $|i - i'| + |j - j'| = 1$.



$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t
 $\mathcal{P}(m, n)$ — perfect matching polytope of $\mathcal{G}(m, n)$

The number $T(m, n, 1)$ of perfect matchings of $\mathcal{G}(m, n)$ can be interpreted as the number of domino tilings of an $m \times n$ board.



Perfect Matchings and Ehrhart Polynomials

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$\mathcal{P}(m, n)$ — perfect matching polytope of $\mathcal{G}(m, n)$

Note that $T(m, n, t) = L_{\mathcal{P}(m, n)}(t)$.

Perfect Matchings and Ehrhart Polynomials

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$\mathcal{P}(m, n)$ — perfect matching polytope of $\mathcal{G}(m, n)$

Note that $T(m, n, t) = L_{\mathcal{P}(m, n)}(t)$.

Theorem (BHS) Assume $m \leq n$. The perfect matching polytope $\mathcal{P}(m, n)$ is Gorenstein (of index k) if and only if one of the following holds:

- (1) $m = 1$ and n is even (in which case \mathcal{P} is a point)
- (2) $m = 2$ (in which case $k = 2$ if $n = 2$, and $k = 3$ for $n > 2$)
- (3) $m = 3$ and n is even (in which case $k = 5$)
- (4) $m = n = 4$ (in which case $k = 4$).

Perfect Matchings and Ehrhart Polynomials

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$\mathcal{P}(m, n)$ — perfect matching polytope of $\mathcal{G}(m, n)$

Note that $T(m, n, t) = L_{\mathcal{P}(m, n)}(t)$.

Theorem (BHS) Assume $m \leq n$. The perfect matching polytope $\mathcal{P}(m, n)$ is Gorenstein (of index k) if and only if one of the following holds:

- (1) $m = 1$ and n is even (in which case \mathcal{P} is a point)
- (2) $m = 2$ (in which case $k = 2$ if $n = 2$, and $k = 3$ for $n > 2$)
- (3) $m = 3$ and n is even (in which case $k = 5$)
- (4) $m = n = 4$ (in which case $k = 4$).

Theorem (BHS) If $\mathcal{P}(m, n)$ is Gorenstein then $\mathcal{P}(m, n)$ has a unimodal Ehrhart h-vector.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

Note that $T(2, n, 1)$ is a shift of the Fibonacci sequence.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

Note that $T(2, n, 1)$ is a shift of the Fibonacci sequence.

Klarner–Pollack (1980) For fixed m , $T(m, n, 1)$ satisfies a linear homogeneous recurrence relation.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

Note that $T(2, n, 1)$ is a shift of the Fibonacci sequence.

Klarner–Pollack (1980) For fixed m , $T(m, n, 1)$ satisfies a linear homogeneous recurrence relation.

Propp (2001) This recurrence relation for $T(m, n, 1)$ satisfies the reciprocity relation

$$T(m, n, 1) = \begin{cases} (-1)^n T(m, -n - 2, 1) & \text{if } m \equiv 2 \pmod{4}, \\ T(m, -n - 2, 1) & \text{otherwise.} \end{cases}$$

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

The natural correspondence between perfect matchings of $\mathcal{G}(m, n)$ and domino tilings begs the question whether there is an analogous construction for magic labellings of $\mathcal{G}(m, n)$.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

The natural correspondence between perfect matchings of $\mathcal{G}(m, n)$ and domino tilings begs the question whether there is an analogous construction for magic labellings of $\mathcal{G}(m, n)$. Here's one option:

A **domino stacking** (of height t) of an $m \times n$ rectangular board is a collection of t domino tilings piled on top of one another.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

The natural correspondence between perfect matchings of $\mathcal{G}(m, n)$ and domino tilings begs the question whether there is an analogous construction for magic labellings of $\mathcal{G}(m, n)$. Here's one option:

A **domino stacking** (of height t) of an $m \times n$ rectangular board is a collection of t domino tilings piled on top of one another.

Note that the number of such domino stackings is $T(m, n, 1)^t$.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

The natural correspondence between perfect matchings of $\mathcal{G}(m, n)$ and domino tilings begs the question whether there is an analogous construction for magic labellings of $\mathcal{G}(m, n)$. Here's one option:

A **domino stacking** (of height t) of an $m \times n$ rectangular board is a collection of t domino tilings piled on top of one another.

Note that the number of such domino stackings is $T(m, n, 1)^t$.

Proposition (BHS) Every magic labelling of sum t of $\mathcal{G}(m, n)$ can be realized as a domino stacking of height t of an $m \times n$ rectangular board.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

$T(m, n, 1)^t$ — number of domino stackings of height t of an $m \times n$ board

Theorem (BHS) For fixed m and t , each of the sequences $(T(m, n, t))_{n \geq 0}$ and $(T(m, n, 1)^t)_{n \geq 0}$ is given by a linear homogeneous recurrence relation.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

$T(m, n, 1)^t$ — number of domino stackings of height t of an $m \times n$ board

Theorem (BHS) For fixed m and t , each of the sequences $(T(m, n, t))_{n \geq 0}$ and $(T(m, n, 1)^t)_{n \geq 0}$ is given by a linear homogeneous recurrence relation.

Corollary (Riordan 1962) (Fixed) powers of Fibonacci numbers can be encoded by a linear homogeneous recurrence relation.

Domino Tilings

$\mathcal{G}(m, n)$ — $m \times n$ grid graph

$T(m, n, t)$ — number of magic labellings of $\mathcal{G}(m, n)$ with sum t

$T(m, n, 1)$ — number of domino tilings of an $m \times n$ board

$T(m, n, 1)^t$ — number of domino stackings of height t of an $m \times n$ board

Theorem (BHS) For fixed m and t , each of the sequences $(T(m, n, t))_{n \geq 0}$ and $(T(m, n, 1)^t)_{n \geq 0}$ is given by a linear homogeneous recurrence relation.

Corollary (Riordan 1962) (Fixed) powers of Fibonacci numbers can be encoded by a linear homogeneous recurrence relation.

Propp's reciprocity relation naturally extends to $(T(m, n, 1)^t)_{n \geq 0}$.

Some Open Problems

- ▶ Do all perfect matching polytopes $\mathcal{P}(m, n)$ have **unimodal** Ehrhart h-vectors? (We can compute Ehrhart h-vectors up to $\mathcal{P}(4, 5)$.)

Some Open Problems

- ▶ Do all perfect matching polytopes $\mathcal{P}(m, n)$ have **unimodal** Ehrhart h-vectors? (We can compute Ehrhart h-vectors up to $\mathcal{P}(4, 5)$.)
- ▶ Which **higher-dimensional grid graphs** have associated Gorenstein polytopes? (The k -dimensional cube graph is one example. We suspect there are not too many others.)

Some Open Problems

- ▶ Do all perfect matching polytopes $\mathcal{P}(m, n)$ have **unimodal** Ehrhart h-vectors? (We can compute Ehrhart h-vectors up to $\mathcal{P}(4, 5)$.)
- ▶ Which **higher-dimensional grid graphs** have associated Gorenstein polytopes? (The k -dimensional cube graph is one example. We suspect there are not too many others.)
- ▶ Do Riordan's recurrence relations for powers of Fibonacci numbers extend to $T(m, n, 1)^t$?

Some Open Problems

- ▶ Do all perfect matching polytopes $\mathcal{P}(m, n)$ have **unimodal** Ehrhart h-vectors? (We can compute Ehrhart h-vectors up to $\mathcal{P}(4, 5)$.)
- ▶ Which **higher-dimensional grid graphs** have associated Gorenstein polytopes? (The k -dimensional cube graph is one example. We suspect there are not too many others.)
- ▶ Do Riordan's recurrence relations for powers of Fibonacci numbers extend to $T(m, n, 1)^t$?
- ▶ Is there a recurrence relation for the number $T(m, n, t)$ of magic labellings of $\mathcal{G}(m, n)$ with sum t when m and n are both allowed to vary?