

- (1) Define a linear map $T \in L(\mathbf{R}^3)$ through $T(x_1, x_2, x_3) = (x_3, 3x_1, 2x_2)$. Compute an isometry S such that $T = S\sqrt{T^*T}$.

Solution. Fix the standard basis of \mathbf{R}^3 ; then T has the matrix form

$$T = \begin{pmatrix} 0 & 0 & 1 \\ 3 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}, \quad \text{and} \quad T^* = \begin{pmatrix} 0 & 3 & 0 \\ 0 & 0 & 2 \\ 1 & 0 & 0 \end{pmatrix}.$$

Thus

$$T^*T = \begin{pmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \sqrt{T^*T} = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and so

$$S = T(\sqrt{T^*T})^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

i.e., $S(x_1, x_2, x_3) = (x_3, x_1, x_2)$ will do the trick. \square

- (2) Prove that $T \in L(V)$ is invertible if and only if there exists a unique isometry S such that $T = S\sqrt{T^*T}$. (Hint for the “if” direction: look at our proof of the polar decomposition theorem.)

Proof. It will be useful (also for later) to know that for $T \in L(V)$,

$$T \text{ is invertible} \iff \sqrt{T^*T} \text{ is invertible}, \quad (\star)$$

so we’ll prove that first. Suppose that $T \in L(V)$ is invertible. Then T^* is also invertible (if $ST = TS = I$ then $T^*S^* = S^*T^* = I$), and thus so is T^*T . But then

$$\sqrt{T^*T}(\sqrt{T^*T}(T^*T)^{-1}) = ((T^*T)^{-1}\sqrt{T^*T})\sqrt{T^*T} = I,$$

i.e., $\sqrt{T^*T}$ is invertible. Conversely, if $\sqrt{T^*T}$ is invertible, then

$$I = \left((\sqrt{T^*T})^{-1} \right)^2 T^*T.$$

So if $\mathbf{v} \in \text{null}(T)$ then

$$\mathbf{v} = \left((\sqrt{T^*T})^{-1} \right)^2 T^*T(\mathbf{v}) = \left((\sqrt{T^*T})^{-1} \right)^2 T^*(\mathbf{0}) = \mathbf{0},$$

i.e., $\text{null}(T) = \{\mathbf{0}\}$, and so T is invertible.

Now for the actual problem: Suppose that T is invertible. Then (as we have just proved) $\sqrt{T^*T}$ is also invertible, and so the isometry S (which is guaranteed by the polar decomposition theorem) equals $T(\sqrt{T^*T})^{-1}$ and is hence unique.

Conversely, suppose there exists a unique isometry S such that $T = S\sqrt{T^*T}$. This means that the isometry $S(\mathbf{u} + \mathbf{w}) = F(\mathbf{u}) + G(\mathbf{w})$ that we constructed in our proof is unique, and thus the isometries

$$F : \text{range}(\sqrt{T^*T}) \rightarrow \text{range}(T) \quad \text{and} \quad G : \left(\text{range}(\sqrt{T^*T}) \right)^\perp \rightarrow (\text{range}(T))^\perp$$

are unique. However, we could have easily replaced G by $-G$, which is still an isometry. Thus G has to be the zero map, and so $(\text{range}(\sqrt{T^*T}))^\perp = \{\mathbf{0}\} = (\text{range}(T))^\perp$. But this means $\text{range}(T) = V$, i.e., T is surjective and thus invertible. \square

- (3) Suppose n is an odd positive integer and $T \in L(\mathbf{R}^n)$ is an isometry. Prove that there exists a nonzero vector $\mathbf{v} \in \mathbf{R}^n$ such that $T^2(\mathbf{v}) = \mathbf{v}$.

Proof. We know that there exists an orthonormal basis with respect to which T has block-diagonal form, with 1×1 blocks (of the form ± 1) and 2×2 blocks. Since n is odd, there must be a 1×1 block, and so there must be an eigenvector \mathbf{v} with eigenvalue $\lambda = \pm 1$. But then

$$T^2(\mathbf{v}) = \lambda^2 \mathbf{v} = \mathbf{v}.$$

□

- (4) Prove that $T \in L(V)$ is
 (a) invertible if and only if 0 is not a singular value of T ;
 (b) an isometry if and only if all the singular values of T are 1.

Proof. (a)

$$\begin{aligned} 0 \text{ is not a singular value of } T &\iff 0 \text{ is not an eigenvalue of } \sqrt{T^*T} \\ &\iff \text{null } \sqrt{T^*T} = \{\mathbf{0}\} \\ &\iff \sqrt{T^*T} \text{ is invertible} \\ &\stackrel{(*)}{\iff} T \text{ is invertible.} \end{aligned}$$

- (b) We proved in class that T is an isometry if and only if $T^*T = I$. This, in turn, holds if and only if $\sqrt{T^*T} = I$, which holds if and only if all eigenvalues of $\sqrt{T^*T}$ are 1, which holds if and only if all singular values of T are 1. □

- (5) Suppose $T \in L(V)$ has smallest singular value s and largest singular value l . Show that for all $\mathbf{v} \in V$

$$s \|\mathbf{v}\| \leq \|T(\mathbf{v})\| \leq l \|\mathbf{v}\|.$$

Proof. Denote the singular values of T by $s = s_1, s_2, \dots, s_n = l$. Suppose $\mathbf{v} \in V$. By the singular-value decomposition theorem, there exist orthonormal bases $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ and $(\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)$ such that

$$T(\mathbf{v}) = \sum_{j=1}^n s_j \langle \mathbf{v}, \mathbf{e}_j \rangle \mathbf{f}_j$$

Since $(\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)$ is orthonormal,

$$\|T(\mathbf{v})\|^2 = \sum_{j=1}^n |s_j \langle \mathbf{v}, \mathbf{e}_j \rangle|^2 = \sum_{j=1}^n s_j^2 |\langle \mathbf{v}, \mathbf{e}_j \rangle|^2$$

and so

$$s^2 \|\mathbf{v}\|^2 = s^2 \sum_{j=1}^n |\langle \mathbf{v}, \mathbf{e}_j \rangle|^2 \leq \|T(\mathbf{v})\|^2 \leq l^2 \sum_{j=1}^n |\langle \mathbf{v}, \mathbf{e}_j \rangle|^2 = l^2 \|\mathbf{v}\|^2.$$

(Here we used the fact that $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ is orthonormal.) □