

Combinatorial Reciprocity Theorems

Matthias Beck

San Francisco State University

math.sfsu.edu/beck/

Joint work with...

Thomas Zaslavsky, Binghamton University (SUNY)

math.CO/0309330

math.CO/0309331

math.CO/0506315

Chromatic polynomials of graphs

$\Gamma = (V, E)$ – graph (without loops)

k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$

Chromatic polynomials of graphs

$\Gamma = (V, E)$ – graph (without loops)

k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$

Proper k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$ such that $x_i \neq x_j$ if there is an edge ij

Chromatic polynomials of graphs

$\Gamma = (V, E)$ – graph (without loops)

k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$

Proper k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$ such that $x_i \neq x_j$ if there is an edge ij

Theorem (Birkhoff 1912, Whitney 1932) $\chi_\Gamma(k) := \#$ (proper k -colorings of Γ) is a monic polynomial in k of degree $|V|$.

Chromatic polynomials of graphs

$\Gamma = (V, E)$ – graph (without loops)

k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$

Proper k -coloring of Γ : mapping $x : V \rightarrow \{1, 2, \dots, k\}$ such that $x_i \neq x_j$ if there is an edge ij

Theorem (Birkhoff 1912, Whitney 1932) $\chi_\Gamma(k) := \#$ (proper k -colorings of Γ) is a monic polynomial in k of degree $|V|$.

Proof: Choose your favorite edge e of Γ and use

$$\chi_\Gamma(k) = \chi_{(\Gamma \setminus e)}(k) - \chi_{(\Gamma \cdot e)}(k)$$

and induction. . .

Stanley's Acyclic-Orientation Theorem

Theorem (Stanley 1973) $(-1)^{|V|}\chi_{\Gamma}(-k)$ equals the number of pairs (α, x) consisting of an acyclic orientation α of Γ and a compatible k -coloring. In particular, $(-1)^{|V|}\chi_{\Gamma}(-1)$ equals the number of acyclic orientations of Γ .

(An orientation α of Γ and a k -coloring x are **compatible** if $x_j \geq x_i$ whenever there is an edge oriented from i to j . An orientation is **acyclic** if it has no directed cycles.)

Flow polynomials

Nowhere-zero A -flow on a graph $\Gamma = (V, E)$: mapping $x : E \rightarrow A \setminus \{0\}$ (A an abelian group) such that for every node $v \in V$

$$\sum_{h(e)=v} x(e) = \sum_{t(e)=v} x(e)$$

$h(e) \quad := \quad \text{head}$
 $t(e) \quad := \quad \text{tail}$ of the edge e in a (fixed) orientation of Γ

Flow polynomials

Nowhere-zero A -flow on a graph $\Gamma = (V, E)$: mapping $x : E \rightarrow A \setminus \{0\}$ (A an abelian group) such that for every node $v \in V$

$$\sum_{h(e)=v} x(e) = \sum_{t(e)=v} x(e)$$

$h(e) :=$ head
 $t(e) :=$ tail of the edge e in a (fixed) orientation of Γ

Nowhere-zero k -flow : \mathbb{Z} -flow with values in $\{1, 2, \dots, k-1\}$

Flow polynomials

Nowhere-zero A -flow on a graph $\Gamma = (V, E)$: mapping $x : E \rightarrow A \setminus \{0\}$ (A an abelian group) such that for every node $v \in V$

$$\sum_{h(e)=v} x(e) = \sum_{t(e)=v} x(e)$$

$h(e) :=$ head
 $t(e) :=$ tail of the edge e in a (fixed) orientation of Γ

Nowhere-zero k -flow : \mathbb{Z} -flow with values in $\{1, 2, \dots, k-1\}$

Theorem

(Tutte 1954) $\overline{\varphi}_{\Gamma}(|A|) := \#(\text{nowhere-zero } A\text{-flows})$ is a polynomial in $|A|$.

(Kochol 2002) $\varphi_{\Gamma}(k) := \#(\text{nowhere-zero } k\text{-flows})$ is a polynomial in k .

Flow polynomials

Nowhere-zero A -flow on a graph $\Gamma = (V, E)$: mapping $x : E \rightarrow A \setminus \{0\}$ (A an abelian group) such that for every node $v \in V$

$$\sum_{h(e)=v} x(e) = \sum_{t(e)=v} x(e)$$

$h(e) :=$ head
 $t(e) :=$ tail of the edge e in a (fixed) orientation of Γ

Nowhere-zero k -flow : \mathbb{Z} -flow with values in $\{1, 2, \dots, k-1\}$

Theorem

(Tutte 1954) $\overline{\varphi}_{\Gamma}(|A|) := \#(\text{nowhere-zero } A\text{-flows})$ is a polynomial in $|A|$.

(Kochol 2002) $\varphi_{\Gamma}(k) := \#(\text{nowhere-zero } k\text{-flows})$ is a polynomial in k .

What about reciprocity?

(Weak) semimagic squares

$H_n(t)$ – number of nonnegative integral $n \times n$ -matrices in which every row and column sums to t

1	1	2
2	1	1
1	2	1

(Weak) semimagic squares

$H_n(t)$ – number of nonnegative integral $n \times n$ -matrices in which every row and column sums to t

1	1	2
2	1	1
1	2	1

Theorem (Ehrhart, Stanley 1973, conjectured by Anand-Dumir-Gupta 1966)
 $H_n(t)$ is a polynomial in t of degree $(n - 1)^2$.

(Weak) semimagic squares

$H_n(t)$ – number of nonnegative integral $n \times n$ -matrices in which every row and column sums to t

1	1	2
2	1	1
1	2	1

Theorem (Ehrhart, Stanley 1973, conjectured by Anand-Dumir-Gupta 1966)
 $H_n(t)$ is a polynomial in t of degree $(n-1)^2$ which satisfies

$$H_n(0) = 1, \quad H_n(-1) = H_n(-2) = \cdots = H_n(-n+1) = 0,$$

and $H_n(-n-t) = (-1)^{n-1} H_n(t)$.

(Weak) semimagic squares

$H_n(t)$ – number of nonnegative integral $n \times n$ -matrices in which every row and column sums to t

1	1	2
2	1	1
1	2	1

Theorem (Ehrhart, Stanley 1973, conjectured by Anand-Dumir-Gupta 1966)
 $H_n(t)$ is a polynomial in t of degree $(n-1)^2$ which satisfies

$$H_n(0) = 1, \quad H_n(-1) = H_n(-2) = \cdots = H_n(-n+1) = 0,$$

and $H_n(-n-t) = (-1)^{n-1} H_n(t)$.

What about “classical” magic squares?

Characteristic polynomials of hyperplane arrangements

$\mathcal{H} \subset \mathbb{R}^d$ – arrangement of affine hyperplanes

$\mathcal{L}(\mathcal{H}) := \{\bigcap \mathcal{S} : \mathcal{S} \subseteq \mathcal{H} \text{ and } \bigcap \mathcal{S} \neq \emptyset\}$, ordered by reverse inclusion

Characteristic polynomials of hyperplane arrangements

$\mathcal{H} \subset \mathbb{R}^d$ – arrangement of affine hyperplanes

$\mathcal{L}(\mathcal{H}) := \{\cap \mathcal{S} : \mathcal{S} \subseteq \mathcal{H} \text{ and } \cap \mathcal{S} \neq \emptyset\}$, ordered by reverse inclusion

Möbius function $\mu(r, s) := \begin{cases} 0 & \text{if } r \not\leq s, \\ 1 & \text{if } r = s, \\ -\sum_{r \leq u < s} \mu(r, u) & \text{if } r < s. \end{cases}$

Characteristic polynomial

$$p_{\mathcal{H}}(\lambda) := \sum_{s \in \mathcal{L}(\mathcal{H})} \mu(\mathbb{R}^d, s) \lambda^{\dim s}$$

Characteristic polynomials of hyperplane arrangements

$\mathcal{H} \subset \mathbb{R}^d$ – arrangement of affine hyperplanes

$\mathcal{L}(\mathcal{H}) := \{\cap \mathcal{S} : \mathcal{S} \subseteq \mathcal{H} \text{ and } \cap \mathcal{S} \neq \emptyset\}$, ordered by reverse inclusion

Möbius function $\mu(r, s) := \begin{cases} 0 & \text{if } r \not\leq s, \\ 1 & \text{if } r = s, \\ -\sum_{r \leq u < s} \mu(r, u) & \text{if } r < s. \end{cases}$

Characteristic polynomial

$$p_{\mathcal{H}}(\lambda) := \sum_{s \in \mathcal{L}(\mathcal{H})} \mu(\mathbb{R}^d, s) \lambda^{\dim s}$$

Theorem (Zaslavsky 1975) If $\mathbb{R}^d \notin \mathcal{H}$ then the number of regions into which a hyperplane arrangement \mathcal{H} divides \mathbb{R}^d is $(-1)^d p_{\mathcal{H}}(-1)$.

Ehrhart (quasi-)polynomials

$\mathcal{P} \subset \mathbb{R}^d$ – convex rational polytope

For $t \in \mathbb{Z}_{>0}$ let $\text{Ehr}_{\mathcal{P}}(t) := \# \left(\mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right)$

Ehrhart (quasi-)polynomials

$\mathcal{P} \subset \mathbb{R}^d$ – convex rational polytope

For $t \in \mathbb{Z}_{>0}$ let $\text{Ehr}_{\mathcal{P}}(t) := \# \left(\mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right)$

Theorem

(Ehrhart 1962) $\text{Ehr}_{\mathcal{P}}(t)$ is a quasipolynomial in t of degree $\dim \mathcal{P}$ with leading term $\text{vol } \mathcal{P}$ (normalized to $\text{aff } \mathcal{P} \cap \mathbb{Z}^d$) and constant term 1.

(A **quasipolynomial** is an expression $c_d(t) t^d + \cdots + c_1(t) t + c_0(t)$ where c_0, \dots, c_d are periodic functions in t .)

Ehrhart (quasi-)polynomials

$\mathcal{P} \subset \mathbb{R}^d$ – convex rational polytope

For $t \in \mathbb{Z}_{>0}$ let $\text{Ehr}_{\mathcal{P}}(t) := \# \left(\mathcal{P} \cap \frac{1}{t} \mathbb{Z}^d \right)$

Theorem

(Ehrhart 1962) $\text{Ehr}_{\mathcal{P}}(t)$ is a quasipolynomial in t of degree $\dim \mathcal{P}$ with leading term $\text{vol } \mathcal{P}$ (normalized to $\text{aff } \mathcal{P} \cap \mathbb{Z}^d$) and constant term 1.

(Macdonald 1971) $(-1)^{\dim \mathcal{P}} \text{Ehr}_{\mathcal{P}}(-t)$ enumerates the interior lattice points in $t\mathcal{P}$.

(A **quasipolynomial** is an expression $c_d(t)t^d + \dots + c_1(t)t + c_0(t)$ where c_0, \dots, c_d are periodic functions in t .)

Shameless plug

M. Beck & S. Robins



Computing the continuous discretely
Integer-point enumeration in polyhedra

To appear (late 2006) in [Springer Undergraduate Texts in Mathematics](#)

Preprint available at math.sfsu.edu/beck



Graph coloring a la Ehrhart

$$\chi_{K_2}(k) = k(k-1) \dots$$




Graph coloring a la Ehrhart

$$\chi_{K_2}(k) = k(k-1) \dots$$



$$\chi_{\Gamma}(k) = \# \left(\left((0, 1)^V \setminus \bigcup \mathcal{H}(\Gamma) \right) \cap \frac{1}{k+1} \mathbb{Z}^V \right)$$

Stanley's Theorem a la Ehrhart




$$\chi_{\Gamma}(k) = \# \left(((0, 1)^V \setminus \bigcup \mathcal{H}(\Gamma)) \cap \frac{1}{k+1} \mathbb{Z}^V \right)$$

Write $(0, 1)^V \setminus \bigcup \mathcal{H}(\Gamma) = \bigcup_j \mathcal{P}_j^{\circ}$, then by Ehrhart-Macdonald reciprocity

$$(-1)^{|V|} \chi_{\Gamma}(-k) = \sum_j \text{Ehr}_{P_j}(k-1)$$

Stanley's Theorem a la Ehrhart



$$\chi_{\Gamma}(k) = \# \left(((0, 1)^V \setminus \bigcup \mathcal{H}(\Gamma)) \cap \frac{1}{k+1} \mathbb{Z}^V \right)$$

Write $(0, 1)^V \setminus \bigcup \mathcal{H}(\Gamma) = \bigcup_j \mathcal{P}_j^{\circ}$, then by Ehrhart-Macdonald reciprocity

$$(-1)^{|V|} \chi_{\Gamma}(-k) = \sum_j \text{Ehr}_{P_j}(k-1)$$

Greene's observation

region of $\mathcal{H}(\Gamma) \iff$ acyclic orientation of Γ

$$x_i < x_j \iff i \longrightarrow j$$

Chromatic polynomials of signed graphs

Σ – signed graph (without loops): each edge is labelled $+$ or $-$

Proper k -coloring of Σ : mapping $x : V \rightarrow \{-k, -k+1, \dots, k\}$ such that, if edge ij has sign ϵ then $x_i \neq \epsilon x_j$



Chromatic polynomials of signed graphs

Σ – signed graph (without loops): each edge is labelled $+$ or $-$

Proper k -coloring of Σ : mapping $x : V \rightarrow \{-k, -k+1, \dots, k\}$ such that, if edge ij has sign ϵ then $x_i \neq \epsilon x_j$



Theorem (Zaslavsky 1982) $\chi_\Sigma(2k+1) := \#$ (proper k -colorings of Σ) and $\chi_\Sigma^*(2k) := \#$ (proper zero-free k -colorings of Σ) are monic polynomials of degree $|V|$. The number of compatible pairs (α, x) consisting of an acyclic orientation α and a k -coloring x of Σ is equal to $(-1)^{|V|} \chi_\Sigma(-(2k+1))$. The number in which x is zero-free equals $(-1)^{|V|} \chi_\Sigma^*(-2k)$. In particular, $(-1)^{|V|} \chi_\Sigma(-1)$ equals the number of acyclic orientations of Σ .

Signed-graph coloring a la Ehrhart



Theorem $\chi_{\Sigma}(2k + 1)$ and $\chi_{\Sigma}^*(2k)$ are two halves of one inside-out quasipolynomial.

Signed-graph coloring a la Ehrhart



Theorem $\chi_{\Sigma}(2k + 1)$ and $\chi_{\Sigma}^*(2k)$ are two halves of one inside-out quasipolynomial.

Open problem Is there a combinatorial interpretation of $\chi_{\Sigma}^*(-1)$?

Flow polynomials revisited

$$\begin{aligned}\varphi_{\Gamma}(k) &:= \# (\text{nowhere-zero } k\text{-flows}) \\ \overline{\varphi}_{\Gamma}(|A|) &:= \# (\text{nowhere-zero } A\text{-flows})\end{aligned}$$

Theorem $(-1)^{|E|-|V|+c(\Gamma)}\varphi_{\Gamma}(-k)$ equals the number of pairs (τ, x) consisting of a totally cyclic orientation τ and a compatible $(k+1)$ -flow x . In particular, the constant term $\varphi_{\Gamma}(0)$ equals the number of totally cyclic orientations of Γ .

(An orientation of Γ is **totally cyclic** if every edge lies in a coherent circle, that is, where the edges are oriented in a consistent direction around the circle. A totally cyclic orientation τ and a flow x are **compatible** if $x \geq 0$ when it is expressed in terms of τ .)

Flow polynomials revisited

$$\begin{aligned}\varphi_{\Gamma}(k) &:= \# (\text{nowhere-zero } k\text{-flows}) \\ \overline{\varphi}_{\Gamma}(|A|) &:= \# (\text{nowhere-zero } A\text{-flows})\end{aligned}$$

Theorem $(-1)^{|E|-|V|+c(\Gamma)}\varphi_{\Gamma}(-k)$ equals the number of pairs (τ, x) consisting of a totally cyclic orientation τ and a compatible $(k+1)$ -flow x . In particular, the constant term $\varphi_{\Gamma}(0)$ equals the number of totally cyclic orientations of Γ .

(An orientation of Γ is **totally cyclic** if every edge lies in a coherent circle, that is, where the edges are oriented in a consistent direction around the circle. A totally cyclic orientation τ and a flow x are **compatible** if $x \geq 0$ when it is expressed in terms of τ .)

Corollary $\varphi_{\Gamma}(0) = (-1)^{|E|-|V|+c(\Gamma)}\overline{\varphi}_{\Gamma}(-1)$

\exists analogous theorems for signed graphs

Open problems

Find a formula for, or a combinatorial interpretation of, the leading coefficient of φ_Γ .

Open problems

Find a formula for, or a combinatorial interpretation of, the leading coefficient of φ_Γ .

Is there a combinatorial interpretation of $\overline{\varphi}_\Gamma(-k)$ for $k \geq 2$?

Open problems


Find a formula for, or a combinatorial interpretation of, the leading coefficient of φ_Γ .

Is there a combinatorial interpretation of $\overline{\varphi}_\Gamma(-k)$ for $k \geq 2$?

For some graphs, both φ_Γ and $\overline{\varphi}_\Gamma$ have integral coefficients and φ_Γ is a multiple of $\overline{\varphi}_\Gamma$. Is there a general reason for these facts?

Inside-out counting functions

Inside-out polytope : $(\mathcal{P}, \mathcal{H})$



Multiplicity of $x \in \mathbb{R}^d$:

$$m_{\mathcal{P}, \mathcal{H}}(x) := \begin{cases} \# \text{ closed regions of } \mathcal{H} \text{ in } \mathcal{P} \text{ that contain } x & \text{if } x \in \mathcal{P}, \\ 0 & \text{if } x \notin \mathcal{P} \end{cases}$$

Closed Ehrhart quasipolynomial $E_{\mathcal{P}, \mathcal{H}}(t) := \sum_{x \in \frac{1}{t}\mathbb{Z}^d} m_{\mathcal{P}, \mathcal{H}}(x)$

Open Ehrhart quasipolynomial $E_{\mathcal{P}, \mathcal{H}}^{\circ}(t) := \# \left(\frac{1}{t}\mathbb{Z}^d \cap [\mathcal{P} \setminus \bigcup \mathcal{H}] \right)$

Basic inside-out results

Theorem If $(\mathcal{P}, \mathcal{H})$ is a closed, full-dimensional, rational inside-out polytope, then $E_{\mathcal{P}, \mathcal{H}}(t)$ and $E_{\mathcal{P}^\circ, \mathcal{H}}(t)$ are quasipolynomials in t of degree $\dim \mathcal{P}$ with leading term $\text{vol } P$, and with constant term $E_{\mathcal{P}, \mathcal{H}}(0)$ equal to the number of regions of $(\mathcal{P}, \mathcal{H})$. Furthermore,

$$E_{\mathcal{P}^\circ, \mathcal{H}}(t) = (-1)^d E_{\mathcal{P}, \mathcal{H}}(-t).$$

Basic inside-out results

Theorem If $(\mathcal{P}, \mathcal{H})$ is a closed, full-dimensional, rational inside-out polytope, then $E_{\mathcal{P}, \mathcal{H}}(t)$ and $E_{\mathcal{P}^\circ, \mathcal{H}}(t)$ are quasipolynomials in t of degree $\dim \mathcal{P}$ with leading term $\text{vol } P$, and with constant term $E_{\mathcal{P}, \mathcal{H}}(0)$ equal to the number of regions of $(\mathcal{P}, \mathcal{H})$. Furthermore,

$$E_{\mathcal{P}^\circ, \mathcal{H}}(t) = (-1)^d E_{\mathcal{P}, \mathcal{H}}(-t).$$

Theorem $(\mathcal{P}, \mathcal{H})$ is a closed, full-dimensional, rational inside-out polytope, then

$$E_{\mathcal{P}^\circ, \mathcal{H}}(t) = \sum_{u \in \mathcal{L}(\mathcal{H})} \mu(\mathbb{R}^d, u) \text{Ehr}_{\mathcal{P} \cap u}(t),$$

and if \mathcal{H} is transverse to \mathcal{P}

$$E_{\mathcal{P}, \mathcal{H}}(t) = \sum_{u \in \mathcal{L}(\mathcal{H})} |\mu(\mathbb{R}^d, u)| \text{Ehr}_{\mathcal{P} \cap u}(t).$$

(\mathcal{H} is **transverse to** \mathcal{P} if every flat $u \in \mathcal{L}(\mathcal{H})$ that intersects \mathcal{P} also intersects \mathcal{P}° , and \mathcal{P} does not lie in any of the hyperplanes of \mathcal{H} .)

(Strong) magic squares

$\text{Mag}_n(t)$ – number of nonnegative integral $n \times n$ -matrices with distinct entries in which every row and column sums to t

4	3	8
9	5	1
2	7	6

(Strong) magic squares

$\text{Mag}_n(t)$ – number of nonnegative integral $n \times n$ -matrices with **distinct** entries in which every row and column sums to t

4	3	8
9	5	1
2	7	6

Corollary $\text{Mag}_n(t)$ is a quasipolynomial in t of degree $n - 2n - 1$.

Open problem Can anything be said about the period of Mag_n ? Even in the weak case, do we ever get a polynomial?

Enumeration of integer points with distinct entries

$\mathcal{P} \subset \mathbb{R}^d$ – rational convex polytope, transverse to

$\mathcal{H} := \mathcal{H}[K_d]^{\text{aff } \mathcal{P}}$ – arrangement corresponding to K_d , induced on $\text{aff } \mathcal{P}$

Enumeration of integer points with distinct entries

$\mathcal{P} \subset \mathbb{R}^d$ – rational convex polytope, transverse to

$\mathcal{H} := \mathcal{H}[K_d]^{\text{aff } \mathcal{P}}$ – arrangement corresponding to K_d , induced on $\text{aff } \mathcal{P}$

Theorem The number $E_{\mathcal{P}^\circ, \mathcal{H}}^\circ(t)$ of integer points in $t\mathcal{P}^\circ$ with distinct entries is a quasipolynomial with constant term equal to the number of permutations of $[d]$ that are realizable in \mathcal{P} . Furthermore, $(-1)^{\dim \mathcal{P}^\circ} E_{\mathcal{P}^\circ, \mathcal{H}}^\circ(-t) = E_{\mathcal{P}, \mathcal{H}}(t) :=$ the number of pairs (x, σ) consisting of an integer point $x \in t\mathcal{P}$ and a compatible \mathcal{P} -realizable permutation σ of $[d]$.

(The point $x \in \mathbb{R}^d$ and the permutation τ are **compatible** if $x_{\tau 1} < x_{\tau 2} < \dots < x_{\tau d}$. τ is **realizable** in X if there exists a compatible $x \in X$.)

Enumeration of integer points with distinct entries

$\mathcal{P} \subset \mathbb{R}^d$ – rational convex polytope, transverse to

$\mathcal{H} := \mathcal{H}[K_d]^{\text{aff } \mathcal{P}}$ – arrangement corresponding to K_d , induced on $\text{aff } \mathcal{P}$

Theorem The number $E_{\mathcal{P}^\circ, \mathcal{H}}^\circ(t)$ of integer points in $t\mathcal{P}^\circ$ with distinct entries is a quasipolynomial with constant term equal to the number of permutations of $[d]$ that are realizable in \mathcal{P} . Furthermore, $(-1)^{\dim s} E_{\mathcal{P}^\circ, \mathcal{H}}^\circ(-t) = E_{\mathcal{P}, \mathcal{H}}(t) :=$ the number of pairs (x, σ) consisting of an integer point $x \in t\mathcal{P}$ and a compatible \mathcal{P} -realizable permutation σ of $[d]$.

(The point $x \in \mathbb{R}^d$ and the permutation τ are **compatible** if $x_{\tau 1} < x_{\tau 2} < \dots < x_{\tau d}$. τ is **realizable** in X if there exists a compatible $x \in X$.)

Applications (strong) magic squares, rectangles, cubes, graphs, ...

Open problems

When does $\mathcal{H}[K_d]$ change the denominator of \mathcal{P} ?

Open problems

When does $\mathcal{H}[K_d]$ change the denominator of \mathcal{P} ?

If \mathcal{P} has integral vertices then $\text{Ehr}_{\mathcal{P}}$ is a polynomial. What conditions on \mathcal{P} ensure that $E_{\mathcal{P}, \mathcal{H}[K_d]}$ is also a polynomial? (It need not be: Consider the line segment \mathcal{P} from $(0, 1)$ to $(1, 0)$ and let $\mathcal{H} = \{x_1 = x_2\}$.)

Open problems

When does $\mathcal{H}[K_d]$ change the denominator of \mathcal{P} ?

If \mathcal{P} has integral vertices then $\text{Ehr}_{\mathcal{P}}$ is a polynomial. What conditions on \mathcal{P} ensure that $E_{\mathcal{P}, \mathcal{H}[K_d]}$ is also a polynomial? (It need not be: Consider the line segment \mathcal{P} from $(0, 1)$ to $(1, 0)$ and let $\mathcal{H} = \{x_1 = x_2\}$.)

The inside-out Ehrhart quasipolynomials for some magic and latin squares have striking symmetries (coefficients alternate in sign, the polynomials factor nicely, etc.). Explain.

Open problems

When does $\mathcal{H}[K_d]$ change the denominator of \mathcal{P} ?

If \mathcal{P} has integral vertices then $\text{Ehr}_{\mathcal{P}}$ is a polynomial. What conditions on \mathcal{P} ensure that $E_{\mathcal{P}, \mathcal{H}[K_d]}$ is also a polynomial? (It need not be: Consider the line segment \mathcal{P} from $(0, 1)$ to $(1, 0)$ and let $\mathcal{H} = \{x_1 = x_2\}$.)

The inside-out Ehrhart quasipolynomials for some magic and latin squares have striking symmetries (coefficients alternate in sign, the polynomials factor nicely, etc.). Explain.

The inside-out Ehrhart quasipolynomials for some magic and latin squares have much lower periods than predicted by their denominators. Explain.

Open problems

When does $\mathcal{H}[K_d]$ change the denominator of \mathcal{P} ?

If \mathcal{P} has integral vertices then $\text{Ehr}_{\mathcal{P}}$ is a polynomial. What conditions on \mathcal{P} ensure that $E_{\mathcal{P}, \mathcal{H}[K_d]}$ is also a polynomial? (It need not be: Consider the line segment \mathcal{P} from $(0, 1)$ to $(1, 0)$ and let $\mathcal{H} = \{x_1 = x_2\}$.)

The inside-out Ehrhart quasipolynomials for some magic and latin squares have striking symmetries (coefficients alternate in sign, the polynomials factor nicely, etc.). Explain.

The inside-out Ehrhart quasipolynomials for some magic and latin squares have much lower periods than predicted by their denominators. Explain.

Compute $\text{Mag}_4, \text{Mag}_5, \dots$ (possibly using LattE and the Möbius function of the intersection lattice of $\mathcal{H}[K_d]$).

Antimagic

$f_1, \dots, f_m \in (\mathbb{R}^d)^*$ – linear forms

$A^\circ(t) := \#$ integer points $x \in (0, t)^d$ such that

$$f_j(x) \neq f_k(x) \quad \text{if } j \neq k$$

Antimagic

$f_1, \dots, f_m \in (\mathbb{R}^d)^*$ – linear forms

$A^\circ(t) := \#$ integer points $x \in (0, t)^d$ such that

$$f_j(x) \neq f_k(x) \quad \text{if } j \neq k$$

Inside-out interpretation: $f(x) := (f_1, \dots, f_m)(x) \notin \bigcup \mathcal{H}[K_m] \subseteq \mathbb{R}^m$

Pullback $\mathcal{H}[K_m]^\# \subseteq \mathbb{R}^d$ obtained from $f^{-1}(h)$ for all $h \in \mathcal{H}[K_m]$

Antimagic : $x \in \mathbb{R}^d \setminus \bigcup \mathcal{H}[K_m]^\#$

Antimagic

$f_1, \dots, f_m \in (\mathbb{R}^d)^*$ – linear forms

$A^\circ(t) := \#$ integer points $x \in (0, t)^d$ such that

$$f_j(x) \neq f_k(x) \quad \text{if } j \neq k$$

Inside-out interpretation: $f(x) := (f_1, \dots, f_m)(x) \notin \bigcup \mathcal{H}[K_m] \subseteq \mathbb{R}^m$

Pullback $\mathcal{H}[K_m]^\# \subseteq \mathbb{R}^d$ obtained from $f^{-1}(h)$ for all $h \in \mathcal{H}[K_m]$

Antimagic : $x \in \mathbb{R}^d \setminus \bigcup \mathcal{H}[K_m]^\#$

Examples : antimagic graphs and relatives (bidirected antimagic graphs, node antimagic, total graphical antimagic), antimagic squares, cubes, etc.

Open problems

Is there a combinatorial interpretation of the regions of $\mathcal{H}[K_m]^\#$?

Open problems

Is there a combinatorial interpretation of the regions of $\mathcal{H}[K_m]^\#$?

What is the intersection-lattice structure of $\mathcal{H}[K_m]^\#$?

Open problems

Is there a combinatorial interpretation of the regions of $\mathcal{H}[K_m]^\#$?

What is the intersection-lattice structure of $\mathcal{H}[K_m]^\#$?

Prove that every graph except K_2 is (strongly) antimagic, i.e., admits an antimagic labelling using the numbers $1, 2, \dots, |E|$.

Open problems

Is there a combinatorial interpretation of the regions of $\mathcal{H}[K_m]^\#$?

What is the intersection-lattice structure of $\mathcal{H}[K_m]^\#$?

Prove that every graph except K_2 is (strongly) antimagic, i.e., admits an antimagic labelling using the numbers $1, 2, \dots, |E|$.

If that's too hard, try trees.