Algebraic and Differential Topology – Lecture Notes

Judith Brinkschulte

Winter Semester 2026/2027

Contents

Index **1**

1 Introduction **3**

2 Homotopy **5**

This script was written by a student and is therefore prone to error. Please direct all typographical errors to the e-mail "hc75nigi@studserv.uni-leipzig.de". For content errors or questions to the content, please ask your lecturer directly.
This is the raw version of the script. Changes to the Layout will happen over time.

1 Introduction

Definition 1.1. A topology on a set X is a collection τ of subsets of X such that:

(i) $\varnothing, X \in \tau$.

(ii) If $U_i \in \tau$ for all $i \in I$, then $\bigcup_{i \in I} U_i \in \tau$.

(iii) If $U_i \in \tau$ for all $i \in I$ where I is finite, then $\bigcap_{i \in I} U_i \in \tau$.

We call the pair (X, τ) a topological space. The elements of τ are called open sets.

Definition 1.2. A **basis** for a topology on X is a collection $\mathcal{B}$ of subsets of X such that:

(i) $\forall x \in X$, $\exists B \in \mathcal{B}$ containing x.

(ii) If $x \in B_1 \cap B_2$ for some $B_1, B_2 \in \mathcal{B}$, then there is a basis element $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

$\mathcal{B}$ generates a topology τ on X as follows:
$U \in \tau$ if $\forall x \in U$, $\exists B \in \mathcal{B}$ such that $x \in B \subseteq U$.

Remark 1.3. It is important that $B \in \mathcal{B}$ is a subset of U in the last statement. If we only required $x \in B \cap U$, we would run into problems. For example consider $X = \mathbb{R}$ and $U = \{0, 1\}$. We know that $\mathcal{B} = \{(a, b) : a, b \in \mathbb{R}, a < b\}$ is a basis for the standard topology on $\mathbb{R}$. Then for $x = 0$, we can choose $B = (-1, 1)$ and for $x = 1$, we can choose $B = (0, 2)$. This would mean that U is open, which should not be the case.

Example 1.4.

- (M, d) is a metric space. Then $\mathcal{B} = \{B_r(x) : x \in M, r > 0\}$ is a basis for the metric topology on M, where $B_r(x) = \{y \in M : d(x, y) < r\}$ is the open ball of radius r around x.
- X, Y topological spaces. The product topology on $X \times Y$ is generated by the basis $\mathcal{B} = \{U \times V : U \subseteq X, V \subseteq Y \text{ open}\}$.
- subspace topology: Let (X, τ) be a topological space, $S \subseteq X$. Then the subspace topology of S is $\tau_S = \{U \cap S : U \in \tau\}$.

Definition 1.5. U is a neighbourhood of x if $\exists V$ open such that $x \in V \subseteq U$.

Definition 1.6. Let X, Y be topological spaces.
A map $f : X \to Y$ is continuous if $\forall U \subseteq Y$ open, $f^{-1}(U)$ is open in X.
If f is a bijection and both f and f^{-1} are continuous, then f is called a homeomorphism. In this case, we say that X and Y are homeomorphic or topologically equivalent ($X \simeq Y$).

Example 1.7.

- $f : [0,1] \to \mathbb{S}^1$ given by $f(t) = (cos(2\pi t), sin(2\pi t))$.
  f is continuous and bijective but f^{-1} is not continuous.
  $f([0,1))$ is not open in $\mathbb{S}^1$.

- $\mathbb{R} \not\cong \mathbb{R}^2$: Assume they are homeomorphic. Then there exists a homeomorphism $f : \mathbb{R} \to \mathbb{R}^2$ with $f(0) = (a,b)$. But $f|_{\mathbb{R}\setminus\{0\}} : \mathbb{R} \setminus \{0\} \to \mathbb{R}^2 \setminus \{(a,b)\}$ is a homeomorphism. But $\mathbb{R} \setminus \{0\}$ is not connected, while $\mathbb{R}^2 \setminus \{(a,b)\}$ is connected. This is a contradiction.

- $\mathbb{R}^2 \not\cong \mathbb{R}^3$: Assume they are diffeomorphic. Then there exists a diffeomorphism $f : \mathbb{R}^2 \to \mathbb{R}^3$. Then $f \circ f^{-1} = id_{\mathbb{R}^3}, f^{-1} \circ f = id_{\mathbb{R}^2}$ so $Df_x Df_y^{-1} = I_3$ and $Df_y^{-1} Df_x = I_2$.

2 Homotopy

Definition 2.1. Let X, Y be topological spaces and $f, g : X \to Y$ be continuous maps. Then f is called homotopic to g if there exists a $F : X \times I \to Y$ continuous s.t. $F(x,0) = f(x)$ and $F(x,1) = g(x)$ for all $x \in X$. We write $f \simeq_F g$ and F is called a homotopy between f and g. If $f \simeq c$ where c is a constant map, then f is called null-homotopic.

Definition 2.2. Let X be a topological space. A continuous map $f : I \to X$ is called a path in X.
Two paths $f, g : I \to X$ are called **path homotopic** if they have the same start and end points, i.e. $f(0) = g(0)$ and $f(1) = g(1)$, and there exists a continuous map $F : I \times I \to X$ s.t. $F(x,0) = f(x)$, $F(x,1) = g(x)$ for all $x \in I$, and $F(0,t) = f(0) = g(0)$, $F(1,t) = f(1) = g(1)$ for all $t \in I$. We write $f \simeq_F g$.

Remark 2.3. This is an equivalence relation:

(i) $f \simeq_F f$ with $F(x,t) = f(x)$ for all $x, t \in I$.

(ii) $f \simeq_F g \implies g \simeq_G f$ with $G(x,t) := F(x, 1-t)$.

(iii) $f \simeq_F g$, $g \simeq_G h \implies f \simeq_H h$ with $\begin{cases} H(x,t) = F(x,2t) & 0 \le t \le \frac{1}{2} \\ H(x,t) = G(x,2t-1) & \frac{1}{2} \le t \le 1 \end{cases}$

If f is a path then we call $[f]$ the path homotopy equivalence class of f.

Example 2.4.

- Any two continuous maps $f, g : X \to \mathbb{R}^2$ are homotopic with $F(x,t) = (1-t)f(x) + tg(x)$.

- Let $X = \mathbb{R}^2 \setminus \{(0,0)\}$ and let $f, g : I \to X$ be defined by $f(t) = e^{\pi i t}$ and $g(t) = e^{-\pi i t}$. Then f and g are not path homotopic in X. Proving this is not trivial. First, we state an equivalent problem which is slightly easier to prove. Instead of two paths, we try to show that the unit circle as a path $h(t) = e^{2\pi i t}$ is not path homotopic to a constant path (w.l.o.g. $(1,0)$).

  Assume there is a path homotopy $F : I \times I \to X$ such that $F(x,0) = h(x)$ and $F(x,1) = (1,0)$. W.l.o.g. we may assume $F : I \times I \to \mathbb{S}^1$: if F' is a path homotopy, then $F := \frac{x}{|x|} \circ F'$ is one too, because $\frac{x}{|x|}$ is continuous on X.

  From here on we use the covering map (which we will define later) $p : \mathbb{R} \to \mathbb{S}^1$, $p(t) = e^{2\pi i t}$. By a lemma which we will prove later, F has a lift $\tilde{F}$, i.e. a continuous map $\tilde{F} : I \times I \to \mathbb{R}$ with $p \circ \tilde{F} = F$. We can normalize $\tilde{F}$ such that $\tilde{F}(0,0) = 0$.

  Because $F(0,t) = F(1,t) = F(x,1) = (1,0)$ are all mapped to 1 (we use $\mathbb{R}^2 \cong \mathbb{C}$ for convenience) and $p^{-1}(1) = \mathbb{Z}$, the maps $\tilde{F}(0,t)$, $\tilde{F}(1,t)$ and $\tilde{F}(x,1)$

are continuous with values in $\mathbb{Z}$, and $\mathbb{Z}$ is discrete. Hence each of them is constant. Since consecutive edges share a corner, the three constants agree, and we get

$$\tilde{F}(0,0) = \tilde{F}(0,1) = \tilde{F}(1,1) = \tilde{F}(1,0) = 0.$$

We have now determined every edge except the bottom one. On the bottom edge, both $x \mapsto \tilde{F}(x,0)$ and $x \mapsto x$ are continuous lifts of h. We see that

$$p\big(\tilde{F}(x,0) - x\big) = \frac{h(x)}{h(x)} = 1,$$

so the difference lies in $\mathbb{Z}$. Since the difference is continuous on the connected set I, it is constant. At $x = 0$ the difference is 0, hence $\tilde{F}(x,0) = x$. But this is a contradiction, because $\tilde{F}(1,0) = 0 \neq 1$.