

LOWER BOUNDS ON BOURGAIN'S CONSTANT FOR HARMONIC MEASURE

MATTHEW BADGER AND ALYSSA GENSCHAW

ABSTRACT. For every $n \geq 2$, Bourgain's constant b_n is the largest number such that the (upper) Hausdorff dimension of harmonic measure is at most $n - b_n$ for every domain in \mathbb{R}^n on which harmonic measure is defined. Jones and Wolff (1988) proved that $b_2 = 1$. When $n \geq 3$, Bourgain (1987) proved that $b_n > 0$ and Wolff (1995) produced examples showing $b_n < 1$. Refining Bourgain's original outline, we prove that

$$b_n \geq c n^{-2n(n-1)/\ln(n)}$$

for all $n \geq 3$, where $c > 0$ is a constant that is independent of n . We further estimate $b_3 \geq 1 \times 10^{-15}$ and $b_4 \geq 2 \times 10^{-26}$.

1. INTRODUCTION

An outstanding problem on the boundary behavior of harmonic functions in space is to identify the maximal *minimal dimension* of a subset of the boundary of a domain through which Brownian motion first exits the domain almost surely [Bis07]. Adopting the parlance of geometric measure theory [Fal97, MMR00], one would like to identify the largest possible (*upper*) *Hausdorff dimension* $\overline{\dim}_H \omega$ of harmonic measure $\omega = \omega_\Omega$ across all connected, open sets $\Omega \subsetneq \mathbb{R}^n$. For planar domains ($n = 2$), the Hausdorff dimension of harmonic measure is at most 1 [JW88] (also see [Øks81, KW85, Car85, Mak85, Wol93]) and this is sharp (e.g. when Ω is a disk). For space domains ($n \geq 3$), the Hausdorff dimension of harmonic measure is at most $n - b_n$ for some undetermined value $0 < b_n < 1$ [Bou87, Wol95] (also see [LVV05]), which we call *Bourgain's constant*. Determining the value of Bourgain's constant is related to understanding certain physical phenomenon (passivation, fouling, poisoning) [FGAS08]. While a conjectural best value $1/(n - 1)$ for b_n has been proposed [Bis92], there appears not to have been any progress in affirming or disproving this conjecture to date.

Very recently, the authors proved an analogue of Bourgain's theorem in the setting of the heat equation [BG21]. In particular, the Hausdorff dimension of caloric measure on any domain in $\mathbb{R}^n \times \mathbb{R}$ (space \times time) with the parabolic distance is at most $n + 2 - \beta_n$ for some $\beta_n > 0$; moreover, $\beta_n \leq b_n$ for all n . In the process of establishing this extension, it was discovered that the demonstration of $b_3 > 0$ in [Bou87] contains an incorrect

Date: May 30, 2022.

2020 Mathematics Subject Classification. Primary 31B15, Secondary 28A75, 31B25, 60J65.

Key words and phrases. harmonic measure, Hausdorff dimension, Bourgain's estimate, Frostman's lemma, net measures.

estimate, leaving a gap in the original proof; see [BG21, Remark 5.4] for a description of the flaw and for a proposed correction. The goal of this note is to record a complete proof that $b_n > 0$ when $n \geq 3$. Further, by refining Bourgain's original outline, we provide quantitative lower bounds on b_n for arbitrary $n \geq 3$ and the first explicit lower bounds on b_n in dimensions $n = 3, 4$.

Theorem 1.1. *There exists $c > 0$ such that $b_n \geq c n^{-2n(n-1)} / \ln(n)$ for all $n \geq 3$.*

Theorem 1.2 (15 decimals, 26 decimals). *We have $b_3 \geq 1 \times 10^{-15}$ and $b_4 \geq 2 \times 10^{-26}$.*

Bourgain's underlying idea to prove estimates of this type is that if the boundary of a domain is spread uniformly in space throughout a cube, then the probability that a Brownian traveler exits the domain near the center of the cube should be smaller than the probability of exiting the domain near the boundary of the cube. This sets up an alternative: at every m -adic location, for some integer $m \gg_n 3$, either the boundary has small $(n - \rho)$ -dimensional content for some $0 < \rho \ll_{m,n} 1$ or the harmonic measure has positive $(n - \lambda)$ -dimensional density on a subset of positive measure for some $0 < \lambda \ll_{m,n} 1$. It follows that $\overline{\dim}_H \omega \leq n - \rho\lambda/(\rho + \lambda)$; see Lemma 2.5 below. This approach exhibits strong dimension dependence in the form of volume concentration near the boundary of balls or cubes in high dimensions. The arguments in [Bou87] and [BG21] use continuity and soft analysis to assert the existence of the parameters m , ρ , and λ to conclude that $b_n > 0$ and $\beta_n > 0$. A challenge in proving Theorem 1.1 is to carry out explicit estimates wherever feasible. For Theorem 1.2, one need only track estimates for a small set of values of m , because $\rho \rightarrow 0$ rapidly as $m \rightarrow \infty$.

In §2, we introduce necessary elements from geometric measure theory, including a tight version of Frostman's lemma for sets in \mathbb{R}^n of Hausdorff dimension $s > n - 1$. In §3, we revisit and sharpen Bourgain's estimate on harmonic measure from below in terms of the relative size of the local m -adic net content. We also record some basic estimates on harmonic measure inside nested rectangles. In §4, we reduce the problem of finding lower bounds on b_n to showing the existence of an admissible set of parameters. Using all of these ingredients, we prove Theorem 1.1 in §5 and Theorem 1.2 in §6.

Acknowledgements. The authors thank Max Engelstein for bringing the work [FGAS08] by Filoche *et al.* to their attention.

2. NET MEASURES, HAUSDORFF DIMENSION, AND FROSTMAN'S LEMMA

Background on Hausdorff measures and related topics can be found in [Mat95, Rog98]. Let \mathbb{R}^n be equipped with the standard Euclidean distance, i.e. $|X - Y| = (\sum_1^n (x_i - y_i)^2)^{1/2}$ for all $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$. We let $\text{diam } E = \sup\{|X - Y| : X, Y \in E\}$ denote the *diameter* of $E \subset \mathbb{R}^n$. For each integer $m \geq 2$, we define the system $\Delta^m(\mathbb{R}^n)$ of *half-open m -adic cubes* to be all sets Q of the form

$$Q = \left[\frac{j_1}{m^k}, \frac{j_1 + 1}{m^k} \right) \times \cdots \times \left[\frac{j_n}{m^k}, \frac{j_n + 1}{m^k} \right) \quad (k, j_1, \dots, j_n \in \mathbb{Z});$$

we say that Q belongs to *generation k* of $\Delta^m(\mathbb{R}^n)$ and has *side length* $\text{side } Q = m^{-k}$ and *volume* $\text{vol } Q = m^{-kn}$. Note that $\text{diam } Q = \sqrt{n} \text{side } Q$ for every $Q \in \Delta^m(\mathbb{R}^n)$.

The cubes in each generation of $\Delta^m(\mathbb{R}^n)$ partition \mathbb{R}^n . Every cube $Q \in \Delta^m(\mathbb{R}^n)$ of generation k is contained in a unique cube $Q^\uparrow \in \Delta^m(\mathbb{R}^n)$ of generation $k-1$; we call Q^\uparrow the *parent* of Q and call Q a *child* of Q^\uparrow . Extending this metaphor, we may refer to the ancestors “above” a cube and descendents “below” a cube. For every $Q \in \Delta^m(\mathbb{R}^n)$, $\# \text{Child}(Q) = m^n$ and $\text{vol } Q = \sum_{R \in \text{Child}(Q)} \text{vol } R$, where $\text{Child}(Q)$ is the set of children.

Definition 2.1. Fix $\Delta = \Delta^m(\mathbb{R}^n)$ and let $s \in [0, \infty)$. For all $\delta \in (0, \infty]$, we define

$$\mathcal{M}_\delta^s(E) = \inf \left\{ \sum_1^\infty (\text{side } E_i)^s : E \subseteq \bigcup_1^\infty E_i, \text{side } E_i \leq \delta, E_i \in \Delta \right\} \quad \text{for all } E \subset \mathbb{R}^n;$$

we call \mathcal{M}_∞^s the *s-dimensional net content* on \mathbb{R}^n . We define the *s-dimensional net measure* on \mathbb{R}^n by $\mathcal{M}^s(E) = \lim_{\delta \downarrow 0} \mathcal{M}_\delta^s(E)$ for all $E \subset \mathbb{R}^n$.

Remark 2.2. The net contents \mathcal{M}_∞^s are outer measures on \mathbb{R}^n and the net measures \mathcal{M}^s are Borel regular outer measures on \mathbb{R}^n . Unlike the Hausdorff measures \mathcal{H}^s and contents \mathcal{H}_∞^s , the net measures and net contents are neither translation nor dilation invariant. Of course, for all $n \geq 1$ and $s > 0$, we have $\mathcal{H}^s \lesssim_n \mathcal{M}^s \lesssim_{n,m} \mathcal{H}^s$ and $\mathcal{H}_\infty^s \lesssim_n \mathcal{M}_\infty^s \lesssim_{n,m} \mathcal{H}_\infty^s$. This allows us to define Hausdorff dimension using net measures in lieu of Hausdorff measures. While the net measures and net contents depend on the choice of the underlying grid $\Delta = \Delta^m(\mathbb{R}^n)$, for simplicity we suppress this from the notation. We choose the letter \mathcal{M} to suggest *m*-adic. An elementary fact is that $\mathcal{M}^s(E) = 0$ if and only if $\mathcal{M}_\infty^s(E) = 0$.

Definition 2.3. Let $E \subset \mathbb{R}^n$. The *Hausdorff dimension* of E is the unique number $\dim_H E \in [0, n]$ where one witnesses a transition from $\mathcal{M}^s(E) = \infty$ for all $s < \dim_H E$ to $\mathcal{M}^s(E) = 0$ for all $s > \dim_H E$.

Definition 2.4. Let μ be a Borel measure on \mathbb{R}^n . The *upper Hausdorff dimension* of μ is defined to be $\overline{\dim}_H \mu = \inf \{ \dim_H E : E \subset \mathbb{R}^n \text{ is Borel, } \mu(\mathbb{R}^n \setminus E) = 0 \}$.

Lemma 2.5. Fix $\Delta = \Delta^m(\mathbb{R}^n)$. Let μ be a Radon measure on \mathbb{R}^n and let $E \subset \mathbb{R}^n$ be a Borel set with $\mu(\mathbb{R}^n \setminus E) = 0$. If there exist constants $0 < \rho < n$ and $\lambda > 0$ such that for every $Q \in \Delta$ with $\text{side } Q \leq 1$,

$$(2.1) \quad \mathcal{M}_{m^{-1} \text{side } Q}^{n-\rho}(E \cap Q) < (\text{side } Q)^{n-\rho}$$

or

$$(2.2) \quad \sum_{R \in \text{Child}(Q)} \mu(R)^{1/2} (\text{vol } R)^{1/2} \leq m^{-\lambda} \mu(Q)^{1/2} (\text{vol } Q)^{1/2},$$

then $\overline{\dim}_H \mu \leq n - \lambda\rho/(\lambda + \rho)$.

Proof. A version of the lemma with the weaker conclusion $\overline{\dim}_H \mu < n$ appears in [Bou87]. The authors supplied a detailed proof with the indicated upper bound, in the setting of $\mathbb{R}^n \times \mathbb{R}$, equipped with the parabolic distance. Repeat the proof of [BG21, Theorem 2.10],

making the following superficial change. Replace each occurrence of $n + 2$ (the Hausdorff dimension of parabolic $\mathbb{R}^n \times \mathbb{R}$) with n (the Hausdorff dimension of \mathbb{R}^n). \square

Lemma 2.6 (Frostman's lemma with better constant). *Fix $\Delta = \Delta^m(\mathbb{R}^n)$. Let $K \subset \mathbb{R}^n$ be a compact set. If $s > n - 1$, then there exists a Radon measure μ on \mathbb{R}^n such that $\mu(Q) \leq (\text{side } Q)^s$ for all $Q \in \Delta$, $\mu(\partial Q) = 0$ for all $Q \in \Delta$, and $\mu(\mathbb{R}^n) = \mu(K) \geq \mathcal{M}_\infty^s(K)$.*

Remark. The “constant” in front of $\mathcal{M}_\infty^s(K)$ is 1.

Proof. Let $K \subset \mathbb{R}^n$ be compact and fix $s > n - 1$. We will modify a standard proof of Frostman's lemma, exploiting the fact that s is greater than the Hausdorff dimension of the boundaries of m -adic cubes. Cover K with a finite list of cubes $Q_1, \dots, Q_l \in \Delta$ with the property that each pair Q_i and Q_j have no common ancestors unless $i = j$. Then $\mathcal{M}_\infty^s(K) = \sum_{i=1}^l \mathcal{M}_\infty^s(K \cap Q_i)$. Suppose that for each i , we can construct a Radon measure μ_i on \mathbb{R}^n such that $\mu_i(Q) \leq (\text{side } Q)^s$ for all $Q \in \Delta$, $\mu_i(\partial Q) = 0$ for all $Q \in \Delta$, and $\mu_i(\mathbb{R}^n) = \mu_i(K \cap \overline{Q_i}) \geq \mathcal{M}_\infty^s(K \cap Q_i)$. (See the next paragraph.) Then $\mu = \mu_1 + \dots + \mu_l$ is our desired measure, because each μ_i vanishes on the boundaries of cubes in Δ and pairwise Q_1, \dots, Q_l have no common ancestors.

Fix a cube $Q_0 \in \Delta$ and assign $E := K \cap Q_0 \subset Q_0$. Following the proof of [Mat95, Theorem 8.8], using m -adic cubes instead of dyadic cubes, one may produce a sequence of Radon measure $(\nu_k)_{k=1}^\infty$ such that

- (i) the support of ν_k belongs to the closure of $\bigcup\{Q \in \Delta : \text{side } Q = m^{-k}, Q \cap E \neq \emptyset\}$;
- (ii) $\nu_k(\partial Q) = 0$ for all $Q \in \Delta$;
- (iii) $\nu_k(Q) \leq (\text{side } Q)^s$ for all $Q \in \Delta$ with $\text{side } Q \geq m^{-k}$; and,
- (iv) for each $X \in E$, there is $Q^X \in \Delta$ with $\text{side } Q^X \geq m^{-k}$ and $\nu_k(Q^X) = (\text{side } Q^X)^s$.

By (i), (ii), and (iii), $\nu_k(\mathbb{R}^n) = \nu_k(\overline{Q_0}) = \nu_k(Q_0) \leq (\text{side } Q_0)^s < \infty$ for all k that are large enough so that $\text{side } Q_0 \geq m^{-k}$. By weak compactness of Radon measures, there exists a subsequence $(\nu_{k_j})_{j=1}^\infty$ and a Radon measure ν such that ν_{k_j} converges weakly to ν as $j \rightarrow \infty$ in the sense of Radon measures. By (i), the support of ν is contained in $K \cap \overline{Q_0}$. By (iv), for each k , there exist (maximal) disjoint cubes Q^{X_1}, \dots, Q^{X_p} such that $E \subset Q^{X_1} \cup \dots \cup Q^{X_p}$ and $\nu_k(\mathbb{R}^n) \geq \sum_{i=1}^p \nu_k(Q^{X_i}) = \sum_{i=1}^p (\text{side } Q^{X_i})^s \geq \mathcal{M}_\infty^s(E)$. Thus,

$$\nu(\mathbb{R}^n) = \nu(\overline{Q_0}) \geq \limsup_{j \rightarrow \infty} \nu_{k_j}(\overline{Q_0}) = \limsup_{j \rightarrow \infty} \nu_{k_j}(\mathbb{R}^n) \geq \mathcal{M}_\infty^s(E).$$

Next, let $Q \in \Delta$. For all $i \geq 1$, a sufficiently small open neighborhood of ∂Q can be covered by $C(n)m^{i(n-1)}$ cubes $R \in \Delta$ with $\text{side } R = m^{-i} \text{side } Q$. By (iii), it follows that

$$\nu(\partial Q) \leq \liminf_{j \rightarrow \infty} \sum_R \nu_{k_j}(R) \leq C(n)m^{i(n-1-s)}(\text{side } Q)^s \quad \text{for all } i \geq 1.$$

Hence $\nu(\partial Q) = 0$, since $s > n - 1$. Consequently, $\nu(Q) = \lim_{j \rightarrow \infty} \nu_{k_j}(Q) \leq (\text{side } Q)^s$. \square

3. ESTIMATES FOR HARMONIC MEASURE

Harmonic measure is perhaps best viewed through several complementary perspectives, including geometric function theory see [GM05], potential theory [Hel09], and stochastic

processes [MP10]. On an unbounded domain $\Omega \subset \mathbb{R}^n$, we let $\partial\Omega$ denote the topological boundary in a one-point compactification of Ω so that $\partial\Omega$ includes the point at infinity. This ensures that for any domain $\Omega \subsetneq \mathbb{R}^n$, $n \geq 3$, harmonic measure ω_Ω^X with pole at $X \in \Omega$ exists and is a Borel probability measure with support in $\partial\Omega$. By Harnack's inequality, ω_Ω^X and ω_Ω^Y are mutually absolutely continuous for all $X, Y \in \Omega$. In particular, the support and Hausdorff dimension of ω_Ω^X are independent of the choice of X .

Given an m -adic grid $\Delta = \Delta^m(\mathbb{R}^n)$, we let

$$\vec{\Delta} := \{Q + m^{-1} \text{side } Q \cdot (j_1, \dots, j_n) : Q \in \Delta, j_1, \dots, j_n \in \mathbb{Z}\}$$

denote the set of translates of m -adic cubes that are aligned with m -adic cubes of the next generation. The following lemma is modeled after [Bou87, Lemma 1]. Results of this type are now collectively referred to as *Bourgain's estimate* and have become a fundamental tool used to study absolute continuity of harmonic measure (see e.g. [HM14, AHM⁺16]). Also see [BZ15, Azz20] for applications of Bourgain's estimate to study the dimension of harmonic measure on special classes of domains.

Lemma 3.1 (Bourgain's estimate with better constants). *Fix $\Delta = \Delta^m(\mathbb{R}^n)$ for some $n \geq 3$ and $m \geq 5$ with $m > \xi_m + 2\sqrt{n}$, where we define $\xi_m := 1$, when m is odd, and $\xi_m := 2$, when m is even. Let $P \in \vec{\Delta}$ and $P_* \in \Delta$ be any cubes such that $\text{side } P_* = m^{-1} \text{side } P$ and $\overline{P_*}$ includes the center of P . Assign $Q := \text{int } P$ and $Q_* := \overline{P_*}$. For all closed sets $E \subset \mathbb{R}^n$, poles $X \in Q_* \setminus E$, and dimensions $n - 1 < s \leq n$,*

$$(3.1) \quad \left(\left(\frac{1}{\sqrt{n}} \right)^{n-2} - \left(\frac{2}{m - \xi_m} \right)^{n-2} \right) \frac{\mathcal{M}_\infty^s(E \cap Q_*)}{(\text{side } Q_*)^s} \leq O(n, m, s) \omega_{Q \setminus E}^X(E \cap Q),$$

where

$$(3.2) \quad O(n, m, s) = \min_{k \in \mathbb{Z}} \left(\left(\frac{2m^k}{(n-2)\sqrt{n}} \right)^{n-2} + \frac{\omega_n}{4} \left(\frac{mn}{n-2} \right)^{n-2} \frac{n^3 m^{k(n-2-s)}}{1 - m^{n-2-s}} \right).$$

Remark. Here and below, we let ω_n denote the volume of the unit ball in \mathbb{R}^n . When m is odd, the cube P_* is uniquely determined by P . When m is even, there are 2^n possibilities for P_* for each P . The gap (distance) between P_* and ∂P is $(m - \xi_m)/2$ times $\text{side } P_*$.

Proof. Let n, m, Q , and Q_* be given with the stated requirements. Let $E \subset \mathbb{R}^n$ be closed. Because Q_* is closed, $K := E \cap Q_*$ is compact. Freeze $n - 1 < s \leq n$. By Lemma 2.6, there exists a finite Borel measure μ on \mathbb{R}^n with support in K such that $\mu(R) \leq (\text{side } R)^s$ for all $R \in \Delta$, $\mu(\partial R) = 0$ for all $R \in \Delta$, and $\mu(\mathbb{R}^n) = \mu(K) \geq \mathcal{M}_\infty^s(K)$. Consider the harmonic function $u(X) = \int_K |X - Y|^{-(n-2)} d\mu(Y)$ defined on $\mathbb{R}^n \setminus K$, which satisfies

$$(E1) \quad u(X) \leq O(n, m, s) (\text{side } Q_*)^{s-(n-2)} \quad \forall X \in \mathbb{R}^n \setminus K,$$

$$(E2) \quad u(X) \geq (\text{diam } Q_*)^{-(n-2)} \mu(K) = \sqrt{n}^{-(n-2)} (\text{side } Q_*)^{-(n-2)} \mu(K) \quad \forall X \in Q_* \setminus K,$$

$$(E3) \quad u(X) \leq (\text{gap}(Q_*, \partial Q))^{-(n-2)} \mu(K) = \left(\frac{m - \xi_m}{2} \text{side } Q_* \right)^{-(n-2)} \mu(K) \quad \forall X \in \partial Q,$$

where $\text{gap}(A, B) = \inf_{a \in A} \inf_{b \in B} |a - b|$ denotes the *gap* between nonempty sets A and B (sometimes referred to as the *distance* between A and B). Of the three estimates, (E2) and (E3) are straightforward; we delay the proof of (E1) to the end of the lemma. Following [Bou87], define an auxiliary harmonic function w on $\mathbb{R}^n \setminus K$ by setting

$$w(X) = \frac{u(X) - \|u\|_{L^\infty(\partial Q)}}{\|u\|_{L^\infty(\mathbb{R}^n \setminus K)}} \quad \text{for all } X \in \mathbb{R}^n \setminus K.$$

By design, $w \leq 0$ on ∂Q , w is continuous on ∂Q , and $w \leq 1$ on all of $\mathbb{R}^n \setminus K$. Hence

$$\limsup_{X \rightarrow X_0} w(X) \leq \chi_K(X_0) \quad \text{for all } X_0 \in \partial(Q \setminus K).$$

Thus, $w(X) \leq \omega_{Q \setminus K}^X(K) \leq \omega_{Q \setminus E}^X(E \cap Q)$ for every $X \in Q \setminus E$ by two applications of the maximum principle. Suppose that $X \in Q_* \setminus E = Q_* \setminus K$. Using (E1) to estimate the denominator in the definition of w and (E2), (E3) to estimate the numerator, we obtain

$$\omega_{Q \setminus E}^X(Q \cap E) \geq \frac{((1/\sqrt{n})^{n-2} - (2/(m - \xi_m))^{n-2}) (\text{side } Q_*)^{-(n-2)} \mathcal{M}_\infty^s(K)}{O(n, m, s) (\text{side } Q_*)^{s-(n-2)}}.$$

This is our desired estimate.

It remains to verify (E1), which is a novel contribution. Fix $X \in \mathbb{R}^n \setminus K$ and let j_0 denote the integer such that $\text{side } Q_* = m^{-j_0}$. Recall that $\mu(R) \leq (\text{side } R)^s$ for all $R \in \Delta$. Let $l \in \mathbb{Z}$ and let $r_l > 0$ (depending on l) to be determined. Let $h \in \mathbb{Z}$ and expand $K = B_{j_0+h}^l \cup \bigcup_{j=j_0+h}^\infty A_j^l$, where

$$B_j^l := \{Y \in K : |X - Y| \geq r_l m^{-j}\}, \quad A_j^l := \{Y \in K : r_l m^{-(j+1)} \leq |X - Y| < r_l m^{-j}\}.$$

On one hand, we may trivially estimate $\mu(B_{j_0+h}^l) \leq \mu(Q_*) \leq m^{-j_0 s}$. On the other hand, let ω_n denote the volume of the unit ball in \mathbb{R}^n . Then A_j^l is covered by $\lfloor \omega_n(r_l m^l + \sqrt{n})^n \rfloor$ or fewer cubes in Δ of side length $m^{-(j+l)}$. (To derive this, let R represent a cube of side length $m^{-(j+l)}$; divide the volume of a ball of radius $r_l m^{-j} + \text{diam } R$ by the volume of R .) Hence $\mu(A_j^l) \leq \omega_n(r_l m^l + \sqrt{n})^n m^{-(j+l)s}$. Therefore,

$$\begin{aligned} u(X) &\leq (1/r_l)^{n-2} m^{(n-2)(j_0+h)} \mu(B_{j_0+h}^l) + \sum_{j=j_0+h}^\infty (1/r_l)^{n-2} m^{(n-2)(j+1)} \mu(A_j^l) \\ &\leq (m^h/r_l)^{n-2} m^{j_0(n-2-s)} + (m/r_l)^{n-2} \omega_n(r_l m^l + \sqrt{n})^n m^{-ls} \sum_{j=j_0+h}^\infty m^{j(n-2-s)} \\ &\leq \underbrace{\left((m^h/r_l)^{n-2} + \omega_n(m/r_l)^{n-2} (r_l m^l + \sqrt{n})^n \frac{m^{-ls+h(n-2-s)}}{1 - m^{n-2-s}} \right)}_I (\text{side } Q_*)^{s-(n-2)}. \end{aligned}$$

The quantity $(m/r_l)^{n-2}(r_l m^l + \sqrt{n})^n$ has a unique critical point and is minimized across all $r_l > 0$ when $r_l = ((n-2)\sqrt{n})/2m^l$. Selecting this value yields

$$\begin{aligned} I &= \left(\frac{2m^{h+l}}{(n-2)\sqrt{n}} \right)^{n-2} + \omega_n \left(\frac{2m^{1+l}}{(n-2)\sqrt{n}} \right)^{n-2} \left(\frac{n\sqrt{n}}{2} \right)^n \frac{m^{-ls+h(n-2-s)}}{1-m^{n-2-s}} \\ &= \left(\frac{2m^{h+l}}{(n-2)\sqrt{n}} \right)^{n-2} + \frac{\omega_n}{4} \left(\frac{mn}{n-2} \right)^{n-2} \frac{n^3 m^{(h+l)(n-2-s)}}{1-m^{n-2-s}}. \end{aligned}$$

Letting $h+l$ range over arbitrary values $k \in \mathbb{Z}$, we arrive at (E1). \square

Corollary 3.2. *For all $n \geq 3$, there exists $C_n > 1$ depending only on n such that for all $m > \xi_m + 2\sqrt{n}$ and $n-1 < s \leq n$,*

$$(3.3) \quad \mathcal{M}_\infty^s(E \cap Q_*) \leq C_n m^{n-2} \omega_{Q \setminus E}^X(E \cap Q) (\text{side } Q_*)^s.$$

Proof. In (3.2), take $k = 0$ or $k = 1$. \square

Corollary 3.3. *For all $\delta > 0$, there exists $n_\delta \geq 3$ such that if n and m are integers with $n \geq n_\delta$ and $m \geq \delta n$, and $n-1/2 < s \leq n$, then*

$$(3.4) \quad \mathcal{M}_\infty^s(E \cap Q_*) \leq \delta(m\sqrt{2\pi e})^{n-2} \omega_{Q \setminus E}^X(E \cap Q) (\text{side } Q_*)^s.$$

Proof. Let $\delta > 0$ be given and fix $\epsilon > 0$ to be specified below. Suppose that n and m are integers with $n \geq 3$ and $m \geq \delta n$. Let us agree to write $c_n \sim d_n$ if c_n is *asymptotic* to d_n as $n \rightarrow \infty$ in the sense that $\lim_{n \rightarrow \infty} c_n/d_n = 1$ and further agree to write $c_n = o(d_n)$ if $\lim_{n \rightarrow \infty} c_n/d_n = 0$. Using Stirling's formula for $\Gamma(x)$ (e.g. [Li07]) and $\omega_n = \pi^{n/2}/\Gamma(\frac{n}{2} + 1)$ (e.g. [ABR01, Appendix A]), we see that

$$\omega_n \sim \frac{1}{\sqrt{\pi n}} \left(\frac{\sqrt{2\pi e}}{\sqrt{n}} \right)^n = \frac{2\sqrt{\pi e}}{\sqrt{n}^3} \left(\frac{\sqrt{2\pi e}}{\sqrt{n}} \right)^{n-2}.$$

By the ‘‘compound interest formula’’, we also have $(n/(n-2))^{n-2} \sim e^2$. Choosing $k = 1$ in (3.2), it follows from the noted asymptotic estimates that for sufficiently large n , depending only on ϵ ,

$$\frac{O(n, m, s)}{(1+\epsilon)e^2} \leq \left(\frac{2m}{n\sqrt{n}} \right)^{n-2} + \frac{\sqrt{\pi e}}{2} \sqrt{n}^3 m^{n-2-s} \left(\frac{m\sqrt{2\pi e}}{\sqrt{n}} \right)^{n-2},$$

where the reader may observe that we also absorbed the factor $(1-m^{n-2-s})^{-1}$ appearing in $O(n, m, s)$ into the error on the left hand side. Because $m \geq \delta n$ and $-2 \leq n-2-s < -3/2$, the factor $\sqrt{n}^3 m^{n-2-s} = o(\max\{\delta^{-2}, \delta^{-3/2}\}) = o(1)$. Also, $(2/n)^{n-2} = o(\sqrt{2\pi e}^{n-2})$. Thus, taking n to be sufficiently large depending only on δ and ϵ , we have

$$(3.5) \quad O(n, m, s) \leq \epsilon(1+\epsilon)e^2 \left(\frac{m\sqrt{2\pi e}}{\sqrt{n}} \right)^{n-2}.$$

Next, because $m \geq \delta n$, we can estimate

$$(3.6) \quad \left(\frac{1}{\sqrt{n}} \right)^{n-2} - \left(\frac{2}{m - \xi_m} \right)^{n-2} \geq (1 + \epsilon)^{-1} \sqrt{n}^{-(n-2)}$$

for all sufficiently large n depending only on ϵ and δ . Combining (3.1), (3.5), and (3.6), we conclude that for all n sufficiently large depending only on ϵ and δ ,

$$\mathcal{M}_\infty^s(E \cap Q_*) \leq \epsilon(1 + \epsilon)^2 e^2 (m \sqrt{2\pi e})^{n-2} \omega_{Q \setminus E}^X(E \cap Q) (\text{side } Q_*)^s.$$

Specifying that $\epsilon(1 + \epsilon)^2 e^2 = \delta$ yields (3.4). \square

Corollary 3.4. *Suppose $n = 3$. For all integers $m \geq 5$ and dimensions $2.999999 \leq s \leq 3$,*

$$(3.7) \quad \left(\frac{1}{\sqrt{3}} - \frac{2}{m - \xi_m} \right) \frac{\mathcal{M}_\infty^s(E \cap Q_*)}{(\text{side } Q_*)^s} \leq \left(\frac{2}{\sqrt{3}} m + 27\pi \frac{m^{-0.999999}}{1 - m^{-1.999999}} \right) \omega_{Q \setminus E}^X(E \cap Q).$$

Proof. In (3.2), take $n = 3$ and $k = 1$. Bound the factor $m^{2-s}/(1 - m^{1-s})$ appearing in $O(3, m, s)$ using the assumption $s \geq 2.999999$. \square

Corollary 3.5. *Suppose $n = 4$. For all integers $m \geq 7$ and dimensions $3.999999 \leq s \leq 4$,*

$$(3.8) \quad \left(\frac{1}{4} - \left(\frac{2}{m - \xi_m} \right)^2 \right) \frac{\mathcal{M}_\infty^s(E \cap Q_*)}{(\text{side } Q_*)^s} \leq \left(\frac{1}{4} m^2 + 32\pi^2 \frac{m^{0.000001}}{1 - m^{-1.999999}} \right) \omega_{Q \setminus E}^X(E \cap Q).$$

Proof. In (3.2), take $n = 4$ and $k = 1$. Bound the factor $m^{4-s}/(1 - m^{2-s})$ appearing in $O(4, m, s)$ using the assumption $s \geq 3.999999$. \square

A closed *rectangle* in \mathbb{R}^n is any set H of the form $H = [x_1, x_1 + s_1] \times \cdots \times [x_n, x_n + s_n]$ with *side lengths* $s_1, \dots, s_n > 0$. Iterating the strong Markov property, we obtain an estimate on harmonic measure of the portion of the boundary lying inside a sequence of nested rectangles. Brownian motion cannot reach the innermost rectangle without passing through the outer rectangles.

Lemma 3.6. *Let $n \geq 3$ and let $\Omega \subsetneq \mathbb{R}^n$ be a domain. Let H_1, \dots, H_k be rectangles in \mathbb{R}^n that are strictly nested in the sense that $H_k \subset \text{int } H_{k-1}$, $H_{k-1} \subset \text{int } H_{k-2}$, \dots , $H_2 \subset \text{int } H_1$. Write $G'_i = \Omega \cap \partial(\Omega \setminus H_i) \subset \partial H_i = G_i$ for each i . If $X \in \Omega \setminus H_1$, then*

$$(3.9) \quad \omega_\Omega^X(H_k) \leq \omega_{\Omega \setminus H_1}^X(G'_1) \left(\sup_{X_1 \in G'_1} \omega_{\Omega \setminus H_2}^{X_1}(G'_2) \right) \cdots \left(\sup_{X_{k-1} \in G'_{k-1}} \omega_{\Omega \setminus H_k}^{X_{k-1}}(G_k) \right).$$

(Except for the final instance, G_k , all instances of a ' G ' in the formula are G'_i .)

Proof. We induct on the number of rectangles. The base case $\omega_\Omega^X(H_1) \leq \omega_{\Omega \setminus H_1}^X(G_1)$ holds by the maximum principle. For the induction step, suppose that the lemma holds with k nested rectangles for some $k \geq 1$. Let H_1, \dots, H_{k+1} be rectangles with $H_{j+1} \subset \text{int } H_j$ for all $1 \leq j \leq k$ and fix $X \in \Omega \setminus H_1$. Note that every $X_1 \in G'_1$ lies outside of H_2 . Thus, the inductive hypothesis applied with H_2, \dots, H_{k+1} guarantees that

$$\omega_\Omega^{X_1}(H_{k+1}) \leq \omega_{\Omega \setminus H_2}^{X_1}(G'_2) \left(\sup_{X_2 \in G'_2} \omega_{\Omega \setminus H_3}^{X_2}(G'_3) \right) \cdots \left(\sup_{X_k \in G'_k} \omega_{\Omega \setminus H_{k+1}}^{X_k}(G_{k+1}) \right).$$

(When $k = 1$, this formula should be read as $\omega_\Omega^{X_1}(H_2) \leq \omega_{\Omega \setminus H_2}^{X_1}(G_2)$.) Since $\Omega \setminus H_1 \subset \Omega$, the strong Markov property (e.g. see [Doo01, p. 117]) ensures that

$$\begin{aligned} \omega_\Omega^X(H_{k+1}) &= \omega_{\Omega \setminus H_1}^X(H_{k+1} \cap \partial\Omega) + \int_{\Omega \cap \partial(\Omega \setminus H_1)} \omega_\Omega^{X_1}(H_{k+1}) d\omega_{\Omega \setminus H_1}^X(X_1) \\ &= \int_{G'_1} \omega_\Omega^{X_1}(H_{k+1}) d\omega_{\Omega \setminus H_1}^X(X_1) \leq \omega_{\Omega \setminus H_1}^X(G'_1) \sup_{X_1 \in G'_1} \omega_\Omega^{X_1}(H_{k+1}), \end{aligned}$$

where $\omega_{\Omega \setminus H_1}^X(H_{k+1} \cap \partial\Omega) = 0$ trivially, since H_{k+1} is contained in the exterior of $\Omega \setminus H_1$. Combining the two displayed equations gives the desired inequality for H_1, \dots, H_{k+1} . \square

A nearly identical argument gives the following dual inequality.

Lemma 3.7. *Let $n \geq 3$ and let $\Omega \subsetneq \mathbb{R}^n$ be a domain. Let H_1, H_2 be rectangles in \mathbb{R}^n with $H_2 \subset \text{int } H_1$. Write $G'_i = \Omega \cap \partial(\Omega \setminus H_i) \subset \partial H_i = G_i$ for all i . If $X \in \Omega \setminus H_1$, then*

$$(3.10) \quad \omega_\Omega^X(H_2) \geq \omega_{\Omega \setminus H_1}^X(G'_1) \left(\inf_{X_1 \in G'_1} \omega_{\Omega \setminus H_2}^{X_1}(G_2) \right).$$

4. BOUNDING BOURGAIN'S CONSTANT FROM BELOW

Recall that Bourgain's constant b_n is the largest value such that the upper Hausdorff dimension of harmonic measure is at most $n - b_n$ for all domains $\Omega \subset \mathbb{R}^n$. The following theorem, based on the (incomplete) demonstration in [Bou87] that $b_3 > 0$, implements the correction identified in [BG21, Remark 5.4]. It reduces the problem of bounding b_n from below to estimation of constants appearing in Bourgain's estimate and selection of parameters m, η, h , and d satisfying the constraint (4.2).

Theorem 4.1. *Let $n \geq 3$ and let $m \geq 5$. Suppose that $\epsilon > 0$ and $\alpha > 0$ are constants such that Bourgain's estimate holds in the sense that for all Q_* and Q , as in Lemma 3.1, for all closed sets $E \subset \mathbb{R}^n$, for all $X \in Q_* \setminus E$, and for all $n - \epsilon < s \leq n$, we have*

$$(4.1) \quad \mathcal{M}_\infty^s(E \cap Q_*) \leq \alpha \omega_{Q \setminus E}^X(E \cap Q) (\text{side } Q_*)^s.$$

Let $\eta > 0$ be any number such that $(2 - m^{-n})\alpha\eta \leq 1 - m^{-n}$. Finally, suppose that $1 \leq h < m/2$ and $d \geq 1$ are integers such that

$$(4.2) \quad \gamma := (1 - (1 - 2h/m)^n)^{1/2} + (1 - 2h/m)^{n/2} \eta^{-1/2} (1 - \eta)^{hm^{d-1}/2} < 1.$$

Then $b_n \geq \lambda\rho/(\lambda + \rho)$, where

$$(4.3) \quad \lambda := -\log_m(\gamma) \quad \text{and} \quad \rho := \min\{\epsilon, 0.914186(1 - \alpha\eta)(1 - m^{-n})m^{-(d+1)n}/\ln(m)\}.$$

In order to prove the theorem, we start with an auxiliary estimate.

Lemma 4.2. *If $n \geq 3$, $m \geq 5$, and $d \geq 1$ are integers and $(2 - m^{-n})a \leq 1 - m^{-n}$, then*

$$(4.4) \quad (m^n - 1)(m^{\rho-n} + m^{2(\rho-n)} + \dots + m^{(d+1)(\rho-n)}) < 1 - am^{(d+1)(\rho-n)}$$

holds for all values of ρ in the range

$$(4.5) \quad 0 \leq \rho \leq 0.914186(1 - a)(1 - m^{-n})m^{-(d+1)n}/\ln(m).$$

Proof. Rewriting (4.4) using the formula for partial geometric series, we want to find $\rho \geq 0$ (bigger is better) such that

$$\frac{(m^\rho - m^{\rho-n})(1 - m^{(d+1)(\rho-n)})}{1 - m^{\rho-n}} < 1 - am^{(d+1)(\rho-n)}.$$

Rearranging, expanding the products, and cancelling like terms, our goal becomes

$$m^\rho(1 - m^{(d+1)(\rho-n)}) + m^{(d+2)(\rho-n)} < 1 - am^{(d+1)(\rho-n)} + am^{(d+2)(\rho-n)}.$$

Set $\rho = \log_m(c)$ with $c \geq 1$, close to 1. Then we would like

$$c - c^{d+2} \underbrace{m^{-(d+1)n}(1 - (1-a)m^{-n} - a/c)}_x < 1.$$

Since $c \geq 1$ and $x > 0$, the inequality $c - c^{d+2}x < 1$ is implied by $c - x < 1$. Hence it suffices to find $c \geq 1$ such that $c - m^{-(d+1)n}(1 - (1-a)m^{-n} - a/c) < 1$. Equivalently,

$$c^2 - (1 + \underbrace{m^{-(d+1)n}(1 - (1-a)m^{-n})}_y) c + \underbrace{am^{-(d+1)n}}_z < 0.$$

Now, $c^2 - (1+y)c + z < 0$ holds at $c = 1$ provided that $z < y$. In our case, we need $(1-a)m^{-n} < 1-a$, which is true since $a < 1$. It follows that we may select c to be any number between 1 and the greater of the two roots of $c^2 - (1+y)c + z = 0$. That is,

$$1 \leq c < \frac{1+y+\sqrt{(1+y)^2-4z}}{2}.$$

Well, $1+(\sqrt{2}-1)(2y-4z) \leq \sqrt{1+2y-4z} < \sqrt{(1+y)^2-4z}$ provided that $0 \leq 2y-4z < 1$. (To verify the first inequality, start by squaring both sides.) In particular, $4z \leq 2y$ as long as $4a \leq 2-2(1-a)m^{-n}$; this holds by our demand that $(2-m^{-n})a \leq 1-m^{-n}$. Hence we can choose

$$\begin{aligned} 1 \leq c &\leq \frac{1+y+1+(\sqrt{2}-1)(2y-4z)}{2} \\ &= 1 + \underbrace{(\sqrt{2}-1/2)m^{-(d+1)n}((1-2(\sqrt{2}-1)a/(\sqrt{2}-1/2)-(1-a)m^{-n}))}_w. \end{aligned}$$

Thus, (4.4) holds if $0 \leq \rho \leq \ln(1+w)/\ln(m)$. Estimating

$$\ln(1+w) \geq w(1-(1/2)w) \geq w(1-(1/2)(\sqrt{2}-1/2) \cdot 5^{-6}) \geq 0.99997w$$

and $1-2(\sqrt{2}-1)a/(\sqrt{2}-1/2)-(1-a)m^{-n} \geq (1-a)(1-m^{-n})$ and checking that $(\sqrt{2}-1/2) \cdot 0.99997 = 0.914186\dots$, we conclude that (4.5) implies (4.4). \square

Proof of Theorem 4.1. Let $n, m, \epsilon, \alpha, \eta, h, d, \gamma, \lambda$, and ρ be given according to the statement of the theorem. Shrinking ρ as needed, we may assume without loss of generality that $\rho < \epsilon$. Let $\Delta = \Delta^m(\mathbb{R}^n)$, let $\Omega \subsetneq \mathbb{R}^n$ be a domain, let $\Omega^c = \mathbb{R}^n \setminus \Omega$, let $X \in \Omega$, and let $\omega = \omega_\Omega^X$. We say (Q, Q_*) is an *admissible pair* if $Q \in \vec{\Delta}$, $Q_* \in \Delta$, $\text{side } Q_* = m^{-1} \text{ side } Q$,

and $\overline{Q_*}$ includes the center of Q . For every admissible pair (Q, Q_*) , the Bourgain type estimate (4.1) with $s = n - \rho$ implies

$$(4.6) \quad \omega_{(\text{int } Q) \setminus \Omega^c}^Z(\Omega^c \cap \text{int } Q) \geq \eta \quad \text{for all } Z \in \overline{Q_*} \setminus \Omega^c,$$

or

$$(4.7) \quad \mathcal{M}_{\infty}^{n-\rho}(\Omega^c \cap Q_*) < \alpha \eta (\text{side } Q_*)^{n-\rho}.$$

To bound $\overline{\dim}_H \omega$ from above by $n - \lambda \rho / (\lambda + \rho)$, we aim to use Lemma 2.5. Because scaling and translating the domain in space and changing the pole has no effect on the Hausdorff dimension of harmonic measure, we may assume without loss of generality that if $P \in \Delta$, $\text{side } P \leq 1$, and $X \in \overline{P}$, then \overline{P} is disjoint from $\partial \Omega$. For any such cube P ,

$$(4.8) \quad \sum_{Q \in \text{Child}(P)} \omega(Q)^{1/2} (\text{vol } Q)^{1/2} = 0 = m^{-\lambda} \omega(P)^{1/2} (\text{vol } P)^{1/2},$$

trivially.

To continue, suppose that $P \in \Delta$ is an m -adic cube with $\text{side } P \leq 1$, for which $X \notin \overline{P}$. For any $j \geq 1$, let $\text{Child}^j(P)$ denote the set of all j -th generation descendents of P in the tree Δ . For example, $\text{Child}^2(P) = \{R \in \Delta : R \subset P, \text{side } R = m^{-2} \text{side } P\}$ is the set of all grandchildren of P . Keeping in mind our goal of checking the hypothesis of Lemma 2.5, we consider two alternatives.

Alternative 1. Suppose that the estimate (4.7) holds for some admissible pair (Q, Q_*) with $Q_* \in \text{Child}^{d+1}(P)$. Let $Q_*^{\uparrow j} \in \Delta$ denote the j -th ancestor of Q_* in Δ . Covering $\Omega^c \cap P$ by $\text{Child}(P) \setminus \{Q_*^{\uparrow d}\}$, $\text{Child}(Q_*^{\uparrow d}) \setminus \{Q_*^{\uparrow d-1}\}, \dots, \text{Child}(Q_*^{\uparrow 1}) \setminus \{Q_*\}$, and $\Omega^c \cap Q_*$, we obtain

$$\mathcal{M}_{m^{-1} \text{side } P}^{n-\rho}(\Omega^c \cap P) < (m^n - 1) ((\text{side } Q_*^{\uparrow d})^{n-\rho} + \dots + (\text{side } Q_*)^{n-\rho}) + \alpha \eta (\text{side } Q_*)^{n-\rho}.$$

Rewriting each side length in terms of $\text{side } P$ and rearranging,

$$\frac{\mathcal{M}_{m^{-1} \text{side } P}^{n-\rho}(\Omega^c \cap P)}{(\text{side } P)^{n-\rho}} \leq (m^n - 1)(m^{-(n-\rho)} + \dots + m^{-(d+1)(n-\rho)}) + \alpha \eta m^{-(d+1)(n-\rho)}.$$

Applying Lemma 4.2 with $a = \alpha \eta$, we conclude that

$$(4.9) \quad \mathcal{M}_{m^{-1} \text{side } P}^{n-\rho}(\Omega^c \cap P) < (\text{side } P)^{n-\rho}.$$

Alternative 2. Suppose that the estimate (4.6) holds for every admissible pair (Q, Q_*) with $Q_* \in \text{Child}^{d+1}(P)$. Partition P into annular rings of d -th generation descendents. Working from the outside to the inside, define $A_0 = \emptyset$, $P_0 = P$,

$$\begin{aligned} A_1 &= \bigcup \{Q \in \text{Child}^d(P) : Q \not\subset A_0, \overline{Q} \cap \partial P_0 \neq \emptyset\}, & P_1 &= P_0 \setminus A_1, \\ A_2 &= \bigcup \{Q \in \text{Child}^d(P) : Q \not\subset A_1, \overline{Q} \cap \partial P_1 \neq \emptyset\}, & P_2 &= P_1 \setminus A_2, \\ &\vdots & &\vdots \\ A_M &= \bigcup \{Q \in \text{Child}^d(P) : Q \not\subset A_{M-1}, \overline{Q} \cap \partial P_{M-1} \neq \emptyset\}, & P_M &= P_{M-1} \setminus A_M, \\ A_{M+1} &= \bigcup \{Q \in \text{Child}^d(P) : Q \not\subset A_M, \overline{Q} \cap \partial P_M \neq \emptyset\} = P_M, & P_{M+1} &= \emptyset, \end{aligned}$$

where $m^d = 2M + 1$, if m is odd, and $m^d = 2M + 2$, if m is even. Next, for each annulus A_i , with $1 \leq i \leq M$, choose a rectangle H_i such that (i) $G_i := \partial H_i \subset A_i$ separates ∂P_{i-1} from ∂P_i and (ii) for any $z \in G_i$, there exists an admissible pair (Q, Q_*) with $Q_* \in \text{Child}^{d+1}(P)$, $z \in Q_*$, $Q \subset A_i$, and $Q \cap A_{i+1} = \emptyset$. There are a continuum of possibilities for each H_i . Further, as in Lemma 3.6, assign $G'_i := \Omega \cap \partial(\Omega \setminus H_i) \subset G_i$ for each i . See Figure 4.1.

Fix any $1 \leq k \leq M$. Later we will choose $k = k(m, h, d)$. Let H_1, \dots, H_k and G_1, \dots, G_k and G'_1, \dots, G'_k be given as above. In addition, by a slight abuse of notation, write $H_{k+1} = \overline{P_k}$ and $G_{k+1} = \partial P_k$. Then $H_{i+1} \subset \text{int } H_i$ for all $1 \leq i \leq k$. Recall that $X \notin \overline{P}$. On the one hand, by Lemma 3.6, the trivial observation $\omega_{\Omega \setminus H_{i+1}}^{X_i}(G'_{i+1}) \leq \omega_{\Omega \setminus H_{i+1}}^{X_i}(G_{i+1})$, and the fact that we are in Alternative 2,

$$\begin{aligned}
 \omega(P_k) &\leq \omega_{\Omega \setminus H_1}^X(G'_1) \prod_{i=1}^k \sup_{X_i \in G'_i \cap \Omega} \omega_{\Omega \setminus H_{i+1}}^{X_i}(G_{i+1}) \\
 (4.10) \quad &\leq \omega_{\Omega \setminus H_1}^X(G'_1) \prod_{i=1}^k \left(1 - \inf_{X_i \in G'_i \cap \Omega} \omega_{\Omega \setminus H_{i+1}}^{X_i}(A_i) \right) \leq \omega_{\Omega \setminus H_1}^X(G'_1) (1 - \eta)^k.
 \end{aligned}$$

To verify the final inequality, fix $z \in G'_i$ and let (Q, Q_*) be the admissible pair given by property (ii) in the definition of G_i . Then $\omega_{\Omega \setminus H_{i+1}}^z(A_i) \geq \omega_{\text{int } Q \setminus \Omega^c}^z(\Omega^c \cap \text{int } Q) \geq \eta$ by the maximum principle and (4.6). On the other hand,

$$(4.11) \quad \omega(P) \geq \omega_{\Omega \setminus H_1}^X(G'_1) \inf_{z \in G'_1 \cap \Omega} \omega_{\Omega}^z(P) \geq \eta \omega_{\Omega \setminus H_1}^X(G'_1)$$

by Lemma 3.7, the maximum principle, and (4.6). Combining (4.10) and (4.11), we obtain

$$(4.12) \quad \omega(P_k) \leq \eta^{-1} (1 - \eta)^k \omega(P).$$

The consequence of this estimate is that if k (hence M , hence d) is sufficiently large, then $\omega(P_k)$ is arbitrarily small relative to $\omega(P)$.

To proceed, note that each union $\tilde{A}_j = A_{1+(j-1)m^{d-1}} \cup \dots \cup A_{m^{d-1}+(j-1)m^{d-1}}$ of m^{d-1} consecutive rings A_i of d -descendants is an annulus formed from children of P . That is, \tilde{A}_1 is the outermost annulus of children, \tilde{A}_2 is the second annulus of children, and so on. Assign $k = hm^{d-1}$, with h given in the hypothesis of the theorem, and note that $k \leq M$, because $h < m/2$. Partition the set of all children of P into two collections:

$$\mathcal{A} = \{Q \in \text{Child}(P) : Q \subset A_1 \cup \dots \cup A_k\} \quad \text{and} \quad \mathcal{B} = \{Q \in \text{Child}(P) : Q \subset P_k\}.$$

Writing $\text{vol}(P_k)/\text{vol}(P) = \delta$, we have

$$\begin{aligned}
 \sum_{Q \in \text{Child}(P)} \omega(Q)^{1/2} (\text{vol } Q)^{1/2} &= \sum_{Q \in \mathcal{A}} \omega(Q)^{1/2} (\text{vol } Q)^{1/2} + \sum_{Q \in \mathcal{B}} \omega(Q)^{1/2} (\text{vol } Q)^{1/2} \\
 &\leq \omega(A_1 \cup \dots \cup A_k)^{1/2} \left(\sum_{Q \in \mathcal{A}} \text{vol } Q \right)^{1/2} + \omega(P_k)^{1/2} (\text{vol } P_k)^{1/2} \\
 &\leq ((1 - \delta)^{1/2} + \delta^{1/2} \eta^{-1/2} (1 - \eta)^{k/2}) \omega(P)^{1/2} (\text{vol } P)^{1/2},
 \end{aligned}$$




FIGURE 4.1. A decomposition $P = \tilde{A}_1 \cup \dots \cup \tilde{A}_h \cup P_{hm^{d-1}}$ when $n = 2$, $m = 9$, $h = 2$, and $d = 1$, where each little square represents a $d + 1$ descendent of P . Brownian motion, started outside of P , cannot reach the inner region $P_{hm^{d-1}}$ without passing through surfaces G_i (not displayed) drawn in the collars of white squares. Increasing h raises the number of annuli \tilde{A}_j of children of P . Increasing d yields a higher density m^{d-1} of separating surfaces per annulus.

where the first inequality holds by Cauchy-Schwarz. To find the value of δ in terms of the parameters n , m , and h , write

$$\delta = \frac{\text{vol } P_k}{\text{vol } P} = \frac{(m^d - 2k)^n}{m^{dn}} = \left(1 - \frac{2k}{m^d}\right)^n = \left(1 - \frac{2h}{m}\right)^n.$$

We have shown that

$$(4.13) \quad \sum_{Q \in \text{Child}(P)} \omega(Q)^{1/2} (\text{vol } Q)^{1/2} \leq m^{-\lambda} \omega(P)^{1/2} (\text{vol } P)^{1/2},$$

where $\lambda = -\log_m(\gamma) = -\log_m((1 - (1 - 2h/m)^n)^{1/2} + (1 - 2h/m)^{n/2} \eta^{-1/2} (1 - \eta)^{hm^{d-1}/2}) > 0$ by (4.2).

Conclusion. By (4.8), (4.9), and (4.13), the harmonic measure $\omega = \omega_\Omega^X$ of Ω satisfies the hypothesis of Lemma 2.5. Therefore, $\overline{\dim}_H \omega \leq n - \lambda\rho/(\lambda + \rho)$. As we let $\Omega \subsetneq \mathbb{R}^n$ be an arbitrary domain, this proves $b_n \geq \lambda\rho/(\lambda + \rho)$. \square

Remark 4.3. Given n, m, η, h , and d , assign

$$(4.14) \quad V := (1 - (1 - 2h/m)^n)^{1/2} \quad \text{and} \quad \Pi := \eta^{-1/2}(1 - \eta)^{hm^{d-1}/2}.$$

Then $\gamma < V + \Pi$, where γ is defined in (4.2). In particular, $V + \Pi < 1$ implies $\gamma < 1$.

Remark 4.4. The structure of the proof of Theorem 4.1 originated in [Bou87]. However, the density parameter d was only recently introduced in [BG21]. See [BG21, Remark 5.4] for an explanation of why the proof that $b_3 > 0$ in [Bou87] (which implicitly uses $d = 1$) is incomplete. To prove $b_n > 0$, one needs the flexibility to choose $d = d(n) > 1$.

5. PROOF OF THEOREM 1.1

Fix $n \geq 3$ (possibly, but not necessarily large). We first verify that $b_n > 0$. Fix a large integer $m > \xi_m + 2\sqrt{n}$ satisfying the stipulations below. By Corollary 3.2, a Bourgain type estimate (4.1) holds with $\alpha = C_n m^{n-2}$ and $\epsilon = 1$. Set $\eta = 1/(3\alpha)$, $h = 1$, and $d = n$. Then certainly $\alpha\eta = 1/3 < (1 - m^{-n})/(2 - m^{-n})$. We claim the quantity γ defined in (4.2) is less than 1 if m is sufficiently large. Indeed, on the one hand,

$$V = (1 - (1 - 2h/m)^n)^{1/2} = (1 - (1 - 2/m)^n)^{1/2} < 1/2$$

for large enough m . On the other hand, $\Pi = \eta^{-1/2}(1 - \eta)^{hm^{d-1}/2} = \eta^{-1/2}(1 - \eta)^{m^{n-1}/2}$. Using the bound $\ln(1 - x) \leq -x$ for all $0 \leq x < 1$, we see that

$$\ln \Pi \leq \frac{1}{2} \ln(3C_n) + \frac{1}{2}(n - 2) \ln(m) - \frac{m^{n-1}}{6C_n m^{n-2}} \rightarrow -\infty \quad \text{as } m \rightarrow \infty.$$

Hence $\Pi < 1/2$ if m is sufficiently large. Thus, $\gamma < V + \Pi < 1$ if m is sufficiently large. Therefore, $b_n > 0$ by Theorem 4.1.

Now, suppose that $n \geq n_{1/3}$, where $n_{1/3}$ is given by Corollary 3.3. Aiming for a quantitative lower bound on b_n that is valid for all sufficiently large n , we may increase the value of n as convenient. Set $m = n$. By Corollary 3.3, a Bourgain type estimate (4.1) holds with $\alpha = (1/3)(n\sqrt{2\pi e})^{n-2}$ and $\epsilon = 1/2$. Set $\eta = (n\sqrt{2\pi e})^{-(n-2)}$, which ensures that $\alpha\eta = 1/3 < (1 - m^{-n})/(2 - m^{-n})$. Set $h = 1$ and $d = 2n - 3$. On the one hand,

$$V = (1 - (1 - 2/n)^n)^{1/2} \sim (1 - e^{-2})^{1/2} = 0.9298\dots$$

On the other hand, $\Pi = \eta^{-1/2}(1 - \eta)^{n^{2n-4}/2}$ satisfies

$$\ln \Pi \leq \frac{1}{2}(n-2) \ln(n\sqrt{2\pi e}) - \frac{1}{2}n^{2n-4}(n\sqrt{2\pi e})^{-(n-2)} = \frac{1}{2}(n-2) \ln(n\sqrt{2\pi e}) - \frac{1}{2} \left(\frac{n}{\sqrt{2\pi e}} \right)^{n-2}.$$

As the latter expression tends to $-\infty$ as n grows, we see that $\Pi < 0.07$ for large n . Thus, $\gamma < V + \Pi < 0.9998$ and $\lambda \geq -\ln(0.9998)/\ln(n)$ for all sufficiently large n . By Theorem 4.1, $b_n \geq \lambda\rho/(\lambda + \rho)$, where $\rho = 0.914186(1 - 1/3)(1 - n^{-n})n^{-2n(n-1)}/\ln(n)$. Since ρ is substantially smaller than λ for large n , it follows that $b_n \approx \rho$. In particular,

since $0.914186(2/3)(1 - n^{-n}) > 0.6$, we may conclude that $b_n \geq 0.6n^{-2n(n-1)}/\ln(n)$ for all sufficiently large n . Therefore, since $b_n > 0$ for all $n \geq 3$, the theorem holds: there exists $c > 0$ such that $b_n \geq cn^{-2n(n-1)}/\ln(n)$ for all $n \geq 3$.

Remark 5.1. For the large n case, one could also choose $d = n - 1 + \theta(n - 2)$ for any $\theta > 0$ by making n large enough depending on θ . For simplicity, we chose $\theta = 1$.

6. PROOF OF THEOREM 1.2

When $n = 3$ and $m = 9$, a Bourgain type estimate (4.1) holds with $\alpha = 60.8979$ and $\epsilon = 0.000001$ by Corollary 3.4. Assign $\eta = 0.0046$, $h = 3$, and $d = 4$. With the aid of a calculator, one can see that $\gamma = (1 - (1 - 2h/m)^n)^{1/2} + (1 - 2h/m)^{n/2}\eta^{-1/2}(1 - \eta)^{hm^{d-1}/2} < 0.9996$. Thus, by Theorem 4.1, $b_3 \geq \lambda\rho/(\lambda + \rho) \geq 1.452... \times 10^{-15}$. See the appendix for details.

When $n = 4$ and $m = 11$, a Bourgain type estimate (4.1) holds with $\alpha = 1660.53$ by Corollary 3.5. When $\eta = 0.00026$, $h = 4$, and $d = 5$, one may check that $\gamma < 0.9995$. Thus, by Theorem 4.1, $b_4 \geq \lambda\rho/(\lambda + \rho) \geq 2.199... \times 10^{-26}$. Once again, see the appendix.

Remark 6.1. The authors do not claim that these bounds are sharp, but do believe that they are likely close to what the method can prove without further improvements to (3.1) or a more complicated case analysis in the proof of Theorem 4.1. Another small optimization available is to use Hölder's inequality with conjugate exponents depending on m, η, h, d instead of the Cauchy-Schwarz inequality and $p = q = 1/2$ in the statement and proof of Lemma 2.5 and in the definition of γ and proof of Theorem 4.1.

APPENDIX A. WOLFRAM LANGUAGE CODE FOR ESTIMATING b_3

We wrote the following code in Mathematica 13 to estimate b_3 using Theorem 4.1 and Corollary 3.4. See Table A.1 for a record of outputs. For each fixed $m \geq 5$, the parameters η , h , and d were optimized by hand. To maximize ρ , the first priority is to minimize d . To rule out small values of d , take $\eta \approx \alpha^{-1}(1 - m^{-3})/(2 - m^{-3})$ and check that $\gamma > 1$ for each integer $1 \leq h < m/2$. Once the optimal value of the integer d is identified, the second priority is to minimize the real-valued parameter η . Using the current best guess for η (keeping $\gamma < 1$), adjust h to minimize γ . One can then test the value of γ against smaller values of η . If $\gamma < 1$ for some smaller value of η , update the best guess for η and repeat (adjust h , test smaller values of η). Halt the search for η once all smaller values of η (up to some predetermined number of decimals) yield $\gamma > 1$. Use the values of λ and ρ associated to m , η , h , and d to bound b_3 from below by $\lambda\rho/(\lambda + \rho)$.

```
(* All formulas use n=3, epsilon=0.000001 *)
bgAlpha[m_] := (LHS = 1/Sqrt[3] - 2/(m - 2 + Mod[m, 2]);
  RHS = m*2/Sqrt[3] + 27*Pi*m^(-0.999999)/(1 - m^(-1.999999));
  Ceiling[10000*RHS/LHS]/10000); (* round up fourth decimal *)

bgMaxEta[m_] := ((1 - m^(-3))/(2 - m^(-3)))/bgAlpha[m];
```

TABLE A.1. Bounding Bourgain's constant for harmonic measure:
 $b_n \geq \lambda\rho/(\lambda + \rho) \approx \rho$ when $\rho \ll \lambda$.

n	m	η	h	d	α	γ	λ	ρ
3	5	0.0005	2	7	303.102	0.9976...	$1.488... \times 10^{-3}$	$8.020... \times 10^{-18}$
3	6	0.0008	2	6	277.560	0.9947...	$2.911... \times 10^{-3}$	$1.801... \times 10^{-17}$
3	7	0.0019	3	5	83.8178	0.9998...	$7.481... \times 10^{-5}$	$2.418... \times 10^{-16}$
3	8	0.0011	3	5	81.9976	0.9965...	$1.678... \times 10^{-3}$	$2.215... \times 10^{-17}$
3	9	0.0046	3	4	60.8979	0.9996...	$1.616... \times 10^{-4}$	$1.452... \times 10^{-15}$
3	10	0.0031	4	4	61.4480	0.9992...	$3.385... \times 10^{-4}$	$3.210... \times 10^{-16}$
3	11	0.0022	4	4	54.2657	0.9984...	$6.516... \times 10^{-4}$	$8.031... \times 10^{-17}$
3	12	0.0016	5	4	55.5835	0.9993...	$2.254... \times 10^{-4}$	$2.174... \times 10^{-17}$
3	13	0.0012	5	4	52.5339	0.9982...	$6.978... \times 10^{-4}$	$6.521... \times 10^{-18}$
3	14	0.0009	5	4	54.1918	0.9988...	$4.385... \times 10^{-4}$	$2.117... \times 10^{-18}$
4	7	0.00006	3	7	2409.54	0.9998...	$7.291... \times 10^{-5}$	$3.637... \times 10^{-28}$
4	8	0.00016	3	6	2425.26	0.9999...	$2.780... \times 10^{-5}$	$1.390... \times 10^{-26}$
4	9	0.00009	3	6	1813.48	0.9978...	$9.801... \times 10^{-4}$	$6.651... \times 10^{-28}$
4	10	0.00005	4	6	1834.77	0.9994...	$2.361... \times 10^{-4}$	$3.605... \times 10^{-29}$
4	11	0.00026	4	5	1660.53	0.9995...	$2.062... \times 10^{-4}$	$2.199... \times 10^{-26}$
4	12	0.00017	5	5	1685.89	0.9999...	$2.779... \times 10^{-5}$	$3.301... \times 10^{-27}$
4	13	0.00012	5	5	1619.82	0.9995...	$1.932... \times 10^{-4}$	$5.289... \times 10^{-28}$
4	14	0.00009	5	5	1649.02	0.9981...	$6.908... \times 10^{-4}$	$9.177... \times 10^{-29}$
4	15	0.00006	6	5	1626.75	0.9997...	$8.531... \times 10^{-5}$	$1.809... \times 10^{-29}$
4	16	0.00005	6	5	1659.76	0.9985...	$5.340... \times 10^{-4}$	$3.816... \times 10^{-30}$

```

bgV[m_,h_] := If[h<m/2, (1 - (1 - 2*h/m)^3)^0.5, 1];

bgEtaProd[m_,eta_,h_,d_] := If[h<m/2,
  (1 - 2*h/m)^1.5 * eta^(-0.5) * ((1-eta)^(0.5*h*m^(d-1))), 1];
(* If h >= m/2, then bgV[m,h]=1 and bgEtaProd[m,eta,h,d]=1 *)

bgGamma[m_,eta_,h_,d_] := bgV[m,h] + bgEtaProd[m,eta,h,d];

bgLambda[m_,eta_,h_,d_] := Max[0,-Log[m,bgGamma[m,eta,h,d]]];
(* If bgGamma[m,eta,h,d]>=1, then bgLambda[m,eta,h,d]=0 *)

bgRho[m_,eta_,d_] := If[eta <= bgMaxEta[m],
  0.914186*(1 - m^(-3))*(1 - bgAlpha[m]*eta)*m^(-3*(d + 1))/Log[m], 0];
(* If eta > bgMaxEta[m], then bgRho[m,eta,d]=0 *)

```



```

bgLowerBound[m_,eta_,h_,d_] := (lambda = bgLambda[m,eta,h,d];
  rho = bgRho[m, eta, d];
  lambda*rho/(lambda + rho));
(* Returns lower bound on b_3 for any admissible (m,eta,h,d) *)
(* For example, bgLowerBound[9,0.0046,3,4] returns 1.45271 * 10^(-15) *)

```

The code used to estimate b_4 (omitted) is similar. It can be reproduced by modifying the definition of `bgAlpha` using Corollary 3.5 instead of Corollary 3.4 and changing $n = 3$ to $n = 4$ in the definitions of `bgMaxEta`, `bgV`, `bgEtaProd`, and `bgRho`.

REFERENCES

- [ABR01] Sheldon Axler, Paul Bourdon, and Wade Ramey, *Harmonic function theory*, second ed., Graduate Texts in Mathematics, vol. 137, Springer-Verlag, New York, 2001. MR 1805196
- [AHM⁺16] Jonas Azzam, Steve Hofmann, José María Martell, Svitlana Mayboroda, Mihalis Mourgoglou, Xavier Tolsa, and Alexander Volberg, *Rectifiability of harmonic measure*, *Geom. Funct. Anal.* **26** (2016), no. 3, 703–728. MR 3540451
- [Azz20] Jonas Azzam, *Dimension drop for harmonic measure on Ahlfors regular boundaries*, *Potential Anal.* **53** (2020), no. 3, 1025–1041. MR 4140087
- [BG21] Matthew Badger and Alyssa Genschaw, *Hausdorff dimension of caloric measure*, preprint, [arXiv:2108.12340](https://arxiv.org/abs/2108.12340), 2021.
- [Bis92] Christopher J. Bishop, *Some questions concerning harmonic measure*, *Partial differential equations with minimal smoothness and applications* (Chicago, IL, 1990), IMA Vol. Math. Appl., vol. 42, Springer, New York, 1992, pp. 89–97. MR 1155854
- [Bis07] Christopher J. Bishop, *Review of Harmonic measure*, *Bull. Amer. Math. Soc.* **44** (2007), no. 2, 267–276.
- [Bou87] J. Bourgain, *On the Hausdorff dimension of harmonic measure in higher dimension*, *Invent. Math.* **87** (1987), no. 3, 477–483. MR 874032
- [BZ15] Athanasios Batakis and Anna Zdunik, *Hausdorff and harmonic measures on non-homogeneous Cantor sets*, *Ann. Acad. Sci. Fenn. Math.* **40** (2015), no. 1, 279–303. MR 3310084
- [Car85] Lennart Carleson, *On the support of harmonic measure for sets of Cantor type*, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **10** (1985), 113–123. MR 802473
- [Doo01] Joseph L. Doob, *Classical potential theory and its probabilistic counterpart*, *Classics in Mathematics*, Springer-Verlag, Berlin, 2001, Reprint of the 1984 edition. MR 1814344
- [Fal97] Kenneth Falconer, *Techniques in fractal geometry*, John Wiley & Sons, Ltd., Chichester, 1997. MR 1449135
- [FGAS08] M. Filoche, D. S. Grebenkov, J. S. Andrade, and B. Sapoval, *Passivation of irregular surfaces accessed by diffusion*, *Proceedings of the National Academy of Sciences* **105** (2008), no. 22, 7636–7640.
- [GM05] John B. Garnett and Donald E. Marshall, *Harmonic measure*, *New Mathematical Monographs*, vol. 2, Cambridge University Press, Cambridge, 2005. MR 2150803
- [Hel09] Lester L. Helms, *Potential theory*, Universitext, Springer-Verlag London, Ltd., London, 2009. MR 2526019
- [HM14] Steve Hofmann and José María Martell, *Uniform rectifiability and harmonic measure I: Uniform rectifiability implies Poisson kernels in L^p* , *Ann. Sci. Éc. Norm. Supér. (4)* **47** (2014), no. 3, 577–654. MR 3239100
- [JW88] Peter W. Jones and Thomas H. Wolff, *Hausdorff dimension of harmonic measures in the plane*, *Acta Math.* **161** (1988), no. 1-2, 131–144. MR 962097

- [KW85] Robert Kaufman and Jang-Mei Wu, *On the snowflake domain*, Ark. Mat. **23** (1985), no. 1, 177–183. MR 800179
- [Li07] Yuan-Chuan Li, *A note on an identity of the gamma function and Stirling's formula*, Real Anal. Exchange **32** (2006/07), no. 1, 267–271. MR 2329236
- [LVV05] John L. Lewis, Gregory C. Verchota, and Andrew L. Vogel, *Wolff snowflakes*, Pacific J. Math. **218** (2005), no. 1, 139–166. MR 2224593
- [Mak85] N. G. Makarov, *On the distortion of boundary sets under conformal mappings*, Proc. London Math. Soc. (3) **51** (1985), no. 2, 369–384. MR 794117
- [Mat95] Pertti Mattila, *Geometry of sets and measures in Euclidean spaces*, Cambridge Studies in Advanced Mathematics, vol. 44, Cambridge University Press, Cambridge, 1995, Fractals and rectifiability. MR 1333890 (96h:28006)
- [MMR00] Pertti Mattila, Manuel Morán, and José-Manuel Rey, *Dimension of a measure*, Studia Math. **142** (2000), no. 3, 219–233. MR 1792606
- [MP10] Peter Mörters and Yuval Peres, *Brownian motion*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, Cambridge, 2010, With an appendix by Oded Schramm and Wendelin Werner. MR 2604525 (2011i:60152)
- [Øks81] Bernt Øksendal, *Brownian motion and sets of harmonic measure zero*, Pacific J. Math. **95** (1981), no. 1, 179–192. MR 631668
- [Rog98] C. A. Rogers, *Hausdorff measures*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1998, Reprint of the 1970 original, With a foreword by K. J. Falconer. MR 1692618
- [Wol93] Thomas H. Wolff, *Plane harmonic measures live on sets of σ -finite length*, Ark. Mat. **31** (1993), no. 1, 137–172. MR 1230270
- [Wol95] Thomas H. Wolff, *Counterexamples with harmonic gradients in \mathbf{R}^3* , Essays on Fourier analysis in honor of Elias M. Stein (Princeton, NJ, 1991), Princeton Math. Ser., vol. 42, Princeton Univ. Press, Princeton, NJ, 1995, pp. 321–384. MR 1315554

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-1009

Email address: `matthew.badger@uconn.edu`

MATHEMATICS DEPARTMENT, MILWAUKEE SCHOOL OF ENGINEERING, MILWAUKEE, WI 53202

Email address: `genschaw@msoe.edu`