

# Quantisation of Constrained Systems

Supervised by Chaolun Wu

Matthew Dunne

Trinity College Dublin

March 27, 2025

# Outline

- 1 Constrained Hamiltonian Systems
- 2 Gauge Symmetry
- 3 Ghosts and Parisi-Sourlas Quantisation

# Constrained Hamiltonian Systems

- Before Dirac, constraints were only analysed on a case by case basis



Paul Dirac

Image: The Nobel Foundation

# Constrained Hamiltonian Systems

- Before Dirac, constraints were only analysed on a case by case basis
- “Lectures on Quantum Mechanics”<sup>[1]</sup> provides a recipe for dealing with general constraints



Paul Dirac

Image: The Nobel Foundation

# Constrained Hamiltonian Systems

- Before Dirac, constraints were only analysed on a case by case basis
- “Lectures on Quantum Mechanics”<sup>[1]</sup> provides a recipe for dealing with general constraints
- Most importantly for this project, the concept of gauge freedom appears more naturally in the Hamiltonian formulation



Paul Dirac

Image: The Nobel Foundation

# Constrained Hamiltonian Systems

- Recall: In setting up the Hamiltonian description of a theory, one defines the canonical momenta

$$p_i = \frac{\partial L}{\partial \dot{q}^i} \quad i = 1, \dots, N$$

# Constrained Hamiltonian Systems

- Recall: In setting up the Hamiltonian description of a theory, one defines the canonical momenta

$$p_i = \frac{\partial L}{\partial \dot{q}^i} \quad i = 1, \dots, N$$

- The theory is then formulated in terms of the  $2N$  coordinates  $(q, p)$  of phase space which obey the Poisson bracket relations

$$\{q^i, p_j\} = \delta^i_j, \quad \{q^i, q^j\} = \{p_i, p_j\} = 0$$

# Constrained Hamiltonian Systems

- Recall: In setting up the Hamiltonian description of a theory, one defines the canonical momenta

$$p_i = \frac{\partial L}{\partial \dot{q}^i} \quad i = 1, \dots, N$$

- The theory is then formulated in terms of the  $2N$  coordinates  $(q, p)$  of phase space which obey the Poisson bracket relations

$$\{q^i, p_j\} = \delta^i_j, \quad \{q^i, q^j\} = \{p_i, p_j\} = 0$$

- When not all of the momentum variables are independent we also get a set of constraints

$$G_a(q, p) = 0 \quad a = 1, \dots, M$$



# Geometry of Constrained Hamiltonian Systems

- If the constraints are independent of each other, they reduce the number of physical degrees of freedom to  $2N - M$

# Geometry of Constrained Hamiltonian Systems

- If the constraints are independent of each other, they reduce the number of physical degrees of freedom to  $2N - M$
- The physical trajectories must be consistent with the constraints so they are confined to the  $(2N - M)$ -dimensional subspace called the physical phase space

# Geometry of Constrained Hamiltonian Systems

- If the constraints are independent of each other, they reduce the number of physical degrees of freedom to  $2N - M$
- The physical trajectories must be consistent with the constraints so they are confined to the  $(2N - M)$ -dimensional subspace called the physical phase space
- In the remaining unphysical sector of phase space, the constraints do not necessarily vanish.

# Geometry of Constrained Hamiltonian Systems

- If the constraints are independent of each other, they reduce the number of physical degrees of freedom to  $2N - M$
- The physical trajectories must be consistent with the constraints so they are confined to the  $(2N - M)$ -dimensional subspace called the physical phase space
- In the remaining unphysical sector of phase space, the constraints do not necessarily vanish.
- For clarity one uses the weak equality  $G_a \approx 0$  as a reminder that the constraints only vanish on the physical phase space.

# Second Class Constraints

- We call our constraints second class if

$$\{G_a, G_b\} \neq 0 \quad a, b = 1, \dots, M$$

# Second Class Constraints

- We call our constraints second class if

$$\{G_a, G_b\} \neq 0 \quad a, b = 1, \dots, M$$

- They induce a structure which can be different from the usual symplectic structure of phase space

# Second Class Constraints

- We call our constraints second class if

$$\{G_a, G_b\} \neq 0 \quad a, b = 1, \dots, M$$

- They induce a structure which can be different from the usual symplectic structure of phase space
- Historically implemented using the Dirac bracket, a method which this project avoids

# First Class Constraints

- We call our constraints first class if

$$\{G_a, G_b\} = U_{ab}{}^c G_c \approx 0 \quad a, b, c = 1, \dots, M$$

where  $U_{ab}{}^c$  are called structure functions



# First Class Constraints

- We call our constraints first class if

$$\{G_a, G_b\} = U_{ab}^c G_c \approx 0 \quad a, b, c = 1, \dots, M$$

where  $U_{ab}^c$  are called structure functions

- First class constraints lead to arbitrary functions of time in the equations of motion.

# First Class Constraints

- We call our constraints first class if

$$\{G_a, G_b\} = U_{ab}{}^c G_c \approx 0 \quad a, b, c = 1, \dots, M$$

where  $U_{ab}{}^c$  are called structure functions

- First class constraints lead to arbitrary functions of time in the equations of motion.
- Specific physical state  $\iff$  Equivalence class of points on physical phase space.

# First Class Constraints

- We call our constraints first class if

$$\{G_a, G_b\} = U_{ab}^c G_c \approx 0 \quad a, b, c = 1, \dots, M$$

where  $U_{ab}^c$  are called structure functions

- First class constraints lead to arbitrary functions of time in the equations of motion.
- Specific physical state  $\iff$  Equivalence class of points on physical phase space.
- Local gauge symmetry: Invariance of physical states under transformations to other points in the equivalence class (gauge transformations)

# First Class Constraints

- We call our constraints first class if

$$\{G_a, G_b\} = U_{ab}^c G_c \approx 0 \quad a, b, c = 1, \dots, M$$

where  $U_{ab}^c$  are called structure functions

- First class constraints lead to arbitrary functions of time in the equations of motion.
- Specific physical state  $\iff$  Equivalence class of points on physical phase space.
- Local gauge symmetry: Invariance of physical states under transformations to other points in the equivalence class (gauge transformations)
- Dirac conjecture: All first class constraints generate gauge transformations

# Gauge Symmetry

- Many theories of physical interest enjoy the property of gauge symmetry: Electrodynamics, Yang-Mills, General Relativity...

# Gauge Symmetry


- Many theories of physical interest enjoy the property of gauge symmetry: Electrodynamics, Yang-Mills, General Relativity...
- Theories which we are very interested in quantising

# Gauge Symmetry

- Many theories of physical interest enjoy the property of gauge symmetry: Electrodynamics, Yang-Mills, General Relativity...
- Theories which we are very interested in quantising
- How do we resolve our redundant description of theories with gauge freedom?

# Gauge Fixing

- Introduce gauge fixing conditions  $F^a(q, p) = 0$




A good choice of gauge and two bad ones [2].



# Gauge Fixing

- Introduce gauge fixing conditions  $F^a(q, p) = 0$
- Must intersect each equivalence class exactly once and satisfy

$$\det[\{F^a, G_b\}] \neq 0$$




A good choice of gauge and two bad ones [2].

# Gauge Fixing

- Introduce gauge fixing conditions  $F^a(q, p) = 0$
- Must intersect each equivalence class exactly once and satisfy

$$\det[\{F^a, G_b\}] \neq 0$$

- Constraints and gauge fixing conditions together form a set of second class constraints!



A good choice of gauge and two bad ones [2].

# The Need for Other Methods

- It is difficult to verify that a gauge choice is a good one (Gribov ambiguity)

# The Need for Other Methods

- It is difficult to verify that a gauge choice is a good one (Gribov ambiguity)
- Even when successful, the standard treatment of second class constraints involving the explicit removal of gauge degrees of freedom can cause essential properties like Lorentz invariance or locality to be lost

# The Need for Other Methods

- It is difficult to verify that a gauge choice is a good one (Gribov ambiguity)
- Even when successful, the standard treatment of second class constraints involving the explicit removal of gauge degrees of freedom can cause essential properties like Lorentz invariance or locality to be lost
- We instead develop methods which make use of some extended symmetry of the gauge phase space in order to obtain a gauge invariant dynamical description

# Dynamics over a Grassmann Algebra

- Anti-commuting variables  $\theta^i$  are odd elements of a Grassmann algebra which satisfy

$$\theta^2 = 0 \qquad \theta^i \theta^j = -\theta^j \theta^i$$

# Dynamics over a Grassmann Algebra

- Anti-commuting variables  $\theta^i$  are odd elements of a Grassmann algebra which satisfy

$$\theta^2 = 0 \qquad \theta^i \theta^j = -\theta^j \theta^i$$

- One may have a dynamical system with  $N$  commuting (bosonic) variables  $q^i(t)$  and  $K$  anti-commuting (fermionic) variables  $\theta^\alpha(t)$

# Dynamics over a Grassmann Algebra

- Anti-commuting variables  $\theta^i$  are odd elements of a Grassmann algebra which satisfy

$$\theta^2 = 0 \qquad \theta^i \theta^j = -\theta^j \theta^i$$

- One may have a dynamical system with  $N$  commuting (bosonic) variables  $q^i(t)$  and  $K$  anti-commuting (fermionic) variables  $\theta^\alpha(t)$
- Both types of variables and functions of them can be taken as elements of the same Grassmann algebra



# Ghosts

- In applications to constrained systems, fermionic variables, when introduced are called ghosts (don't worry they're not that scary)

# Ghosts

- In applications to constrained systems, fermionic variables, when introduced are called ghosts (don't worry they're not that scary)
- Ghosts have properties that can be exploited to perform gauge fixing without eliminating unphysical degrees of freedom at intermediate stages of analysis.

# Ghosts

- In applications to constrained systems, fermionic variables, when introduced are called ghosts (don't worry they're not that scary)
- Ghosts have properties that can be exploited to perform gauge fixing without eliminating unphysical degrees of freedom at intermediate stages of analysis.
- Famously Fadeev-Popov ghosts were used to obtain a gauge fixed path-integral for Yang-Mills

# Ghosts

- In applications to constrained systems, fermionic variables, when introduced are called ghosts (don't worry they're not that scary)
- Ghosts have properties that can be exploited to perform gauge fixing without eliminating unphysical degrees of freedom at intermediate stages of analysis.
- Famously Fadeev-Popov ghosts were used to obtain a gauge fixed path-integral for Yang-Mills
- In my project I am exploring the application of one such method called Parisi-Sourlas quantisation<sup>[3]</sup>

# Parisi-Sourlas Formalism

- Suppose that we want to formulate a path-integral for a theory of  $N$  dynamical variables with 2 constraints without explicitly reducing the action to a lower dimensional form. Then by analytic continuation we can imagine writing the reduced path integral in the equivalent form

$$\int d^{N-2}x f(x^2) \sim \lim_{d \rightarrow -2} \int d^N x d^d y f(x^2 + y^2)$$

# Parisi-Sourlas Formalism

- Suppose that we want to formulate a path-integral for a theory of  $N$  dynamical variables with 2 constraints without explicitly reducing the action to a lower dimensional form. Then by analytic continuation we can imagine writing the reduced path integral in the equivalent form

$$\int d^{N-2}x f(x^2) \sim \lim_{d \rightarrow -2} \int d^N x d^d y f(x^2 + y^2)$$

# Parisi-Sourlas Formalism

- Suppose that we want to formulate a path-integral for a theory of  $N$  dynamical variables with 2 constraints without explicitly reducing the action to a lower dimensional form. Then by analytic continuation we can imagine writing the reduced path integral in the equivalent form

$$\int d^{N-2}x f(x^2) \sim \lim_{d \rightarrow -2} \int d^N x d^d y f(x^2 + y^2)$$

- The “negative dimensional coordinates” of the above integral can then be realised by anti-commuting variables

# Ghosts as Negative Dimensional Coordinates: Parisi-Sourlas Supersymmetry



# References



P.A.M. Dirac.

*Lectures on Quantum Mechanics.*

Belfer Graduate School of Science, monograph series. Dover Publications, 2001.



Bengtsson. I.

Constrained hamiltonian systems.

*U. Stockholm, Unknown Year.*



Antti J. Niemi.

Pedagogical Introduction to BRST.

*Phys. Rept.*, 184:147–165, 1989.

# Thank You!