Homework #5

BSTA 519 - Fall 2021

Applied Longitudinal Data Analysis Due - Monday 11/15/2021 by midnight

Linear mixed effects models, Problem 1

Chapter 8, problem 8.1.

The data is provided in Exercise.txt. If you open up the dataset using WordPad, it will show the correct format. This is the dataset we used to illustrate the covariance pattern models.

Linear mixed effects models, Problem 2

The Childhood Asthma Management Program study

The Childhood Asthma Management Program (CAMP) was a clinical trial carried out in children with asthma. The trial was designed to determine the long-term effects of 3 treatments (budesonide, nedocromil, or placebo) on pulmonary function for up to 10 years period. The design of CAMP was a multicenter, masked, placebo-controlled, randomized trial. A total of 695 children aged 5-12 years were enrolled between December of 1993 and September of 1995.

For this analysis, you will only look at the data from the treatment group of nedocromil and the outcome variable is, FFratio, the ratio of FEV1 to forced vital capacity (FVC) where FEV1 is the Forced Expiratory Volume at 1 second. The data for this analysis is provided in the file *CAMP HWK5.xlsx*.

Dataset Variables

id Randomized participant ID age_rz Age in years at Randomization

GENDER m=male, f=female

ETHNIC w=white, b=black, h=hispanic, o=other

FFratio FEV1/FVC ratio %

Fdays Days since randomization

1) Create an appropriate time plot to show the trend of how FFratio changes over time and describe the trend.

Homework #5

BSTA 519 - Fall 2021

Applied Longitudinal Data Analysis Due - Monday 11/15/2021 by midnight

- 2) Fit a random intercept model to evaluate both the cross-sectional association between age at randomization and FFratio, and the longitudinal association between age and FFratio.
 - i. Write out the model you fit to evaluate both associations.
 - ii. Is there a significant cross-sectional association? Is there a significant longitudinal association? Interpret your results.
 - iii. Obtain the estimated covariance and correlation matrix. What is the estimated intra-class correlation coefficient in this case? Is there a strong correlation among the repeated measures within the same subject?
 - iv. Based on this model, how do the subject-specific response curves differ among the individuals?
 - v. Plot the marginal mean response over time based on the fitted model.
 - vi. Fit a marginal model using the same mean response with an exchangeable correlation matrix, and the REML method. How do the results from the marginal model compare with the results above?
- 3) Fit a random intercept and random slope model with the above specification for the mean response, with an unstructured covariance matrix for the random effects.
 - i. Write out the model you fit to evaluate both associations.
 - ii. How does the resulted variance-covariance matrix differ from the one in 1)? Provide the estimated covariance and correlation matrix.
 - iii. Does this model provide a better fit to the data, compared to the model in 1)?
 - iv. Based on this model, how do the subject-specific response curves differ from each other? Could you provide the range of the slopes for the middle 95% subjects?
- 4) Fit a random intercept and random slope model to evaluate whether the longitudinal association between age and FFratio differs by gender after adjusting for ethnicity and age at randomization, and interpret your results.