

# Derivative Rules

## Advanced



The Derivative tells us the slope of a function at any point.

There are **rules** we can follow to find many derivatives.

For example:

- The slope of a **constant** value (like 3) is always 0
- The slope of a **line** like  $2x$  is 2, or  $3x$  is 3 etc
- and so on.

Here are useful rules to help you work out the derivatives of many functions (with examples below). Note: the little mark ' means "Derivative of".



### Common Functions

Constant

$c$

0

Line

$x$

1

$ax$

$a$

Square

$x^2$

$2x$

Square Root

$\sqrt{x}$

$(\frac{1}{2})x^{-\frac{1}{2}}$

Exponential

$e^x$

$e^x$

$a^x$

$\ln(a) a^x$

Logarithms

$\ln(x)$

$1/x$

$\log_a(x)$

$1 / (x \ln(a))$

Trigonometry ( $x$  is in radians)

$\sin(x)$

$\cos(x)$

$\cos(x)$

$-\sin(x)$

$\tan(x)$

$\sec^2(x)$

Inverse Trigonometry

$$\sin^{-1}(x)$$

$$1/\sqrt{1-x^2}$$

$$\cos^{-1}(x)$$

$$-1/\sqrt{1-x^2}$$

$$\tan^{-1}(x)$$

$$1/(1+x^2)$$

### Rules

### Function

### Derivative

Multiplication by constant

$$cf$$

$$cf'$$

[Power Rule](#)

$$x^n$$

$$nx^{n-1}$$

Sum Rule

$$f + g$$

$$f' + g'$$

Difference Rule

$$f - g$$

$$f' - g'$$

Product Rule

$$fg$$

$$f g' + f' g$$

Quotient Rule

$$f/g$$

$$(f' g - g' f)/g^2$$

Reciprocal Rule

$$1/f$$

$$-f'/f^2$$

Chain Rule

(as ["Composition of Functions"](#)).

$$f \circ g$$

$$(f' \circ g) \times g'$$

Chain Rule (using ' )

$$f(g(x))$$

$$f'(g(x))g'(x)$$

Chain Rule (using  $\frac{d}{dx}$  )

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

"The derivative of" is also written  $\frac{d}{dx}$

So  $\frac{d}{dx}\sin(x)$  and  $\sin(x)'$  both mean "The derivative of  $\sin(x)$ "

## Examples

Example: what is the derivative of  $\sin(x)$  ?

From the table above it is listed as being  **$\cos(x)$**

It can be written as:

$$\frac{d}{dx} \sin(x) = \cos(x)$$

Or:

$$\sin(x)' = \cos(x)$$

## Power Rule

Example: What is  $\frac{d}{dx} x^3$  ?

The question is asking "what is the derivative of  $x^3$  ?"

We can use the [Power Rule](#), where  $n=3$ :

$$\frac{d}{dx} x^n = nx^{n-1}$$

$$\frac{d}{dx} x^3 = 3x^{3-1} = \mathbf{3x^2}$$

(In other words the derivative of  $x^3$  is  $3x^2$ )

So it is simply this:

$$\begin{array}{c} x^3 \\ \swarrow \searrow \\ 3x^2 \end{array}$$

"multiply by power  
then reduce power by 1"

It can also be used in cases like this:

Example: What is  $\frac{d}{dx} (1/x)$  ?

$1/x$  is also  $x^{-1}$

We can use the Power Rule, where  $n = -1$ :

$$\frac{d}{dx} x^n = nx^{n-1}$$

$$\frac{d}{dx} x^{-1} = -1x^{-1-1} = -x^{-2}$$

So we just did this:

$$x^{-1} \rightarrow -1x^{-2}$$

which simplifies to  $-x^{-2}$

## Multiplication by constant

Example: What is  $\frac{d}{dx} 5x^3$  ?

the derivative of  $cf = cf'$

the derivative of  $5f = 5f'$

We know (from the Power Rule):

$$\frac{d}{dx} x^3 = 3x^{3-1} = 3x^2$$

So:

$$\frac{d}{dx} 5x^3 = 5 \frac{d}{dx} x^3 = 5 \times 3x^2 = \mathbf{15x^2}$$

## Sum Rule

Example: What is the derivative of  $x^2 + x^3$  ?

The Sum Rule says:

the derivative of  $f + g = f' + g'$

So we can work out each derivative separately and then add them.

Using the Power Rule:

- $\frac{d}{dx}x^2 = 2x$
- $\frac{d}{dx}x^3 = 3x^2$

And so:

$$\text{the derivative of } x^2 + x^3 = \mathbf{2x + 3x^2}$$

## Difference Rule

It doesn't have to be **x**, we can differentiate with respect to, for example, **v**:

Example: What is  $\frac{d}{dv}(v^3 - v^4)$  ?

The Difference Rule says

$$\text{the derivative of } f - g = f' - g'$$

So we can work out each derivative separately and then subtract them.

Using the Power Rule:

- $\frac{d}{dv}v^3 = 3v^2$
- $\frac{d}{dv}v^4 = 4v^3$

And so:

$$\text{the derivative of } v^3 - v^4 = \mathbf{3v^2 - 4v^3}$$

## Sum, Difference, Constant Multiplication And Power Rules

Example: What is  $\frac{d}{dz}(5z^2 + z^3 - 7z^4)$  ?

Using the Power Rule:

- $\frac{d}{dz}z^2 = 2z$

- $\frac{d}{dz} z^3 = 3z^2$
- $\frac{d}{dz} z^4 = 4z^3$

And so:

$$\frac{d}{dz} (5z^2 + z^3 - 7z^4) = 5 \times 2z + 3z^2 - 7 \times 4z^3 = \mathbf{10z + 3z^2 - 28z^3}$$

## Product Rule

Example: What is the derivative of  $\cos(x)\sin(x)$  ?

The Product Rule says:

$$\text{the derivative of } fg = f g' + f' g$$

In our case:

- $f = \cos$
- $g = \sin$

We know (from the table above):

- $\frac{d}{dx} \cos(x) = -\sin(x)$
- $\frac{d}{dx} \sin(x) = \cos(x)$

So:

$$\text{the derivative of } \cos(x)\sin(x) = \cos(x)\cos(x) - \sin(x)\sin(x)$$

$$= \mathbf{\cos^2(x) - \sin^2(x)}$$

## Reciprocal Rule

Example: What is  $\frac{d}{dx}(1/x)$  ?

The Reciprocal Rule says:

$$\text{the derivative of } 1/f = -f'/f^2$$

**With  $f(x) = x$ , we know that  $f'(x) = 1$**

So:

$$\text{the derivative of } 1/x = -1/x^2$$

Which is the same result we got above using the Power Rule.

## Chain Rule

Example: What is  $\frac{d}{dx}\sin(x^2)$  ?

**$\sin(x^2)$**  is made up of  **$\sin()$**  and  **$x^2$** :

- $f(g) = \sin(g)$
- $g(x) = x^2$

The Chain Rule says:

$$\text{the derivative of } f(g(x)) = f'(g(x))g'(x)$$

The individual derivatives are:

- $f'(g) = \cos(g)$
- $g'(x) = 2x$

So:

$$\begin{aligned}\frac{d}{dx}\sin(x^2) &= \cos(g(x)) (2x) \\ &= 2x \cos(x^2)\end{aligned}$$

Another way of writing the Chain Rule is:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

Let's do the previous example again using that formula:

Example: What is  $\frac{d}{dx} \sin(x^2)$  ?

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Have  $u = x^2$ , so  $y = \sin(u)$ :

$$\frac{d}{dx} \sin(x^2) = \frac{d}{du} \sin(u) \frac{d}{dx} x^2$$

Differentiate each:

$$\frac{d}{dx} \sin(x^2) = \cos(u) (2x)$$

Substitute back  $u = x^2$  and simplify:

$$\frac{d}{dx} \sin(x^2) = 2x \cos(x^2)$$

Same result as before (thank goodness!)

Another couple of examples of the Chain Rule:

Example: What is  $\frac{d}{dx} (1/\cos(x))$  ?

**$1/\cos(x)$**  is made up of  **$1/g$**  and  **$\cos()$** :

- $f(g) = 1/g$
- $g(x) = \cos(x)$

The Chain Rule says:

$$\text{the derivative of } f(g(x)) = f'(g(x))g'(x)$$

The individual derivatives are:



- $f'(g) = -1/(g^2)$
- $g'(x) = -\sin(x)$

So:

$$\begin{aligned} (1/\cos(x))' &= -1/(g(x))^2 \times -\sin(x) \\ &= \sin(x)/\cos^2(x) \end{aligned}$$

Note:  $\sin(x)/\cos^2(x)$  is also  $\tan(x)/\cos(x)$ , or many other forms.

Example: What is  $\frac{d}{dx}(5x-2)^3$  ?

The Chain Rule says:

$$\text{the derivative of } f(g(x)) = f'(g(x))g'(x)$$

$(5x-2)^3$  is made up of  $g^3$  and  $5x-2$ :

- $f(g) = g^3$
- $g(x) = 5x-2$

The individual derivatives are:

- $f'(g) = 3g^2$  (by the Power Rule)
- $g'(x) = 5$

So:

$$\frac{d}{dx}(5x-2)^3 = 3g(x)^2 \times 5 = 15(5x-2)^2$$

[Question 1](#) [Question 2](#) [Question 3](#) [Question 4](#) [Question 5](#) [Question 6](#)  
[Question 7](#) [Question 8](#) [Question 9](#) [Question 10](#) [Question 11](#) [Question 12](#)  
[Question 13](#)

Copyright © 2017 MathsIsFun.com