

# Reconstruction Algorithms for Blind Ptychographic Imaging

**Matthew K. Tam**

School of Mathematical and Physical Sciences  
University of Newcastle Australia

With generous support from the AustMS Student Support Scheme



Joint work with R. Hesse, D.R. Luke and S. Sabach



59th Annual Meeting of the Australian Mathematical Society  
Flinders University (Adelaide), Sept. 28–Oct. 1, 2015

# Scanning Ptychography: A Crash Course

- An unknown **specimen** is illuminated by a **localized illumination function** resulting in an **exit-wave** whose intensity is observed.
- A **ptychography dataset** is a series of these observations, each is obtained by shifting the illumination function to a different position relative to the specimen. **Neighbouring illumination regions overlap**.
- Given a ptychographic dataset, the **blind ptychography problem** is to simultaneously reconstruct the specimen and illumination function.




Figure : An illumination function (left), specimen (center), and exit-wave (right).

# Scanning Ptychography: A Crash Course

- An unknown **specimen** is illuminated by a **localized illumination function** resulting in an **exit-wave** whose intensity is observed.
- A **ptychography dataset** is a series of these observations, each is obtained by shifting the illumination function to a different position relative to the specimen. **Neighbouring illumination regions overlap**.
- Given a ptychographic dataset, the **blind ptychography problem** is to simultaneously reconstruct the specimen and illumination function.




Figure : An illumination function (left), specimen (center), and exit-wave (right).

# Scanning Ptychography: A Crash Course

- An unknown **specimen** is illuminated by a **localized illumination function** resulting in an **exit-wave** whose intensity is observed.
- A **ptychography dataset** is a series of these observations, each is obtained by shifting the illumination function to a different position relative to the specimen. **Neighbouring illumination regions overlap**.
- Given a ptychographic dataset, the **blind ptychography problem** is to simultaneously reconstruct the specimen and illumination function.




Figure : An illumination function (left), specimen (center), and exit-wave (right).

# Scanning Ptychography: A Crash Course

The mathematical model is:

- The unknown **illumination function**:  $x \in \mathbb{C}^{n \times n}$ ,
- The unknown **specimen**:  $y \in \mathbb{C}^{n \times n}$ ,
- An  $m$ -tuple of **diffraction patterns**:  $\mathbf{z} = (z_1, \dots, z_m) \in (\mathbb{C}^{n \times n})^m$ ,
- The *shift map*  $S_j : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$  moves  $x$  to the position corresponding to the  $j^{\text{th}}$  diffraction pattern measurement.
- The elements of the triple  $(x, y, \mathbf{z})$  are related by:

$$S_j(x) \odot y = z_j \quad \text{for } j = 1, 2, \dots, m.$$




Figure : An example of  $S_j(x) \odot y = z_j$  with  $S_j$  localising “ $x$ ” to the  $j^{\text{th}}$  position.

# Scanning Ptychography: A Crash Course

The mathematical model is:

- The unknown **illumination function**:  $x \in \mathbb{C}^{n \times n}$ ,
- The unknown **specimen**:  $y \in \mathbb{C}^{n \times n}$ ,
- An  $m$ -tuple of **diffraction patterns**:  $\mathbf{z} = (z_1, \dots, z_m) \in (\mathbb{C}^{n \times n})^m$ ,
- The *shift map*  $S_j : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$  moves  $x$  to the position corresponding to the  $j^{\text{th}}$  diffraction pattern measurement.
- The elements of the triple  $(x, y, \mathbf{z})$  are related by:

$$S_j(x) \odot y = z_j \quad \text{for } j = 1, 2, \dots, m.$$




Figure : An example of  $S_j(x) \odot y = z_j$  with  $S_j$  localising “ $x$ ” to the  $j^{\text{th}}$  position.

# Scanning Ptychography: A Crash Course

The mathematical model is:

- The unknown **illumination function**:  $x \in \mathbb{C}^{n \times n}$ ,
- The unknown **specimen**:  $y \in \mathbb{C}^{n \times n}$ ,
- An  $m$ -tuple of **diffraction patterns**:  $\mathbf{z} = (z_1, \dots, z_m) \in (\mathbb{C}^{n \times n})^m$ ,
- The *shift map*  $S_j : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$  moves  $x$  to the position corresponding to the  $j^{\text{th}}$  diffraction pattern measurement.
- The elements of the triple  $(x, y, \mathbf{z})$  are related by:

$$S_j(x) \odot y = z_j \quad \text{for } j = 1, 2, \dots, m.$$




Figure : An example of  $S_j(x) \odot y = z_j$  with  $S_j$  localising “ $x$ ” to the  $j^{\text{th}}$  position.

# Scanning Ptychography: A Crash Course

In a ptychography experiment we observe  $m$  non-negative matrices:

$$b_1, \dots, b_m \in \mathbb{R}_+^{n \times n}.$$

Each corresponds to a different illumination region on the specimen:

$$b_j \equiv |\mathcal{F}(z_j)| \quad \text{where} \quad z_j = S_j(x) \odot y,$$


where  $\mathcal{F}$  is the *2D Fourier transform*, and  $|\cdot|$  is taken element-wise.

The **blind ptychography problem** can now be stated:

Given  $b_1, b_2, \dots, b_m \in \mathbb{R}_+^{n \times n}$  reconstruct the triple  $(x, y, z)$ .

# Two Algorithms in the Literature

Maiden & Rodenburg proposed:




Update functions are of the form:

$$x^{k+1} = x^k + \alpha \frac{S_j^{-1}(y^k \odot \bar{y}^k)}{\|S_j^{-1}(y^k \odot \bar{y}^k)\|_\infty} (z_j^{k+1} - z_j^k).$$

# Two Algorithms in the Literature

Maiden & Rodenburg proposed:



Thibault *et al.* proposed:

$$P_F(\mathbf{z}) : z_j \rightarrow P_F(z_j) \text{ for all } j = 1, 2, \dots, m. \quad (4)$$

$$P_O(\mathbf{z}) : z_j \rightarrow P_O(z_j) \equiv S_j(x) \odot y. \quad (6)$$

$$\mathbf{z}^{k+1} = \mathbf{z}^k + P_F(2P_O(\mathbf{z}^k) - \mathbf{z}^k) - P_O(\mathbf{z}^k). \quad (9)$$

On computing (6):

$$\hat{y}^k = \frac{\sum_{j=1}^m S_j(x^k) \odot z_j^k}{\|\sum_{j=1}^m x^k \odot \bar{x}^k\|}. \quad (7)$$

$$\hat{x}^k = \frac{\sum_{j=1}^m S_j^{-1}(y^k \odot z_j^k)}{\|\sum_{j=1}^m y^k \odot \bar{y}^k\|}. \quad (8)$$

In the event the probe  $P$  is already known, the overlap projection is given by (6), where  $\hat{O}$  is computed with Eq. (7). If  $\hat{P}$  also needs to be retrieved, both Eqs. (7) and (8) need to be simultaneously solved. While the system cannot be decoupled analytically, applying the two equations in turns for a few iterations was observed to be an efficient procedure to find the minimum. Within the reconstruction scheme, initial guesses for  $\hat{P}$  and  $\hat{O}$  are readily available from the previous iteration—apart

Update functions are of the form:

$$x^{k+1} = x^k + \alpha \frac{S_j^{-1}(y^k \odot \bar{y}^k)}{\|S_j^{-1}(y^k \odot \bar{y}^k)\|_\infty} (z_j^{k+1} - z_j^k).$$

# Our Framework

- We considered the following optimisation problem:

$$\begin{aligned} \min \quad & F(x, y, z) := \sum_{j=1}^m \|S_j(x) \odot y - z_j\|^2 \\ \text{s.t.} \quad & x \in X := \{x : \|x\|_\infty \leq M_x, x_{ij} = 0, \forall (i, j) \notin \mathbb{I}_x\}, \\ & y \in Y := \{y : \|y\|_\infty \leq M_y\}, \\ & z \in Z := \{z : |\mathcal{F}(z_j)| = b_j \text{ for } j = 1, 2, \dots, m\}, \end{aligned} \tag{P}$$

where  $M_x, M_y \in \mathbb{R}$  are bounds, and  $\mathbb{I}_x$  is an index set (**support** of  $x$ ).

- Separable constraint sets coupled through an objective function.
  - However, all constraints are  $X, Y, Z$  are closed semi-algebraic sets.
  - $F$  is a continuous semi-algebraic function (graph  $F$  is semi-algebraic).
- (P) is equivalent to the formally unconstrained problem:

$$\min \Psi(x, y, z) := F(x, y, z) + \iota_X(x) + \iota_Y(y) + \iota_Z(z).$$

---

A set  $S \subseteq \mathbb{R}^d$  is **semi-algebraic** if there exists finitely many polynomials  $p_{ij}, q_{ij} : \mathbb{R}^d \rightarrow \mathbb{R}$  such that

$$S = \bigcup_{j=1}^N \bigcap_{i=1}^K \left\{ u \in \mathbb{R}^d : p_{ij}(u) = 0, q_{ij}(u) < 0 \right\}.$$

# Our Framework

- We considered the following optimisation problem:

$$\begin{aligned} \min \quad & F(x, y, z) := \sum_{j=1}^m \|S_j(x) \odot y - z_j\|^2 \\ \text{s.t.} \quad & x \in X := \{x : \|x\|_\infty \leq M_x, x_{ij} = 0, \forall (i, j) \notin \mathbb{I}_x\}, \\ & y \in Y := \{y : \|y\|_\infty \leq M_y\}, \\ & z \in Z := \{z : |\mathcal{F}(z_j)| = b_j \text{ for } j = 1, 2, \dots, m\}, \end{aligned} \tag{P}$$

where  $M_x, M_y \in \mathbb{R}$  are bounds, and  $\mathbb{I}_x$  is an index set (**support** of  $x$ ).

- Separable constraint sets coupled through an objective function.
  - However, all constraints are  $X, Y, Z$  are closed **semi-algebraic sets**.
  - $F$  is a continuous **semi-algebraic function** (**graph**  $F$  is semi-algebraic).
- (P) is equivalent to the formally unconstrained problem:

$$\min \Psi(x, y, z) := F(x, y, z) + \iota_X(x) + \iota_Y(y) + \iota_Z(z).$$

---

A set  $S \subseteq \mathbb{R}^d$  is **semi-algebraic** if there exists finitely many polynomials  $p_{ij}, q_{ij} : \mathbb{R}^d \rightarrow \mathbb{R}$  such that

$$S = \bigcup_{j=1}^N \bigcap_{i=1}^K \left\{ u \in \mathbb{R}^d : p_{ij}(u) = 0, q_{ij}(u) < 0 \right\}.$$

# Our Framework

- We considered the following optimisation problem:

$$\begin{aligned} \min \quad & F(x, y, z) := \sum_{j=1}^m \|S_j(x) \odot y - z_j\|^2 \\ \text{s.t.} \quad & x \in X := \{x : \|x\|_\infty \leq M_x, x_{ij} = 0, \forall (i, j) \notin \mathbb{I}_x\}, \\ & y \in Y := \{y : \|y\|_\infty \leq M_y\}, \\ & z \in Z := \{z : |\mathcal{F}(z_j)| = b_j \text{ for } j = 1, 2, \dots, m\}, \end{aligned} \tag{P}$$

where  $M_x, M_y \in \mathbb{R}$  are bounds, and  $\mathbb{I}_x$  is an index set (**support** of  $x$ ).

- Separable constraint sets coupled through an objective function.
  - However, all constraints are  $X, Y, Z$  are closed **semi-algebraic sets**.
  - $F$  is a continuous **semi-algebraic function** (graph  $F$  is semi-algebraic).
- (P) is equivalent to the formally unconstrained problem:

$$\min \Psi(x, y, z) := F(x, y, z) + \iota_X(x) + \iota_Y(y) + \iota_Z(z).$$

---

A set  $S \subseteq \mathbb{R}^d$  is **semi-algebraic** if there exists finitely many polynomials  $p_{ij}, q_{ij} : \mathbb{R}^d \rightarrow \mathbb{R}$  such that

$$S = \bigcup_{j=1}^N \bigcap_{i=1}^K \left\{ u \in \mathbb{R}^d : p_{ij}(u) = 0, q_{ij}(u) < 0 \right\}.$$

# A Naïve Algorithm: Alternating Minimisation

Alternating Minimisation Algorithm (over three blocks):

**Initialization.** Choose  $(x^0, y^0, z^0) \in X \times Y \times Z$ .

**General Step.** ( $k = 0, 1, \dots$ )

1. Select  $x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k),$

2. Select  $y^{k+1} \in \arg \min_{y \in Y} F(x^{k+1}, y, z^k),$

3. Select  $z^{k+1} \in \arg \min_{z \in Z} F(x^{k+1}, y^{k+1}, z).$

What's involved? Roughly speaking, to compute Step 1 we need to minimise terms of the form  $\|S_j(x) \odot y^k - z_j^k\|^2$ . To do so:

$$S_j(x) \odot y^k \approx z_j^k \implies \underbrace{S_j(x) \approx z_j^k \oslash y_k}_{\text{pointwise division } X} \implies \underbrace{x \approx S_j^{-1}(z_j^k \oslash y_k)}_{\text{un-shift operator } \checkmark}.$$

# A Naïve Algorithm: Alternating Minimisation

Alternating Minimisation Algorithm (over three blocks):

**Initialization.** Choose  $(x^0, y^0, z^0) \in X \times Y \times Z$ .

**General Step.** ( $k = 0, 1, \dots$ )

1. Select  $x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k),$

2. Select  $y^{k+1} \in \arg \min_{y \in Y} F(x^{k+1}, y, z^k),$

3. Select  $z^{k+1} \in \arg \min_{z \in Z} F(x^{k+1}, y^{k+1}, z).$

**What's involved?** Roughly speaking, to compute Step 1 we need to minimise terms of the form  $\|S_j(x) \odot y^k - z_j^k\|^2$ . To do so:

$$S_j(x) \odot y^k \approx z_j^k \implies \underbrace{S_j(x) \approx z_j^k \oslash y_k}_{\text{pointwise division } x} \implies \underbrace{x \approx S_j^{-1}(z_j^k \oslash y_k)}_{\text{un-shift operator } \checkmark}.$$

# PHeBIE: Proximal Block Implicit-Explicit Algorithm

From the previous slide, recall our naïve Step 1:

$$x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k).$$

Replace the objective function  $F$  with a better behaved regularisation:

$$x^{k+1} \in \arg \min_{x \in X} \left( F(x, y^k, z^k) \right)$$

- \* No longer requires any ill-conditioned or unstable operations!

---

Given a set  $C$ , its (nearest point) projection,  $P_C$ , is given by

$$P_C(w) := \arg \min_{u \in C} \|u - w\|.$$

# PHeBIE: Proximal Block Implicit-Explicit Algorithm

From the previous slide, recall our naïve Step 1:

$$x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k).$$

Replace the objective function  $F$  with a better behaved regularisation:

$$x^{k+1} \in \arg \min_{x \in X} \left( \underbrace{F(x^k, y^k, z^k) + \langle x - x^k, \nabla_x F(x^k, y^k, z^k) \rangle}_{\text{linearisation of } F \text{ at } (x^k, y^k, z^k)} \right)$$

- \* No longer requires any ill-conditioned or unstable operations!

---

Given a set  $C$ , its (nearest point) projection,  $P_C$ , is given by

$$P_C(w) := \arg \min_{u \in C} \|u - w\|.$$

# PHeBIE: Proximal Block Implicit-Explicit Algorithm

From the previous slide, recall our naïve Step 1:

$$x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k).$$

Replace the objective function  $F$  with a better behaved regularisation:

$$x^{k+1} \in \arg \min_{x \in X} \left( \underbrace{F(x^k, y^k, z^k) + \langle x - x^k, \nabla_x F(x^k, y^k, z^k) \rangle}_{\text{linearisation of } F \text{ at } (x^k, y^k, z^k)} + \underbrace{\frac{\alpha^k}{2} \|x - x^k\|^2}_{\text{proximal term}} \right)$$

- \* No longer requires any ill-conditioned or unstable operations!

---

Given a set  $C$ , its (nearest point) projection,  $P_C$ , is given by

$$P_C(w) := \arg \min_{u \in C} \|u - w\|.$$

# PHeBIE: Proximal Block Implicit-Explicit Algorithm

From the previous slide, recall our naïve Step 1:

$$x^{k+1} \in \arg \min_{x \in X} F(x, y^k, z^k).$$

Replace the objective function  $F$  with a better behaved regularisation:

$$\begin{aligned} x^{k+1} &\in \arg \min_{x \in X} \left( \underbrace{F(x^k, y^k, z^k) + \langle x - x^k, \nabla_x F(x^k, y^k, z^k) \rangle}_{\text{linearisation of } F \text{ at } (x^k, y^k, z^k)} + \underbrace{\frac{\alpha^k}{2} \|x - x^k\|^2}_{\text{proximal term}} \right) \\ &= P_X \left( x^k - \frac{2}{\alpha^k} \sum_{j=1}^m S_j^{-1}(\bar{y}^k) \odot S_j^{-1}(y^k - z_j^k) \right). \end{aligned}$$

- ★ No longer requires any ill-conditioned or unstable operations!

---

Given a set  $C$ , its (nearest point) projection,  $P_C$ , is given by

$$P_C(w) := \arg \min_{u \in C} \|u - w\|.$$

# PHeBIE: Proximal Block Implicit-Explicit Algorithm

## Proximal Heterogeneous Block Implicit-Explicit Algorithm:

**Initialization.** Choose  $\alpha, \beta, \gamma > 0$  and  $(x^0, y^0, z^0) \in X \times Y \times Z$ .

**General Step.** ( $k = 0, 1, \dots$ )

1. Choose  $\alpha^k > \alpha$  and select

$$x^{k+1} \in P_X \left( x^k - \frac{2}{\alpha^k} \sum_{j=1}^m S_j^{-1}(\bar{y^k}) \odot S_j^{-1}(y^k - z_j^k) \right),$$

2. Choose  $\beta^k > \beta$  and select

$$y^{k+1} \in P_Y \left( y^k - \frac{2}{\beta^k} \sum_{j=1}^m S_j(\bar{x^{k+1}}) \odot (S_j(x^{k+1}) - z_j^k) \right),$$

3. Choose  $\gamma^k > \gamma$  and select

$$z^{k+1} \in P_Z \left( \left[ \frac{2}{2 + \gamma_k} S_j(x^{k+1}) \odot y^{k+1} + \frac{\gamma_k}{2 + \gamma_k} z_j^k \right]_{j=1}^m \right).$$

For convergence we choose:  $\alpha^k \geq L_x(y^k, z^k)$  and  $\beta^k \geq L_y(x^{k+1}, z^k)$  where  $L_x(y^k, z^k)$  and  $L_y(x^{k+1}, z^k)$  denote the partial Lipschitz constant of  $\nabla_x F(\cdot, y^k, z^k)$  and  $\nabla_y F(x^{k+1}, \cdot, z^k)$ .

# PHeBIE: Convergence Theorem

Theorem (Hesse–Luke–Sabach–T, 2015)

Let  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  be a sequence generated by the PHeBIE algorithm for the blind ptychography problem. Then the following hold.

- ① The sequence  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  has finite length. That is,

$$\sum_{k=1}^{\infty} \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\| < \infty.$$

- ② The sequence  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  converges to point  $(x^*, y^*, z^*)$  which is a critical point of the function  $\Psi$ . That is,

$$0 \in \partial\Psi(x, y, z) = \nabla F(x^*, y^*, z^*) + \partial\iota_X(x^*) + \partial\iota_Y(y^*) + \partial\iota_Z(z^*),$$

where  $\partial(\cdot)$  denotes the limiting Fréchet subdifferential.

For  $u$  in the domain of  $f$ , the limiting Fréchet subdifferential is given by

$$\partial f(u) := \left\{ v : \exists u^k \rightarrow u, f(u^k) \rightarrow f(u), v^k \rightarrow v, v^k \in \widehat{\partial}f(u^k) \right\}, \text{ where } \widehat{\partial}f(u) = \left\{ v : \liminf_{\substack{w \neq u \\ w \rightarrow u}} \frac{f(w) - f(u) - \langle v, w - u \rangle}{\|w - u\|} \geq 0 \right\}.$$

# PHeBIE: Example

# PHeBIE: Convergence Theorem (cont.)

*Proof Sketch.*

The proof has three steps:

- ① (Sufficient decrease) Use properties of the algorithm to establish that the sequence  $\{F(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  is decreasing, converges to some  $F^* > -\infty$ , and that

$$\sum_{k=1}^{\infty} \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|^2 < \infty.$$

- ② (Subdifferential bound) Use properties of the algorithm to show that

$$\|w^{k+1}\| \leq \kappa \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|,$$

for some  $w^{k+1} \in \partial\Psi(x^{k+1}, y^{k+1}, z^{k+1})$  and  $\kappa > 0$ .

- ③ To establish convergence of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  to a critical point, uses the fact that  $\Psi$  satisfied the so-called Kurdyka–Łojasiewicz (KL) Property which gives Cauchyness of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$ . □

---

The proof strategy is based on a paper of Bolte, Sabach Teboulle (2014).

# PHeBIE: Convergence Theorem (cont.)

*Proof Sketch.*

The proof has three steps:

- ① (Sufficient decrease) Use properties of the algorithm to establish that the sequence  $\{F(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  is decreasing, converges to some  $F^* > -\infty$ , and that

$$\sum_{k=1}^{\infty} \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|^2 < \infty.$$

- ② (Subdifferential bound) Use properties of the algorithm to show that

$$\|w^{k+1}\| \leq \kappa \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|,$$

for some  $w^{k+1} \in \partial\Psi(x^{k+1}, y^{k+1}, z^{k+1})$  and  $\kappa > 0$ .

- ③ To establish convergence of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  to a critical point, uses the fact that  $\Psi$  satisfied the so-called Kurdyka–Łojasiewicz (KL) Property which gives Cauchyness of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$ . □

---

The proof strategy is based on a paper of Bolte, Sabach Teboulle (2014).

# PHeBIE: Convergence Theorem (cont.)

*Proof Sketch.*

The proof has three steps:

- ① (Sufficient decrease) Use properties of the algorithm to establish that the sequence  $\{F(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  is decreasing, converges to some  $F^* > -\infty$ , and that

$$\sum_{k=1}^{\infty} \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|^2 < \infty.$$

- ② (Subdifferential bound) Use properties of the algorithm to show that

$$\|w^{k+1}\| \leq \kappa \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|,$$

for some  $w^{k+1} \in \partial\Psi(x^{k+1}, y^{k+1}, z^{k+1})$  and  $\kappa > 0$ .

- ③ To establish convergence of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  to a critical point, uses the fact that  $\Psi$  satisfied the so-called Kurdyka–Łojasiewicz (KL) Property which gives Cauchyness of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$ . □

The proof strategy is based on a paper of Bolte, Sabach Teboulle (2014).

# PHeBIE: Convergence Theorem (cont.)

*Proof Sketch.*

The proof has three steps:

- ① (Sufficient decrease) Use properties of the algorithm to establish that the sequence  $\{F(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  is decreasing, converges to some  $F^* > -\infty$ , and that

$$\sum_{k=1}^{\infty} \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|^2 < \infty.$$

- ② (Subdifferential bound) Use properties of the algorithm to show that

$$\|w^{k+1}\| \leq \kappa \|(x^{k+1}, y^{k+1}, z^{k+1}) - (x^k, y^k, z^k)\|,$$

for some  $w^{k+1} \in \partial\Psi(x^{k+1}, y^{k+1}, z^{k+1})$  and  $\kappa > 0$ .

- ③ To establish convergence of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$  to a critical point, uses the fact that  $\Psi$  satisfied the so-called Kurdyka–Łojasiewicz (KL) Property which gives Cauchyness of  $\{(x^k, y^k, z^k)\}_{k \in \mathbb{N}}$ . □

The proof strategy is based on a paper of Bolte, Sabach Teboulle (2014).

# The Kurdyka-Łojasiewicz (KL) Property

Roughly speaking, functions which satisfy the **KL-property** can be made “sharp” by a reparametrisation of their range by an increasing function. A simple example: the function  $f(x) = x^2$  can be reparametrised by  $\varphi(x) = \sqrt{|x|}$ :



- The **utility of the KL-property** is that it holds for many real world optimisation problems which lack convexity (e.g., ptychography!).

## Theorem (Bolte–Danillidis–Lewis 2006)

Every proper, lower semi-continuous, **semi-algebraic** function satisfies the KL-property throughout its domain.

Let  $f : \mathbb{R}^d \rightarrow (-\infty, +\infty]$  be proper. For  $\eta \in (0, +\infty]$  define

$$\mathcal{C}_\eta \equiv \{\varphi : [0, \eta] \rightarrow \mathbb{R}_+ : \varphi(0) = 0, \varphi'(s) > 0 \text{ for all } s \in (0, \eta)\}.$$

The function  $f$  has **KL property** at  $\bar{u} \in \text{dom } \partial f$  if there exists a neighbourhood  $U$  of  $\bar{u}$  and a function  $\varphi \in \mathcal{C}_\eta$ , such that, for all  $u \in U : f(\bar{u}) < f(u) < f(\bar{u}) + \eta$ , it holds that

$$\underbrace{\varphi'(f(u) - f(\bar{u}))}_{\text{Think: minimum norm element of } \partial(\varphi \circ g)} \text{dist}(0, \partial f(u)) \geq 1.$$

Think: minimum norm element of  $\partial(\varphi \circ g)$  where  $g = f - f(\bar{u})$ .

# Interpreting Current State-of-the-Art Algorithms

The update steps all of the algorithms: PHeBIE algorithm, Madien & Rodenburg, and Thibault *et al.* are **more-or-less the same**.

- The PHeBIE algorithm:
  - Minimises w.r.t. three blocks  $X, Y, Z$  in cyclic order.
  - Each  $x$ -update/ $y$ -update uses **all  $m$  diffraction patterns**. In Step 1, the weight  $\alpha^k$  given by partial Lipschitz constant of  $\nabla_x F(\cdot, y^k, z^k)$  is

$$L_x(y^k, z^k) = 2 \left\| \left( \sum_{j=1}^m S_j^*(\bar{y}^k \odot y^k) \right) \right\|_\infty.$$

- Madien & Rodenburg method:
  - Can be viewed as minimisation w.r.t. the blocks  $X, Y$  and  $Z$ .
  - Each  $x$ -update/ $y$ -update uses **only a single diffraction pattern**. In Step 1, the weight when updating with the  $j$ th diffraction pattern is

$$2 \left\| S_j^*(\bar{y}^k \odot y^k) \right\|_\infty.$$

- Thibault *et al.* method:
  - Minimise w.r.t. three blocks  $X, Y, Z$ , but **many  $X, Y$  updates are performed between  $Z$  updates**.
  - Updates embedded within a “difference map” algorithmic structure.

# Interpreting Current State-of-the-Art Algorithms

The update steps all of the algorithms: PHeBIE algorithm, Madien & Rodenburg, and Thibault *et al.* are **more-or-less the same**.

- The PHeBIE algorithm:
  - Minimises w.r.t. three blocks  $X, Y, Z$  in cyclic order.
  - Each  $x$ -update/ $y$ -update uses **all  $m$  diffraction patterns**. In Step 1, the weight  $\alpha^k$  given by partial Lipschitz constant of  $\nabla_x F(\cdot, y^k, z^k)$  is

$$L_x(y^k, z^k) = 2 \left\| \left( \sum_{j=1}^m S_j^*(\bar{y}^k \odot y^k) \right) \right\|_\infty.$$

- Madien & Rodenburg method:
  - Can be viewed as minimisation w.r.t. the blocks  $X, Y$  and  $Z$ .
  - Each  $x$ -update/ $y$ -update uses **only a single diffraction pattern**. In Step 1, the weight when updating with the  $j$ th diffraction pattern is

$$2 \left\| S_j^*(\bar{y}^k \odot y^k) \right\|_\infty.$$

- Thibault *et al.* method:
  - Minimise w.r.t. three blocks  $X, Y, Z$ , but **many  $X, Y$  updates are performed between  $Z$  updates**.
  - Updates embedded within a “difference map” algorithmic structure.

# Interpreting Current State-of-the-Art Algorithms

The update steps all of the algorithms: PHeBIE algorithm, Madien & Rodenburg, and Thibault *et al.* are **more-or-less the same**.

- The PHeBIE algorithm:
  - Minimises w.r.t. three blocks  $X, Y, Z$  in cyclic order.
  - Each  $x$ -update/ $y$ -update uses **all  $m$  diffraction patterns**. In Step 1, the weight  $\alpha^k$  given by partial Lipschitz constant of  $\nabla_x F(\cdot, y^k, z^k)$  is

$$L_x(y^k, z^k) = 2 \left\| \left( \sum_{j=1}^m S_j^*(\bar{y}^k \odot y^k) \right) \right\|_\infty.$$

- Madien & Rodenburg method:
  - Can be viewed as minimisation w.r.t. the blocks  $X, Y$  and  $Z$ .
  - Each  $x$ -update/ $y$ -update uses **only a single diffraction pattern**. In Step 1, the weight when updating with the  $j$ th diffraction pattern is

$$2 \left\| S_j^*(\bar{y}^k \odot y^k) \right\|_\infty.$$

- Thibault *et al.* method:
  - Minimise w.r.t. three blocks  $X, Y, Z$ , but **many  $X, Y$  updates are performed between  $Z$  updates**.
  - Updates embedded within a “difference map” algorithmic structure.

simultaneously solved. While the system cannot be decoupled analytically, applying the two equations in turns for a few iterations was observed to be an efficient procedure to find the minimum. Within the reconstruction scheme, initial guesses for  $\hat{P}$

# Concluding Remarks and Ongoing Work

In summary:

- We have presented an algorithm for scanning ptychography within a clear mathematical optimisation framework.
- Under practically verifiable assumptions, the algorithm is **provably convergent** to **critical points** of the function  $\Psi \equiv F + \iota_X + \iota_Y + \iota_Z$ .
- The framework is flexible and allows for an interpretations of current state-of-the-art ptychography algorithms.

Ongoing and future work:

- Can the critical points of  $\Psi$  be characterised in a meaningful way?
- What happens when the data is noisy? Our convergence theorem holds independently of the presence of noise in the data.

---

Further details can be found in:

**Proximal Heterogeneous Block Implicit-Explicit Method and Application to Blind Ptychographic Diffraction Imaging** with R. Hesse, D.R. Luke and S. Sabach. *SIAM J. on Imaging Sciences*, 8(1):426–457 (2015).