

Complex Analysis

Matthew M. Jones

October 2019

Contents

1	Complex Numbers	1
1.1	Complex numbers	1
1.2	The field of complex numbers	3
1.3	The geometry of complex numbers	5
1.4	The metric space of complex numbers	8
1.5	Power series	11
1.6	Differentiation	12
2	Complex Differentiation	19
2.1	Holomorphic functions	19
2.2	The Cauchy-Riemann equations	24
2.3	Holomorphic functions as vector fields	29
2.4	Power series	30
3	Special Functions	41
3.1	Introduction	41
3.2	The exponential function	41
3.3	The logarithm	44
3.4	Complex powers	50
3.5	Mappings of given domains	53
3.6	Trigonometric functions	53
4	Complex Integration	57

4.1	Integration of complex functions	57
4.2	Integration along curves	58
4.3	Integration with respect to arc-length	63
4.4	Some topological notions	64
4.5	Cauchy's theorem	67
4.6	The weak form of Cauchy's Theorem	69
4.7	Proof of Cauchy's theorem	70
5	Applications of Cauchy's Theorem	81
5.1	Doubly connected domains	81
5.2	Cauchy's integral formula	83
5.3	Holomorphic functions are analytic	86
5.4	Zeros of holomorphic functions	89
5.5	Liouville's theorem	91
6	The Residue Theorem	97
6.1	Singularities	97
6.2	The Residue Theorem	101
6.3	Simple poles	105
6.4	Applications of the Residue Theorem to real integrals	107
6.4.1	The Cauchy principal value	109
6.4.2	Integrals involving trigonometric functions	113
7	Mapping Properties	119
7.1	The maximum modulus	119
7.2	The argument principle	122

Preparation and Other Reading

“I am always doing that which I can not do, in order that I may learn how to do it.”

Pablo Picasso

These notes will introduce complex analysis. This subject is fundamental to many areas of mathematics and has a long history dating back to the mid 19th century. Many important mathematicians including Cauchy and Riemann initiated the work and you will see their names aligned to results throughout these notes.

There are many textbooks on complex analysis that might help you to understand concepts and delve deeper into areas not covered, the main recommendations are:

- Garling, D. J. H., “A Course in Mathematical Analysis, volume III”, [3].

This is a new textbook; it covers all of the topics, and extends some of what we will go over. This is a more modern book but it does avoid some of the geometric insights of some of the concepts. This is largely because the author is not a native complex analyst like the books below. But at least he is still alive.

- Ahlfors, L. V., “Complex Analysis”, [1].

This is considered one of the classics in complex analysis. Lars Ahlfors was one of the original Fields medal winners and he revolutionised many areas of complex analysis by introducing a geometric intuition to many of the topics. He is considered to be one of the greatest complex analysts of the 20th century. This book covers all of the module and is written by a master.

- Rudin, W., “Real and Complex Analysis”, [8].

This is another classic, including two sections covering real and complex analysis separately. This book departs a little from what we learn in this module and includes areas of current research.

- Remmert, R, “Theory of Complex Functions”, [7]. I’ll not reference this book very much but there’s a copy in the library. It covers everything we will do in this module and includes quite a lot of detail (and some history of the authors of much of the work).

Chapter 1

Complex Numbers

“I have also learned not to take glory in the difficulty of a proof: difficulty means we have not understood. The idea is to be able to paint a landscape in which the proof is obvious.”

Alexander Grothendieck, quoted by Pierre Deligne, *Notices of the AMS* 63 (2016), p. 250

1.1 Complex numbers

The field of complex numbers is represented by the symbol \mathbb{C} . Elements of \mathbb{C} are normally written in the form

$$a + bi \quad i^2 = -1$$

where $a, b \in \mathbb{R}$. If we treat i as a variable then we may define arithmetic in \mathbb{C} in the same way as we define arithmetic of polynomials in $\mathbb{R}[i]$, with the additional assumption that $i^2 = -1$. In particular we have

$$(a + bi) + (c + di) = (a + c) + (b + d)i, \quad \text{and} \\ (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

The value of a in the complex number $a+bi$ is called the **real part** and denoted $Re(a+bi)$. The value of b is called the **imaginary part** and denoted $Im(a + bi)$.

Given a complex number we can describe it using the exponential form

$$a + bi = re^{i\theta}$$

where r and θ are related to a and b in the following way

$$a = r \cos \theta \quad b = r \sin \theta.$$

These follow from Euler's formula

$$e^{i\theta} = \cos \theta + i \sin \theta \quad (1.1)$$

The value r is often written as $|\cdot|$ and called the modulus or absolute value, for example $|1 + i| = \sqrt{2}$. The value of θ is called the argument and denoted by \arg . It should be noted here that the argument is not uniquely defined since any multiple of 2π can be added or subtracted to produce the argument of the same complex number.

This representation is often called the exponential representation and is more convenient in many ways. For example we have the product formula:

$$(r_1 e^{i\theta_1})(r_2 e^{i\theta_2}) = r_1 r_2 e^{i(\theta_1 + \theta_2)}.$$

For example it follows that

$$\frac{1}{re^{i\theta}} = \frac{1}{r} e^{-i\theta}.$$

We will normally reserve the letters z and w for complex numbers and x and y for real numbers.

Finally, if $z = a + ib$ then the complex conjugate is $\bar{z} = a - ib$. In exponential form

$$\overline{re^{i\theta}} = re^{-i\theta}.$$

Geometrically the conjugate of a complex number represents the reflection in the real axis of the Argand diagram.

We collect here properties of the conjugate and modulus of complex numbers.




Figure 1.1: The exponential form of a complex number

Proposition 1.1

Suppose $z, w \in \mathbb{C}$ then the following hold.

1. $\operatorname{Re} z = \frac{1}{2}(z + \bar{z}), \operatorname{Im} z = \frac{1}{2i}(z - \bar{z})$
2. $\overline{z + w} = \bar{z} + \bar{w}, \overline{zw} = \bar{z}\bar{w}$
3. $|z|^2 = z\bar{z}$
4. $|z + w| \leq |z| + |w|$ (the Triangle inequality)
5. $-|z| \leq \operatorname{Re} z \leq |z|$
6. $-|z| \leq \operatorname{Im} z \leq |z|$

1.2 The field of complex numbers

The complex number field first came about as the smallest splitting field over \mathbb{R} . This means that every polynomial with real coefficients can be factorised over the complex numbers. This follows from the following important theorem which we will prove at the end of the course as a corollary to Liouville's theorem.

Theorem 1.2 – The fundamental theorem of algebra

Let $p(x) \in \mathbb{C}[x]$ have $\deg p \geq 1$. Then there is $\alpha \in \mathbb{C}$ such that $p(\alpha) = 0$ and p can be factorised as follows

$$p(x) = (x - \alpha)q(x)$$

where $q(x) \in \mathbb{C}[x]$.

The value α is called a root of the polynomial. Although this theorem only guarantees a single factor, it can be proved by induction that a polynomial $p(x)$ with $\deg p = n$ has exactly n roots (not necessarily distinct).

The solutions to equations of the form $z^n - a = 0$ are of particular importance in complex analysis. Finding solutions to this equation is equivalent to finding the n th roots of the number a . The process of finding n th roots starts with finding the roots of unity, these are the solutions to the equation $z^n - 1$. The n distinct roots of unity are

$$\{1, \omega, \omega^2, \dots, \omega^{n-1}\} \quad \text{where } \omega = e^{2\pi i/n}. \quad (1.2)$$

An example of these are plotted on the left in Figure 1.2.




Figure 1.2: (Left) roots of unity with $n = 7$ and (right) cube roots of -8 .

Example 1.3. Find the 3rd roots of unity.

Here $n = 3$ so $\omega = e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$, and

$$\omega^2 = e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i.$$

In general it is possible to find the n th roots of any number using the same technique. If $z = re^{i\theta}$ then $\eta = r^{1/n}e^{i\theta/n}$ is the **principle n th root of z** . The n th roots of z are then given by

$$\{\eta, \eta\omega, \eta\omega^2, \dots, \eta\omega^{n-1}\},$$

where ω is the n th root of unity.

Example 1.4. Solve $z^3 + 8 = 0$.

First we write $-8 = 8e^{\pi i}$. Then $\eta = 2e^{\pi i/3}$ is the principle 3rd root of -8 . The 3rd roots of -8 are then

$$\{\eta, \eta\omega, \eta\omega^2\}.$$

These can be calculated as

$$\left\{ \frac{1}{2} + \frac{\sqrt{3}}{2}i, -1, \frac{1}{2} - \frac{\sqrt{3}}{2}i \right\}.$$

In the right-hand diagram in Figure 1.2 these roots are plotted. They are in a familiar pattern of an equilateral triangle.

1.3 The geometry of complex numbers

Complex analysis relies very much on geometric intuition, indeed the domain of definition of a function – as you will shortly see – is given much higher status than it is in real analysis and functions are studied much more in terms of their mapping properties. In this section we'll look at some important algebraic definitions of sets in \mathbb{C} .

Example 1.5. Sketch the set $B(i, 1) = \{z \in \mathbb{C} : |z - i| < 1\}$.

The quantity $|z - i|$ represents the distance from the variable z to the point i on the complex plane. This set consists of all the points closer than 1 to i and is the (open) disk drawn on the left in Figure 1.3.




Figure 1.3: The sets in Examples 1.5 and 1.6

In general a set of the form

$$B(a, r) = \{z \in \mathbb{C} : |z - a| < r\}$$

represents the open disk of radius r centred at the point a .

Related to the disk, the following are the **circle** and the **exterior of a disk**, respectively:

$$\partial B(a, r) = \{z \in \mathbb{C} : |z - a| = r\}, \quad \text{and} \quad B(a, r)^* = \{z \in \mathbb{C} : |z - a| > r\}.$$

Example 1.6. Sketch the set $B(0, 1/3)^* = \{z \in \mathbb{C} : |z| > 1/3\}$

This time this set is the exterior of the disk of radius $1/3$ centred at the origin. A sketch of this is given on the right in Figure 1.3, it is the unbounded region external to the disk.

It is possible to combine these sets to form other examples. In Figure 1.4 a number of examples are given, they are respectively

$$A = \{z \in \mathbb{C} : 1 < |z - i| < 2\}$$

$$C = \{z \in \mathbb{C} : |z| < 1\} \setminus \{z \in \mathbb{C} : |z - 1/2| < 1/2\}$$

It will be useful to define straight lines in \mathbb{C} . The easiest way to do this is to simply note that a set such as

$$\{xe^{i\theta} : x \in \mathbb{R}\}$$





Figure 1.4: Other sets in \mathbb{C} : an annulus and a crescent

defines a straight line extending, in both directions, towards infinity. The line passes through the origin and is in the same direction as $e^{i\theta}$. If we want to define a line in general that passes through a point $w \in \mathbb{C}$ in the direction $e^{i\theta}$ we have

$$L = \{w + xe^{i\theta} : x \in \mathbb{R}\}.$$

Note that the direction $e^{i\theta}$ can be replaced by $e^{-i\theta}$, i.e.


$$\{w + xe^{i\theta} : x \in \mathbb{R}\} = \{w + xe^{-i\theta} : x \in \mathbb{R}\}.$$



Following from the definition of a line above we can define a ray, that is a half-line pointing in a particular direction. For example the line emanating from the point $1 + 2i$

in the direction $\pi/4$ is drawn in the following diagram. We can write this as

$$\{z: \arg(z - (1 + 2i)) = \pi/4\}$$



In general a ray starting at a point a and heading in the direction θ can be described as

$$\{z: \arg(z - a) = \theta\}.$$

1.4 The metric space of complex numbers

The complex numbers \mathbb{C} form a metric space with the metric

$$d(z, w) = |z - w|.$$

We define the mapping

$$\begin{aligned} J: \mathbb{C} &\rightarrow \mathbb{C} \\ a + ib &\mapsto \begin{pmatrix} a \\ b \end{pmatrix}. \end{aligned} \tag{1.3}$$

The mapping J is a metric space isometry from the metric space (\mathbb{C}, d) to the metric space (\mathbb{R}^2, d_2) where d_2 is the standard Euclidean metric. The definitions of open sets, closed sets, continuous functions etc. on (\mathbb{C}, d) can be thought of in terms of the metric space (\mathbb{R}^2, d_2) . In particular, by the Heine-Borel theorem, a set A in \mathbb{C} is compact if and only if it is closed and bounded.

For example an ϵ -neighbourhood of a point $a \in \mathbb{C}$ is the set $B(a, \epsilon)$ and is an open disk centred at a with radius $\epsilon > 0$.

A sequence of complex numbers z_0, z_1, z_2, \dots will be denoted by $(z_n)_{n=0}^\infty$. A sequence is said to converge to the limit z if for each $\epsilon > 0$ there is a $N > 0$ such that

$$n > N \quad \Rightarrow \quad |z_n - z| < \epsilon.$$

Familiar rules of real sequences apply to complex sequences. For example limits are unique, the limit of the sum of two (or more) sequences is the sum of the limits. Furthermore a convergent sequence is bounded. Recall that a set is bounded in a metric space if it is contained in an open ball $B(a, r)$. In the complex metric space this is equivalent to the existence of a $r > 0$ such that each element of the set has absolute value less than r .

Proposition 1.7

If (z_n) is a convergent complex-valued sequence then it is bounded.

Proof. Let $f(z) = |z| = d(0, z)$ then f is a continuous function from \mathbb{C} to \mathbb{R} by standard results from metric spaces. It follows that if $z_n \rightarrow z$ as $n \rightarrow \infty$ then

$$|z_n| \rightarrow |z| \quad \text{as } n \rightarrow \infty.$$

Since this is a real-valued convergent sequence it is bounded and hence (z_n) is bounded. \square

Other common results transfer from real analysis via the next result.

Proposition 1.8

Suppose $z_n = x_n + iy_n$ for all $n \geq 0$. The sequence (z_n) converges if and only if the sequences (x_n) and (y_n) converge. When the limits exist

$$\lim_{n \rightarrow \infty} z_n = \lim_{n \rightarrow \infty} x_n + i \lim_{n \rightarrow \infty} y_n.$$

The sequences (x_n) and (y_n) can be described using the isometry (1.3) as

$$\begin{pmatrix} x_n \\ y_n \end{pmatrix} = J(z_n) \quad \text{for all } n.$$

Proof. Suppose that (z_n) converges to $z = x + iy$. Then since $\operatorname{Re} w \leq |\operatorname{Re} w| \leq |w|$ for any $w \in \mathbb{C}$ we have that

$$\begin{aligned} d(x_n, x) &= |x_n - x| \\ &= |\operatorname{Re}(z_n - z)| \\ &\leq |z_n - z| \\ &= d(z_n, z) \end{aligned}$$

It follows that if $d(z_n, z) \rightarrow 0$ then $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$. The same argument with the imaginary part instead shows that $d(y_n, y) \leq d(z_n, z)$ and so, again, if $d(z_n, z) \rightarrow 0$ then $d(y_n, y) \rightarrow 0$ as $n \rightarrow \infty$.

To prove the converse is easier. Suppose that

$$x_n \rightarrow x, \quad \text{and} \quad y_n \rightarrow y$$

as $n \rightarrow \infty$. Then, with $z = x + iy$,

$$\begin{aligned} d(z_n, z) &= |x_n + iy_n - (x + iy)| \\ &= |(x_n - x) + i(y_n - y)| \\ &\leq |(x_n - x)| + |i| |y_n - y| \\ &= d(x_n, x) + d(y_n, y). \end{aligned}$$

Since $d(x_n, x) \rightarrow 0$ and $d(y_n, y) \rightarrow 0$ as $n \rightarrow \infty$ we have that $d(z_n, z) \rightarrow 0$ and the lemma is proved. \square

Lastly we have the following result that will be used throughout.

Proposition 1.9

A complex sequence (z_n) tends to L if and only if the real sequence $(|z_n - L|)$ tends to 0.

Proof. This is simply a restatement of the definition. \square

It is not possible to define the **lim sup** and **lim inf** of a complex sequence since \mathbb{C} does not have an ordering by \leq like the set of real numbers does and therefore there is no concept of a supremum or infimum for \mathbb{C} .

A series of complex numbers $\sum_{n=0}^{\infty} a_n$ is said to converge if the sequence of partial sums

$$s_N = \sum_{n=0}^N a_n$$

converge as a sequence. A series is said to converge absolutely if

$$\sum_{n=0}^{\infty} |a_n|$$

converges. Since the latter is a real-valued series the tests of convergence that you have come across elsewhere can be used to determine if a given series converges absolutely or not. Although we won't prove it in this module it can be shown quite easily that if a series converges absolutely then it converges.

1.5 Power series

Power series are particularly important in complex analysis, these are functions of the form

$$\sum_{n=0}^{\infty} a_n (z - a)^n$$

A power series converges for the value z whenever the limit

$$\lim_{N \rightarrow \infty} \sum_{n=0}^N a_n (z - a)^n$$

exists.

In any introductory analysis course you learn that given a power series with real coefficients there is a value $R \geq 0$ such for which the series converges absolutely for all $|z - a| < R$. The number R is defined by

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} |a_n|^{1/n},$$

and is called the **radius of convergence** of the power series. It is possible for the limit above to be 0 in which case we say that $R = \infty$ and the power series converges for all values of z . Similarly it is possible for the limit to be infinite in which case the power series does not converge absolutely for any value of z (except a trivially).

The set in which a power series converges is therefore a disk

$$B(a, R) = \{z : |z - a| < R\}$$

(hence the use of 'radius' to describe R), Figure 1.5.




Figure 1.5: The disk centred at a with radius $R > 0$

1.6 Differentiation

The main object of study in complex analysis is a complex differentiable function. We will see that the implications of complex differentiability are more restrictive than those of real differentiability. It is worth discussing briefly the definitions of both before we study them in more detail.

Suppose that $\mathbf{f}(\mathbf{x}) = (f_1(\mathbf{x}), f_2(\mathbf{x}))^T$ is a vector field from some open set $A \subset \mathbb{R}^2$ into \mathbb{R}^2 . Here

$$\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}.$$

If the components of \mathbf{f} are continuously differentiable at each point in A then the derivative of \mathbf{f} at \mathbf{x} is given by

$$\nabla \mathbf{f}(\mathbf{x}) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(\mathbf{x}) & \frac{\partial f_1}{\partial y}(\mathbf{x}) \\ \frac{\partial f_2}{\partial x}(\mathbf{x}) & \frac{\partial f_2}{\partial y}(\mathbf{x}) \end{pmatrix}$$

Recall that this means that

$$\lim_{\mathbf{h} \rightarrow 0} \frac{|\mathbf{f}(\mathbf{x}_0 + \mathbf{h}) - \mathbf{f}(\mathbf{x}_0) - \nabla \mathbf{f}(\mathbf{x}_0)\mathbf{h}|}{|\mathbf{h}|} = 0$$

In other words the expression

$$\mathbf{f}(\mathbf{x}_0) + \nabla \mathbf{f}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)$$

is the best linear approximation to $\mathbf{f}(\mathbf{x})$ in a neighbourhood of \mathbf{x}_0 .

On the other hand suppose instead that we have a function

$$\begin{aligned} f: \mathbb{C} &\rightarrow \mathbb{C} \\ z &\mapsto f(z). \end{aligned}$$

We define the complex derivative of f at the point z_0 as

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

whenever this limit exists.

This is equivalent to the condition that

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0) - f'(z_0)h}{h} = 0.$$

These two definitions of differentiation are not the same.

Example 1.10. Let $f(z) = |z|$. Find the real derivative of z at 1 and the complex derivative.

Given the isometry J defined in (1.3) we can always define the equivalent vector field $\mathbf{f}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$\mathbf{f} = J \circ f \circ J^{-1}.$$

In this example we have that

$$\mathbf{f}(\mathbf{x}) = (|\mathbf{x}|, 0)^T.$$

We need to calculate this derivative at $\mathbf{x} = (1, 0)^T = J(1)$.

Now the partial derivatives of this exist and are continuous at $(1, 0)^T$ and so by Theorem 3.17 of Nick's notes we can write

$$\nabla \mathbf{f} = \begin{pmatrix} x(x^2 + y^2)^{-1/2} & y(x^2 + y^2)^{-1/2} \\ 0 & 0 \end{pmatrix}.$$

Evaluated at $(1, 0)^T$ we have that

$$\nabla \mathbf{f}((1, 0)^T) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

On the other hand to calculate the complex derivative we need to find the limit as $h \rightarrow 0$ of the fraction

$$\frac{f(1 + h) - f(1)}{h} = \frac{|1 + h| - 1}{h}.$$

This limit does not exist. To see this suppose first that $h \rightarrow 0$ along the real line, then $|1 + h| = 1 + h$ and

$$\frac{|1 + h| - 1}{h} = \frac{1 + h - 1}{h} = 1.$$

On the other hand if $h \rightarrow 0$ along the imaginary axis, then for we can write $h = ih'$. Now

$$\frac{|1 + h| - 1}{h} = \frac{(1 + (h')^2)^{1/2} - 1}{ih'}.$$

The limit as $h' \rightarrow 0$ of the fraction on the right is 0 which is different to the limit we found when z converged to 1 along the real line. Therefore the complex derivative doesn't exist.

This example begins to highlight the difficulty with defining complex differentiation. Namely the limit $h \rightarrow 0$ can approach from any direction. You encountered this type of issue when you looked at directional derivatives with Nick.

It is for this reason that complex analysis places a higher emphasis on the **domain** of a function than real analysis does. A set A in \mathbb{C} is said to be **connected** if there is a path from any point to any other point in A .

Definition 1.11

A set Ω in \mathbb{C} is a **domain** if it is open and connected.

Throughout these complex analysis notes, unless otherwise stated, the set Ω will be a domain. It is usual to define complex-valued functions from a domain to the complex plane. For example

$$\begin{aligned} f: \Omega &\rightarrow \mathbb{C} \\ z &\mapsto f(z). \end{aligned}$$

Things to know

- Complex numbers – addition, multiplication and division;
- Polar and exponential forms, and how to convert a complex number into any given form;
- Complex conjugates;

- The statement of Proposition 1.1;
- How to calculate the n th roots of a given complex number;
- Geometric descriptions of sets such as $\{z \in \mathbb{C} : |z - a| = r\}$;
- Definitions of convergence of complex sequences and series, how to find the limit of common sequences;
- The statements of Propositions 1.4 and 1.5.

Further reading

There is very little in the way of introductory material in [3]. Ahlfors, [1, Chapter 1, §1 - 2], provides a nice description of most of the properties of complex numbers, Problem 4 is taken from the Exercises in this chapter. For a more detailed discussion of complex numbers see Remmert, [7, Chapter 0]. There is, in particular, a more detailed discussion of Problem 4 there that will give you a good grounding for the Cauchy-Riemann equations that we'll look at in the next chapter.

Problems

1. If $z = 1 + 2i$ and $w = 2 - i$ find

$$\frac{3}{z} + \frac{1}{w}.$$

2. Use Euler's formula to show that

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}.$$

Deduce a formula for $\tan \theta$ in terms of $e^{i\theta}$.

3. Draw the region defined by the following

(a) $\left| \frac{z-3}{z+3} \right| = 2$

(b) $\left| \frac{z-3}{z+3} \right| < 2$

(c) $|z| = |z-2|$

(d) $1 < |z-i| < 2$

4. (a) Find the 5th roots of -1

- (b) Find the cube roots of i

- (c) Find the cube roots of $-i$

- (d) Find the 4th roots of $1+i$

- (e) Find the cube roots of $2 + 2\sqrt{3}i$

5. Sketch the following regions in \mathbb{C}

(a) $\{z \in \mathbb{C} : |z+1| < 2\}$

(b) $\{z \in \mathbb{C} : \operatorname{Re}(z - 1/2) > 0\}$

(c) $\{z \in \mathbb{C} : \operatorname{Re}(z - i) > 0\}$

(d) $\{z \in \mathbb{C} : |z-i| > |z+i|\}$

(e) $\{z \in \mathbb{C} : |z-i| > 2|z+i|\}$

6. Find the solution of $z^2 + 3iz + i = 0$. Why are the roots **not** complex conjugates?

7. Decide whether the following sequences converge or diverge:

(a) $\left(\frac{i^n}{n}\right);$

(b) $\left(\frac{(1+i)^n}{n}\right).$

8. Prove that if $z_n \rightarrow z$ then $|z_n| \rightarrow |z|$ as $n \rightarrow \infty$. Give a counterexample to show that the converse is **not** true.

9. Show that whenever $|z| < 1$

$$1 + z + z^2 + \dots = \frac{1}{1 - z}.$$

* 10. Show that \mathbb{C} is isomorphic to $\mathbb{R}[x]/(x^2 + 1)$, the field of polynomials with real coefficients factored by $x^2 + 1$.

* 11. Let \mathcal{C} be the set of real-valued matrices of the form

$$\begin{pmatrix} a & b \\ -b & a \end{pmatrix}.$$

(a) Show that \mathcal{C} is a vector space over \mathbb{R} (show that it's closed under addition, and scalar multiplication)

(b) Show that the matrices

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

form a basis for \mathcal{C} and hence that \mathcal{C} is 2-dimensional.

(c) Show that the map $H: \mathcal{C} \rightarrow \mathbb{C}$ defined by

$$H(aI + bJ) = a + bi$$

is bijective.

* 12. Show that if we think of z and w as vectors in \mathbb{C} then the dot product is given by $\operatorname{Re} z \bar{w}$.

Chapter 2

Complex Differentiation

“Excellence is for the few but dedication is something everybody can have – should have – and without it life is not worth living.”
I want to be a mathematician, Paul Halmos

2.1 Holomorphic functions

Complex analysis can be summarised as a study of complex-differentiable functions (or, more accurately, holomorphic functions). In the last part you saw that complex-differentiation is different to real-differentiation. We will start this topic by comparing complex and real-differentiation via the Cauchy-Riemann equations but end up highlighting the differences between the two.

Definition 2.1 – Holomorphic functions

A complex-valued function f defined on a domain Ω is **holomorphic** if the derivative

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

exists for all values $z_0 \in \Omega$. We denote the set of holomorphic functions on Ω as $H(\Omega)$. A function $f \in H(\mathbb{C})$ is called **entire**.

It is important that whenever we discuss complex functions we state or assume that

the function is defined on a domain (in the sense of Definition 1.11). In the words of Lars V. Ahlfors, [1]:

“It is important to stress that the open set Ω is part of the definition. As a rule one should avoid speaking of a [holomorphic] function $f(z)$ without referring to a specific open set Ω on which it is defined.”

Example 2.2. *If p is a polynomial then $p \in H(\mathbb{C})$. So p is entire.*

You will study this example later.

Example 2.3. *The function*

$$f(z) = \frac{1}{z}$$

is in $H(\Omega)$ where

$$\Omega = \{z: 1 < |z| < 2\}.$$

Here the derivative is

$$f'(z) = -\frac{1}{z^2}.$$

Example 2.4. *The function*

$$f(z) = \frac{1}{z-1}$$

is holomorphic on any domain that does not contain the point 1.

For example we could take the unit disc centred at the origin:

$$\Omega = B(0, 1) = \{z: |z| < 1\}.$$

The derivative for this function is

$$f'(z) = -\frac{1}{(z-1)^2}.$$

For the remainder of these notes we will often refer to complex-differentiability simply as differentiability unless it is not clear whether I am talking about real or complex-differentiation.

Theorem 2.5

Suppose that f is defined in a neighbourhood of a point z_0 , $B(z_0, \epsilon)$. Then f is differentiable at z_0 if and only if there is a complex valued function E defined on $B(0, \epsilon) \setminus \{0\}$ with $E(h)/h \rightarrow 0$ as $h \rightarrow 0$ and a $d \in \mathbb{C}$ such that

$$f(z_0 + h) = f(z_0) + dh + E(h).$$

In this case we have that $f'(z_0) = d$.

We will use this result frequently as a characterisation of differentiability. It is very useful not only for deciding if a function is differentiable but also for finding its derivative.

Proof. Suppose that f is differentiable at z_0 then let

$$e(h) = \frac{f(z_0 + h) - f(z_0)}{h} - f'(z_0). \quad (2.1)$$

This is defined for all $h \in B(0, \epsilon) \setminus \{0\}$. Furthermore $\lim_{h \rightarrow 0} e(h) = 0$ since f is differentiable at z_0 . Letting

$$E(h) = he(h)$$

and rearranging (2.1) we get

$$f(z_0 + h) = f(z_0) + f'(z_0)h + E(h)$$

and $E(h)/h = e(h) \rightarrow 0$ as $h \rightarrow 0$.

Suppose, conversely, that

$$f(z_0 + h) = f(z_0) + dh + E(h)$$

where $E(h)$ is defined in a neighbourhood $B(0, \epsilon) \setminus \{0\}$ and $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. Then again we can rearrange this so that

$$\frac{f(z_0 + h) - f(z_0)}{h} = d + \frac{E(h)}{h}.$$

It follows that

$$\begin{aligned} f'(z_0) &= \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \\ &= \lim_{h \rightarrow 0} \left(d + \frac{E(h)}{h} \right) \\ &= d. \end{aligned}$$

□

In general the standard rules for differentiation in calculus have equivalent counterparts in the complex plane, for example if $f(z) = z^n$ for some $n \in \mathbb{C}$ then $f'(z) = nz^{n-1}$. The product rule, chain rule and quotient rule hold when we replace real-differentiation with complex-differentiation. The examples above illustrate the importance of the domain as a crucial element of the definition of a holomorphic function and the same is true when we use the rules of differentiation. We need to make sure we are aware that the rules are only true as long as our functions are holomorphic.

Theorem 2.6 – Product Rule

Suppose $f \in H(\Omega_1)$ and $g \in H(\Omega_2)$ then for all $z_0 \in \Omega_1 \cap \Omega_2$

$$\frac{d}{dz}(fg)(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0).$$

Proof. Since both $f'(z_0)$ and $g'(z_0)$ exist we have, by Theorem 2.5, functions $E(h)$ and $F(h)$ such that

$$\begin{aligned} f(z_0 + h) &= f(z_0) + f'(z_0)h + E(h) \\ g(z_0 + h) &= g(z_0) + g'(z_0)h + F(h), \end{aligned}$$

where $\lim_{h \rightarrow 0} E(h)/h = \lim_{h \rightarrow 0} F(h)/h = 0$.

Now

$$\begin{aligned} f(z_0 + h)g(z_0 + h) &= (f(z_0) + f'(z_0)h + E(h))(g(z_0) + g'(z_0)h + F(h)) \\ &= f(z_0)g(z_0) + f(z_0)g'(z_0)h + f(z_0)F(h) \\ &\quad + f'(z_0)g(z_0)h + f'(z_0)g'(z_0)h^2 + f'(z_0)F(h)h \\ &\quad + E(h)g(z_0) + E(h)g'(z_0)h + E(h)F(h) \\ &= f(z_0)g(z_0) + (f(z_0)g'(z_0) + f'(z_0)g(z_0))h \\ &\quad + f(z_0)F(h) + f'(z_0)g'(z_0)h^2 \\ &\quad + f'(z_0)F(h)h + E(h)g(z_0) + E(h)g'(z_0)h + E(h)F(h) \\ &= f(z_0)g(z_0) + (f(z_0)g'(z_0) + f'(z_0)g(z_0))h + G(h) \end{aligned}$$

where

$$G(h) = f(z_0)F(h) + E(h)g(z_0) + h(f'(z_0)F(h) + E(h)g'(z_0)) + f'(z_0)g'(z_0)h^2 + E(h)F(h).$$




Figure 2.1: The Chain Rule

But

$$\frac{G(h)}{h} = f(z_0) \frac{F(h)}{h} + g(z_0) \frac{E(h)}{h} + (f'(z_0)F(h) + E(h)g'(z_0)) + f'(z_0)g'(z_0)h + \frac{E(h)}{h}F(h).$$

Since each term converges to 0 in this expression, we have that $\lim_{h \rightarrow 0} G(h)/h = 0$ and by Theorem 2.5 the result is proved. \square

We will also state here without proving it the chain rule.

Theorem 2.7 – Chain Rule

Suppose that $f: \Omega_1 \rightarrow \Omega_2$ is holomorphic, as is $g: \Omega_3 \rightarrow \Omega_1$. Then $f \circ g \in H(\Omega_3)$ and

$$\frac{d}{dz} f \circ g(z_0) = g'(z_0) f'(g(z_0)), \quad z_0 \in \Omega_3.$$

Example 2.8. The derivative of z^n is nz^{n-1} whenever $n \in \mathbb{N}$.

Let $h(z) = z^n$. To prove this note that when $n = 1$, $h(z) = z$ and so it follows immediately from Theorem 2.5 that $h'(z) = 1$.

Suppose that the result is true for k and let $h(z) = z^{k+1}$ then we write

$$h(z) = z^{k+1} = f(z)g(z)$$

where $f(z) = z^k$ and $g(z) = z$. Then by the inductive step $f'(z) = kz^{k-1}$ and we have that $g'(z) = 1$. So by the Product Rule

$$f'(z) = kz^{k-1} \times z + z^k \times 1 = (k+1)z^k$$

as required.

Finally we have the following result.

Theorem 2.9 – Quotient Rule

Let Ω be a domain. Suppose $f, g \in H(\Omega)$ and $g(z) \neq 0$ for any $z \in \Omega$. Then $F(z) = f(z)/g(z)$ is holomorphic in Ω and for all $z_0 \in \Omega$

$$\frac{F'(z_0)}{F(z_0)} = \frac{f'(z_0)}{f(z_0)} - \frac{g'(z_0)}{g(z_0)}.$$

Example 2.10. The function $F(z) = \frac{z+1}{z-1}$ is holomorphic throughout the domain $B(0, 1)$. Indeed by the quotient rule we may write $f(z) = z+1$ and $g(z) = z-1$ to find that

$$\begin{aligned} \frac{F'(z)}{F(z)} &= \frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \\ &= \frac{1}{z+1} - \frac{1}{z-1}, \\ &= -\frac{2}{(z-1)(z+1)} \end{aligned}$$

so that,

$$\begin{aligned} F'(z) &= -F(z) \frac{2}{(z-1)(z+1)} \\ &= -\frac{z+1}{z-1} \frac{2}{(z-1)(z+1)} \\ &= -\frac{2}{(z-1)^2} \end{aligned}$$

2.2 The Cauchy-Riemann equations

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ and define $\mathbf{f} = J \circ f \circ J^{-1}$ where J was defined in (1.3).

We saw in the last section that f is complex-differentiable at $z_0 = x_0 + iy_0$ if

$$f(z_0 + h) \approx f(z_0) + dh \quad (2.2)$$

for some $d \in \mathbb{C}$ and h close enough to 0 .

Similarly \mathbf{f} is real-differentiable at $\mathbf{x}_0 = (x_0, y_0)^T$ if

$$\mathbf{f}(\mathbf{x}_0 + \mathbf{h}) \approx \mathbf{f}(\mathbf{x}_0) + D\mathbf{h} \quad (2.3)$$

for some 2×2 square matrix D , and \mathbf{h} close enough to $\mathbf{0}$.

If we write $d = a + ib$ in (2.2) then define

$$D = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}.$$

A calculation shows that

$$J(dz) = DJ(z).$$

So if f satisfies (2.2) with derivative $d \in \mathbb{C}$ then \mathbf{f} satisfies (2.3) with derivative D . That is, if f is complex-differentiable then \mathbf{f} is real-differentiable. The converse is not, however, true: if \mathbf{f} is real-differentiable it does not mean that f is complex-differentiable. In this section we will prove the Cauchy-Riemann equations that demonstrate that this relationship holds throughout the domain of definition of a holomorphic function.

Theorem 2.11 – The Cauchy-Riemann equations

Suppose that $f: \Omega \rightarrow \mathbb{C}$ is written as

$$f(x + iy) = u(x, y) + iv(x, y).$$

Then

1. If f is holomorphic then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad (2.4)$$

at each point in Ω .

2. Conversely if u and v are continuously differentiable functions in Ω that satisfy (2.4) then f is holomorphic.

Proof. Since Ω is open we only need to prove the theorem in a neighbourhood of a given point $z_0 = x_0 + iy_0$.

We prove part 1 first. Since $f'(z_0)$ exists we have that

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

where the limit is taken in any direction. In particular if we restrict h to the real numbers, $h \in \mathbb{R}$, then $z_0 + h = (x_0 + h) + iy_0$ and

$$\begin{aligned} f'(z_0) &= \lim_{h \rightarrow 0} \frac{(u(x_0 + h, y_0) + iv(x_0 + h, y_0)) - (u(x_0, y_0) + iv(x_0, y_0))}{h} \\ &= \lim_{h \rightarrow 0} \frac{u(x_0 + h, y_0) - u(x_0, y_0)}{h} + i \lim_{h \rightarrow 0} \frac{v(x_0 + h, y_0) - v(x_0, y_0)}{h} \\ &= \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0). \end{aligned}$$

Similarly we can take the limit along the imaginary axis so that $h = ik$ for $k \in \mathbb{R}$. Then $z_0 + h = x_0 + i(y_0 + k)$ and

$$\begin{aligned} f'(z_0) &= \lim_{k \rightarrow 0} \frac{(u(x_0, y_0 + k) + iv(x_0, y_0 + k)) - (u(x_0, y_0) + iv(x_0, y_0))}{ik} \\ &= \lim_{k \rightarrow 0} \frac{u(x_0, y_0 + k) - u(x_0, y_0)}{ik} + i \lim_{k \rightarrow 0} \frac{v(x_0, y_0 + k) - v(x_0, y_0)}{ik} \\ &= -i \lim_{k \rightarrow 0} \frac{u(x_0, y_0 + k) - u(x_0, y_0)}{k} + \lim_{k \rightarrow 0} \frac{v(x_0, y_0 + k) - v(x_0, y_0)}{k} \end{aligned}$$

Since $1/i = -i$,

$$= -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}.$$

These two limits are the same since $f'(z_0)$ exists and hence we have that

$$\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}.$$

The Cauchy-Riemann equations (2.4) follow by equating real and imaginary parts.

To prove the converse recall that since u and v are continuously differentiable throughout Ω ,

$$\nabla u = \left(\frac{\partial u}{\partial x} \quad \frac{\partial u}{\partial y} \right), \quad \nabla v = \left(\frac{\partial v}{\partial x} \quad \frac{\partial v}{\partial y} \right).$$

Moreover

$$\lim_{\mathbf{h} \rightarrow \mathbf{0}} \frac{|u(\mathbf{x}_0 + \mathbf{h}) - u(\mathbf{x}_0) - \nabla u \mathbf{h}|}{|\mathbf{h}|} = 0, \quad (2.5)$$

and

$$\lim_{\mathbf{h} \rightarrow \mathbf{0}} \frac{|v(\mathbf{x}_0 + \mathbf{h}) - v(\mathbf{x}_0) - \nabla v \mathbf{h}|}{|\mathbf{h}|} = 0, \quad (2.6)$$

where \mathbf{h} and \mathbf{x}_0 are vector quantities in \mathbb{R}^2 .

Let $h \in \mathbb{C}$ and define

$$E(h) = f(z_0 + h) - f(z_0) - dh$$

where

$$d = \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0).$$

By Theorem 2.5 the result will follow if we show that $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. Let us write $h = h_1 + ih_2$. We first take the real part of $E(h)$,

$$\begin{aligned} \operatorname{Re} E(h) &= \operatorname{Re}(f(z_0 + h) - f(z_0) - dh) \\ &= \operatorname{Re} \left(u(x_0 + h_1, y_0 + h_2) + iv(x_0 + h_1, y_0 + h_2) - (u(x_0, y_0) + iv(x_0, y_0)) \right. \\ &\quad \left. - \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right) (h_1 + ih_2) \right) \\ &= u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0) - \operatorname{Re} \left(\left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right) (h_1 + ih_2) \right) \\ &= u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0) - \left(\frac{\partial u}{\partial x} h_1 - \frac{\partial v}{\partial x} h_2 \right) \\ &= u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0) - \frac{\partial u}{\partial x} h_1 + \frac{\partial v}{\partial x} h_2 \\ &= u(x_0 + h_1, y_0 + h_2) - u(x_0, y_0) - \frac{\partial u}{\partial x} h_1 - \frac{\partial u}{\partial y} h_2 \end{aligned}$$

by (2.4).

Now if we let $\mathbf{x}_0 = J(z_0) = (x_0, y_0)^T$ and $\mathbf{h} = J(h) = (h_1, h_2)^T$ then we may rewrite this as

$$\operatorname{Re} E(h) = u(\mathbf{x}_0 + \mathbf{h}) - u(\mathbf{x}_0) - \nabla u \mathbf{h}$$

then

$$\lim_{h \rightarrow 0} \left| \frac{\operatorname{Re} E(h)}{h} \right| = \lim_{|\mathbf{h}| \rightarrow 0} \frac{|u(\mathbf{x}_0 + \mathbf{h}) - u(\mathbf{x}_0) - \nabla u \mathbf{h}|}{|\mathbf{h}|} = 0$$

by (2.5).

If instead we calculate the imaginary part of $E(h)$ above then we see that

$$\operatorname{Im} E(h) = v(\mathbf{x}_0 + \mathbf{h}) - v(\mathbf{x}_0) - \nabla v \mathbf{h}$$

and it follows by (2.6) that

$$\lim_{h \rightarrow 0} \left| \frac{\operatorname{Im} E(h)}{h} \right| = \lim_{|h| \rightarrow 0} \frac{v(\mathbf{x}_0 + \mathbf{h}) - v(\mathbf{x}_0) - \nabla v \mathbf{h}}{|h|} = 0.$$

Since $\operatorname{Re} E(h)/h \rightarrow 0$ and $\operatorname{Im} E(h)/h \rightarrow 0$ as $h \rightarrow 0$, it follows by Proposition 1.4 that $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. So f is differentiable at \mathbf{z}_0 by Theorem 2.5. Since Ω is open this is true at all points. It follows that the function f is holomorphic. \square

A number of consequences of the Cauchy-Riemann equations will demonstrate why they are important, the first being the following.

Corollary 2.12. *Suppose that $f(x + iy) = u(x, y) + iv(x, y)$ is holomorphic in a neighbourhood then*

$$\begin{aligned} f'(z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ f'(z) &= -i \left(\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) \end{aligned}$$

The first description of the derivative in this corollary gives us an easy way to find the derivative of a function since we can think of it as

$$f'(z) = \frac{\partial}{\partial x} f(z) \tag{2.7}$$

where $z = x + iy$. Similarly the second can be written as

$$f'(z) = -i \frac{\partial}{\partial y} f(z). \tag{2.8}$$

Example 2.13. *Find the derivative of z^2 .*

Here we write this as

$$(x + iy)^2 = x^2 + 2ixy - y^2$$

then

$$\begin{aligned} \frac{d}{dz} z^2 &= \frac{\partial}{\partial x} (x^2 + 2ixy - y^2) \\ &= 2x + 2iy = 2(x + iy) \\ &= 2z. \end{aligned}$$

2.3 Holomorphic functions as vector fields

Let f be a holomorphic function and define

$$\mathbf{f} = J \circ f \circ J^{-1}$$

be the equivalent vector field $\mathbf{f}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. With the notation of Theorem 2.11 we may write this function as

$$\begin{aligned} \mathbf{f}: \mathbb{R}^2 &\rightarrow \mathbb{R}^2 \\ (x, y)^T &\mapsto (u(x, y), v(x, y))^T. \end{aligned}$$

The derivative is

$$\nabla \mathbf{f} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix}.$$

However, note that the Cauchy-Riemann equations state that we may write this as

$$\nabla \mathbf{f} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ -\frac{\partial u}{\partial y} & \frac{\partial u}{\partial x} \end{pmatrix}. \quad (2.9)$$

We therefore have

$$\begin{aligned} \det \nabla \mathbf{f} &= \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \\ &= |f'(z)|^2. \end{aligned}$$

We have proved the following.

Corollary 2.14. *Let f and \mathbf{f} be defined as above and $z = x + iy$. If f is holomorphic then*

$$\det \nabla \mathbf{f}(x, y) = |f'(z)|^2.$$

Before moving on we will, on occasion, need the Cauchy-Riemann equations in polar form. To find these require some elementary work using the equations

$$x = r \cos \theta \quad y = r \sin \theta.$$

The chain rule for partial derivatives then gives the following

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial u}{\partial \theta} \quad \frac{\partial u}{\partial \theta} = -\frac{1}{r} \frac{\partial u}{\partial r}. \quad (2.10)$$

2.4 Power series

A power series is an expression of the form

$$\sum_{n=0}^{\infty} a_n (z - a)^n$$

for $a \in \mathbb{C}$ and a sequence of complex numbers $(a_n)_{n=0}^{\infty}$. The complex number a is called the centre of the power series.

Power series are important in complex analysis and we will spend quite some time discussing them in detail. There are three main themes:

1. For which values of z does the series converge?
2. What can be said about the function defined by a power series?
3. Under what conditions can a function be represented as a power series?

The first of these questions has a familiar answer.

The **radius of convergence** of a power series is a number R defined by the relationship

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} |a_n|^{1/n},$$

where R can be 0 or ∞ , when the upper limit on the right-hand-side is ∞ or 0 respectively.

If z satisfies $|z - a| < R$ the series converges absolutely. This follows from the root test.

Example 2.15. *The Taylor series of a function $f(z)$ is the power series*

$$f(z) = \sum_{n=0}^{\infty} a_n (z - a)^n,$$

with

$$a_n = \frac{f^{(n)}(a)}{n!}.$$

A function for which the Taylor series exists has to be very smooth in real analysis, indeed it has to be differentiable infinitely many times in a neighbourhood of a .




Figure 2.2: The graph of $f(x) = 1/(1+x^2)$.

Example 2.16. Consider the power series

$$\sum_{n=0}^{\infty} (-1)^n z^{2n}.$$

The terms of this series are 0 for odd values of n , for even values they alternate between ± 1 . So we have

$$|a_n| = \begin{cases} 1, & n \text{ even;} \\ 0, & n \text{ odd.} \end{cases}$$

Therefore

$$\limsup_{n \rightarrow \infty} |a_n|^{1/n} = 1$$

and the power series converges for $|z| < 1$.

In real analysis it is not always clear why certain power series have finite radii of convergence. The example above highlights this. The power series is in fact the Taylor series

$$f(z) = \sum_{n=0}^{\infty} (-1)^n z^{2n} = \frac{1}{1+z^2}.$$

The real-valued function $1/(1+x^2)$ is defined for all $x \in \mathbb{R}$ and is infinitely differentiable there, so it is not clear why the Taylor series does not converge for $|x| \geq 1$. The graph of $1/(1+x^2)$ is shown in Figure 2.2.

The answer lies in considering this as a complex-valued function:

$$\frac{1}{1+z^2} = \frac{1}{(z-i)(z+i)}.$$




Figure 2.3: The radius of convergence for $f(z) = 1/(1 + z^2)$.

It is clear that this function is not defined at the points $z = \pm i$ and these lie on the boundary of the disk $\{z: |z| < 1\}$, see Figure 2.3. It is for this reason that the power series fails to converge for $|z| \geq 1$.

Theorem 2.17

Let $f(z) = \sum_{n=0}^{\infty} a_n(z - a)^n$ have radius of convergence $R > 0$. Then

$$\sum_{n=1}^{\infty} n a_n (z - a)^{n-1} \quad (2.11)$$

also has radius of convergence R and

$$f'(z) = \sum_{n=1}^{\infty} n a_n (z - a)^{n-1}, \quad |z - a| < R.$$

To prove this requires some manipulation of binomial series. We will use the following estimate of binomial coefficients.

Lemma 2.18. For $n \geq 2$ and $r = 0, 1, \dots, n$,

$$\binom{n}{r+2} \leq n^2 \binom{n-2}{r}.$$

Proof. We have

$$\binom{n}{r+2} = \frac{n!}{(n-r-2)!(r+2)!}, \quad \binom{n-2}{r} = \frac{(n-2)!}{(n-r-2)!r!}.$$

Therefore

$$\begin{aligned} \binom{n}{r+2} / \binom{n-2}{r} &= \frac{n!}{(n-2)!} \frac{r!}{(r+2)!} \\ &= \frac{n(n-1)}{(r+2)(r+1)} \\ &\leq n(n-1) \leq n^2. \end{aligned}$$

□

Next we will show that the radii of convergence are the same for our two series.

Lemma 2.19. *The series $\sum_{n=0}^{\infty} a_n(z-a)^n$ and (2.11) have the same radius of convergence.*

Proof. We first re-index (2.11) as follows:

$$\sum_{n=0}^{\infty} (n+1)a_{n+1}(z-a)^n.$$

Then we need to show that

$$\limsup_{n \rightarrow \infty} ((n+1)|a_{n+1}|)^{1/n} = \limsup_{n \rightarrow \infty} |a_n|^{1/n}.$$

Now $(n+1)^{1/n} \rightarrow 1$ as $n \rightarrow \infty$. Therefore

$$\begin{aligned} \limsup_{n \rightarrow \infty} ((n+1)|a_{n+1}|)^{1/n} &= \limsup_{n \rightarrow \infty} |a_{n+1}|^{1/n} \\ &= \limsup_{n \rightarrow \infty} (|a_{n+1}|^{1/(n+1)})^{\frac{n+1}{n}} \\ &= \limsup_{n \rightarrow \infty} |a_{n+1}|^{1/(n+1)}, \end{aligned}$$

since $(n+1)/n \rightarrow 1$ as $n \rightarrow \infty$,

$$= \limsup_{n \rightarrow \infty} |a_n|^{1/n},$$

as required.

□

Proof of Theorem 2.4. We will show that $g(z) = \sum_{n=1}^{\infty} n a_n (z - a)^{n-1}$ is the derivative of $f(z)$. To simplify notation we assume without loss of generality that $a = 0$, otherwise we can simply consider the functions $g(z + a)$ and $f(z + a)$. Let

$$E(h) = f(z + h) - f(z) - g(z)h$$

then by Theorem 2.5 we need to show that $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. Now

$$\begin{aligned} E(h) &= f(z + h) - f(z) - g(z)h \\ &= \sum_{n=0}^{\infty} a_n (z + h)^n - \sum_{n=0}^{\infty} a_n z^n - h \sum_{n=1}^{\infty} n a_n z^{n-1} \\ &= \sum_{n=0}^{\infty} a_n ((z + h)^n - z^n - n z^{n-1} h) \\ &= \sum_{n=0}^{\infty} a_n d_n(h) \end{aligned}$$

where

$$d_n(h) = (z + h)^n - z^n - n z^{n-1} h.$$

In order to show that $E(h)/h \rightarrow 0$ as $h \rightarrow 0$ we want to estimate the size of $|a_n|$ and the size of $|d_n(h)|$.

Now $d_0(h) = 0$ and $d_1(h) = 0$ for all h . For $n \geq 2$ the binomial theorem tells us

$$(z + h)^n = \sum_{k=0}^n \binom{n}{k} h^k z^{n-k} = z^n + n z^{n-1} h + \sum_{k=2}^n \binom{n}{k} h^k z^{n-k},$$

and therefore

$$d_n(h) = \sum_{k=2}^n \binom{n}{k} h^k z^{n-k}, \quad n \geq 2.$$

Choose h with $|h|$ small enough so that we may find r_1 and r_2 with

$$|z| + |h| < r_1 < r_2 < R.$$

Now $r_2 \in B(0, R)$ and so the power series $f(z)$ converges for $z = r_2$. That is $\sum_{n=0}^{\infty} a_n r_2^n$ converges. It follows that $a_n r_2^n \rightarrow 0$ as $n \rightarrow \infty$, and the sequence $(a_n r_2^n)$ is bounded. Hence we may find a $M > 0$ such that $|a_n| r_2^n \leq M$, or

$$|a_n| \leq \frac{M}{r_2^n} \quad \forall n.$$

Now, using the estimate in Lemma 2.18,

$$\begin{aligned} |d_n(h)| &= \left| \sum_{k=2}^n \binom{n}{k} h^k z^{n-k} \right| \leq \sum_{k=2}^n \binom{n}{k} |h|^k |z|^{n-k} \\ &= \sum_{i=0}^{n-2} \binom{n}{i+2} |h|^{i+2} |z|^{n-2-i}, \end{aligned}$$

where we are changing the indexing in the sum, $k = i + 2$,

$$\begin{aligned} &= |h|^2 \sum_{i=0}^{n-2} \binom{n}{i+2} |h|^i |z|^{n-2-i} \\ &\leq |h|^2 \sum_{i=0}^{n-2} n^2 \binom{n-2}{i} |h|^i |z|^{n-2-i}, \end{aligned}$$

by Lemma 2.18,

$$= |h|^2 n^2 (|h| + |z|)^{n-2}.$$

Since $|z| + |h| < r_1$ we can estimate this by

$$|d_n(h)| \leq |h|^2 n^2 r_1^{n-2}$$

and so using this and our estimate for $|a_n|$ we have that

$$\begin{aligned} |E(h)| &= \left| \sum_{n=0}^{\infty} a_n d_n(h) \right| \leq \sum_{n=0}^{\infty} |a_n| |d_n(h)| \\ &\leq \sum_{n=2}^{\infty} \frac{M}{r_2^n} n^2 |h|^2 r_1^{n-2} = M |h|^2 \sum_{n=2}^{\infty} n^2 \frac{r_1^{n-2}}{r_2^n}. \end{aligned}$$

The sum on the right hand side converges by the ratio test. Therefore we have shown that $|E(h)| \leq C |h|^2$ for some constant $C > 0$. Hence $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. By Theorem 2.5 we have shown that

$$f'(z) = g(z)$$

as required. □

This result highlights the importance of power series as a way of defining functions. It also allows us to find the derivative of a function.

Example 2.20. The geometric series tells us that, for $|z| < 1$,

$$\frac{1}{1-z} = 1 + z + z^2 + \dots.$$

Differentiating both sides tells us that, again for all $|z| < 1$,

$$\frac{1}{(1-z)^2} = 1 + 2z + 3z^2 + \dots.$$

Definition 2.21 – Analytic functions

A function on a domain Ω that can be written as a power series

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n$$

with positive radius of convergence, $R > 0$, at each point $a \in \Omega$ is called an **analytic function**. The set of analytic functions on a domain Ω is written $A(\Omega)$.

By Theorem 2.4 an analytic function is differentiable at each point in Ω and so we have $A(\Omega) \subset H(\Omega)$. We will show, eventually,

$$A(\Omega) = H(\Omega)$$

but in order to do that we will need to develop complex integration. We will, in preparation, note a couple of consequences of Theorem 2.4.

Corollary 2.22. Suppose that $f \in A(\Omega)$ then $f^{(k)} \in A(\Omega)$ for each $k \geq 0$.

This follows directly from the theorem since, if $\sum_{n=0}^{\infty} a_n(z-a)^n$ is analytic then so is its derivative $\sum_{n=0}^{\infty} n a_n(z-a)^{n-1}$, therefore so is its derivative and so on. Furthermore the radius of convergence is the same for each derivative. In fact it can be seen from the theorem that the k th derivative is

$$f^{(k)}(z) = \sum_{n=0}^{\infty} \frac{(n+k)!}{n!} a_{n+k} (z-a)^n. \quad (2.12)$$

Letting $z = a$ in this equation gives the Taylor coefficients

$$f^{(k)}(a) = k! a_k.$$

So if $f \in A(\Omega)$ then at each point $a \in \Omega$ $f(z)$ is equal to its Taylor series centred at a in some disk $B(a, r)$. We state this as the following corollary that will be useful later on.

Corollary 2.23. Suppose that $f \in A(\Omega)$ and $a \in \Omega$ then there is a $r > 0$ such that

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (z - a)^n$$

for all $z \in B(a, r)$.

Things to know

- The definition of complex differentiation;
- The product, chain and quotient rules;
- The definition of a holomorphic function, and simple examples;
- The statement of Theorem 2.5;
- The Cauchy-Riemann equations;
- The definition of a power series and how to calculate its radius of convergence;
- Understand the statement of Theorem 2.4;
- The definition of an analytic function;

Further reading

First it should be noted that as we said holomorphic functions and analytic functions are the same thing in complex analysis (although we've not proved that yet). It is only modern textbooks that distinguish the two of these concepts. Ahlfors, [1], only uses 'Analytic', for example, whereas Remmert, [7], uses 'Holomorphic'. The reason we distinguish these nowadays is (I believe) because there is a similar concept in real analysis and these are not the same as real-differentiable functions.

All of the material in this chapter can be found in Chapter 1 of [3]. Remmert, [7], deals with the topic in a lot of detail but might be worth reading to get to grips with these ideas more. Ahlfors, [1, Chapter 2], also includes a standard cover of these ideas.

The notion of a conformal mapping is an entire subject in its own right and lots of textbooks are devoted to this material. However you should only embark on these once you have finished this course as the concepts are quite advanced.

Problems

1. By calculating $E(h)$, use Theorem 2.5 to show directly that the derivative of $f(z) = 1/z$ is

$$f'(z) = -\frac{1}{z^2}, \quad z \neq 0.$$

2. Use Theorem 2.5 to find the derivatives of the following functions

- (a) $f(z) = 3z^2 - 2z + 4$;
- (b) $f(z) = (1 - 4z^2)^3$;
- (c) $f(z) = \frac{z-1}{2z+1}$, $z \neq -1/2$.

3. Use the Cauchy-Riemann equations to decide whether or not the following functions are holomorphic in \mathbb{C} .

- (a) $(x^2 + y^2) + 2ixy$
- (b) $(x^2 - y^2) - 2ixy$
- (c) $(x^2 + 2x - y^2) + 2iy(1 + x)$

4. Use the Cauchy-Riemann equations to show that the function $f(z) = \operatorname{Re} z$ is not holomorphic at any point.

5. Use the Cauchy-Riemann equations to show that the function $f(z) = \operatorname{Im} z$ is not holomorphic at any point.

6. Use the Cauchy-Riemann equations to show that the function $f(z) = |z|$ is not holomorphic at any point.

7. Use the Cauchy-Riemann equations to show that if $\operatorname{Re} f(z) = 0$ in a domain then $f(z)$ is constant throughout the domain.

8. Find the radius of convergence for the following power series:

- (a) $\sum_{n=1}^{\infty} \frac{n^2}{a^n} z^n$ for $a \in \mathbb{C}$;
- (b) $\sum_{n=1}^{\infty} \frac{i^n n^2}{a^n} z^n$ for $a \in \mathbb{C}$;

(c) $\sum_{n=1}^{\infty} 2^n z^{2^n}$ for $a \in \mathbb{C}$;

(d) $\sum_{n=1}^{\infty} \frac{i^n - 1}{n} z^n$

9. Suppose that the sequence $(|a_n|)$ is bounded. Show that $\sum_{n=0}^{\infty} a_n z^n$ has radius of convergence **at least 1**.

10. Let $0 < R < \infty$. Give an example of a power series whose radius of convergence is R .

* 11. Use Theorem 2.5 to show that if $P(z) = a_0 + a_1 z + \cdots + a_n z^n$ is a polynomial then it has derivative at all points equal to

$$P'(z) = a_1 + 2a_2 z^{n-1} + \cdots + na_n z^n.$$

* 12. Suppose $f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n$ is analytic in $B(a, r)$ for some $r > 0$. Show that there is a $K > 0$ such that $|a_n| \leq K^n$ for all n .

* 13. Suppose $f(z) = \sum_{n=0}^{\infty} a_n z^n$ is a power series defined in $B(0, 1)$. Show that if $|a_n| \leq 1$ for all n then

$$|f(z)| \leq \frac{1}{1 - |z|}, \quad z \in B(0, 1).$$

Chapter 3

Special Functions

“Mathematics, rightly viewed, possesses not only truth, but supreme beauty – a beauty cold and austere, like that of sculpture, without appeal to any part of our weaker nature, without the gorgeous trappings of painting or music, yet sublimely pure, and capable of a stern perfection such as only the greatest art can show.”

Bertrand Russell

3.1 Introduction

Functions that you are familiar with from real analysis take on a different, more geometric, form in complex analysis. Often they display some interesting geometric behaviour in addition to their analytic definitions. In this chapter we will collect together definitions of some important functions. We will both define them and look at how they behave geometrically.

3.2 The exponential function

We begin with the exponential function. In real analysis this is the function $\exp x = e^x$ where $e = 2.71828 \dots$ is Euler’s number. This only really makes sense when $x \in \mathbb{Q}$ since $\exp p/q$ can be defined as the q th root of e^p . Or similarly the smallest positive real

solution to the equation

$$y^q - e^p = 0.$$

In the last chapter we discussed power series. This is a more flexible and robust way of defining the exponential function. We define

$$\begin{aligned} \exp : \mathbb{C} &\rightarrow \mathbb{C} \\ z &\mapsto \sum_{n=0}^{\infty} \frac{z^n}{n!}. \end{aligned}$$

This definition makes sense for all $z \in \mathbb{C}$ since the radius of convergence R , satisfies

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} |n!|^{-1/n} = 0$$

and so $R = \infty$. By Theorem 2.4 \exp is differentiable everywhere and hence is holomorphic in \mathbb{C} . This is, therefore, an entire function. Note that if x is real then $\exp x$ is our usual definition of the real-valued exponential function.

Proposition 3.1

The exponential function satisfies the following:

1. $\exp(z + w) = \exp z \exp w$;
2. $\frac{1}{\exp z} = \exp(-z)$;
3. $\exp(z + 2\pi i) = \exp z$;
4. If $\exp z = \exp w$ then $z = w + 2\pi i k$ for some $k \in \mathbb{Z}$;
5. $\frac{d}{dz} \exp z = \exp z$.

Lemma 3.2 (The Cauchy product). *Suppose that $\sum_{i=0}^{\infty} a_i$ and $\sum_{j=0}^{\infty} b_j$ are two absolutely convergent complex series. Then*

$$\left(\sum_{i=0}^{\infty} a_i \right) \left(\sum_{j=0}^{\infty} b_j \right) = \sum_{k=0}^{\infty} c_k$$

where

$$c_k = \sum_{l=0}^k a_l b_{k-l}$$

Proof of Theorem 3.2. 1. By the Binomial theorem

$$\frac{1}{n!}(z+w)^n = \frac{1}{n!} \sum_{m=0}^n \binom{n}{m} z^m w^{n-m} = \sum_{k=0}^n \frac{z^m w^{n-m}}{m!(n-m)!}.$$

So

$$\exp(z+w) = \sum_{n=0}^{\infty} \sum_{m=0}^n \frac{z^m w^{n-m}}{m!(n-m)!}. \quad (3.1)$$

On the other hand, using the Cauchy product and the fact that the exponential series is entire and hence converges absolutely for all values,

$$\begin{aligned} \exp z \exp w &= \left(\sum_{i=0}^{\infty} \frac{z^i}{i!} \right) \left(\sum_{j=0}^{\infty} \frac{w^j}{j!} \right) \\ &= \sum_{k=0}^{\infty} \sum_{l=0}^k \frac{z^l}{l!} \frac{w^{k-l}}{(k-l)!}. \end{aligned}$$

Comparing this series with (3.1) proves that

$$\exp(z+w) = \exp z \exp w$$

as required.

2. This follows from part 1 with $w = -z$, since $\exp 0 = 1$.

3. By part 1 we have that

$$\exp(z + 2\pi i) = \exp z \exp(2\pi i).$$

However by Euler's formula, (1.1), $\exp 2\pi i = \cos 2\pi + i \sin 2\pi = 1$.

4. By parts 1 and 2 if $\exp z = \exp w$ then $\exp(z-w) = 1$. So it is enough to show that if $\exp z = 1$ then $z = 2\pi i k$ for some $k \in \mathbb{Z}$.

Suppose $z = a + ib$ satisfies $\exp z = 1$. Then

$$1 = \exp z = e^a e^{ib}.$$

Taking absolute values of both sides gives $1 = e^a$ and so we may deduce $a = 0$. Therefore we have

$$1 = e^{ib} = \cos b + i \sin b.$$

Taking real parts of both sides gives $1 = \cos b$ and so $b = 2\pi i k$ for some $k \in \mathbb{Z}$.

5. Finally given Theorem 2.4 we have that

$$\begin{aligned}\frac{d}{dz} \exp z &= \frac{d}{dz} \sum_{n=0}^{\infty} \frac{z^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{nz^{n-1}}{n!} \\ &= \sum_{n=1}^{\infty} \frac{z^{n-1}}{(n-1)!} \\ &= \exp z.\end{aligned}$$

□

The exponential function is relatively simple to understand. If $z = x + iy$ then by part 1 of Proposition 3.2

$$\exp z = e^x e^{iy}.$$

Thus if we consider a line with constant real part x_0 , then this is mapped onto a circle of radius e^{x_0} . Similarly the line with constant imaginary part y_0 is mapped onto the ray with the argument, y_0 . In Figure 3.1 the complex planes on the right show how the exponential function maps a rectangle and an infinite strip. The rectangle at the top is of the form

$$\{z = x + iy : 1 < x < 2, -2\pi/3 < y < 2\pi/3\}.$$

It is mapped by the exponential function to the set on the right:

$$\{z = re^{i\theta} : e < r < e^2, -2\pi/3 < \theta < 2\pi/3\}.$$

The diagram at the bottom of Figure 3.1 shows the strip $\{z = x + iy : 1 < x < 2\}$. This domain is mapped onto the annulus

$$\{z \in \mathbb{C} : e < |z| < e^2\}.$$

The rectangle at the top of the diagram in Figure 3.1 is mapped **bijectively** onto the domain on the right. However in the bottom diagram the infinite strip is not bijectively mapped onto the annulus on the right.

3.3 The logarithm

The natural way to define the logarithm is to define it as the inverse function of the exponential function, **exp**. In order to do that though we need **exp** to be bijective and




Figure 3.1: The mapping of the exponential function




Figure 3.2: The strip S in Theorem 3.3

this is not true in general. From Theorem 3.2 part 3, for any $k \in \mathbb{Z}$

$$\exp z = \exp(z + 2k\pi i).$$

Therefore \exp is not injective. This is similar to the argument and we treat it in the same way: we restrict the possible values to numbers $z \in \mathbb{C}$ with $\operatorname{Im} z \in [0, 2\pi)$. Given any $w \in \mathbb{C}$ we can choose a suitable k and a number $z \in \mathbb{C}$ with $\operatorname{Im} z \in [0, 2\pi)$ so that

$$\exp z = \exp w.$$

This is used to prove the following, see Figure 3.2.

Theorem 3.3

Let $S = \{z \in \mathbb{C} : 0 \leq \operatorname{Im} z < 2\pi\}$ then \exp is a bijective function from S to $\mathbb{C} \setminus \{0\}$.

Proof. Suppose that $\exp z = \exp w$ for $z, w \in S$. Then $z = w + 2k\pi i$ for some $k \in \mathbb{Z}$ by part 4 of Proposition 3.2. Since $z, w \in S$ we have that

$$0 \leq \operatorname{Im} z < 2\pi, \quad 0 \leq \operatorname{Im} w < 2\pi.$$

So

$$-2\pi < \operatorname{Im}(z - w) < 2\pi.$$

It follows that if $\operatorname{Im}(z - w) = 2k\pi$ then $k = 0$ since this is the only possibility in this range. Therefore $z = w$ and \exp is injective.

To prove that \exp is surjective take any $w \in \mathbb{C} \setminus \{0\}$. Let $r = |w|$ and $\theta = \arg w$ be chosen so that $\theta \in [0, 2\pi)$. Letting $z = \log |w| + i\theta$ we have that

$$\begin{aligned} \exp z &= \exp(\log |w| + i\theta) \\ &= \exp(\log |w|) \exp(i\theta) \\ &= |w| \exp(i\theta) = w \end{aligned}$$

and so \exp is surjective. □

The inverse of \exp defined in the theorem above is called the **principle logarithm**:

$$\text{Log } w = \log |w| + i \text{Arg } w \quad (3.2)$$

where Arg represents the principle argument of w , i.e. the argument contained in $[0, 2\pi)$.

Example 3.4. *The principle log of $1 + i$.*

Here $|1 + i| = \sqrt{2}$ and the $\text{Arg}(1 + i) = \pi/4$ so

$$\text{Log}(1 + i) = \frac{1}{2} \log 2 + i \frac{\pi}{4}.$$

An important point to note here is that whereas $\log x$ is not defined in real analysis when $x < 0$ it is possible to define it in complex analysis.

Example 3.5. *The principle log of -2 .*

Here we can write $-2 = 2 \exp(i\pi)$. Then

$$\text{Log}(-2) = \log 2 + i\pi.$$

Some books consider the principle argument to consist of arguments in the interval $[-\pi, \pi)$. There is no substantive difference in doing this. In fact our choice of strip S in the theorem and hence the choice of interval is arbitrary. The proof can be modified quite easily to show the following.

Corollary 3.6. *Suppose that $\alpha \in \mathbb{R}$. Let $S_\alpha = \{z \in \mathbb{C} : \alpha \leq \text{Im } z < \alpha + 2\pi\}$. Then \exp is a bijective mapping from S_α to $\mathbb{C} \setminus \{0\}$.*

We define the inverse of $\exp : S_\alpha \rightarrow \mathbb{C} \setminus \{0\}$ as \log_α , the α -logarithm. The function \log_α is called a **branch of the logarithm**.

The logarithm is not holomorphic everywhere, there is a problem when the argument changes from α to $\alpha + 2\pi$. At this point it is no longer continuous and therefore cannot be differentiable. It is holomorphic everywhere else though.

Theorem 3.7

The α -logarithm is holomorphic on $\mathbb{C} \setminus \{re^{i\alpha} : r \geq 0\}$. Its derivative is $1/z$ there.

Proof. We will simply verify the polar form of the Cauchy-Riemann equations, (2.10). Since

$$\mathbb{C} \setminus \{re^{i\alpha} : r \geq 0\}$$

is open this will suffice by Theorem 2.11.

To prove this we check the Cauchy-Riemann equations in polar form, (2.10). Here $u(r, \theta) = \log r$ and $v(r, \theta) = \theta$, where $\theta \in [\alpha, \alpha + 2\pi)$. It follows that

$$\begin{aligned} \frac{\partial u}{\partial r} &= \frac{1}{r}, & \frac{\partial u}{\partial \theta} &= 0, \\ \frac{\partial v}{\partial r} &= 0, & \frac{\partial v}{\partial \theta} &= 1. \end{aligned}$$

Therefore

$$\frac{\partial u}{\partial r} = \frac{1}{r} = \frac{1}{r} \frac{\partial v}{\partial \theta},$$

and

$$\frac{\partial u}{\partial \theta} = 0 = -\frac{1}{r} \frac{\partial v}{\partial r}.$$

Therefore these satisfy the polar form of the Cauchy-Riemann equations, (2.10), and so by Theorem 2.11 \log_α is holomorphic. To find the derivative simply note that

$$\exp(\log_\alpha z) = z$$

for any α , we can then use the chain rule and inverse function theorem. The left-hand-side becomes

$$\exp(\log_\alpha z) \frac{d}{dz}(\log_\alpha z) = z \frac{d}{dz} \log_\alpha z$$

and the right-hand-side becomes 1. □

Although in general formulas like

$$\log_\alpha zw = \log_\alpha z + \log_\alpha w$$

are not true it is possible to come up relationships of this form as long as we understand that the right-hand side may need to add or subtract $2\pi i$. These follow from Proposition 3.2.

Since the logarithm is not defined everywhere in a consistent manner, we will consider the mapping properties of the α -logarithm.

Given a ϕ_1 and ϕ_2 with $0 \leq \phi_2 - \phi_1 \leq 2\pi$ we can find $\alpha \notin (\phi_1, \phi_2)$ such that \log_α maps the sector $S(\phi_1, \phi_2)$ bijectively onto the infinite strip

$$\{z \in \mathbb{C} : \phi_1 < \operatorname{Im} z < \phi_2\}.$$

As a particularly useful example $S(0, \pi)$ is the upper half-plane. This can be mapped by the principle logarithm (where $\alpha = 0$) onto the strip

$$\{z \in \mathbb{C} : 0 < \operatorname{Im} z < \pi\}.$$




Figure 3.3: The mapping of the logarithm function

We end this section with the following result that will be used elsewhere. If f is holomorphic it's not necessarily true that $\log f$ makes sense at all. For $\log f$ to exist we can't have $f(z) = 0$ for any z for example. We state the following result without proof. We can't prove it here because we haven't done enough yet (we need to prove Cauchy's theorem). The proof can be found in [3, Theorem 22.71] but don't look at it unless this is the second time you're reading these notes.

Proposition 3.8

Let $f \in H(B(a, r))$ for some $a \in \mathbb{C}$ and $r > 0$. Suppose $f(z) \neq 0$ for any $z \in B(a, r)$. Then there is a branch of the logarithm such that $\log_\alpha f(z)$ is holomorphic in $B(a, r)$ and has derivative

$$\frac{f'(z)}{f(z)}.$$

3.4 Complex powers

Suppose $\alpha \in \mathbb{C}$. We define

$$z^\alpha = \exp(\alpha \log z)$$

where \log is any branch of the logarithm.

If α is an integer then this has its usual meaning (multiplying z by itself α times). The function $z \mapsto z^\alpha$, in this case, is said to be **single-valued** since there is only one possible value.

If $\alpha = p/q$ for $p, q \in \mathbb{Z}$ ($q \neq 0$) then $z^{p/q}$ has q distinct values and the function $z \mapsto z^\alpha$ is said to be q -valued or just multiple-valued. This should be familiar from the real-valued function $x \mapsto \sqrt{x}$. You have learned that \sqrt{x} means the positive square root, however the square root is in reality either positive or negative. When x is real we can simplify this by always using the positive square root. This is no longer possible in complex analysis since there is no way of preferring one root over another.

This may seem like a problem but is not in reality. Riemann considered this in great depth and realised that it could be overcome if instead of treating the roots as different numbers, treating them as the **same** number. So that the $z \mapsto z^\alpha$ maps one value to one of these abstracted values. This led to the notion of a **Riemann surface** which is a generalisation of the complex plane. We will not go through this but it would make a good topic for a project.

For other values of α the function is infinite-valued. They are not particularly interesting though — they're just annoying and difficult.

The functions z^α where $\alpha \in \mathbb{R}$ and $\alpha > 0$ are particularly interesting geometrically.

Example 3.9. Let $f(z) = z^2$.

If we consider the grid shown on the top-left of Figure 3.4. This is the square of side length $3/2$. The function $f(z)$ takes the grid in the first quadrant and spins it around into the first and second quadrant. This is because if you think of it in polar form:

$$re^{i\theta} \mapsto r^2 e^{2i\theta}$$

then you can see that the argument is being doubled under this mapping whilst the absolute value is being squared. A sector, for example, like the one at the bottom of Figure 3.4, is mapped onto a sector twice the angle.

Suppose that $\alpha > 0$ is real. Then $f(z) = z^\alpha$ is not in general single-valued in \mathbb{C} . A consequence of this is that we cannot define it easily. We can define it on a sector




Figure 3.4: The mapping $f(z) = z^2$ on a grid and a sector.




Figure 3.5: The mapping $f(z) = z^\alpha$ where $\alpha > 0$.

of opening less than $2\pi/\alpha$ at 0 though and in this domain it is holomorphic. In [1] Ahlfors describes this nicely on page 44 where he introduces the sectors $S(\phi_1, \phi_2) = \{z \in \mathbb{C} : \phi_1 < \arg z < \phi_2\}$. The mapping $f(z) = z^\alpha$ is injective as long as $0 \leq \phi_2 - \phi_1 \leq 2\pi/\alpha$. Then f maps this sector onto $S(\alpha\phi_1, \alpha\phi_2)$.

One thing to note here is that if $S(0, \phi)$ is a sector then the mapping

$$z \mapsto z^{\pi/\phi}$$

maps this sector onto $S(0, \pi)$ and this is the upper half-plane $\{z \in \mathbb{C} : \operatorname{Im} z > 0\}$. This can be useful for ironing out corners in domains.

We end this section with the following consequence of Proposition 3.3.

Corollary 3.10. *Suppose Ω is a simply connected domain. Let $f \in H(\Omega)$. If $f(z) \neq 0$ for any $z \in \Omega$ then for $p > 0$, $f(z)^p$ defines a holomorphic function in Ω .*

3.5 Mappings of given domains

The mappings described above can be combined or modified to include mappings of many domains onto each other. For example the function

$$z \mapsto \text{Log } z^2$$

maps the first quadrant in \mathbb{C} onto the strip $\{z \in \mathbb{C} : 0 < \text{Im } z < \pi\}$ bijectively. The question of whether or not there is a bijective, holomorphic mapping from one domain onto another is a complicated one and the solution is remarkable and unique to complex analysis. It is the content of probably the most important result in complex analysis.

Theorem 3.11 – The Riemann mapping theorem

Suppose that Ω is a simply connected domain whose boundary consists of more than one point. Then there is a bijective, holomorphic mapping from Ω to the disk $B(0, 1)$.

Of course one can combine functions to find mappings from any such domain onto any other.

3.6 Trigonometric functions

Before we end this chapter we will define trigonometric functions. The reason this is not done earlier is that they don't display any useful geometric mapping properties. However they are very useful in understanding the exponential function a little more.

$$\cos z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}, \quad \sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}.$$

These functions are also entire functions and therefore we can rearrange terms without hesitation to deduce the identities

$$\cos z = \frac{\exp(iz) + \exp(-iz)}{2}, \quad \text{and} \quad \sin z = \frac{\exp(iz) - \exp(-iz)}{2i}.$$

Proposition 3.12

The following identities hold

1. $\exp iz = \cos z + i \sin z$ for $z \in \mathbb{C}$.
2. $\cos^2 z + \sin^2 z = 1$
3. $\frac{d}{dz} \cos z = -\sin z$, and $\frac{d}{dz} \sin z = \cos z$

Proof. The proof is left as an exercise. □

Things to know

- The definitions of the standard functions as power series;
- The logarithm and branches of logarithms.

Further reading

If you want to learn more about the geometric properties of mappings then you should go straight to [1, Chapter 3]. Most of the material in this chapter is inspired by Ahlfors' exposition. You can find more about the definitions of these special functions in [3, §20.4] but not much in the way of the geometric intuition. In [7, Chapter 2] the concepts are extended further and are worth reading if you feel so inspired.

Problems

1. Let $f(z) = az + b$ be a linear function. Describe geometrically what this mapping does for different a and b (using terms like 'rotate' and 'shift' and 'scale').
2. Showed that if $z \in \mathbb{C}$ satisfies $|z| < 1$ then

$$\operatorname{Re} \frac{1+z}{1-z} > 0.$$

What does this tell you about the mapping properties of $\frac{1+z}{1-z}$? What does this tell you about the mapping properties of $i\frac{1+z}{1-z}$? How about $\operatorname{Log} \frac{1+z}{1-z}$?

3. Prove the following:
 - (a) $\exp iz = \cos z + i \sin z$ for $z \in \mathbb{C}$;
 - (b) $\cos^2 z + \sin^2 z = 1$;
 - (c) $\frac{d}{dz} \cos z = -\sin z$, and $\frac{d}{dz} \sin z = \cos z$.
4. By writing $z = x + iy$ and using Corollary 2.12 show that

$$\frac{d}{dz} \exp z = \exp z.$$

5. Show that

$$\exp iz = \cos z + i \sin z$$

6. The **hyperbolic trigonometric functions**, \sinh and \cosh are defined, for complex z , as follows

$$\sinh z = \frac{1}{2}(\exp z - \exp -z), \quad \cosh z = \frac{1}{2}(\exp z + \exp -z).$$

- (a) Find the power series representations for both these functions
- (b) Find the derivatives of these functions and show that

$$\frac{d}{dz} \sinh z = \cosh z, \quad \frac{d}{dz} \cosh z = \sinh z.$$

- (c) Show that

$$\sinh z = -i \sin iz, \quad \cosh z = \cos iz.$$

7. Show that $\text{Log}(zw) = \text{Log } z + \text{Log } w$ whenever $\text{Im } z > 0$ and $\text{Im } w > 0$. Give an example of points $z, w \in \mathbb{C}$ such that $\text{Log}(zw) \neq \text{Log } z + \text{Log } w$.
8. Describe a set on which we can define $\log(z - i)$
9. Suppose that $p(z)$ is a polynomial of degree n . Explain why p is analytic in \mathbb{C} .
- * 10. Show that the exponential function never takes the value 0 , i.e. there is no z such that $\exp z = 0$.
- * 11. Suppose that $p(z) = \prod_{i=1}^n (z - \alpha_i)$ is a polynomial. Prove that

$$\frac{p'(z)}{p(z)} = \sum_{i=1}^n \frac{1}{z - \alpha_i}.$$

Chapter 4

Complex Integration

“Mathematics is not a careful march down a well-cleared highway, but a journey into a strange wilderness, where the explorers often get lost. Rigour should be a signal to the historian that the maps have been made, and the real explorers have gone elsewhere.”

W. S. Anglin, *Mathematics and History*, Mathematical Intelligencer v. 4, no. 4

4.1 Integration of complex functions

We begin with the familiar notion of integration. Suppose that $f(t)$ is a complex-valued function on an interval (a, b) . We write this as

$$f(t) = u(t) + iv(t)$$

then we can define the integral of f (assuming it exists) as

$$\int_a^b f(t)dt = \int_a^b u(t)dt + i \int_a^b v(t)dt.$$

This integral inherits a number of properties from the real-integral counterparts. We will collect two here for use elsewhere.

Proposition 4.1

Let $f(t) = u(t) + iv(t)$ be a complex-valued function on an interval (a, b) such that both u and v are integrable.

1. For $c \in \mathbb{C}$,

$$\int_a^b c f(t) dt = c \int_a^b f(t) dt;$$

2. The triangle inequality holds,

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt.$$

Proof. The first property follows on writing $c = d + ie$ and expanding the integral. To prove the second property assume that the integral is non-zero. Let

$$\theta = \arg \int_a^b f(t) dt$$

then, with $c = e^{-i\theta}$, we have that

$$\begin{aligned} \left| \int_a^b f(t) dt \right| &= \operatorname{Re} \left(e^{-i\theta} \int_a^b f(t) dt \right) \\ &= \int_a^b \operatorname{Re} (e^{-i\theta} f(t)) dt \\ &\leq \int_a^b |f(t)| dt \end{aligned}$$

Since $\operatorname{Re}(e^{-i\theta} f(t)) \leq |e^{-i\theta} f(t)| = |f(t)|$. □

4.2 Integration along curves

Recall the following definitions from calculus.




Figure 4.1: The curve $C = \{(\sin 2t, \cos t)^T : t \in [0, \pi]\}$

Definition 4.2 – Curves and paths

- A **path** is a continuous function $p : [a, b] \rightarrow \mathbb{C}$;
- The **trace** of a path is the set $\text{tr } p = \{z \in \mathbb{C} : z = p(t), t \in [a, b]\}$;
- A set $C \subseteq \mathbb{C}$ is a **curve** if it is the trace of some path.

We will always assume that a path p is piecewise continuously differentiable. If $\text{tr } p = C$ then we say C is parametrised by p .

For example the curve C in \mathbb{R}^2 parametrised by

$$p(t) = (\sin 2t, \cos t)^T \quad t \in [0, \pi]$$

is drawn in Figure 4.1. We normally include an arrow on a curve to represent its orientation. In this example $t = 0$ is mapped to the top of the contour $(0, 1)^T$, as t increases the parametrisation represents points moving along the curve heading towards the point at the bottom, $(0, -1)^T$, which corresponds to $t = \pi$.

In this section we will develop, in the same way that you did in real analysis, the technique of finding the integral of a complex-valued function along a curve. In the complex plane we have the advantage of being able to define a curve as a function of complex numbers. For example the curve C of Figure 4.1 could just as easily have been defined as

$$p(t) = \sin 2t + i \cos t, \quad t \in [0, \pi].$$




Figure 4.2: The curve $C = \{2 \cos t + i \sin t : t \in [0, 2\pi]\}$

Example 4.3. The curve C parametrised by

$$p(t) = 2 \cos t + i \sin t, \quad t \in [0, 2\pi]$$

is an ellipse with major axis 2 and minor axis 1. See Figure 4.2.

For a piecewise continuously differentiable curve C of finite length we can define the integral of a continuous function f along C as the limit of Riemann sums in the following way. Let P be a partition of C , i.e. suppose that

$$P = \{z_0, z_1, \dots, z_n\}$$

where z_0 and z_n are the endpoints of C . Then we let

$$S(f, P) = \sum_{k=0}^{n-1} f(z_k)(z_{k+1} - z_k).$$

Let $I \in \mathbb{C}$. If for each $\epsilon > 0$ there is a partition P such that

$$|S(f, P) - I| < \epsilon$$

then we say I is the integral along C of f and write

$$\int_C f(z) dz = I.$$

In general we don't need this limiting argument, since we can define the integral using a parametrisation $p(t)$ as follows.

Definition 4.4 – Integral along a curve

Let $p(t)$, $a \leq t \leq b$, be a path. Suppose that $p'(t)$ exists almost everywhere and f is a complex-valued function such that $f(p(t))$ is integrable. Then

$$\int_C f(z)dz = \int_a^b f(p(t))p'(t)dt.$$

Example 4.5. Suppose that $f(z) = 2z$ and C is parametrised by $p(t) = t + t^2i$ for $-1 \leq t \leq 1$. Find

$$\int_C f(z)dz.$$

Here $f(p(t)) = 2(t + t^2i) = 2t + 2t^2i$, and $p'(t) = 1 + 2ti$. Therefore

$$\begin{aligned}\int_C f(z)dz &= \int_{-1}^1 f(p(t))p'(t)dt = \int_{-1}^1 (2t + 2t^2i)(1 + 2ti)dt \\ &= \int_{-1}^1 -4t^3 + 2t + 6it^2 dt \\ &= \int_{-1}^1 -4t^3 + 2t dt + i \int_{-1}^1 6t^2 dt \\ &= 4i.\end{aligned}$$

When you learned about integrals along curves in calculus a number of important properties were proved that have direct analogues in complex integration. Firstly if $C = C_1 \cup C_2$ then

$$\int_C f(z)dz = \int_{C_1} f(z)dz + \int_{C_2} f(z)dz.$$

For any curve C defined by $p(t)$, $a \leq t \leq b$, we may define the curve $-C$ as the same curve but in the opposite direction. So $-C$ is parametrised by

$$q(t) = p(-t) \quad -b \leq t \leq -a.$$

Then

$$\int_{-C} f(z)dz = - \int_C f(z)dz.$$

We won't prove these. Instead we prove the following that relates the complex integral along a curve to the real integral along a curve that you learned about in calculus.

Proposition 4.6

Suppose that $f: \mathbb{C} \rightarrow \mathbb{C}$. Write

$$f(x + iy) = u(x, y) + iv(x, y).$$

Let $\mathbf{a}, \mathbf{b}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be vector fields defined by

$$\begin{aligned}\mathbf{a}(x, y) &= (u(x, y) \quad -v(x, y))^T; \\ \mathbf{b}(x, y) &= (v(x, y) \quad u(x, y))^T,\end{aligned}$$

and suppose that C is a curve. Then

$$\int_C f(z)dz = \int_C \mathbf{a} \cdot d\mathbf{r} + i \int_C \mathbf{b} \cdot d\mathbf{r}$$

Proof. Suppose that $p: [c, d] \rightarrow \mathbb{C}$ is a path parametrising C . Then write $p(t) = p_1(t) + ip_2(t)$.

$$\begin{aligned}\int_C f(z)dz &= \int_c^d f(p(t))p'(t)dt \\ &= \int_c^d \left((u(p(t)) + iv(p(t)))(p_1'(t) + ip_2'(t)) \right) dt \\ &= \int_c^d (u(p(t))p_1'(t) - v(p(t))p_2'(t) + iv(p(t))p_1'(t) + iu(p(t))p_2'(t))dt \\ &= \int_c^d (u(p(t))p_1'(t) - v(p(t))p_2'(t))dt + i \int_c^d (v(p(t))p_1'(t) + u(p(t))p_2'(t))dt \\ &= \int_c^d \mathbf{a}(p(t)) \cdot p'(t)dt + i \int_c^d \mathbf{b}(p(t)) \cdot p'(t)dt \\ &= \int_C \mathbf{a} \cdot d\mathbf{r} + i \int_C \mathbf{b} \cdot d\mathbf{r}.\end{aligned}$$

□

The purpose of this proposition is so that we may use the previously proved results on integration of vector fields along curves in the definition of a complex integral over a curve. For example as a consequence of this result that two different parameterisations of the curve C lead to the same value for the integral, since this was proved for vector fields.

4.3 Integration with respect to arc-length

Recall that the length of a curve C , parametrised by $p(t)$, $a \leq t \leq b$, is given by the formula

$$\text{length}(C) = \int_a^b |p'(t)| dt.$$

For example the circle C_r of radius r can be defined by $p(t) = re^{it}$ for $0 \leq t \leq 2\pi$. The length of C_r is

$$\begin{aligned} \text{length}(C_r) &= \int_0^{2\pi} |ire^{it}| dt = \int_0^{2\pi} r dt \\ &= 2\pi r. \end{aligned}$$

This is an example of an integral with respect to arc-length. In general we have the following definition.

Definition 4.7 – Integral with respect to arc-length

Suppose that C is a piecewise continuously differentiable curve with parametrisation $p(t)$, $a \leq t \leq b$. The integral with respect to arc-length of an integrable function f is

$$\int_C f(z) |dz| = \int_a^b f(p(t)) |p'(t)| dt.$$

So we could write

$$\text{length}(C) = \int_C |dz|.$$

Integration with respect to arc-length is different to integration along a curve, and is useful in complex analysis since it provides a triangle inequality for integrals.

Proposition 4.8

Suppose that f is a complex-valued function defined on a piecewise continuously differentiable curve C . Then the following inequality holds.

$$\left| \int_C f(z) dz \right| \leq \int_C |f(z)| |dz|.$$

Proof. Using the triangle inequality, Proposition 4.1 part 2, we have that

$$\begin{aligned} \left| \int_C f(z) dz \right| &= \left| \int_a^b f(p(t)) p'(t) dt \right| \\ &\leq \int_a^b |f(p(t))| |p'(t)| dt \\ &= \int_C |f(z)| |dz|. \end{aligned}$$

□

You should make sure you understand what each of these quantities means.

As a consequence we have the following useful estimate of the contour integral over an arc of finite length.

$$\left| \int_C f(z) dz \right| \leq \text{length}(C) \sup_{z \in C} |f(z)|. \quad (4.1)$$

4.4 Some topological notions

Complex integration depends on curves. As such we need to collect together some ideas that help us describe correctly curves and other considerations.

First remember the following definitions.

Definition 4.9 – L

Let C be a curve parameterised by $p(t)$ for $a \leq t \leq b$.


- If $p(a) = p(b)$ then C is called a **closed curve**.
- If p is injective then C is called a **simple curve**.

Informally a closed curve is one that loops round and joins up at its endpoints. A simple curve is one that does not cross itself. See Figure 4.3.


Theorem 4.10 – The Jordan Curve Theorem

Let C be a simple closed curve in \mathbb{C} . Then $\mathbb{C} \setminus C$ consists of two components, a bounded component and an unbounded component.


A closed curve



A simple curve



A simple closed curve



A non-simple curve




Figure 4.3: Types of curves




Figure 4.4: The Jordan curve theorem

Drawing a diagram of this makes it look like a trivial result. Figure 4.4 does that. However it is not trivial and is in fact quite difficult to prove. A simple closed curve is often called a **Jordan curve** because of this theorem.

Jordan curves need not be well-behaved like the one in Figure 4.4. The snowflake curve for example is a Jordan curve. It is possible for Jordan curves to have positive area, [6].

In the long tradition of courses in complex analysis I will not go through this proof, it's rather long and involves many interesting ideas. In [1] it is merely mentioned as a side-comment. The full proof can be found in §21.3 of [3, pages 661-667].

Previously we said that a **domain** is a set in \mathbb{C} that is open and connected, see Definition 1.11. You spent a considerable time last year learning about open sets in metric spaces and these are quite familiar in \mathbb{C} . A set Ω is connected (also called path-connected) if for any two points $z, w \in \Omega$ there is a path $p: [0, 1] \rightarrow \Omega$ with $p(0) = z$ and $p(1) = w$.

We will also make use of the following definition (Definition 5.6 from the calculus notes).




Figure 4.5: Simply and multiply connected domains

Definition 4.11 – Simply connected domains

A set $\Omega \subset \mathbb{C}$ is **simply connected** if

- Ω is path-connected;
- the interior of any closed curve $C \subset \Omega$ is contained in Ω .

Informally a simply connected domain is one without holes in it. Figure 4.5 shows two domains the first is simply connected and the second is not — a domain that is not simply connected is called **multiply connected**. The red dashed curve is a simple closed curve whose interior is not a subset of the domain.

4.5 Cauchy's theorem

It doesn't matter how we parameterise a curve, the integral of a function over that curve is the same. This follows from Proposition 4.2 and the equivalent result for vector fields. In this section we will show that under some restrictions on f it doesn't matter what the curve looks like either, as long as the end-points are the same.

This is one of the most important theorems in complex analysis. Under fairly mild assumptions of complex-differentiability and curves of finite length it turns out that integrating along a curve depends only on the endpoints.

The entirety of this section will be devoted to proving the following.




Figure 4.6: A consequence of Cauchy's theorem

Theorem 4.12 – Cauchy's theorem

Suppose that $f(z)$ is holomorphic in the simply connected domain Ω . Then for any simple closed curve C in Ω of finite length

$$\int_C f(z)dz = 0.$$

In order to understand the importance of this theorem, consider a function $f(z)$ holomorphic in the domain Ω and suppose that a and b are two points in Ω . Let C_1 and C_2 be two curves joining a to b , see Figure 4.6. Then

$$\int_{C_1} f(z)dz = \int_{C_2} f(z)dz.$$

This follows because we can define the simple closed curve $C = C_1 \cup (-C_2)$. Then by Cauchy's theorem

$$\begin{aligned} 0 &= \int_C f(z)dz = \int_{C_1} f(z)dz + \int_{-C_2} f(z)dz \\ &= \int_{C_1} f(z)dz - \int_{C_2} f(z)dz. \end{aligned}$$

This argument only follows if C_1 and C_2 do not intersect, but even if they do the conclusion is still true. In general as long as $f(z)$ is holomorphic on a simply connected domain containing both arcs the integrals will be the same.

4.6 The weak form of Cauchy's Theorem

It is possible to prove a weak form of Cauchy's theorem using the Cauchy-Riemann equations and Green's theorem, via Proposition 4.2 that we proved earlier. Since it is so much more simple we provide it here before moving on to our stronger form.

Theorem 4.13 – Weak form of Cauchy's theorem

Suppose that $f(z)$ is holomorphic in the simply connected domain Ω . Let $f(x + iy) = u(x, y) + iv(x, y)$ and suppose that u and v are continuously differentiable. Then for any simple closed curve C in Ω of finite length

$$\int_C f(z)dz = 0.$$

The difference is in the assumption of continuous derivatives in the hypothesis. This allows us to use Green's theorem. This is the form of the theorem that Cauchy proved in 1825. It wasn't until 1900 that the mathematician Goursat devised a method for dispensing with the requirement that the derivative be continuous.

Proof of the weak form of Cauchy's theorem. With the notation of Proposition 4.2 we need only show that

$$\int_C \mathbf{a} \cdot d\mathbf{r} = 0, \quad \text{and} \quad \int_C \mathbf{b} \cdot d\mathbf{r} = 0.$$

First $\mathbf{a} = (u \ -v)^T$ and so, since these functions are continuously differentiable, by Green's theorem

$$\int_C \mathbf{a} \cdot d\mathbf{r} = \iint_U \left(-\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy,$$

where U is the interior of C .

However, since f is holomorphic, the Cauchy-Riemann equations (2.4) hold and

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}.$$

The integrand on the right is therefore 0 throughout U and so the integral is 0.

Similarly, again using Green's theorem,

$$\int_C \mathbf{b} \cdot d\mathbf{r} = \iint_U \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) dx dy$$

and, again, by the Cauchy-Riemann equations the integral on the right is 0. \square




Figure 4.7: Breaking a triangle into four sub-triangles

4.7 Proof of Cauchy's theorem

We will prove Cauchy's theorem in three steps, the first has already been done by proving the weak form in the last section. The next step is to prove the theorem for the perimeter of a triangle. This is based on an idea of Pringsheim. Goursat originally proved the theorem with rectangles instead of triangles. I prefer this proof, based on [4], since it involves a direct proof without any additional sidesteps or excursions.

Theorem 4.14 – Cauchy's theorem for a triangle

Suppose that f is holomorphic in a simply connected domain Ω , and T is the perimeter of a triangle contained in Ω . Then

$$\int_T f(z)dz = 0.$$

Proof. We divide the triangle T into four similar triangles, T_1 , T_2 , T_3 and T_4 by joining the midpoints of each side of T as indicated in Figure 4.7. The common sides of the triangles cancel each other when integrating since they are oriented in opposite directions. So

$$\int_T f(z)dz = \int_{T_1} f(z)dz + \int_{T_2} f(z)dz + \int_{T_3} f(z)dz + \int_{T_4} f(z)dz.$$

By the triangle inequality we have that

$$\left| \int_T f(z) dz \right| \leq \left| \int_{T_1} f(z) dz \right| + \left| \int_{T_2} f(z) dz \right| + \left| \int_{T_3} f(z) dz \right| + \left| \int_{T_4} f(z) dz \right|.$$

Therefore at least one of the integrals on the right hand side satisfies

$$\left| \int_{T_i} f(z) dz \right| \geq \frac{1}{4} \left| \int_T f(z) dz \right|.$$

We rename T_i as S_1 and continue this process with this triangle: Subdivide the triangle S_1 into four similar sub-triangles, with side-length half of the side-length of S_1 . It follows again that we may choose one of these smaller triangles, that we will call S_2 , such that

$$\left| \int_{S_2} f(z) dz \right| \geq \frac{1}{4} \left| \int_{S_1} f(z) dz \right| \geq \frac{1}{4^2} \left| \int_T f(z) dz \right|.$$

We continue in this way finding a sequence of triangles S_k , $k = 1, 2, \dots$ each smaller than the last, with side-length halved each time, and satisfying

$$\left| \int_{S_k} f(z) dz \right| \geq \frac{1}{4^k} \left| \int_T f(z) dz \right|. \quad (4.2)$$

Let \tilde{S}_k denote the union of S_k and its interior. We claim that the intersection of these triangles is a set containing a single point, z_0 . To see this let us define

$$\text{diam } A = \sup_{a, b \in A} |a - b|$$

to be the diameter of a set A . Since \tilde{S}_k is a triangle $\text{diam } \tilde{S}_k$ is the length of the largest side. Therefore we have that

$$\text{diam } \tilde{S}_{k+1} = \frac{1}{2} \text{diam } \tilde{S}_k, \quad k = 1, 2, \dots$$

It follows by induction that

$$\text{diam } \tilde{S}_k = \frac{1}{2^k} \text{diam } T.$$

Therefore $\text{diam } \tilde{S}_k \rightarrow 0$ as $k \rightarrow \infty$ and there is a z_0 such that

$$\cap_{k=1}^{\infty} \tilde{S}_k = \{z_0\}.$$

Now f is holomorphic at z_0 and so by Theorem 2.5 there is a function $E(h)$ such that

$$f(z_0 + h) = f(z_0) + f'(z_0)h + E(h)$$

and $E(h)/h \rightarrow 0$ as $h \rightarrow 0$. Write $z = z_0 + h$ then this becomes

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + E(z - z_0).$$

Since $f(z_0) + f'(z_0)(z - z_0)$ is a linear polynomial it has continuous derivatives of all orders, therefore by the weak form of Cauchy's theorem we have that

$$\int_{S_k} (f(z_0) + f'(z_0)(z - z_0))dz = 0.$$

It follows that for each k

$$\begin{aligned} \int_{S_k} f(z)dz &= \int_{S_k} (f(z_0) + f'(z_0)(z - z_0))dz + \int_{S_k} E(z - z_0)dz \\ &= \int_{S_k} E(z - z_0)dz. \end{aligned}$$

Let $\epsilon > 0$. Since $E(z - z_0)/(z - z_0) \rightarrow 0$ as $z \rightarrow z_0$ we may find n so that for all $z \in \tilde{S}_n$

$$|E(z - z_0)|/|z - z_0| < \epsilon. \quad (4.3)$$

Hence

$$\begin{aligned} \left| \int_{S_k} f(z)dz \right| &= \left| \int_{S_k} E(z - z_0)dz \right| \\ &\leq \int_{S_k} |E(z - z_0)||dz|, \end{aligned}$$

by Proposition 4.3,

$$\leq \epsilon \int_{S_k} |z - z_0||dz|.$$

by (4.3).

Now $z, z_0 \in S_k$, so

$$|z - z_0| \leq \text{diam } \tilde{S}_k = \frac{1}{2^k} \text{diam } T.$$

Furthermore the length of S_k is

$$\text{length}(S_k) = \frac{\text{length}(T)}{2^k}.$$

Let $L = \text{length}(T)$ and $D = \text{diam } T$. Then

$$\begin{aligned} \left| \int_{S_k} f(z) dz \right| &\leq \epsilon \int_{S_k} |z - z_0| |dz| \\ &\leq \epsilon \frac{D}{2^k} \int_{S_k} |dz| \\ &\leq \epsilon \frac{D}{2^k} \frac{L}{2^k} = \epsilon \frac{DL}{4^k}. \end{aligned}$$

Combining this with (4.2) we find that for any $\epsilon > 0$ have

$$\begin{aligned} \left| \int_T f(z) dz \right| &\leq 4^k \left| \int_{S_k} f(z) dz \right| \\ &\leq 4^k \epsilon \frac{DL}{4^k} \\ &= \epsilon DL. \end{aligned}$$

Since $\epsilon > 0$ was arbitrary,

$$\int_T f(z) dz = 0$$

as required. □

Any polygon, such as the one drawn in Figure 4.8, can be decomposed into triangles – a process called triangulation. Integrating along a polygon can be written as the sum of a finite number of integrals along the triangles. As we have seen before sides of triangles appear twice in this decomposition but oriented in opposite directions, so their integrals along the common edges cancel. Therefore a consequence of Theorem 4.14 is the following.

Corollary 4.15. *Suppose that f is holomorphic in a simply connected domain Ω , and C is the perimeter of a polygon contained in Ω . Then*

$$\int_C f(z) dz = 0.$$




Figure 4.8: Cauchy's theorem for a polygonal curve

It remains to prove Cauchy's theorem in full generality.

The idea is to approximate the curve C by a sequence of polygons sufficiently well that the integrals along the polygons converge to the integral along C . The only thing we have to worry about is the convergence and so we need to be careful about how we define the polygon. Integral convergence is relatively weak and so we only need to employ the continuity of f , not the fact that it is holomorphic.

We will require the following result that strengthens the notion of continuity in metric spaces.

Lemma 4.16. *Let (X, d_X) and (Y, d_Y) be metric spaces and suppose that $f: X \rightarrow Y$ is a continuous function. Let E be a compact set in (X, d_X) then for all $\epsilon > 0$ there is a $\delta > 0$ such that for all $x, y \in E$ and $d_X(x, y) < \delta$ we have that $d(f(x), f(y)) < \epsilon$.*

The important thing here is that δ is independent of x and y as long as they are in the compact set E .

Proof. Suppose that $\epsilon > 0$. Since f is continuous we can find, for each $x \in E$ a $\delta(x)$ such that

$$f(B(x, \delta(x))) \subset B(f(x), \epsilon/2).$$

Now the family of open sets $\{B(x, \delta(x)/2) : x \in E\}$ is an open cover of E and so there is a finite sub cover

$$\{B(x_i, \delta(x_i)/2) : i = 1, 2, \dots, N\}.$$

Let $\delta = \min\{\delta(x_i) : i = 1, 2, \dots, N\}$.

Suppose that $x, y \in E$ are such that $d_X(x, y) < \delta$. Let i be chosen so that $x \in B(x_i, \delta(x_i)/2)$ then

$$\begin{aligned} d_X(y, x_i) &\leq d_X(y, x) + d_X(x, x_i) \\ &< \frac{\delta}{2} + \frac{\delta(x_i)}{2} \\ &\leq \frac{\delta(x_i)}{2} + \frac{\delta(x_i)}{2} = \delta(x_i). \end{aligned}$$

Therefore $y \in B(x_i, \delta(x_i))$ for some i . But then

$$\begin{aligned} d_Y(f(x), f(y)) &\leq d_Y(f(x), f(x_i)) + d_Y(f(x_i), f(y)) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

The result is proved. □

Proof of Cauchy's theorem. Since C is a simple closed curve in Ω we can find a compact set R such that $C \subset R \subset \Omega$. By Lemma 4.16 there is a $\delta > 0$ so that whenever $z, w \in R$,

$$|z - w| < \delta \quad \Rightarrow \quad |f(z) - f(w)| < \frac{\epsilon}{2L}$$

where $L = \text{length}(C)$.

Suppose that C is positively oriented and $P_\epsilon = \{z_0, z_1, \dots, z_n\}$ is a partition of C ordered so that z_{k+1} follows z_k on C for each $k = 0, 1, \dots, n-1$, P_ϵ is such that

$$\left| \int_C f(z) dz - \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) \right| < \frac{\epsilon}{2}, \quad (4.4)$$

where we adopt the convention $z_{n+1} = z_0$ to avoid repetition.

We can choose P_ϵ so that the following hold, see Figure 4.9:

1. for each $k = 0, 1, \dots, n$ the line segment joining z_k to z_{k+1} is contained in R ;
2. the polygon, Π , whose edges are the line segments z_k to z_{k+1} , $k = 0, 1, \dots, n$, does not intersect itself
3. for each k , $|z_{k+1} - z_k| < \delta$.




Figure 4.9: The proof of Cauchy's theorem

Only the second of these is controversial. It is possible of course that Π does intersect itself, but in this case we are left with a finite number of polygons, each of which satisfies the properties above. Since we only need the integral along Π to be 0 we can restrict the remainder of the proof to these finitely many polygons individually if needed.

Let Π_k be the part of Π consisting of the line segment from z_k to z_{k+1} , $k = 0, 1, \dots, n$. We have that

$$\int_{\Pi} f(z) dz = \sum_{k=0}^n \int_{\Pi_k} f(z) dz.$$

We also have

$$\int_{\Pi_k} f(z_k) dz = f(z_k)(z_{k+1} - z_k).$$

Therefore

$$\begin{aligned} \left| \int_{\Pi} f(z) dz - \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) \right| &= \left| \sum_{k=0}^n \int_{\Pi_k} (f(z) - f(z_k)) dz \right| \\ &\leq \sum_{k=0}^n \int_{\Pi_k} |f(z) - f(z_k)| |dz|, \end{aligned}$$

by Proposition 4.3, and the estimate above

$$\begin{aligned} &\leq \frac{\epsilon}{2L} \sum_{k=0}^n \int_{\Pi_k} |dz| \\ &\leq \frac{\epsilon}{2L} L = \frac{\epsilon}{2} \end{aligned}$$

since the length of Π is less than or equal to the length of C .

It follows by the triangle inequality that

$$\begin{aligned} \left| \int_C f(z) dz - \int_{\Pi} f(z) dz \right| &= \left| \int_C f(z) dz - \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) + \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) - \int_{\Pi} f(z) dz \right| \\ &\leq \left| \int_C f(z) dz - \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) \right| + \left| \int_{\Pi} f(z) dz - \sum_{k=0}^n f(z_k)(z_{k+1} - z_k) \right| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

However by Corollary 4.15

$$\int_{\Pi} f(z) dz = 0$$

and so this reduces to

$$\left| \int_C f(z) dz \right| < \epsilon.$$

Since $\epsilon > 0$ was arbitrary the theorem is proved. \square

Things to know

- How to find a parameterisation of standard examples of curves, such as a line segment or an arc of a circle;
- How to distinguish between the different types of curves as shown in Figure 4.3;
- How to integrate a given function along a curve;
- How to integrate a given function along a curve with respect to arc-length, especially the calculation of the length of a curve;
- The triangle inequality for integrals along a curve;

- The statement but not the proof of Cauchy's theorem. You should know in particular the important subtleties of the statement — namely that Ω has to be simply connected and (very importantly) that f is not required to have continuous partial derivatives.

Further reading

Cauchy's theorem is fundamental and so can be found in any and all books on undergraduate complex analysis. Each of our standard references treats it differently and proves it differently (many for example use Goursat's original proof of Cauchy's theorem in a rectangle instead of a triangle). If you can get your hands on [4] and [5] then they treat this topic very well.

There are some nice examples to work through at the end of [7, Chapter 6, §1]. Unfortunately not many textbooks give practice examples of finding parameterisations in \mathbb{C} – so use the ones I've given you.

Problems

1. To parameterise a line segment from a to b you can use

$$p(t) = a + t(b - a), \quad 0 \leq t \leq 1.$$

Find a parameterisation for the following line segments.

- (a) The line from 1 to i
 - (b) The line above in the opposite direction
 - (c) The line from $1 + i$ to $-1 - i$
 - (d) The line above in the opposite direction
2. To parameterise an arc of the boundary of $B(a, r)$ in the anti-clockwise direction

$$p(t) = a + re^{it}, \quad \theta_1 \leq t \leq \theta_2.$$

Find the parameterisations of the following arcs in the anti-clockwise direction.

- (a) Centred at i from π to $3\pi/2$, radius 1;
 - (b) Centred at $-i$ from 0 to π , radius 1.
3. To parameterise an arc of the boundary of $B(a, r)$ in the clockwise direction

$$p(t) = a + re^{i(\theta_1 + \theta_2 - t)}, \quad \theta_1 \leq t \leq \theta_2.$$

Find the parameterisations of the following arcs in the clockwise direction.

- (a) Centred at i from π to $3\pi/2$;
 - (b) Centred at $-i$ from 0 to π .
4. Find a parameterisation of the simple closed curve from -1 to 1 and then along a semi-circle in the upper half-plane (above the real axis) back to -1 .
5. Sketch the following curves and find a parameterisation for each of them
- (a) The semi-circle of radius 3 , anticlockwise
 - (b) The semi-circle of radius 3 , clockwise
 - (c) The line segment from 0 to 2

- (d) The line segment from 0 to $1 + i$
6. Find the integral of $f(z) = z^2$ along the line segment joining $-i$ to i
 7. Find the integral of $f(z) = z^3$ along the portion of the circle of radius 1 centred at the origin in the first quadrant, in the anti-clockwise direction
 8. Find the integral of $f(z) = 1/z$ along the circle of radius 2 centred at the origin in the clockwise direction
 9. Find the length of the curve parameterised by $p(t) = 3e^{2it} + 2$ for $-\pi \leq t \leq \pi$
 10. Find the length of the curve parameterised by $p(t) = e^t \cos t + ie^t \sin t$ for $0 \leq t \leq 2\pi$
 11. Find the length of the curve C parameterised by $p(t) = 10e^{it} - 2e^{5it}$, $0 \leq t \leq 2\pi$.
 12. Verify Cauchy's theorem for the functions

(a) $3z^2 + iz - 4$

(b) $\exp z$

if C is the square with vertices $1 \pm i$ and $-1 \pm i$.

- * 13. Suppose C_r is the circle of radius r taken in the anti-clockwise direction, and $|f(z)| \leq M$ for all $|z| \leq 1$. Use the triangle inequality to show

$$\lim_{r \rightarrow 0} \int_{C_r} f(z) dz = 0.$$

Cauchy's theorem proof

- * 14. Show that for any **bounded** open set $A \subset \mathbb{C}$, there are two limit points $a, b \in A'$ such that

$$\text{diam } A = |a - b|.$$

Deduce that the diameter of a triangle T is the length of the largest side.

- * 15. Check that for the line segment Π_k joining points z_k to z_{k+1} we have

$$\int_{\Pi_k} f(z_k) dz = f(z_k)(z_{k+1} - z_k)$$

- * 16. Construct a simple closed curve C and a polygon Π as in the proof of Cauchy's theorem such that Π intersects itself. Why does this not affect the proof?

Chapter 5

Applications of Cauchy's Theorem

“[The student] should understand mathematics as a living organism, a growing body of learning, fed by the efforts of many workers. If possible, he should acquire a sense of veneration for those who built up the structure that he is studying.”

Analytic Function Theory, Vol I by Einar Hille.

5.1 Doubly connected domains

Cauchy's theorem has a number of important applications in complex analysis. In this chapter we will explore a few of them to extend what you have already learned about holomorphic functions.

Our first application of Cauchy's theorem is its extension to doubly connected domains. These are domains such as the one in Figure 5.1. Informally, they are domains with a hole in them. Cauchy's theorem is no longer necessarily true if our holomorphic function is defined only in a doubly connected domain.

However we can prove the following. First note that the Jordan curve theorem lets us assign an interior and exterior to a simple closed curve.

Lemma 5.1. *Let C_1 and C_2 denote simple closed curves with C_2 in the interior of C_1 . Let Ω denote the doubly connected domain with boundary $C_1 \cup C_2$. If a function f is holomorphic in a domain containing Ω , then*




Figure 5.1: A doubly connected domain

$$\int_{C_1} f(z)dz = \int_{C_2} f(z)dz.$$




Figure 5.2: Proof of Lemma 5.1

Proof. Let L be a curve that joins C_2 to C_1 and lies entirely inside the doubly connected domain Ω . And suppose the end-points of the L are the points z_1 and z_2 , as shown in Figure 5.2.

Let K be the curve starting at z_1 and travelling around $-C_2$, then along L to z_2 and along C_1 , and finally back to z_1 along L . Note K is a simple closed curve with simply connected interior, so that by Cauchy's theorem

$$\int_K f(z)dz = 0.$$

Hence,

$$\begin{aligned} 0 &= \int_K f(z)dz = \int_{-C_2} f(z)dz + \int_L f(z)dz + \int_{-L} f(z)dz + \int_{C_1} f(z)dz \\ &= - \int_{C_2} f(z)dz + \int_{C_1} f(z)dz \end{aligned}$$

□

5.2 Cauchy's integral formula

We begin with the following remarkable result.

Theorem 5.2 – Cauchy's integral formula

Suppose that $f \in H(\Omega)$ where Ω is a simply connected domain. Let C be a simple closed curve in Ω with ζ a point in its interior. Then

$$f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z - \zeta} dz.$$

Lemma 5.3. Suppose that $f \in H(\Omega)$ and C a simple closed curve with ζ a point in its interior. Then

$$\int_C \frac{1}{z - \zeta} dz = 2\pi i.$$

Proof. First, by Lemma 5.1,

$$\int_C \frac{1}{z - \zeta} dz = \int_D \frac{1}{z - \zeta} dz,$$

where D is the boundary of the disk $B(\zeta, r)$ and $r > 0$ is chosen so that $B(\zeta, r)$ is interior to C , see Figure 5.3




Figure 5.3: The proof of Lemma 5.3

Next, we parameterise $\partial B(\zeta, R)$ as $p(t) = \zeta + re^{it}$ for $0 \leq t < 2\pi$. Then

$$\begin{aligned} \int_{\partial B(\zeta, R)} \frac{1}{z - \zeta} dz &= \int_0^{2\pi} \frac{p'(t)}{p(t) - \zeta} dt \\ &= \int_0^{2\pi} \frac{ire^{it}}{re^{it}} dt \\ &= \int_0^{2\pi} i dt = 2\pi i. \end{aligned}$$

□

Proof of Theorem 5.2. Since $(f(z) - f(\zeta))/(z - \zeta)$ is holomorphic in any sub-domain that does not contain ζ , by Lemma 5.1,

$$\int_C \frac{f(z) - f(\zeta)}{z - \zeta} dz = \int_{\partial B(\zeta, r)} \frac{f(z) - f(\zeta)}{z - \zeta} dz \quad (5.1)$$

for $r > 0$ chosen so that $B(\zeta, r)$ is in the interior of C .

Next we show that this integral is 0. Since f is continuous at ζ we have that for any $\epsilon > 0$ there is a $\delta > 0$ such that

$$|z - \zeta| < \delta \quad \Rightarrow \quad |f(z) - f(\zeta)| < \epsilon.$$

For $r < \delta$

$$\begin{aligned} \left| \int_{\partial B(\zeta, r)} \frac{f(z) - f(\zeta)}{z - \zeta} dz \right| &\leq \int_{\partial B(\zeta, r)} \frac{|f(z) - f(\zeta)|}{|z - \zeta|} |dz| \\ &< \epsilon \int_{\partial B(\zeta, r)} \frac{1}{|z - \zeta|} |dz| \\ &= \epsilon \frac{1}{r} \int_{\partial B(\zeta, r)} |dz|, \end{aligned}$$

since $|z - \zeta| = r$ on $\partial B(\zeta, r)$,

$$\begin{aligned} &= \epsilon \frac{1}{r} \text{length}(\partial B(\zeta, r)) = \epsilon \frac{1}{r} 2\pi r \\ &= 2\pi\epsilon. \end{aligned}$$

By (5.1) we have that

$$\left| \int_C \frac{f(z) - f(\zeta)}{z - \zeta} dz \right| = \left| \int_{\partial B(\zeta, r)} \frac{f(z) - f(\zeta)}{z - \zeta} dz \right| < 2\pi\epsilon$$

and since ϵ was arbitrary we have shown that

$$\int_C \frac{f(z) - f(\zeta)}{z - \zeta} dz = 0.$$

Now

$$\frac{1}{2\pi i} \int_C \frac{f(z)}{z - \zeta} dz - f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z - \zeta} dz - \frac{1}{2\pi i} f(\zeta) \int_C \frac{1}{z - \zeta} dz,$$

by Lemma 5.3,

$$\begin{aligned} &= \frac{1}{2\pi i} \int_C \frac{f(z)}{z - \zeta} dz - \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{z - \zeta} dz \\ &= \frac{1}{2\pi i} \int_C \frac{f(z) - f(\zeta)}{z - \zeta} dz = 0, \end{aligned}$$

and the result is proved. □

5.3 Holomorphic functions are analytic

When we defined analytic functions we noted that since you can differentiate power series, an analytic function is holomorphic. The converse is also true. A function that is holomorphic in a domain Ω is equal to its Taylor series in a neighbourhood of each point in Ω . In particular a holomorphic function is analytic, making the two definitions equivalent.

In this section we will prove this result. It is based on Cauchy's integral formula.

Theorem 5.4

Suppose that f is holomorphic in $B(z_0, r)$ for $z_0 \in \mathbb{C}$ and $r > 0$. Then $f(z)$ is equal to its Taylor series centred at z_0 for all $z \in B(z_0, r)$.

Lemma 5.5. Let $z, z_0, h \in \mathbb{C}$ then

$$\frac{1}{z - (z_0 + h)} = \frac{1}{z - z_0} + \frac{h}{(z - z_0)^2} + \cdots + \frac{h^n}{(z - z_0)^{n+1}} + \frac{h^{n+1}}{(z - z_0)^{n+1}(z - (z_0 + h))} \quad (5.2)$$

Proof. We begin with the geometric series formula:

$$(1 - w)(1 + w + \cdots + w^n) = 1 - w^{n+1}.$$

If we substitute $w = h/(z - z_0)$ and rearrange, then

$$\left(1 - \frac{h}{z - z_0}\right) \left(1 + \frac{h}{z - z_0} + \cdots + \frac{h^n}{(z - z_0)^n}\right) + \frac{h^{n+1}}{(z - z_0)^{n+1}} = 1.$$

It follows that

$$\left(\frac{z - (z_0 + h)}{z - z_0}\right) \left(1 + \frac{h}{z - z_0} + \cdots + \frac{h^n}{(z - z_0)^n}\right) + \frac{h^{n+1}}{(z - z_0)^{n+1}} = 1$$

and so

$$(z - (z_0 + h)) \left(\frac{1}{z - z_0} + \frac{h}{(z - z_0)^2} + \cdots + \frac{h^n}{(z - z_0)^{n+1}}\right) + \frac{h^{n+1}}{(z - z_0)^{n+1}} = 1.$$

Dividing by $z - (z_0 + h)$ then gives the result. □

Lemma 5.6. Suppose that $d > 0$, $z \in \partial B(z_0, d)$ and $\zeta \in B(z_0, d)$ then

$$d - |\zeta - z_0| \leq |z - \zeta| \leq |\zeta - z_0| + d.$$




Figure 5.4: The proof of Lemma 5.6

Proof. The diagram in Figure 5.4 demonstrates this. Thinking of z_0 and ζ as fixed points and z as moving around the circle, you should convince yourself that the distance from z to ζ is minimised and maximised precisely when z is at the opposite ends of the diameter passing through ζ . \square

Proof of Theorem 5.4. We only need to show that f can be written as a power series with radius of convergence r . The fact that it is then its Taylor series follows from Corollary 2.23.

Let $d < r$. Then the closure of the ball $B(z_0, d)$ is contained in $B(z_0, r)$. We will show that $f(\zeta)$ can be written as a power series whenever $\zeta = z_0 + h \in B(z_0, d)$.

First, by (5.2),

$$\frac{1}{z - \zeta} = \frac{1}{z - z_0} + \frac{h}{(z - z_0)^2} + \cdots + \frac{h^n}{(z - z_0)^{n+1}} + \frac{h^{n+1}}{(z - z_0)^{n+1}(z - \zeta)}.$$

Multiplying by $f(z)/(2\pi i)$ and integrating along the curve $C = \partial B(z_0, d)$ in the positive

direction gives

$$\begin{aligned} f(\zeta) &= \frac{1}{2\pi i} \int_C \frac{f(z)}{z - \zeta} dz = \sum_{k=0}^n \left(\frac{1}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{k+1}} dz \right) h^k + R_n(h) \\ &= \sum_{k=0}^n \left(\frac{1}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{k+1}} dz \right) (\zeta - z_0)^k + R_n(h) \end{aligned}$$

where

$$R_n(h) = \frac{h^{n+1}}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{n+1}(z - \zeta)} dz.$$

Now

$$\begin{aligned} |R_n(h)| &\leq \frac{|h|^{n+1}}{2\pi} \int_C \frac{|f(z)|}{|z - z_0|^{n+1}|z - \zeta|} |dz| \\ &\leq \frac{|h|^{n+1}}{2\pi} \frac{M}{d^{n+1}} \frac{1}{d - |h|} \text{length}(C) \\ &= \frac{Md}{d - |h|} \left(\frac{|h|}{d} \right)^{n+1}, \end{aligned}$$

where $M = \sup_{z \in C} |f(z)|$ and we have used Lemma 5.6 to estimate $1/|z - \zeta| \leq 1/(d - |h|)$, and $\text{length}(C) = 2\pi d$.

It follows that $R_n(h) \rightarrow 0$ as $n \rightarrow \infty$ and the series converges to $f(\zeta)$:

$$f(\zeta) = \sum_{k=0}^{\infty} \left(\frac{1}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{k+1}} dz \right) (\zeta - z_0)^k. \quad (5.3)$$

This is true for all $\zeta \in B(z_0, d)$ and so since $d < r$ was arbitrary this means $f(z)$ can be represented by its Taylor series, centred at z_0 , for all $z \in B(z_0, r)$. \square

A number of corollaries follow from this result.

Corollary 5.7. *Suppose that $\Omega \subset \mathbb{C}$ is a domain then $H(\Omega) = A(\Omega)$.*

Proof. Suppose that $z_0 \in \Omega$ is arbitrary. Then since Ω is open there is a $r > 0$ such that $B(z_0, r) \subset \Omega$. By the theorem $f(z)$ is equal to its Taylor series at all points in $B(z_0, r)$ and hence it is analytic at z_0 . \square

Combining this with Corollary 2.22 we get the following incredible result.

Corollary 5.8. Suppose that f is complex-differentiable in $B(z_0, r)$ for some z_0 and $r > 0$ then $f^{(k)}$ exists at all points of $B(z_0, r)$ for all k

In other words if a function is differentiable then it is infinitely differentiable!

If we now compare the description of the Taylor coefficients in Corollary 2.23 to (5.3) and rearrange we have the following consequence of the result, which is an extension of Cauchy's integral formula

Theorem 5.9 – Cauchy's integral formula for derivatives

Suppose that $f \in H(\Omega)$ where Ω is a simply connected domain. Let C be a simple closed curve in Ω with ζ a point in its interior. Then

$$f^{(k)}(\zeta) = \frac{k!}{2\pi i} \int_C \frac{f(z)}{(z - \zeta)^{k+1}} dz.$$

And finally we put to bed the question of why the Taylor series of $1/(1+x^2)$ has radius of convergence 1 when it has continuous derivatives of all orders (see Example 2.16).

Corollary 5.10. Suppose that the power series $\sum_{n=0}^{\infty} a_n(z-a)^n$ has radius of convergence $R > 0$. Then there is at least one point of $\partial B(a, R)$ at which $f(z)$ fails to be holomorphic.

5.4 Zeros of holomorphic functions

A zero of a holomorphic function is a point z such that $f(z) = 0$. For example the function $z^2 + 1$ has zeros at $\pm i$. There are a couple of important results on the zeros of holomorphic functions that are worth mentioning here. They again highlight the difference between real analysis and complex analysis.

Theorem 5.11

Suppose $f \in H(\Omega)$ and (z_m) is a sequence of zeros of f , i.e. for each m , $f(z_m) = 0$. If $z_m \rightarrow a$ as $m \rightarrow \infty$ and $a \in \Omega$ then $f(z) = 0$ for all $z \in \Omega$.

Proof. Since $a \in \Omega$, by Theorem 5.4 we may define $f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n$, in $B(a, R)$, for $R > 0$. We will prove by induction that $a_n = 0$ for all n .

Since f is holomorphic it is continuous at a . Therefore $f(a) = f(\lim_{m \rightarrow \infty} z_m) = \lim_{m \rightarrow \infty} f(z_m) = 0$. So

$$0 = f(a) = a_0.$$

Now suppose then that a_0, a_1, \dots, a_k are all 0. Then

$$\begin{aligned} f(z) &= a_{k+1}(z-a)^{k+1} + a_{k+2}(z-a)^{k+2} + \dots \\ &= (z-a)^{k+1}(a_{k+1} + a_{k+2}(z-a) + \dots) \\ &= (z-a)^{k+1}g(z). \end{aligned}$$

Here $g(z) = a_{k+1} + a_{k+2}(z-a) + \dots$ is a holomorphic function in $B(a, R)$ since it can be written as a power series. Note also that

$$g(z_m) = \frac{f(z_m)}{(z_m - a)^{k+1}} = 0$$

for all m and so by continuity $a_{k+1} = g(a) = \lim_{m \rightarrow \infty} g(z_m) = 0$.

It follows by induction that $a_n = 0$ for all n . □

Results about zeros always lead to results about uniqueness – i.e. results that give conditions under which two holomorphic functions are equal.

Corollary 5.12. *Suppose that $f, g \in H(\Omega)$. If $z_m \rightarrow a \in \Omega$, as $m \rightarrow \infty$, is such that $f(z_m) = g(z_m)$ for all m then $f = g$.*

Another important consequence is the following

Corollary 5.13. *Suppose $f \in H(\Omega)$ and $f(a) = 0$. Then there is a $R > 0$ such that $f(z) \neq 0$ for any $z \in B(a, R)$ except for a .*

Finally we have the following characterisation of zeros of holomorphic functions.

Corollary 5.14. *Suppose that $f \in H(\Omega)$ and $f(a) = 0$. Then there is a $k \geq 1$ and $R > 0$ such that $f(z) = (z-a)^k g(z)$ where $g \in H(B(a, R))$ and $g(z) \neq 0$ for all $z \in B(a, R)$.*

Proof. As in the proof of Theorem 5.11 if $f(a) = 0$ then we may write $f(z)$ as a power series

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n$$

in a neighbourhood $B(a, R)$. By the previous corollary we also know that we can choose R small enough so that $f(z)$ has no other zeros in $B(a, R)$.

Since $f(a) = 0$ we have that $a_0 = 0$. If we let k be the smallest positive integer such that $a_k \neq 0$ then $a_0 = a_1 = \dots = a_{k-1} = 0$ and

$$f(z) = \sum_{n=k}^{\infty} a_n (z-a)^n = (z-a)^k \sum_{n=k}^{\infty} a_n (z-a)^{n-k}.$$

The corollary now follows since $g(z) = \sum_{n=k}^{\infty} a_n (z-a)^{n-k}$ is holomorphic and non-zero in $B(a, R)$. \square

Definition 5.15 – Multiplicity

The number k in the previous corollary is called the multiplicity of the zero a .

5.5 Liouville's theorem

A final consequence of Cauchy's theorem, or more precisely a consequence of Cauchy's integral formula, is an important estimate of entire functions that is surprisingly useful. It is a deep theorem of Liouville that is used broadly throughout mathematics.

Theorem 5.16 – Liouville's theorem

Suppose that f is an entire function and there is a $M > 0$ such that for all $z \in \mathbb{C}$,

$$|f(z)| \leq M.$$

Then $f(z)$ is constant throughout \mathbb{C} .

Proof. Since $f(z)$ is entire we can write it as a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ with infinite radius of convergence.

Let $k \geq 1$ be arbitrary. We will show that $a_k = 0$. To do this let

$$C_R = \{z \in \mathbb{C} : |z| = R\}.$$

By Cauchy's integral formula, Theorem 5.9,

$$\begin{aligned}
 |a_k| &= \left| \frac{f^{(k)}(0)}{k!} \right| = \left| \frac{1}{2\pi i} \int_{C_R} \frac{f(z)}{z^{k+1}} dz \right| \\
 &\leq \frac{1}{2\pi} \int_{C_R} \frac{|f(z)|}{|z|^{k+1}} |dz| \\
 &\leq \frac{M}{2\pi} \frac{1}{R^{k+1}} \text{length}(C_R) \\
 &= M \frac{R}{R^{k+1}} = \frac{M}{R^k}, \quad \text{since } \text{length}(C_R) = 2\pi R.
 \end{aligned}$$

But since $R^{-k} \rightarrow 0$ as $R \rightarrow \infty$ the last expression can be made arbitrarily small. Therefore we have shown $a_k = 0$ for all $k \geq 1$. Hence $f(z) = a_0$ as required. \square

An immediate consequence of this is the fundamental theorem of algebra.

Theorem 5.17 – The fundamental theorem of algebra

Let $p(z) \in \mathbb{C}[z]$ have $\deg p \geq 1$. Then there is $\alpha \in \mathbb{C}$ such that $p(\alpha) = 0$ and p can be factorised as follows

$$p(z) = (z - \alpha)q(z)$$

where $q(z) \in \mathbb{C}[z]$.

The proof is a direct application of Liouville's theorem. The idea is that if $p(z) \neq 0$ then $1/p(z)$ is an entire function. If we can show it is bounded then we can deduce it is constant, contradicting the hypothesis that $\deg p \geq 1$.

Proof. Writing $p(z) = a_0 + a_1 z + \dots + a_n z^n$ where $a_n \neq 0$ and $n > 0$ we have

$$p(z) = z^n \left(a_n + \frac{a_{n-1}}{z} + \dots + \frac{a_0}{z^n} \right) = z^n q(z).$$

Now $\lim_{z \rightarrow \infty} q(z) = a_n$ and so as $z \rightarrow \infty$, $|p(z)| \rightarrow \infty$. It follows that there is a $R > 0$ such that $|p(z)| \geq 1$ for z such that $|z| \geq R$. Now let $f(z) = 1/p(z)$. Then $|f(z)| \leq 1$ for $|z| \geq R$.

Suppose now that $p(z) \neq 0$ for any $z \in \mathbb{C}$. Then $f(z)$ is holomorphic on the compact set $\overline{B(0, R)}$ and so we also have that $|f(z)|$ is bounded, by $M > 1$ say, on this set. Therefore, combining these two bounds, $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Hence, by Liouville's theorem,

$f(z)$ is constant. It follows that p is constant – a contradiction. The existence of α follows, as does the factorisation. \square

Entire functions are different in many ways to functions that are holomorphic on a domain Ω strictly contained in \mathbb{C} . Complex analysis can be split between entire functions and non-entire functions. Two important theorems that illustrate this will be given here, they are unique to entire functions. For more information the classic book of Boas, [2], although difficult to find these days, is an excellent introduction.

Theorem 5.18

A non-constant entire function comes arbitrarily close to every complex number.

In other words given any $w \in \mathbb{C}$ and any $\epsilon > 0$ there is a z such that $|w - f(z)| < \epsilon$.

Proof. Suppose to the contrary that $|f(z) - w| \geq \epsilon$ for all $z \in \mathbb{C}$. Then $g(z) = 1/(f(z) - w)$ is entire and bounded by $1/\epsilon$. Therefore $g(z)$ is constant and, rearranging, it follows that $f(z)$ is constant, contradicting the hypothesis. \square

Of course this doesn't say that an entire function takes every complex value, it just says it comes arbitrarily close to them. However we state here the following fundamental theorem that says that there can only be one point that an entire function doesn't take.

Theorem 5.19 – Picard's theorem

A non-constant entire function assumes each complex value, with one possible exception.

Things to know

- Cauchy's integral formula, both the standard one and the one for derivatives, and how to apply them to find a given integral;
- The notion that Holomorphic functions and Analytic functions are the same, and the consequences of that;
- The properties of the zeros of a holomorphic function, in particular from Theorem 5.11, Corollary 5.13 and Corollary 5.14;

- The definition of multiplicity;
- The statement and proof of Liouville's theorem;
- Applications of Liouville's theorem to uniqueness problems in complex analysis.

Further reading

The proof of Cauchy's integral formula is based on the treatment in [3]. The proof can be done with a more general definition of a curve – I've just insisted that it has finite length but in fact all the results can be proved with a more technical assumption of **rectifiability**. The two notions are similar but the latter adds an extra distracting level of technicality to proofs. The results can be found in full generality in [3] and also in [1] and [7].

Zeros of holomorphic functions are best discussed in [3]. Ahlfors, [1, Chapter 4, §3], covers these ideas briefly as well.

Liouville's theorem and its applications can be found in [3] and all other textbooks. It is classic.

Problems

1. State the following integrals (no calculation is needed). In each case C is the boundary of the unit disk $B(0, 1)$.

(a) $\frac{1}{2\pi i} \int_C \frac{1}{z} dz;$

(b) $\frac{1}{2\pi i} \int_C \frac{1}{z^2} dz;$

(c) $\frac{1}{2\pi i} \int_C \frac{z^3}{z - 1/2} dz;$

(d) $\frac{1}{2\pi i} \int_C \frac{z^3}{(z - 1/2)^2} dz.$

2. Suppose C is the closed curve parameterised by $p(t) = e^{it}$ for $0 \leq t \leq 4\pi$. Show that

$$\frac{1}{2\pi i} \int_C \frac{1}{z} dz = 2.$$

Why isn't this a contradiction of Cauchy's integral formula?

3. Verify Cauchy's integral formula for the function $f(z) = z^n$ in the disk $B(0, 1)$. That is show that

$$\zeta^n = \frac{1}{2\pi i} \int_C \frac{z^n}{z - \zeta} dz.$$

Hint: write $z^n = z^n - \zeta^n + \zeta^n$, split the integral and use Cauchy's theorem and Lemma 4.2.

4. Prove that if $f(z)$ is holomorphic in $B(0, R)$ for some $R > 1$, then the coefficients a_n of its power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ satisfy

$$a_n = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-in\theta} d\theta.$$

5. Suppose $f(z)$ is entire and $|f(z)| \leq |\exp z| = e^{\operatorname{Re} z}$ for all $z \in \mathbb{C}$. Show that there is a λ with $|\lambda| = 1$ such that $f(z) = \lambda \exp z$.
6. Extend Liouville's theorem to the following. Suppose that $|f(z)| \leq M|z|^n$ for all $z \in \mathbb{C}$ and some $M > 0$. Then $f(z)$ is a polynomial and $\deg f(z) \leq n$.

- * 7. Show that if $f(z)$ is entire and $\operatorname{Re} f(z) \leq M$ for all z then $f(z)$ is constant.
- * 8. Suppose $f(z)$ and $g(z)$ are entire and there is a $a > 0$ such that $\operatorname{Re} f(z) \leq a \operatorname{Im} g(z)$ for all z . Show that

$$f(z) = \alpha g(z) + \beta, \quad \text{for some } \alpha, \beta \in \mathbb{C}.$$

- * 9. Show that if $f(z)$ is entire and $(\operatorname{Re} f(z))^2 \leq (\operatorname{Im} f(z))^2$ for all z then $f(z)$ is constant.

Chapter 6

The Residue Theorem

“The shortest route between two truths in the real domain passes through the complex domain.”

Jacques Salomon Hadamard (1865-1963)

6.1 Singularities

In much of what we’ve done so far we have been studying holomorphic functions in a domain Ω . Applications of Cauchy’s theorem require functions to be holomorphic.

In this chapter we’re going to extend some of the ideas to functions that are not holomorphic everywhere but have isolated points where they are not holomorphic. We’ve already seen an example of this when we saw Cauchy’s integral formula:

$$f^{(n)}(\zeta) = \frac{n!}{2\pi i} \int_C \frac{f(z)}{(z - \zeta)^{n+1}} dz, \quad n \geq 0.$$

The integrand here, $\frac{f(z)}{(z - \zeta)^{n+1}}$ is holomorphic in a domain containing C **except** at the point ζ .

Example 6.1. Let $f(z) = 1/z^2$ defined in $B(0, 1)$.

In the unit disk $B(0, 1)$ the function is defined and holomorphic everywhere except at the point 0. At 0 the denominator has a zero of multiplicity 2.

It turns out that this example perfectly illustrates the possibilities when we have a function $f(z)$ that is defined everywhere except possibly at **isolated** singularities. You may notice a similarity between these **singularities** and zeros. In the same way as we used power series to characterise the multiplicity of zeros we will use an extension of the Taylor series to define and characterise singularities.

Definition 6.2 – Laurent series

If $f(z)$ is holomorphic in the annulus $\{z \in \mathbb{C} : r < |z - a| < R\}$ then

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - a)^n, \quad a_n = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z - \zeta)^{n+1}} dz$$

and $C = \{z \in \mathbb{C} : |z - a| = \rho \text{ for some } r < \rho < R\}$.

The series is called a Laurent series, it is seemingly the same as a Taylor series except that the terms include negative indexes as well. In practise there are other ways of finding the series rather than using the integral.

Example 6.3. Let $f(z) = \frac{1}{z^2} \exp z$. Find the Laurent series for $f(z)$ centred at 0.

Here

$$\begin{aligned} f(z) &= \frac{1}{z^2} \left(1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots \right) \\ &= \frac{1}{z^2} + \frac{1}{z} + \frac{1}{2!} + \frac{z}{3!} \\ &= \sum_{n=-2}^{\infty} \frac{1}{(n+2)!} z^n. \end{aligned}$$

Note here we have that $f(z)$ is analytic everywhere in the complex plane except the point 0. So our annulus here is

$$\{z \in \mathbb{C} : 0 < |z| < \infty\}.$$

The point 0 is an example of an **isolated singularity**.

Definition 6.4 – Isolated singularities

Suppose that $f(z)$ is defined by its Laurent series in the annulus $\{z \in \mathbb{C} : 0 < |z - a| < R\}$ for $R \leq \infty$. Then a is an **isolated singularity** for $f(z)$.

The nature of the Laurent series determines the behaviour of the singularity. There are three distinct possibilities here.

1. **No negative powers.**

In this case we have that $f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n$. If we merely define $f(a) = a_0$ here then we can extend our function from the annulus to the disk $B(a, R)$. Furthermore the function is now holomorphic since it can be defined in terms of a power series. So in this case the point a is not a problem at all and can be dealt with quite easily, we call these kinds of points **removable singularities**.

2. **A finite number of negative powers.**

This time we have $a_n = 0$ for all $n < -m$ but $a_{-m} \neq 0$, for some $m > 0$. So we may write the Laurent series as

$$f(z) = \frac{a_{-m}}{(z-a)^m} + \frac{a_{-m+1}}{(z-a)^{m-1}} + \cdots + \frac{a_{-1}}{z-a} + a_0 + a_1(z-a) + \cdots.$$

Here our function tends to infinity as $z \rightarrow a$ and so it cannot be dealt with as we did with the last case. We say that $f(z)$ has a **pole of order m** at a .

Since it is not always possible or efficient to write a function as its Laurent series we prefer a different definition of a pole. Note that if $f(z)$ has a pole of order m then

$$(z-a)^m f(z) = a_{-m} + a_{-m+1}(z-a) + a_{-m+2}(z-a)^2 + \cdots.$$

The following definition of a pole is, then, equivalent.

Definition 6.5 – S

ppose that $f(z)$ is holomorphic in $\{z \in \mathbb{C} : 0 < |z-a| < R\}$ for some $R > 0$. Then $f(z)$ is said to have a pole of order m if $(z-a)^m f(z)$ has a removable singularity at a and m is the smallest such integer.

Example 6.6. Let $f(z) = \frac{\sin z}{z}$ in $\{z \in \mathbb{C} : 0 < |z| < \infty\}$.

This time

$$\begin{aligned} f(z) &= \frac{1}{z} \left(\sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!} \right) \\ &= \frac{1}{z} \left(z - \frac{z^3}{3!} + \frac{z^5}{5!} + \dots \right) \\ &= 1 - \frac{z^2}{3!} + \frac{z^4}{5!} + \dots. \end{aligned}$$

This function has a removable singularity at 0.

Example 6.7. Let $f(z) = \frac{\cos z}{z^2}$ in $\{z \in \mathbb{C} : 0 < |z| < \infty\}$.

Here

$$\begin{aligned} f(z) &= \frac{1}{z^2} \left(\sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!} \right) \\ &= \frac{1}{z^2} - \frac{1}{2!} + \frac{z^2}{4!} + \dots. \end{aligned}$$

This function has a pole of order 2 at 0.

Note for this function we could also note that

$$zf(z) \rightarrow \infty, \quad z \rightarrow 0$$

but $z^2 f(z) = \cos z$ has a removable singularity at 0. So $f(z)$ has a pole of order 2 at 0.

Example 6.8. Let $f(z) = \frac{1}{1+z^2}$ for $\{z \in \mathbb{C} : 0 < |z-i| < 2\}$.

Here it is less easy to define $f(z)$ as a Laurent series. Instead note that

$$(z-i)f(z) = (z-i) \frac{1}{(z+i)(z-i)} = \frac{1}{z+i} \rightarrow \frac{1}{2i}, \quad z \rightarrow i$$

and so $f(z)$ has a pole of order 1 at i .

3. Infinitely many negative terms.

This case is the worst in terms of behaviour of the function. We say that $f(z)$ has an **essential singularity** at a .

Example 6.9. Let $f(z) = \exp(1/z)$ for $\{z \in \mathbb{C} : 0 < |z| < \infty\}$.

This time

$$\begin{aligned} f(z) &= \sum_{n=0}^{\infty} \frac{(1/z)^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{1}{z^n}. \end{aligned}$$

The function $f(z)$ has an essential singularity at 0.

Definition 6.10 – Meromorphic functions

A function f defined in a domain Ω except possibly at isolated points where it has poles is called a **meromorphic function**.

A meromorphic function can always be written as the ratio of two holomorphic functions. Meaning meromorphic functions and holomorphic functions have the same relationship as rational numbers and integers.

6.2 The Residue Theorem

We would like to extend Cauchy's theorem to meromorphic functions. We can't expect to have an integral equating to 0 here but we might expect something that doesn't depend on the simple closed curve C . We will begin with the following lemma.

Lemma 6.11. Suppose that C is a simple closed curve, positively oriented. Suppose also that a is a point in the interior of C . Then

$$\int_C \frac{1}{(z-a)^n} dz = \begin{cases} 2\pi i, & n = 1; \\ 0, & n \neq 1. \end{cases}$$

Proof. By Lemma 5.1 we replace C with the boundary of a disk centred at a , $B(a, r)$ say, where $r > 0$ is chosen so that $B(a, r)$ is in the interior of C .

We parameterise $\partial B(a, r)$ by $p(t) = a + re^{it}$ for $0 \leq t \leq 2\pi$ then $p'(t) = rie^{it}$. Therefore,

$$\begin{aligned} \int_{\partial B(a, r)} \frac{1}{(z-a)^n} dz &= \int_0^{2\pi} \frac{1}{re^{int}} rie^{it} dt \\ &= \int_0^{2\pi} ie^{i(1-n)t} dt. \end{aligned}$$

If $n = 1$ then the integrand reduces to i and so the integral is $2\pi i$. If $n \neq 1$ then we get

$$\int_0^{2\pi} i e^{i(1-n)t} dt = \left[\frac{1}{1-n} e^{i(1-n)t} \right]_0^{2\pi} = \frac{1}{1-n} (1 - e^{2(1-n)\pi i}) = 0.$$

□

Consider a Laurent series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n.$$

Let C be a simple closed curve, positively oriented, with a in its interior (but no other singularities). The assuming that we can swap the summation and the integral around we should get

$$\begin{aligned} \int_C f(z) dz &= \int_C \left(\sum_{n=-\infty}^{\infty} a_n (z-a)^n \right) dz \\ &= \sum_{n=-\infty}^{\infty} \left(\int_C a_n (z-a)^n dz \right) \\ &= 2\pi i a_{-1} \end{aligned}$$

by Lemma 6.11.

The value a_{-1} is called the **Residue** of $f(z)$ at a . This equation gives us a method for evaluating the integral of a meromorphic function with a single pole.

Example 6.12. Let C be simple closed curve, positively oriented, with 0 in its interior. Find the integral along C of the function $f(z) = \frac{\cos z}{z^2}$.

We previously found that

$$f(z) = \frac{1}{z^2} - \frac{1}{2!} + \frac{z^2}{4!} + \dots$$

and so this time $a_{-1} = 0$ and so

$$\int_C f(z) dz = 2\pi i \times 0 = 0.$$

A better method for finding residues is needed in most practical situations. Suppose that $f(z)$ has a pole of order m at a then

$$f(z) = \sum_{n=-m}^{\infty} a_n (z-a)^n$$

and a_{-1} is the residue at a . Now

$$(z - a)^m f(z) = \sum_{k=0}^{\infty} a_{n-m}(z - a)^n = g(z)$$

and then a_{-1} is the $(m - 1)$ th coefficient of the power series for $g(z)$. So

$$a_{-1} = \frac{1}{(m - 1)!} g^{(m-1)}(a).$$

This is the basis for the following definition when a is a pole.

Definition 6.13 – Residues for poles

Suppose that f is a meromorphic function with a pole of order m at a . Then the residue at a is

$$\text{Res}_a f(z) = \frac{1}{(m - 1)!} \lim_{z \rightarrow a} \frac{d^{m-1}}{dz^{m-1}} (z - a)^m f(z).$$

We will illustrate the use of this formula with the following example.

Example 6.14. Let $f(z) = (z^4 - z^3 - 17z + 2)/(z - 1)^3$.

Here we have a pole at 1 of order 3 (be careful here! You should check that $(z - 1)$ is not a factor of the numerator).

$$\begin{aligned} \text{Res}_1 f(z) &= \frac{1}{2!} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} (z - 1)^3 f(z) \\ &= \frac{1}{2} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} (z^4 - z^3 - 17z + 2) \\ &= \frac{1}{2} \lim_{z \rightarrow 1} (12z^2 - 6z) \\ &= \frac{1}{2} \times 6 = 3. \end{aligned}$$

Therefore if C is any simple closed curve, positively oriented, and containing 1 in its interior, then

$$\int_C f(z) dz = 2\pi i \times 3 = 6\pi i.$$

Some functions have many poles of different orders. We can extend the method above to this scenario.




Figure 6.1: Proof of Cauchy's Residue Theorem

Theorem 6.15 – Cauchy's Residue Theorem

Suppose f is meromorphic function in a domain Ω . Let C be a simple closed curve in Ω . If the poles of f in the interior of C are written z_1, z_2, \dots, z_n then

$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}_{z_k} f(z).$$

Proof. We will only sketch the proof. Let r_k be chosen small enough so that the disks $B(z_k, r_k)$ do not overlap and are small enough to be in the interior of C . See Figure 6.1. Let $C_k = \partial B(z_k, r_k)$ then it is possible to show, extending Lemma 5.1, that

$$\int_C f(z) dz = \sum_{k=1}^n \int_{C_k} f(z) dz.$$

Then

$$\int_{C_k} f(z) dz = 2\pi i \text{Res}_{z_k} f(z)$$

and the theorem is proved. □

6.3 Simple poles

A pole of order 1 is called a **simple pole**. For simple poles the calculation of the residue is more straightforward. Here we have

$$\text{Res}_a f(z) = \lim_{z \rightarrow a} (z - a)f(z).$$

Example 6.16. Find the residue at $2i$ of $f(z) = (z^2 - 4)/(z^2 + 4)$.

This is a simple pole so

$$\begin{aligned} \text{Res}_{2i} f(z) &= \lim_{z \rightarrow 2i} (z - 2i) \frac{(z^2 - 4)}{(z^2 + 4)} \\ &= \lim_{z \rightarrow 2i} (z - 2i) \frac{(z^2 - 4)}{(z + 2i)(z - 2i)} \\ &= \lim_{z \rightarrow 2i} \frac{(z^2 - 4)}{(z + 2i)} \\ &= \frac{(2i)^2 - 4}{2i + 2i} \\ &= \frac{-8}{4i} = 2i. \end{aligned}$$

Example 6.17. Find the residues of $f(z) = 1/(1 + z^3)$ at each of its poles.

The poles are the cube roots of -1 which are

$$\omega_k = e^{i(2\pi k + \pi/2)/3}, \quad k = 0, 1, 2.$$

We will factorise $(1 + z^3) = (z - \omega_0)(z - \omega_1)(z - \omega_2)$ then

$$\text{Res}_{\omega_k} f(z) = \lim_{z \rightarrow \omega_k} (z - \omega_k) \prod_{i=0}^2 \frac{1}{z - \omega_i} = \prod_{\substack{i=1 \\ i \neq k}}^2 \frac{1}{\omega_k - \omega_i}.$$

An useful tool for calculating residues, and limits of ratios in general, is l'Hopital's rule. This is a standard tool in calculus but has a particularly nice statement in complex analysis.

Theorem 6.18

Suppose that $f(z)$ and $g(z)$ are holomorphic in a disk $B(a, r)$ and a is a zero of multiplicity m for both $f(z)$ and $g(z)$. Then

$$\lim_{z \rightarrow a} \frac{f(z)}{g(z)} = \frac{f^{(m)}(a)}{g^{(m)}(a)}.$$

Proof. By Corollary 5.14 we may write

$$f(z) = (z - a)^m \sum_{n=0}^{\infty} a_{n+m}(z - a)^n, \quad g(z) = (z - a)^m \sum_{n=0}^{\infty} b_{n+m}(z - a)^n.$$

By Corollary 2.23,

$$a_k = \frac{f^{(k)}(a)}{k!}, \quad b_k = \frac{g^{(k)}(a)}{k!}, \quad k = m, m + 1, \dots$$

Therefore

$$\begin{aligned} \lim_{z \rightarrow a} \frac{f(z)}{g(z)} &= \lim_{z \rightarrow a} \frac{(z - a)^m \sum_{n=0}^{\infty} a_{n+m}(z - a)^n}{(z - a)^m \sum_{n=0}^{\infty} b_{n+m}(z - a)^n} \\ &= \lim_{z \rightarrow a} \frac{\sum_{n=0}^{\infty} a_{n+m}(z - a)^n}{\sum_{n=0}^{\infty} b_{n+m}(z - a)^n} \\ &= \frac{a_m}{b_m} = \frac{f^{(m)}(a)/m!}{g^{(m)}(a)/m!} \\ &= \frac{f^{(m)}(a)}{g^{(m)}(a)}. \end{aligned}$$

□

Example 6.19. Consider again the residues of $f(z) = 1/(1 + z^3)$ at each of its poles.

We'll use l'Hopital's rule this time. First

$$\begin{aligned} \operatorname{Res}_{\omega_k} f(z) &= \lim_{z \rightarrow \omega_k} (z - \omega_k) f(z) \\ &= \lim_{z \rightarrow \omega_k} \frac{z - \omega_k}{1 + z^3} \\ &= \frac{1}{3\omega_k^2}. \end{aligned}$$

We can simplify this even further by noticing that $-1 = \omega_k^3 = \omega_k^2 \omega_k$ so that

$$\operatorname{Res}_{\omega_k} f(z) = -\frac{1}{3} \omega_k.$$




Figure 6.2: The closed contour in Example 6.20

6.4 Applications of the Residue Theorem to real integrals

One of the most commonly seen applications of the residue theorem is in the evaluation of previously difficult real integrals. The best way to learn about this is to see an example.

Example 6.20. Find the value of the integral

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx.$$

We will integrate $f(z) = 1/(1+z^2)$ along the curve consisting of the semi-circle of radius $R > 0$, centred at 0 in the upper half-plane, and the line-segment from $-R$ to R , see Figure 6.2. Call this curve C . As long as $R > 1$ we can see from the diagram that C contains only the simple pole at i . Therefore by the residue theorem

$$\begin{aligned} \int_C \frac{1}{1+z^2} dz &= 2\pi i \operatorname{Res}_i f(z) \\ &= 2\pi i \lim_{z \rightarrow i} \frac{z-i}{1+z^2} \\ &= 2\pi i \lim_{z \rightarrow i} \frac{1}{2z} = 2\pi i \frac{1}{2i} \\ &= \pi. \end{aligned}$$

Now this integral does not depend on R so we can choose R to be anything at all. In particular we could let $R \rightarrow \infty$. The reason we might want to do this is that the integral

along the curve C can be split between the integral along the line segment I from $-R$ to R , and the semi-circle S . Then letting $R \rightarrow \infty$

$$\int_I \frac{1}{1+z^2} dz = \int_{-R}^R \frac{1}{1+x^2} dx \rightarrow \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx.$$

We'll need to be careful about this convergence but for the moment let's just accept that it all makes sense.

We will show that

$$\lim_{R \rightarrow \infty} \int_S \frac{1}{1+z^2} dz = 0.$$

As we have done a number of times we will do this using the triangle inequality.

$$\begin{aligned} \left| \int_S \frac{1}{1+z^2} dz \right| &\leq \int_S \frac{1}{|1+z^2|} |dz| \\ &\leq \int_S \frac{1}{|z|^2 - 1} |dz|, \end{aligned}$$

since $|z^2 + 1| \geq |z|^2 - 1$ by the triangle inequality,

$$\begin{aligned} &= \frac{1}{R^2 - 1} \int_S |dz| = \frac{1}{R^2 - 1} \text{length}(S) \\ &= \frac{\pi R}{R^2 - 1}. \end{aligned}$$

Therefore since the last expression tends to 0 as $R \rightarrow \infty$ we have that

$$\lim_{R \rightarrow \infty} \int_S \frac{1}{1+z^2} dz = 0$$

as required.

Collecting this together we have that for all $R > 1$

$$\pi = \int_C \frac{1}{1+z^2} = \int_{-R}^R \frac{1}{1+x^2} dx + \int_S \frac{1}{1+z^2} dz.$$

If we now take the limit as $R \rightarrow \infty$ we get

$$\pi = \lim_{R \rightarrow \infty} \pi = \lim_{R \rightarrow \infty} \left(\int_{-R}^R \frac{1}{1+x^2} dx + \int_S \frac{1}{1+z^2} dz \right) = \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx.$$

So

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \pi.$$




Figure 6.3: The curve in Example 6.21.

6.4.1 The Cauchy principal value

For this all to make sense we need to understand what we mean by integrals over the entire real line. We will explain this using the integral

$$\int_{-\infty}^{\infty} x dx.$$

First we might calculate,

$$\int_R^S x^3 dx = \frac{1}{2}(S^2 - R^2),$$

and let $S \rightarrow \infty$ and $R \rightarrow -\infty$ independently. This leads to a problem since the right-hand side won't converge if we let $S \rightarrow \infty$ first. What if S depends somehow on R , i.e. S is a function of R , for example suppose $S = \sqrt{R^2 + 4}$ then

$$\frac{1}{2}(S^2 - R^2) = 1.$$

So,

$$\lim_{R \rightarrow -\infty, S \rightarrow \infty} \int_R^S x^3 dx = 1.$$

You should check that by replacing 4 in the equation of S in terms of R you can make this limit any non-negative number at all. You might want to try to think how you could make it any negative value at all.

To avoid this problem we define the **Cauchy principal value** of an improper integral like this to mean the following:

$$PV \int_{-\infty}^{\infty} f(x)dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x)dx.$$

We often drop the “PV” here with the understanding that we’re always talking about the Cauchy principal value. Notice how this lines up nicely with the way it’s understood in the last example.

As a final remark here note that this is all academic if the integrand is a function in $L^1(\mathbb{R})$, the Lebesgue space, because the dominated convergence theorem guarantees no matter R and S tend to minus or plus infinity the integral converges.

Now let’s have another example.

Example 6.21. *Using the residue theorem find*

$$\int_0^{\infty} \frac{x^2}{1+x^4} dx.$$

This time we are integrating from 0 to infinity, however notice that $x^2/(1+x^4)$ is an even function and so

$$\int_0^{\infty} \frac{x^2}{1+x^4} dx = \frac{1}{2} \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx.$$

We can therefore find this integral by considering a curve like the one we had in the last example, see Figure 6.3.

The poles of $z^2/(1+z^4)$ are the fourth roots of -1 which are

$$\omega_k = e^{i(2\pi k + \pi)/4}, \quad k = 0, 1, 2, 3,$$

also drawn in Figure 6.3. Here only

$$\omega_0 = \frac{1}{\sqrt{2}}(1+i), \quad \omega_1 = \frac{1}{\sqrt{2}}(-1+i)$$

are contained in the interior of the curve C . Therefore by the residue theorem

$$\int_C \frac{z^2}{1+z^4} dz = 2\pi i \left(\text{Res}_{\omega_0} f(z) + \text{Res}_{\omega_1} f(z) \right).$$

Both these are simple poles so we can use l'Hopital's rule to find them.

$$\begin{aligned}
 \operatorname{Res}_{\omega_0} f(z) &= \lim_{z \rightarrow \omega_0} (z - \omega_0) f(z) \\
 &= \lim_{z \rightarrow \omega_0} \frac{z^2(z - \omega_0)}{1 + z^4} \\
 &= \frac{3\omega_0^2 - \omega_0(2\omega_0)}{4\omega_0^3} \\
 &= \frac{\omega_0^2}{4\omega_0^3} \\
 &= \frac{1}{4\omega_0} = \frac{\sqrt{2}}{4(1+i)} \\
 &= \frac{\sqrt{2}}{8}(1-i).
 \end{aligned}$$

Similarly

$$\begin{aligned}
 \operatorname{Res}_{\omega_1} f(z) &= \lim_{z \rightarrow \omega_1} (z - \omega_1) f(z) \\
 &= \frac{3\omega_1^2 - \omega_1(2\omega_1)}{4\omega_1^3} \\
 &= \frac{1}{4\omega_1} = \frac{\sqrt{2}}{4(-1+i)} \\
 &= \frac{\sqrt{2}}{8}(-1-i).
 \end{aligned}$$

Therefore

$$\operatorname{Res}_{\omega_0} f(z) + \operatorname{Res}_{\omega_1} f(z) = \frac{\sqrt{2}}{8}(1-i) + \frac{\sqrt{2}}{8}(-1-i) = -i\frac{\sqrt{2}}{4}.$$

It follows that

$$\int_C \frac{z^2}{1+z^4} dz = 2\pi i \left(-i\frac{\sqrt{2}}{4} \right) = \pi \frac{\sqrt{2}}{2}.$$

As before we now have to show that

$$\lim_{R \rightarrow \infty} \int_C \frac{z^2}{1+z^4} dz = \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx.$$

First if I is the line segment joining $-R$ to R then

$$\lim_{R \rightarrow \infty} \int_I f(z) dz = \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx.$$

Second let S be the semi-circular part of C . Then

$$\begin{aligned} \left| \int_S f(z) dz \right| &\leq \int_S \left| \frac{z^2}{1+z^4} \right| |dz| \\ &\leq \int_S \frac{|z|^2}{|z|^4 - 1} |dz| \\ &= \frac{R^2}{R^4 - 1} \pi R \\ &= \frac{\pi R^3}{R^4 - 1}. \end{aligned}$$

And it follows that

$$\lim_{R \rightarrow \infty} \int_S f(z) dz = 0.$$

So as before we have

$$\frac{\sqrt{2}}{2} \pi = \lim_{R \rightarrow \infty} \int_C \frac{z^2}{1+z^4} dz = \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx.$$

Finally we have

$$\int_0^{\infty} \frac{x^2}{1+x^4} dx = \frac{1}{2} \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx = \frac{\sqrt{2}}{4} \pi.$$

This method can be used for finding real integrals of the form

$$\int_{-\infty}^{\infty} \frac{p(x)}{q(x)} dx$$

when p and q are polynomials with $\deg p \leq \deg q - 2$ (i.e. the degree of p is at least two less than the degree of q) as long as this integral exists and $q(x) \neq 0$ for any $x \in \mathbb{R}$. The strategy is to integrate over a closed curve like the one in Figure 6.2, calculate the residues and show that the integral along the semi-circle tends to 0 as $R \rightarrow \infty$.

Residues can also be used to calculate other types of real integrals. In general the choice of closed curve needs to be chosen wisely so that the 'complex part' of the integral tends to 0. There are a number of general methods for doing this.

6.4.2 Integrals involving trigonometric functions

We will consider, in this section, integrals of the form

$$\int_0^{2\pi} G(\cos t, \sin t) dt$$

We will use the standard relationships

$$\cos t = \frac{e^{it} + e^{-it}}{2}, \quad \sin t = \frac{e^{it} - e^{-it}}{2i}$$

to rewrite integrals of this form as an integral over a circular path. We already know that we can parameterise these integrals using the parameterisation

$$p(t) = ae^{it}, \quad 0 \leq t \leq 2\pi,$$

so our aim is to rewrite the integral in the form

$$\int_0^{2\pi} f(p(t))p'(t)dt = \int_C f(z)dz.$$

We will demonstrate the technique with the following example.

Example 6.22. Evaluate the integral

$$\int_0^{2\pi} \frac{1}{2 + \sin t} dt.$$

Let $p(t) = e^{it}$ then we have

$$\cos t = \frac{e^{it} + e^{-it}}{2} = \frac{p(t) + 1/p(t)}{2},$$

and

$$\sin t = \frac{e^{it} - e^{-it}}{2i} = \frac{p(t) - 1/p(t)}{2i}.$$

Here we have used the fact that $e^{-it} = 1/e^{it} = 1/p(t)$.

The denominator in the integrand is

$$\begin{aligned} 2 + \sin t &= 2 + \frac{p(t) - 1/p(t)}{2i} \\ &= \frac{1}{p(t)} \left(2p(t) + \frac{1}{2i}(p(t)^2 - 1) \right). \end{aligned}$$

Now $p'(t) = ie^{it} = ip(t)$ so that $p(t) = p'(t)/i$ and so we can rewrite the integrand above as

$$\begin{aligned}\frac{1}{2 + \sin t} &= \frac{p(t)}{\left(2p(t) + \frac{1}{2i}(p(t)^2 - 1)\right)} \\ &= \frac{p'(t)}{i\left(2p(t) + \frac{1}{2i}(p(t)^2 - 1)\right)} \\ &= \frac{2p'(t)}{p(t)^2 + 4ip(t) - 1}.\end{aligned}$$

The reason we rearrange the integral as we have done is so that we may now write this as

$$\int_0^{2\pi} \frac{2p'(t)}{p(t)^2 + 4ip(t) - 1} dt = \int_C \frac{2}{z^2 + 4iz - 1} dz$$

where C is the boundary of the unit disk $B(0, 1)$. We can use the quadratic formula (which holds for complex quadratics as well) to find that the poles are where the denominator has zeros. They are simple poles at

$$a = (-2 + \sqrt{3})i, \quad b = (-2 - \sqrt{3})i.$$

Only the first is in $B(0, 1)$ and its residue is

$$\begin{aligned}\text{Res}_a \frac{2}{z^2 + 4iz - 1} &= \text{Res}_a \frac{2}{(z - a)(z - b)} \\ &= \frac{2}{a - b} = \frac{2}{(-2 + \sqrt{3})i - (-2 - \sqrt{3})i} \\ &= \frac{1}{\sqrt{3}i}.\end{aligned}$$

It follows by the Residue Theorem that

$$\int_0^{2\pi} \frac{1}{2 + \sin t} dt = \int_C \frac{2}{z^2 + 4iz - 1} dz = 2\pi i \frac{1}{\sqrt{3}i} = \frac{2\pi}{\sqrt{3}}.$$

Again the technique is fairly straightforward – even if it is quite length. And the important thing to note is that it allows us to find the integrals of many functions that we previously couldn't.

Things to know

- The definition of the Laurent series and how to find it for a standard example;
- The definition of an isolated singularity;
- The classification of isolated singularities as removable singularities, poles and essential singularities, and how to classify the singularities of a given function;
- The definition of a residue and how to find it for a given function, especially including applying l'Hopital's rule for simple poles;
- The statement and proof of Lemma 6.11;
- The statement of the Residue Theorem and how to use it to find the integral of a function;
- The application of the Residue Theorem to find integrals of real-valued integrals.

Further reading

The residue theorem is important in complex analysis – not just as a way of calculating real integrals. Other applications are the proof that $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$. This and other applications can be found in [3, Chapter 24]. Some of the material in this chapter are modelled on Ahlfors treatment, [1, Chapter 4, §5.3].

Problems

- Find and classify the poles of the following functions and calculate the residue of each of them.

(a) $\frac{z^2 + 2}{z - 1}$

(b) $\left(\frac{z}{2z + 1}\right)^3$

(c) $\frac{\exp z}{z^2 + \pi^2}$

(d) $\frac{a - z}{1 - az}$ where $a \in \mathbb{R}$ and $a \neq 0$

- Find and classify all the poles of $\frac{\exp z + 1}{\exp z - 1}$. For each of them find its residue.

- Let C be the simple closed curve consisting of the line segment from -2 to 2 and then the semi-circle in the upper half-plane centred at 0 with radius 2 . Use the Residue Theorem to find

$$\int_C \frac{1}{1 + z^6} dz.$$

- Suppose $f(z)$ and $g(z)$ both have a simple pole at a . Prove that $f(z)/g(z)$ has a removable singularity at a and hence can be extended to a holomorphic function in a neighbourhood of a .
- If $f(z)$ is holomorphic in Ω except at poles a_i , $i = 1, 2, \dots, N$ of order m_i . Show that

$$f(z) = g(z) \prod_{i=1}^N \frac{1}{(z - a_i)^{m_i}}$$

where $g(z)$ can be extended to a holomorphic function in Ω .


- Use residues to find

(a) $\int_0^\infty \frac{1}{1 + x^4} dx$. Ans: $\pi/(2\sqrt{2})$.

(b) $\int_0^\infty \frac{x^2}{(x^2 + 1)(x^2 + 4)} dx$. Ans: $\pi/6$.

7. Use the curve shown below, where $R > 1$, to show that

$$\int_0^{\infty} \frac{1}{1+x^3} dx = \frac{2\pi}{3\sqrt{3}}.$$



Chapter 7

Mapping Properties

“The beauty of mathematics only shows itself to more patient followers.”
Maryam Mirzakhani

7.1 The maximum modulus

In calculus and real-analysis in general it is often useful to consider the problem of finding the maximum of a function in a domain. In complex analysis of course you can't maximise a function because \mathbb{C} doesn't have an ordering, i.e. there is no notion of one complex number being larger than another. However we can maximise the real-valued function $|f(z)|$ in a domain and this is the problem we will look at in this section.

You've already seen one example of this when we looked at Liouville's theorem, Theorem 5.16. We can think of this theorem as saying that an entire function cannot map \mathbb{C} into a bounded domain Ω .

We will begin with the following result that may take some time to digest.

Theorem 7.1 – Maximum modulus theorem

Suppose that $f(z)$ is a non-constant holomorphic function in a domain Ω . Then $|f(z)|$ does not attain its supremum in Ω .

If we let $M = \sup_{z \in \Omega} |f(z)|$ then this means that there is no point in Ω where $|f(z)| = M$. Remember that Ω is an open set so that this makes sense. To prove this important theorem we need the following.

Lemma 7.2 (Gauss' mean value theorem). *Suppose $f(z)$ is holomorphic in a domain containing the closed disk $\overline{B(a, r)} = \{z \in \mathbb{C} : |z - a| \leq r\}$. Then*

$$f(a) = \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{it}) dt.$$

This looks suspiciously like Cauchy's integral formula. That is because it's simply a restatement. It should be noted then that this property holds for a much wider class of functions called **harmonic** functions and there the proof is more difficult.

Proof. We let $C = \{z \in \mathbb{C} : |z - a| = r\}$. We parameterise this by

$$p(t) = a + re^{it}, \quad 0 \leq t \leq 2\pi.$$

Then by Cauchy's integral formula

$$\begin{aligned} f(a) &= \frac{1}{2\pi i} \int_C \frac{f(z)}{z - a} dz \\ &= \frac{1}{2\pi i} \int_0^{2\pi} f(p(t)) p'(t) dt = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(a + re^{it})}{re^{it}} ire^{it} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{it}) dt \end{aligned}$$

□

We also need the following uniqueness result.

Lemma 7.3. *Suppose that $f(z)$ is holomorphic on a domain containing $B(a, r)$ and $|f(z)| = m$ for all $z \in B(a, r)$. Then $f(z)$ is constant throughout Ω .*

Proof. First if $m = 0$ then by Theorem 5.11 $f(z) = 0$ for all $z \in \Omega$. If $m > 0$ then since $f(z)$ is continuous there is a $\rho > 0$ such that $f(z) \neq 0$ for all $z \in B(a, \rho)$. So by Proposition 3.3 there is a branch of the logarithm for which $g(z) = \log_\alpha f(z)$ is holomorphic in $B(a, \rho)$. Now

$$g(z) = \log |f(z)| + i \arg_\alpha z$$

and this function has constant real part. Therefore $g(z)$ is constant by the Cauchy-Riemann equations. So $f(z) = \exp g(z)$ is also constant. □

Proof of Theorem 7.1. We prove this by contradiction. Suppose that there is a point $a \in \Omega$ such that $|f(a)| \geq |f(z)|$ for all $z \in \Omega$. Let $R > 0$ be chosen so that $\overline{B(a, R)}$ is contained in Ω . For each $0 < r \leq R$ we have, by Lemma 7.2,

$$f(a) = \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{it}) dt.$$

Now $|f(z)|$ is not constant in $B(a, r)$ by Lemma 7.3 so we can find a $\rho < R$ and a $\phi \in [0, 2\pi)$ such that $|f(a + \rho e^{i\phi})| < |f(a)|$. Since f is continuous, there is an interval I with $\phi \in I \subset [0, 2\pi)$ such that

$$|f(a + \rho e^{it})| < |f(a)|, \quad t \in I.$$

We therefore have

$$\begin{aligned} |f(a)| &\leq \frac{1}{2\pi} \int_0^{2\pi} |f(a + \rho e^{it})| dt = \frac{1}{2\pi} \left(\int_I |f(a + \rho e^{it})| dt + \int_{I^c} |f(a + \rho e^{it})| dt \right) \\ &< \frac{1}{2\pi} \int_I |f(a)| dt + \frac{1}{2\pi} \int_{I^c} |f(a)| dt, \quad \text{where } I^c = [0, 2\pi) \setminus I, \\ &= \frac{1}{2\pi} \int_0^{2\pi} |f(a)| dt = |f(a)|. \end{aligned}$$

This is of course a contradiction and so the theorem is proved. \square

This is an important property of holomorphic functions. One consequence is that if $f(z)$ is holomorphic in a domain containing $\overline{B(0, r)}$ for all $r < R$ for some $R > 0$ then the maximum value of $|f(z)|$ in the disk $B(0, r)$ is taken at a point on the circle $\{z \in \mathbb{C} : |z| = r\}$. This justifies the following definition.

Definition 7.4 – The maximum modulus function

Suppose that $f(z)$ is a complex valued function. The **maximum modulus function** is

$$M(r; f) = \max_{\theta \in [0, 2\pi)} |f(re^{i\theta})|.$$

By Theorem 7.1, if f is holomorphic in a domain containing $\overline{B(a, r)}$, the value of $M(r; f)$ is the maximum value taken by $|f(z)|$ in the whole of $B(0, r)$. Therefore we have that $M(r; f)$ is a strictly increasing function of r whenever f is a non-constant holomorphic function.

For non-constant entire functions we know by Liouville's theorem that $M(r; f) \rightarrow \infty$ as $r \rightarrow \infty$. For functions only holomorphic on a smaller domain we have the following.

Theorem 7.5 – Schwarz’s lemma

Suppose that f is holomorphic on $B(0, 1)$, $|f(z)| \leq 1$ and $f(0) = 0$ then

$$|f(z)| \leq |z|, \quad \text{for all } z \in B(0, 1).$$

If equality occurs at **any** value of z then $f(z) = \lambda z$ for some λ with $|\lambda| = 1$.

Proof. Since $f(0) = 0$ we can write $f(z) = \sum_{n=m}^{\infty} a_n z^n$ where $m > 0$ is the multiplicity of the zero at 0. In particular we can write

$$g(z) = \frac{f(z)}{z}$$

and $g(z)$ is holomorphic in $B(0, 1)$.

Now if $|z| \leq r < 1$ then by the maximum modulus theorem, Theorem 7.1,

$$|g(z)| \leq \max_{\theta \in [0, 2\pi)} \frac{|f(re^{i\theta})|}{|re^{i\theta}|} \leq \frac{1}{r}.$$

This is true for all $r \in (0, 1)$ and so is also true if we let $r \rightarrow 1$. Therefore for all $z \in B(0, 1)$, $|g(z)| \leq 1$. Hence $|f(z)| \leq |z|$ for all $z \in B(0, 1)$.

To prove the second part suppose that $|f(z)| = |z|$ for some $z \in B(0, 1)$, then $|g(z)| = 1$ for some $z \in B(0, 1)$. But this is the maximum value taken by $|g(z)|$ and so by the Maximum-Modulus theorem, Theorem 7.1, the function g must be constant. It follows that $f(z)/z = \lambda$ where $|\lambda| = 1$ as required. \square

Note this means that for f ,

$$M(r; f) \leq r, \quad 0 < r < 1.$$

Thinking of f as a mapping, Schwarz’s lemma says that if f maps $B(0, 1)$ into $B(0, 1)$ and $f(0) = 0$ then f maps $B(0, r)$ into $B(0, r)$ for any $0 < r < 1$. So f can be thought of as *contracting* the ball $B(0, r)$.

7.2 The argument principle

Suppose that C is a simple closed curve of finite length then a holomorphic function maps C onto another closed curve D of finite length. However it is no longer necessarily

true that D is simple, it may wind its way around any number of times. For example the function $f(z) = z^2$ maps the circle $\{z \in \mathbb{C} : |z| = 1\}$ onto itself but the resulting curve is covered twice. In this section we prove the following result.

Theorem 7.6 – The argument principle

Let $f(z)$ be a function holomorphic in a domain Ω except possibly at a set of isolated singularities. Suppose C is a simple closed curve of finite length such that C and its interior are contained in Ω . Then

$$\frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz = N - P$$

where N is the number of zeros and P the number of poles in the interior of C counted according to their multiplicity.

When we say that zeros or poles are *counted according to their multiplicity* in the statement above this means that if a zero, for example, has multiplicity 3 then it contributes 3 to the value of N .

Note: the requirement that the interior of C also be in Ω ensures that the function $f(z)$ is holomorphic in this interior. Consider for example the annulus $\Omega = \{1 < |z| < 3\}$. The function $f(z) = 1/z$ is holomorphic throughout Ω but we can't invoke Cauchy's theorem to find the integral along the path $\{|z| = 2\}$ since Ω is not simply connected.

Proof. Suppose a is a zero of order m . Then $f(z) = (z - a)^m g(z)$ where $g(z)$ has a removable singularity at a . Now

$$\frac{f'(z)}{f(z)} = \frac{m(z - a)^{m-1} g(z) + (z - a)^m g'(z)}{(z - a)^m g(z)} = \frac{m}{z - a} + \frac{g'(z)}{g(z)}.$$

Now if b is a pole of order n then $g(z) = (z - b)^{-n} h(z)$ and we may perform the same calculation to show that

$$\frac{f'(z)}{f(z)} = \frac{m}{z - a} + \frac{-n}{z - b} + \frac{h'(z)}{h(z)}.$$

We can apply this calculation to all zeros and poles of $f(z)$. If z_1, z_2, \dots, z_k are the zeros of $f(z)$ interior to C and a_1, a_2, \dots, a_l are the poles then

$$\frac{f'(z)}{f(z)} = \sum_{i=1}^k \frac{m_i}{z - z_k} + \sum_{j=1}^l \frac{-p_j}{z - a_j} + \frac{G'(z)}{G(z)}.$$

Here $G(z)$ has no zeros or poles, m_i is the multiplicity of the zero z_i , and p_j the order of the pole at a_j .

Hence $f'(z)/f(z)$ has residues m_i at each z_i , and $-p_j$ at each a_j . Since the interior of C is contained in Ω the Residue Theorem implies that

$$\frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz = \sum_{i=1}^k m_i - \sum_{j=1}^l p_j = N - P.$$

□




Figure 7.1: The mapping of the circle $\partial B(0, 2)$ under the mapping $z \mapsto z^2$

The reason this is called the argument principle is the interpretation of the integral. The integrand $f'(z)/f(z)$ is the logarithmic derivative of $f(z)$, i.e. $\frac{d}{dz} \log f(z)$. Suppose that $f(z)$ is holomorphic and non-zero for all z in the interior of C . Then there is a branch of the logarithm for which $\log_\alpha f(z)$ is defined and is holomorphic by Proposition 3.3. If C is parameterised by $p(t)$ for $a \leq t \leq b$ then

$$\begin{aligned} \int_C \frac{f'(z)}{f(z)} dz &= \int_C \frac{d}{dz} \log_\alpha f(z) dz \\ &= \int_a^b p'(t) \frac{d}{dz} \log f(p(t)) dt \\ &= \int_a^b \frac{d}{dt} \log f(p(t)) dt \\ &= \left[\log |f(p(t))| + i \arg_\alpha f(p(t)) \right]_a^b. \end{aligned}$$

The curve C is closed so $|f(p(a))| = |f(p(b))|$. Thus the integral above is $i(\arg f(p(b)) - \arg f(p(a)))$. This is $2\pi i$ times the **winding number** of the curve $f(C)$. The winding number of $f(C)$ is the number of times a curve winds around the origin before closing. In Figure 7.1 the image of the circle on the left is the curve on the right which has winding number 2. This corresponds to the fact that the function $f(z) = z^2$ has two zeros. The winding number of a closed curve D is denoted by $n(D, 0)$ where we highlight 0 as the point around which D is winding. The argument principle is thus

$$n(f(C), 0) = N - P.$$

A classic application of the argument principle is the following important theorem.

Theorem 7.7 – Rouché's theorem

Let C be a simple closed curve in a domain Ω such that the interior of C is also in Ω . Suppose that $f(z)$ and $g(z)$ are holomorphic in Ω and for all $z \in C$

$$|f(z) - g(z)| < |f(z)|.$$

Then $f(z)$ and $g(z)$ have the same number of zeros.

I will follow the proof given in [1, page 153] since it is an elegant, geometric proof. It is based on the observation that if D is a simple closed curve contained in a disk that does not contain the origin then $n(D, 0) = 0$.

Proof. By assumption $f(z)$ and $g(z)$ are non-zero on C . Furthermore

$$\left| \frac{g(z)}{f(z)} - 1 \right| < 1, \quad z \in C.$$

Therefore $F(z) = g(z)/f(z)$ maps C into the disk $B(1, 1)$. It follows that $n(F(C), 0) = 0$. Now

$$\frac{F'(z)}{F(z)} = \frac{g'(z)}{g(z)} - \frac{f'(z)}{f(z)}$$

so that

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_C \frac{F'(z)}{F(z)} dz = \frac{1}{2\pi i} \int_C \frac{g'(z)}{g(z)} dz - \frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz \\ &= N_g - N_f \end{aligned}$$

by the argument principle, where N_f and N_g are the number of zeros of $f(z)$ and $g(z)$. □

A continuous mapping does not necessarily map open sets to open sets. For example the real function $f(x) = x^2$ maps $(-1, 1)$ to $[0, 1)$. However for holomorphic functions this is an important and useful property. We can prove this as an application of Rouché's theorem.

Theorem 7.8 – The open mapping theorem

A non-constant holomorphic function maps open sets to open sets.

Such a function is called an **open mapping** – hence the name of the theorem. Recall that an open set is a set in which every point contains an open neighbourhood. So to prove that $f(A)$ is open when A is open we need to show that f maps open neighbourhoods of each point into another open neighbourhood.

Proof. Suppose $f \in H(\Omega)$ for some domain Ω and let $z_0 \in \Omega$ be arbitrary. By Corollary 5.14 let $\delta > 0$ be chosen so that $B(z_0, \delta) \subset \Omega$ and there is only one zero of $f(z) - f(z_0)$ in $\overline{B(z_0, \delta)}$, namely z_0 itself.

Let ϵ be the minimum value of $|f(z) - f(z_0)|$ on the circle $\partial B(z_0, \delta) = \{z \in \mathbb{C} : |z - z_0| = \delta\}$. We will show that the image of $B(z_0, \delta)$ contains the disk $B(f(z_0), \epsilon)$.

Choose $w \in B(f(z_0), \epsilon)$. Then on $\partial B(z_0, \delta)$ we have

$$|f(z_0) - w| < \epsilon \leq |f(z) - f(z_0)|.$$

Therefore by Rouché's theorem

$$(f(z) - f(z_0)) + (f(z_0) - w) = f(z) - w$$

has the same number of zeros in $B(z_0, \delta)$ as $f(z) - f(z_0)$ does. Since $f(z) - f(z_0)$ has one zero (at $z = z_0$), the function $f(z) - w$ also has a zero. That is, $f(z) = w$ for some $z \in B(z_0, \delta)$. Since w was arbitrary the image of $B(z_0, \delta)$ must contain all the points of $B(f(z_0), \epsilon)$.

We have shown that every point in the image of Ω contains a neighbourhood, the image $f(\Omega)$ is also open. \square

Things to know

- The different behaviour of standard functions in terms of their mapping properties;

- The statement but not the proof of the Maximum Modulus Theorem and the definition of $M(r; f)$;
- The statement and proof of Gauss' mean value theorem, Lemma 7.2;
- The statement and proof of Schwarz's lemma;
- The statement of the argument principle in terms of winding numbers: $n(f(C), 0) = N - P$;
- The statement of Rouché's Theorem and the Open Mapping Theorem.

Further reading

The topics in this chapter head towards more advanced concepts in complex analysis. About the behaviour of holomorphic functions in a domain or locally. This is one main branch of higher level complex analysis and is worth reading about. The best introduction to these advanced topics is [1, Chapters 5-8]. You can also see the entire book of Hille, [5].

Problems

1. If $f(z)$ is entire show that $\lim_{r \rightarrow \infty} M(r; f) = \infty$
2. Suppose $f(z) = \frac{1+z}{1-z}$. Find $M(r; f)$ for $0 < r < 1$.
3. Use Rouché's Theorem to locate zeros of $f(z) = z^7 - 4z^3 + z - 1$ as follows:
 - (a) Define $g(z) = -4z^3$, so that $f(z) - g(z) = z^7 + z - 1$. Show that if $|z| = 1$ then

$$|f(z) - g(z)| \leq 3;$$
 - (b) deduce that for $|z| = 1$

$$|f(z) - g(z)| < |g(z)|;$$
 - (c) apply Rouché's theorem to show that $p(z)$ has exactly three zeros in $B(0, 1)$, counting multiplicity.
4. Find the number of zeros, counting multiplicities, of $z^6 - 5z^4 + z^3 - 2z$ in $B(0, 1)$.
5. Suppose $f(z)$ had a zero of multiplicity $m > 1$ at a . Use the Argument Principle to deduce that $f(z)$ is not injective in $B(a, R)$ for any $R > 0$.
- * 6. Suppose $f(z)$ is holomorphic in $B(0, R)$ and $r_1 < r_2 < R$. Show that if $M(r_1; f) = M(r_2; f)$ then $f(z)$ is constant.
- * 7. (Another proof of the fundamental theorem of algebra). Suppose $n \geq 1$ and

$$f(z) = a_0 + a_1 z + \cdots + a_{n-1} z^{n-1} + a_n z^n.$$

- (a) Let $g(z) = a_n z^n$. Show that if $R = |z| > 1$ then

$$|f(z) - g(z)| \leq M(n-1)R^{n-1}, \quad M = \max\{|a_0|, |a_1|, \dots, |a_{n-1}|\};$$

- (b) show that there is a $R > 0$ for which

$$M(n-1)R^{n-1} < |a_n|R^n = |g(z)|;$$

- (c) deduce from Rouché's theorem that $f(z)$ has exactly n roots.

Bibliography

- [1] Lars V. Ahlfors. *Complex analysis*. McGraw-Hill Book Co., New York, third edition, 1978. An introduction to the theory of analytic functions of one complex variable, International Series in Pure and Applied Mathematics.
- [2] Ralph Philip Boas, Jr. *Entire functions*. Academic Press Inc., New York, 1954.
- [3] D. J. H. Garling. *A course in mathematical analysis. Vol. III*. Cambridge University Press, Cambridge, 2014. Complex analysis, measure and integration.
- [4] Einar Hille. *Analytic function theory. Vol. 1*. Introduction to Higher Mathematics. Ginn and Company, Boston, 1959.
- [5] Einar Hille. *Analytic function theory. Vol. II*. Introductions to Higher Mathematics. Ginn and Co., Boston, Mass.-New York-Toronto, Ont., 1962.
- [6] William F. Osgood. A Jordan curve of positive area. *Trans. Amer. Math. Soc.*, 4(1):107–112, 1903.
- [7] Reinhold Remmert. *Theory of complex functions*, volume 122 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. Translated from the second German edition by Robert B. Burckel, Readings in Mathematics.
- [8] Walter Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.