

The Global Sparsity and Regulatory Rigidity of Freshwater Fecal Indicator Bacteria Monitoring

[Author list...]¹

¹[Affiliations...]

Key Points:

- We harmonized 5.3 million freshwater fecal indicator bacteria measurements from ten public sources into one global dataset.
- Openly shared monitoring is concentrated in the United States, Europe, and Canada, revealing a data sharing gap as much as a collection gap.
- Only about seven percent of monitored sites are sampled often enough to capture the rapid changes in fecal contamination.

Corresponding author: [TBD], [corresponding author contact]

Abstract

Fecal indicator bacteria, especially *Escherichia coli*, are the primary tool used worldwide to judge whether inland fresh waters are safe from waterborne pathogens, yet the monitoring record that underpins these decisions has never been characterized at a harmonized global scale. We compiled, harmonized, and analyzed 5,314,791 freshwater fecal indicator observations from ten public sources, standardized to a common schema, indicator set, and unit. Two structural weaknesses emerge. First, openly shared data are spatially unequal: the United States, Europe, and Canada account for roughly 89% of records, while much of the Global South is sparse and large monitoring nations are absent from open repositories, indicating that apparent data scarcity is in part an open-data sharing barrier rather than an absence of monitoring. Second, the record is temporally rigid: only about 7% of multi-sample sites are sampled at a daily or near-daily frequency, while contamination is a transient, weather-driven condition. Using the rare high-frequency sites, including three contrasting case studies, we show that low-frequency sampling misrepresents safety in both episodically and chronically contaminated waters. We argue for wider adoption of open-data practices and for higher-frequency, event-based, and predictive monitoring to make fecal indicator data actionable for public health. Key uncertainties include uneven coverage of non-*E. coli* indicators, a geographic bias of high-frequency sites toward North America and Europe, and an approximate unit conversion between enumeration methods.

Plain Language Summary

Bacteria such as *E. coli* are measured around the world to tell whether rivers and lakes are safe for swimming and for drinking-water supply. We gathered more than five million of these measurements from ten public sources in many countries and combined them into a single comparable dataset. We found two problems. First, almost all of the freely shared data come from the United States, Europe, and Canada; many other countries monitor their waters but do not publish the results in open, reusable form, so the world looks emptier of data than it really is. Second, most places are tested too infrequently (often only monthly or a few times a year) to catch the short bursts of pollution that follow rainfall and that pose the greatest health risk. At the few sites tested almost daily, we show that infrequent testing gives a misleading picture of safety, both at beaches that are usually clean and at rivers that are chronically polluted. Making water-quality data more openly shared, and testing more frequently where it matters, would help protect public health.

1 Introduction

Fecal indicator bacteria, and *Escherichia coli* in particular, are the primary tool used worldwide to judge whether fresh waters are safe from waterborne pathogens. The public-health stakes are large: inadequate water, sanitation, and hygiene remain responsible for a substantial global disease burden (Prüss-Ustün et al., 2014; Prüss-Ustün et al., 2019; Wolf et al., 2023), fecal contamination of drinking water is widespread across low- and middle-income settings (Bain, Cronk, Wright, et al., 2014; Bain, Cronk, Hosain, et al., 2014), and contact with fecally contaminated recreational water is an established cause of gastrointestinal and other illness (Gerdes et al., 2023; T. Wade et al., 2022; Zmirou et al., 2003). Because directly enumerating the many waterborne pathogens is impractical for routine monitoring, agencies instead measure culturable indicator organisms whose presence signals fecal contamination. A long line of epidemiological studies has linked indicator concentrations to swimmer illness and established *E. coli* and intestinal enterococci as the standard indicators for fresh and marine waters, respectively (Cabelli et al. (1982); subsequent reviews and cohort studies refined these relationships (Prüss, 1998; T. Wade et al., 2003; T. J. Wade et al., 2005; T. Wade et al., 2010), and

E. coli is widely regarded as the most reliable bacterial indicator of fecal contamination in fresh water (Edberg et al., 2000; Boehm & Sassoubre, 2014; Korajkic et al., 2018). These relationships underpin regulatory thresholds such as the United States Environmental Protection Agency recreational criteria and the European Bathing Water Directive (U.S. Environmental Protection Agency, 2012; European Parliament and Council of the European Union, 2006; World Health Organization, 2021; Boehm et al., 2008).

Despite this central role, the global record that underpins freshwater safety decisions has never been assembled and characterized at a harmonized, global scale. Monitoring data are dispersed across dozens of national and supranational portals with incompatible schemas, units, and indicator vocabularies. A growing body of work aggregates water-quality observations to enable large-sample analysis: in the United States, the Water Quality Portal unifies federal, state, and tribal records (Read et al., 2017), although reusing such multi-source data requires substantial harmonization and quality control (Sprague et al., 2017; Bousquin & Mullin, 2024). At continental to global scale, compilations such as AquaSat, the Global River Chemistry database, the Global River Water Quality Archive, large-sample chemistry extensions of catchment datasets, and global salinity syntheses have assembled physical and chemical variables for thousands of sites (Ross et al., 2019; Hartmann et al., 2014; Virro et al., 2021; Sterle et al., 2024; Thorslund & van Vliet, 2020), and recent work has begun to argue for global microbial water-quality data infrastructure (Rose et al., 2023). These resources are invaluable, but they are typically organized around water chemistry or single-nation coverage and do not characterize the spatial sharing structure or the temporal sampling cadence of fecal indicators specifically.

These two dimensions matter because fecal contamination is a transient, weather-driven condition. Numerous site studies document that fecal indicator concentrations vary by orders of magnitude over hours to days, exhibiting diurnal cycles, rainfall- and runoff-driven spikes, and rapid sunlight-driven decay (Whitman & Nevers, 2004; Whitman et al., 2004; Traister & Anisfeld, 2006; Meays et al., 2006; Desai & Rifai, 2013; Nguyen et al., 2022). This variability makes infrequent grab samples unreliable for classifying recreational safety (Kim & Grant, 2004), and it has motivated high-frequency sampling campaigns and statistical or data-driven nowcasting models that predict water quality from rainfall, turbidity, and discharge (Searcy et al., 2023; Searcy & Boehm, 2021; Heasley et al., 2021). Regulatory thresholds, however, are defined for compliance statistics and implicitly assume representative sampling — an assumption that weekly, monthly, or quarterly designs, often confined to a summer bathing season, routinely violate.

A complementary literature helps interpret why openly shared data are so unevenly distributed. The FAIR principles articulate norms for findable, accessible, interoperable, and reusable data (Wilkinson et al., 2016), yet technical, institutional, and social barriers impede open sharing of water-quality and other environmental monitoring data (Brasier et al., 2017; Sugg, 2021; Hazell et al., 2023; Heacock et al., 2022), and monitoring capacity itself varies systematically with national development and the flow of information through regulatory programs (Kirschke et al., 2020; Kumpel et al., 2020). Together these dynamics imply that an apparent absence of data in global repositories often reflects where data are *shared* rather than where they are *collected*.

Here we compile, harmonize, and analyze a global freshwater fecal indicator database of 5,314,791 observations drawn from ten public sources and standardized to a common schema, indicator vocabulary, and unit. Using this dataset we (i) map the global distribution of freshwater fecal indicator monitoring and show that openly shared data are concentrated in the United States, Europe, and Canada; (ii) characterize the indicator and ecosystem composition of the record; (iii) quantify per-site sampling cadence and show that only a small minority of sites are sampled at a frequency capable of resolving transient contamination; and (iv) use the rare high-frequency sites, including three contrasting case studies, to demonstrate that low-frequency sampling misrepresents safety

Table 1. Freshwater fecal indicator observations contributed by each source after cleaning and restriction to inland fresh waters.

Source	Region	Observations	Access
US Water Quality Portal	United States	3,321,490	API
Eionet (EEA bathing water)	Europe	763,440	Web scrape
GEMStat	Global (ex-US)	399,706	Manual
DataStream	Canada	296,913	API
UK Open WIMS	United Kingdom	189,485	API
NMMP	South Africa	161,467	Manual
Hub'Eau (Naiades)	France	146,650	API
LAWA	New Zealand	35,248	Manual
WPdx	Global	392	Manual
ANA (annual summaries)	Brazil	14,853	Manual

in both episodically and chronically contaminated waters. We interpret the resulting geography as evidence that apparent data sparsity is, in substantial part, an open-data sharing barrier, and we argue for both wider adoption of FAIR data practices and a shift toward higher-frequency, event-based, and predictive monitoring.

2 Data and Methods

2.1 Data sources

We aggregated freshwater fecal indicator observations from nine public point-observation networks and one annual-summary source, listed in Table 1. The sources span national and supranational programs retrieved either through programmatic interfaces or by manual download of curated releases. The United States Water Quality Portal contributed the majority of records, and the GEMStat compilation contributed globally distributed observations from many countries.

2.2 Harmonization and quality control

All concentrations were standardized to colony forming units per 100 mL. Most-probable-number values were treated as colony forming units, an approximation, and qPCR-equivalent and non-convertible units were dropped. Censored values were assigned a direction and filled at one half the lower detection limit (left-censored) or at the upper limit (right-censored). Source-specific location descriptors were mapped to a controlled vocabulary of water-body types. Repeated observations were averaged within each source and then de-duplicated across sources using a fuzzy key with approximately 1 km coordinate tolerance to remove records that appear in multiple portals.

2.3 Freshwater classification and cleaning

Cleaning retained records with a valid indicator, plausible date and coordinates, and a non-negative, bounded value, reducing 9,055,019 harmonized records to 9,044,108. Each record was then classified as marine, groundwater, or freshwater. Marine status was assigned where the location type was ocean or estuary, or where the bathing-water category indicated coastal or transitional waters; this category overrode an ambiguous location type and prevented coastal sites from being miscounted as inland. Groundwater was excluded as a distinct exposure pathway. Removing 3,409,529 marine and 26,284 groundwater records, reclassifying 212,554 records by bathing-water category, and dropping 293,504




Figure 1. Global density of freshwater fecal indicator bacteria observations, aggregated into hexagonal bins and shaded on a logarithmic scale. Monitoring is heavily concentrated in the United States and Europe, with sparse coverage across most of the Global South and an almost complete absence over Russia and China. Observations for India include fecal and total coliform, but exclude *E. coli*.

records that could not be verified as inland yielded the final analysis dataset of 5,314,791 freshwater observations.

2.4 Sampling-cadence and safety metrics

For each site we computed the number of observations, the temporal span, the mean return interval, and the median gap between consecutive samples. We grouped multi-sample sites into return-interval bands (daily, weekly, monthly, quarterly, yearly, and longer). High-frequency sites were defined as those with a median consecutive sampling gap of at most three days and at least 30 samples. To assess recreational safety we used a single-sample threshold of 235 colony forming units per 100 mL for *E. coli*, consistent with widely applied freshwater criteria (U.S. Environmental Protection Agency, 2012; Boehm et al., 2008), and computed the fraction of samples at or below this value.

3 Results

3.1 Geographic distribution and the sharing gap

Freshwater fecal indicator monitoring is highly spatially unequal (Figure 1). As summarized in Table 2, the United States accounts for 62.9% of records, Europe for 20.7%, and Canada for 5.7%, so that these regions together hold 89.3% of all observations; the remaining 10.7% is distributed across the rest of the world, led by Mexico, South Africa, India, and New Zealand. Large monitoring nations including China and Russia are effectively absent from the open, harmonized record, and Brazil appears only through publicly released annual summaries despite the existence of sub-annual sampling. Because several of the absent or thinly represented regions are known to operate national monitoring programs, we interpret much of this sparsity as an open-data sharing barrier rather than an absence of monitoring.

Table 2. Geographic distribution of the freshwater analysis dataset.

Region	Records	Share
United States	3,342,758	62.9%
Europe	1,099,575	20.7%
Canada	305,432	5.7%
Rest of world	567,026	10.7%

Table 3. Indicator and ecosystem composition of the freshwater dataset.

Category	Records	Share
<i>E. coli</i>	2,308,643	43.4%
Fecal coliform	1,692,665	31.8%
Total coliform	560,355	10.5%
Intestinal enterococci	479,053	9.0%
Fecal streptococci	274,075	5.2%
River or stream	3,848,418	72.4%
Lake	1,199,766	22.6%
Other freshwater	266,607	5.0%

3.2 Indicator and ecosystem composition

No single indicator dominates the dataset: *E. coli* accounts for 43.4% of records, fecal coliform for 31.8%, total coliform for 10.5%, intestinal enterococci for 9.0%, and fecal streptococci for 5.2% (Table 3). Coverage of non-*E. coli* indicators is uneven across sources; intestinal enterococci, for example, derive almost entirely from European inland bathing waters. By ecosystem, the data are dominated by rivers and streams (72.4%) and lakes (22.6%), with reservoirs, canals, ditches, springs, wetlands, and ponds making up the remainder.

3.3 The actionability gap in sampling frequency

Most sites are sampled too sparsely, and too infrequently, to resolve transient contamination (Figure 3). Of 133,328 distinct sites, only 10.8% have at least 100 observations and just 0.7% have at least 500. Among the 116,489 sites with two or more samples, only 7.0% are sampled at a daily or near-daily cadence, whereas 29.9% are sampled monthly and 30.7% quarterly (Table 4). The contrast in monitoring objectives is illustrated in Figure 2, which compares three Wisconsin sites whose designs differ by an order of magnitude in sampling frequency.

3.4 High-frequency safety and case studies

High-frequency monitoring is rare and geographically concentrated in North America and Europe (Figure 5). Across 631 high-frequency sites comprising 235,398 *E. coli* samples, 77.6% of observations were at or below the safe threshold and 22.4% exceeded it, but per-site exceedance rates vary widely, from sites that are consistently safe to sites in agricultural or urban catchments that are chronically above the threshold. The three 2023 case studies in Figure 4 and Table 5 make the implication concrete: an enclosed Lake Erie beach in the United States was safe most of the time (18.0% exceedance) with short-lived spikes that recovered within days, whereas the River Lune in the United Kingdom and the Taruheru River in New Zealand exceeded the safe threshold for the major-




Figure 2. Three Wisconsin freshwater sites with contrasting sampling cadences. The locator map marks a daily-sampled swimming beach, a monthly regulatory river site, and a quarterly trend site; the panels show their 2022–2023 *E. coli* observations, with 101, 19, and 17 samples respectively. Sampling frequency, rather than concentration trend, distinguishes the three designs.




Figure 3. Quantifying the actionability gap. Left: the percentage of sites retained as the minimum observation count per site increases. Right: the distribution of mean sampling return intervals among multi-sample sites, with the near-daily category highlighted.

Table 4. Distribution of per-site sampling return intervals among the 116,489 sites with at least two samples.

Return interval band	Share of multi-sample sites
Daily (at most 1.5 days)	7.0%
Weekly (1.5 to 8 days)	9.1%
Monthly (8 to 31 days)	29.9%
Quarterly (31 to 92 days)	30.7%
Yearly (92 to 366 days)	19.2%
Longer (over 366 days)	4.2%

Table 5. High-frequency case-study sites in 2023.

Site	Country	Samples	Exceedance	Median	Maximum
Euclid Beach	United States	128	18.0%	41	2,420
River Lune	United Kingdom	43	83.7%	580	100,000
Taruheru River	New Zealand	40	62.5%	465	11,000

^a Median and maximum concentrations in colony forming units per 100 mL.

ity of the year (83.7% and 62.5% exceedance). In the first case, a single monthly grab sample taken during a transient spike would have closed an otherwise-safe beach for weeks; in the latter two, a favorably timed monthly sample could have certified as safe a river that was unsafe most of the year. Low-frequency sampling therefore misrepresents safety in both regimes, and higher-frequency data improve management in both.

4 Discussion

4.1 Collection gap versus sharing gap

The geography of the freshwater record (Figure 1) is best understood as the joint product of two distinct gaps. A genuine collection gap exists in resource-limited regions where monitoring capacity is constrained and varies with national development (Kirschke et al., 2020). But the near-absence of large, industrialized monitoring nations from the open record points to a second, sharing gap: data are collected but not published in open, machine-readable, harmonized form. This interpretation is consistent with a literature documenting persistent technical, institutional, and social barriers to sharing environmental and water-quality data (Brasier et al., 2017; Sugg, 2021; Hazell et al., 2023; Heacock et al., 2022; Kumpel et al., 2020), and it implies that funding new stations alone will not close the gap. Wider adoption of the FAIR principles (Wilkinson et al., 2016), together with data-sharing norms comparable to those that govern international exchange of meteorological data, would convert existing but invisible monitoring into a usable global resource.

4.2 Regulatory rigidity and management implications

Our cadence results quantify a mismatch between how fecal indicators are monitored and how they behave. Because contamination is transient (Whitman & Nevers, 2004; Traister & Anisfeld, 2006; Meays et al., 2006), calendar-based low-frequency sampling is structurally blind to acute risk, and because culture-based assays return results only after roughly a day, advisories are inherently retrospective (Kim & Grant, 2004). The high-frequency case studies show that this rigidity cuts both ways: infrequent sam-




Figure 4. High-frequency *E. coli* monitoring at three sites on three continents in 2023, plotted against the 235 colony forming units per 100 mL safe threshold with a shaded safe zone. The Lake Erie beach (United States) remained largely safe with short-lived spikes, whereas the River Lune (United Kingdom) and Taruheru River (New Zealand) exceeded the threshold for most of the year.




Figure 5. Global safety of high-frequency *E. coli* monitoring sites. The map plots sites with at least near-daily sampling, colored by the share of samples exceeding the safe threshold; the donut summarizes the overall split of 77.6 percent safe versus 22.4 percent unsafe across 631 sites and 235,398 samples.

pling can both over-restrict an otherwise-safe beach after a single spike and certify as safe a chronically polluted river. Higher-frequency, event-based sampling, and predictive nowcasting models that exploit rainfall, turbidity, and discharge (Searcy & Boehm, 2021; Searcy et al., 2023; Heasley et al., 2021), therefore offer substantial gains in both directions. Scaling such approaches beyond the handful of intensively studied sites we identify is, in our view, the central operational opportunity.

4.3 Limitations

Several limitations bound our conclusions. The treatment of most-probable-number values as colony forming units is an approximation, and datetimes and time zones were not harmonized globally. Coverage of non-*E. coli* indicators is uneven across sources, so cross-indicator comparisons of total monitoring effort should be made cautiously. High-frequency sites are geographically biased toward North America and Europe, so the 77.6% safe fraction is conditional on those sites and is not a globally representative safety rate. Finally, the case studies are individual sites chosen to illustrate distinct regimes rather than a representative sample.

5 Conclusions

We assembled and characterized the first harmonized, global, freshwater-focused dataset of fecal indicator bacteria, comprising 5,314,791 observations from ten public sources. The record exhibits two structural weaknesses for public health: a spatial concentration of openly shared data in a few wealthy regions that reflects an open-data sharing barrier as much as a collection gap, and a temporal rigidity in which only a small minority of sites are sampled often enough to resolve the transient dynamics of fecal contamination. High-frequency sites demonstrate that low-frequency monitoring misrepresents safety in both episodically and chronically contaminated waters. Closing these gaps will require both open, FAIR data sharing and a shift toward higher-frequency, event-based, and predictive monitoring.

Open Research Section

The harmonized freshwater fecal indicator dataset and the data-cleaning and figure-generation code that support the findings of this study will be archived in a public repository with a persistent identifier prior to publication [repository DOI — to be completed]; data are not available solely from the authors. The underlying observations were obtained from the following public sources, each of which should be cited in the final reference list with its access point: the US Water Quality Portal (Read et al., 2017), the European Environment Agency Eionet bathing-water reporting system, the UN GEMS/Water GEM-Stat database, DataStream, the UK Environment Agency Water Quality Archive, the South African National Microbial Monitoring Programme, the French Hub'Eau Naiades service, Land Air Water Aotearoa, the Water Point Data Exchange, and the Brazilian National Water Agency. [Per-source data citations with persistent identifiers to be completed.]

Conflict of Interest disclosure

The authors declare no conflicts of interest for this manuscript.

Acknowledgments

[Funding sources and acknowledgments to be completed.]

References

- Bain, R., Cronk, R., Hossain, R., Bonjour, S., Onda, K., Wright, J. A., ... Bartram, J. (2014). Global assessment of exposure to faecal contamination through drinking water based on a systematic review. *Tropical medicine & international health*. doi: 10.1111/tmi.12334
- Bain, R., Cronk, R., Wright, J., Yang, H., Slaymaker, T., & Bartram, J. (2014). Fecal contamination of drinking-water in low- and middle-income countries: A systematic review and meta-analysis. *PLoS Medicine*. doi: 10.1371/journal.pmed.1001644
- Boehm, A. B., Ashbolt, N. J., Colford, J. M., Dunbar, L. E., Fleming, L. E., Gold, M., ... Weisberg, S. B. (2008). A sea change ahead for recreational water quality criteria. *Journal of Water and Health*. doi: 10.2166/wh.2009.122
- Boehm, A. B., & Sassoubre, L. M. (2014). *Enterococci as indicators of environmental fecal contamination*.
- Bousquin, J., & Mullin, C. (2024). harmonize-wq: Standardize, clean and wrangle waterquality portal data into more analytic-ready formats. *The Journal of Open Source Software*. doi: 10.21105/joss.07305
- Brasier, K. J., Jalbert, K., Kinchy, A., Brantley, S. L., & Unroe, C. (2017). Barriers to sharing water quality data: experiences from the shale network. *Journal of Environmental Planning and Management*. doi: 10.1080/09640568.2016.1276435
- Cabelli, V. J., Dufour, A., McCabe, L. J., & Levin, M. A. (1982). Swimming-associated gastroenteritis and water quality. *American Journal of Epidemiology*. doi: 10.1093/oxfordjournals.aje.a113342
- Desai, A. M., & Rifai, H. S. (2013). *Escherichia coli* concentrations in urban watersheds exhibit diurnal sag: Implications for water-quality monitoring and assessment. *JAWRA Journal of the American Water Resources Association*. doi: 10.1111/jawr.12039
- Edberg, S. C., Rice, E. W., Karlin, R. J., & Allen, M. J. (2000). *Escherichia coli*: the best biological drinking water indicator for public health protection. *Journal of Applied Microbiology*. doi: 10.1111/j.1365-2672.2000.tb05338.x
- European Parliament and Council of the European Union. (2006). *Directive 2006/7/ec concerning the management of bathing water quality and repealing directive 76/160/ee*. Official Journal of the European Union L 64.
- Gerdes, M. E., Miko, S., Kunz, J. M., Hannapel, E., Hlavsa, M., Hughes, M., ... Collier, S. (2023). Estimating waterborne infectious disease burden by exposure route, united states, 2014. *Emerging Infectious Diseases*. doi: 10.3201/eid2907.230231
- Hartmann, J., Lauerwald, R., & Moosdorf, N. (2014). *A brief overview of the global river chemistry database, glorich*. doi: 10.1016/J.PROEPS.2014.08.005
- Hazell, P., Novitzky, P., & Oord, S. V. D. (2023). Socio-technical system analysis of responsible data sharing in water systems as critical infrastructure. *Frontiers in Big Data*. doi: 10.3389/fdata.2022.1057155
- Heacock, M., Lopez, A. R., Amolegbe, S. M., Carlin, D., Henry, H., Trottier, B., ... Suk, W. (2022). Enhancing data integration, interoperability, and reuse to address complex and emerging environmental health problems. *Environmental Science and Technology*. doi: 10.1021/acs.est.1c08383
- Heasley, C., Sánchez, J., Tustin, J., & Young, I. (2021). Systematic review of predictive models of microbial water quality at freshwater recreational beaches. *PLoS ONE*. doi: 10.1371/journal.pone.0256785
- Kim, J. H., & Grant, S. B. (2004). Public mis-notification of coastal water quality: A probabilistic evaluation of posting errors at huntington beach, california. *Environmental Science & Technology*. doi: 10.1021/es034382v
- Kirschke, S., Avellán, T., Bärlund, I., Bogárdi, J. J., Carvalho, L., Chapman, D. V., ... Warner, S. (2020). Capacity challenges in water quality monitoring: un-

- derstanding the role of human development. *Environmental Monitoring and Assessment*. doi: 10.1007/s10661-020-8224-3
- Korajkic, A., McMinn, B., & Harwood, V. (2018). Relationships between microbial indicators and pathogens in recreational water settings. *International Journal of Environmental Research and Public Health*. doi: 10.3390/ijerph15122842
- Kumpel, E., MacLeod, C., Stuart, K., Cock-Esteb, A., Khush, R., & Peletz, R. (2020). From data to decisions: understanding information flows within regulatory water quality monitoring programs. *npj Clean Water*. doi: 10.1038/s41545-020-00084-0
- Meays, C. L., Broersma, K., Nordin, R., Mazumder, A., & Samadpour, M. (2006). Diurnal variability in concentrations and sources of escherichia coli in three streams. *Canadian Journal of Microbiology*. doi: 10.1139/w06-066
- Nguyen, K., Smith, S. M., Roundtree, A., Feistel, D. J., Kirby, A. E., Levy, K., & Mattioli, M. C. (2022). Fecal indicators and antibiotic resistance genes exhibit diurnal trends in the chattahoochee river: Implications for water quality monitoring. *Frontiers in Microbiology*. doi: 10.3389/fmicb.2022.1029176
- Prüss, A. (1998). Review of epidemiological studies on health effects from exposure to recreational water. *International Journal of Epidemiology*. doi: 10.1093/ije/27.1.1
- Prüss-Ustün, A., Bartram, J., Clasen, T., Colford, J., Cumming, O., Curtis, V., ... Cairncross, S. (2014). Burden of disease from inadequate water, sanitation and hygiene in low- and middle-income settings: a retrospective analysis of data from 145 countries. *Tropical medicine & international health*. doi: 10.1111/tmi.12329
- Prüss-Ustün, A., Wolf, J., Bartram, J., Clasen, T., Cumming, O., Freeman, M. C., ... Johnston, R. B. (2019). Burden of disease from inadequate water, sanitation and hygiene for selected adverse health outcomes: An updated analysis with a focus on low- and middle-income countries. *International Journal of Hygiene and Environmental Health*. doi: 10.1016/j.ijheh.2019.05.004
- Read, E. K., Carr, L., Cicco, L. A. D., Dugan, H. A., Hanson, P. C., Hart, J. A., ... Winslow, L. (2017). Water quality data for national-scale aquatic research: The water quality portal. *Water Resources Research*. doi: 10.1002/2016wr019993
- Rose, J. B., Hofstra, N., Hollmann, E., Katsivelis, P., Medema, G., Murphy, H., ... Verbyla, M. E. (2023). Global microbial water quality data and predictive analytics: Key to health and meeting sdg 6. *PLOS Water*. doi: 10.1371/journal.pwat.0000166
- Ross, M., Topp, S., Appling, A., Yang, X., Kuhn, C., Butman, D., ... Pavelsky, T. (2019). Aquasat: A data set to enable remote sensing of water quality for inland waters. *Water Resources Research*. doi: 10.1029/2019WR024883
- Searcy, R. T., & Boehm, A. (2021). A day at the beach: Enabling coastal water quality prediction with high-frequency sampling and data-driven models. *Environmental Science and Technology*. doi: 10.1021/acs.est.0c06742
- Searcy, R. T., Phaneuf, J. R., & Boehm, A. B. (2023). High-frequency fecal indicator bacteria (fib) observations to assess water quality drivers at an enclosed beach. *PLoS ONE*. doi: 10.1371/journal.pone.0286029
- Sprague, L., Oelsner, G., & Argue, D. M. (2017). Challenges with secondary use of multi-source water-quality data in the united states. *Water Research*. doi: 10.1016/j.watres.2016.12.024
- Sterle, G., Perdrial, J., Kincaid, D. W., Underwood, K. L., Rizzo, D. M., Haq, I. U., ... Harpold, A. A. (2024). Camels-chem: augmenting camels (catchment attributes and meteorology for large-sample studies) with atmospheric and stream water chemistry data. *Hydrology and earth system sciences*. doi: 10.5194/hess-28-611-2024
- Sugg, Z. (2021). Social barriers to open (water) data. *WIREs Water*. doi: 10.1002/

- wat2.1564
- Thorslund, J., & van Vliet, M. V. (2020). A global dataset of surface water and groundwater salinity measurements from 1980–2019. *Scientific Data*. doi: 10.1038/s41597-020-0562-z
- Traister, E., & Anisfeld, S. C. (2006). Variability of indicator bacteria at different time scales in the upper hoosic river watershed. *Environmental Science & Technology*. doi: 10.1021/es0601437
- U.S. Environmental Protection Agency. (2012). *Recreational water quality criteria* (Tech. Rep. No. EPA 820-F-12-058). U.S. Environmental Protection Agency, Office of Water.
- Virro, H., Amatulli, G., Kmoch, A., Shen, L. Q., & Uuemaa, E. (2021). Grqa: Global river water quality archive. *Earth system science data*. doi: 10.5194/essd-13-5483-2021
- Wade, T., Arnold, B., Schiff, K., Colford, J., Weisberg, S., Griffith, J., & Dufour, A. (2022). Health risks to children from exposure to fecally-contaminated recreational water. *PLoS ONE*. doi: 10.1371/journal.pone.0266749
- Wade, T., Pai, N., Eisenberg, J., & Colford, J. (2003). Do u.s. environmental protection agency water quality guidelines for recreational waters prevent gastrointestinal illness? a systematic review and meta-analysis. *Environmental Health Perspectives*. doi: 10.1289/ehp.6241
- Wade, T., Sams, E., Brenner, K., Haugland, R., Chern, E., Beach, M., ... Dufour, A. (2010). Rapidly measured indicators of recreational water quality and swimming-associated illness at marine beaches: a prospective cohort study. *Environmental Health*. doi: 10.1186/1476-069X-9-66
- Wade, T. J., Calderon, R. L., Sams, E., Beach, M. J., Brenner, K. P., Williams, A. H., & Dufour, A. (2005). Rapidly measured indicators of recreational water quality are predictive of swimming-associated gastrointestinal illness. *Environmental Health Perspectives*. doi: 10.1289/ehp.8273
- Whitman, R. L., & Nevers, M. B. (2004). *Escherichia coli* sampling reliability at a frequently closed chicago beach: Monitoring and management implications. *Environmental Science & Technology*. doi: 10.1021/es034978i
- Whitman, R. L., Nevers, M. B., Korinek, G. C., & Byappanahalli, M. N. (2004). Solar and temporal effects on *escherichia coli* concentration at a lake michigan swimming beach. *Applied and Environmental Microbiology*. doi: 10.1128/aem.70.7.4276-4285.2004
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., ... Mons, B. (2016). The fair guiding principles for scientific data management and stewardship. *Scientific Data*. doi: 10.1038/sdata.2016.18
- Wolf, J., Johnston, R., Ambelu, A., Arnold, B., Bain, R. E. S., Brauer, M., ... Cumming, O. (2023). Burden of disease attributable to unsafe drinking water, sanitation, and hygiene in domestic settings: a global analysis for selected adverse health outcomes. *The Lancet*. doi: 10.1016/S0140-6736(23)00458-0
- World Health Organization. (2021). *Guidelines on recreational water quality. volume 1: Coastal and fresh waters* (Tech. Rep.). World Health Organization.
- Zmirou, D., Pena, L., Ledrans, M., & LeTertre, A. (2003). Risks associated with the microbiological quality of bodies of fresh and marine water used for recreational purposes: Summary estimates based on published epidemiological studies. *Archives of Environmental Health An International Journal*. doi: 10.3200/aeoh.58.11.703-711