

Temporal Equivalence Principle: Suppressed Density Scaling in Globular Cluster Pulsars

Matthew Lukin Smawfield

Version: v0.10 (Caracas)

First published: 9 January 2026 | Last updated: 17 October 2026

DOI: 10.5281/zenodo.18165798

Code Availability: github.com/matthewsmawfield/TEP-COS

Abstract

A spatially stratified spin-down anomaly is reported in 199 globular-cluster millisecond pulsars compared with 351 field controls. Cluster pulsars show a 0.63 dex raw excess and a 0.40 dex controlled residual, with covariance-aware significance of 5.6σ. The signal exhibits suppressed density scaling: a mixed-effects slope $\Gamma = 0.39 \pm 0.08$ dex/dex compared with a Newtonian ensemble baseline $\Gamma = 0.75$, rejecting the baseline at 4.1σ. Field binaries provide a null control, while cluster binaries are quieter than isolated cluster pulsars; corrected CMC reproduces this binary direction, so the binary result functions as a comparator check rather than the primary failure mode. An adversarial dynamics model accounting for line-of-sight acceleration, jerk, Shklovskii terms, mass segregation, and cluster mass profiles cannot explain the observed suppression. An N-body baseline derived from CMC synthetic pulsars (20 cluster selections mapping to 14 unique Kremer et al. 2020 realizations; 9,526 evolved neutron stars, 6,617 unique) predicts a structurally rescaled excess 3.2× larger than observed (13.2σ nominal model-data tension, not detection significance; 13.0σ model-aware, 8.6σ under 20-cluster ensemble-propagated uncertainty, and 4.1σ under the 100%-correlated conservative quadrature ceiling), and a mock pulsar-timing-array detection pipeline confirms that the missing high-acceleration population cannot be explained by the tested observational-selection models. These results are interpreted as a time-domain test of the TEP observable response coefficient κ_{MSP}, not as a measurement of a microscopic scalar coupling.

Code Availability: All data and analysis code required to reproduce the results presented in this work, including the full pulsar catalog compilation, are available in the public repository at https://github.com/matthewsmawfield/TEP-COS.

Keywords: temporal equivalence principle, pulsar timing, globular clusters, time dilation, screening transition, modified gravity

1. Introduction: Time-Domain Tests of Modified Gravity

1.1 The Intermediate-Scale Problem

General Relativity has passed every precision test in the Solar System. Yet at intermediate and cosmological scales, persistent discrepancies arise—rotation curves, cluster dynamics, cosmic acceleration—that conventionally require invisible mass or exotic energy to resolve. A fundamental question follows: Is gravitational time dilation scale-dependent? This work explores the hypothesis that these anomalies reflect not missing matter but modified temporal structure: a scale-dependent amplification of gravitational time dilation beyond the predictions of standard General Relativity.

The Temporal Equivalence Principle (TEP) formalizes this possibility within a two-metric framework (see Section 2), predicting that the rate of proper time accumulation is environment-dependent at intermediate scales while remaining consistent with precision tests in the screened Solar System regime. The central prediction is that *rate-dependent* physical processes—pulsar spin-down, photon arrival times, clock frequencies—should exhibit anomalies in deep gravitational potentials, while *fossil* observables that integrate over formation timescales remain insensitive.

1.2 Why Time-Domain Tests Are Critical

TEP modifies the proper time relation: $d\tilde{\tau}/d\tau_g = A(\varphi)$ where $A(\varphi) = \exp(\beta\varphi/M_{Pl})$, $\tilde{\tau}$ is the TEP-modified proper time, τ_g is the GR proper time, and $A(\varphi)$ is the universal conformal factor. This creates two classes of observables with quantitatively different TEP sensitivity:

Observable Class	Examples	TEP Sensitivity	Rationale
Time-Domain (Rates)	Pulsar $\dot{P}$, Clock frequencies	HIGH	Measures present-tense clock rate
Fossil (Archaeology)	Stellar ages, [α/Fe], colors, SFH	LOW	Integrates over ~Gyr formation history

The expected TEP differential (~10 kyr over cosmic time) is $O(10^{-6})$ of the formation timescale spread (~Gyr) for stellar populations. Fossil observables are unlikely to distinguish TEP from standard astrophysical processes at practical significance levels. This paper therefore focuses exclusively on time-domain tests: pulsar spin-down rates.

1.3 Central Results

Theoretical Framework

The TEP framework uses observational data to constrain the *class* of viable modified gravity theories through the observable pulsar response coefficient κ_{MSP} (absorbing stellar physics and environment, not the microscopic coupling β) and the topological transition scale R_T, following the same strategy as the PPN (Parameterized Post-Newtonian) framework used to test GR in the Solar System. The effective screening framework motivates the response model tested here; candidate transfer models are discussed separately and are not required for the empirical inference. Section 2.1.1 presents the screening framework, making this manuscript self-contained and independently evaluable.

Observable	Status	Result	Details
Pulsar Timing	Cluster Spin-down Residual	Anomaly Detection	0.63 dex raw excess; 0.40 dex controlled residual; null in field
Field Binary Control	Binary vs Isolated (Field)	Null Control	p = 0.78 (supports environmental origin)
Binary Comparator	Binary vs Isolated (Cluster)	Comparator Check	Binaries −0.33 dex quieter than isolated; corrected CMC predicts −0.27 dex (3rc filter), consistent direction
Spatial Stratification	Core vs Outskirts	Suggestive	−0.30 dex (inner, p=0.074) vs −0.14 dex (outer, p=0.41)
Suppressed Density Scaling	Does the signal track dynamical noise (ρ^2) or potential (Φ)?	Consistency test	Observed slope = 0.39 vs Newtonian slope = 0.75 (4.1σ rejection)

The pulsar signal satisfies three independent criteria consistent with TEP: (i) Spatial Resolution: The spin-down anomaly is concentrated in cluster cores (−0.30 dex for inner binaries, p = 0.074) and absent in the outskirts (−0.14 dex, p = 0.41), directly tracking the local temporal-field environment. (ii) Environmental Isolation: The Field Binary Control supports an environmental rather than intrinsic origin—the binary vs isolated difference vanishes in the galactic field (p = 0.78). (iii) Suppressed Density

Scaling: While standard dynamics predicts residuals scaling strongly with density (ensemble slope ≈ 0.75), the observed slope is only 0.39 ± 0.08—a 4.1σ rejection. Leave-one-cluster-out validation confirms this result is stable (3.7% relative instability, STABLE assessment). All 15 clusters with sufficient statistics (N ≥ 4) show positive controlled residuals (+0.009 to +0.695 dex), consistent with a broad environmental enhancement among the well-sampled clusters that saturates rather than scaling with density. Two of the remaining 10 lower-N clusters show marginally negative residuals, but these carry large uncertainties and do not alter the primary inference.

1.4 The Screening Hierarchy and ρ_T

A central requirement of TEP phenomenology is that intermediate-scale signals coexist with stringent Solar System bounds. Local light-propagation tests constrain the screened local gradient/source-charge sector and any disformal cone tilt. The pulsar analysis instead fits an observable response coefficient, $\kappa_{\rm MSP}$, in globular-cluster environments. $\kappa_{\rm MSP}$ should not be compared directly to PPN couplings without a solved transfer function.

The screening hierarchy itself is governed by the non-linear superposition of field gradients, termed Temporal Shear. While the asymptotic saturation of screening occurs at the Temporal Topology saturation proximity scale $\rho_T \approx 20$ g/cm³, the onset of modulation is sensitive to the gradient coherence length. In compact systems like globular clusters ($\rho \sim 10^{-18}$ g/cm³), the gradient is set by the diffuse embedding halo, allowing TEP effects to remain fully active despite high local densities. In contrast, distributed galactic environments ($\rho \sim 10^{-23}$ g/cm³) matching the gradient scale exhibit partial suppression, as observed in the distance-ladder modulation of Paper 11.

Within this weakly screened regime, the TEP clock response exhibits topological flattening rather than scaling indefinitely with potential depth. This produces a characteristic signature: residuals that do not track density as strongly as Newtonian dynamics predicts. The observed suppressed density scaling (4.1σ rejection of ρ² dynamics, with all 15 clusters meeting the N ≥ 4 threshold showing positive controlled residuals) is consistent with this continuous geometric screening behavior.

1.5 Paper Structure

The analysis is organized to prioritize empirical evidence from time-domain probes:

1. Section 2 establishes the theoretical framework: the TEP modification and spin-down predictions for pulsars.
2. Section 3 presents the primary detection: pulsar timing in globular clusters using 550 MSPs (199 GC, 351 field) with measured Ṗ, including the Suppressed Density Scaling test, Spatial Stratification, and Field Binary Control. A conservative period-matched analysis with 400 MSPs serves as robustness confirmation.
3. Section 4 discusses the unified picture, falsification criteria, and implications.
4. Section 5 concludes.

The Joint Pattern

The TEP framework is supported by the joint pattern: a robust GC–field environmental excess, suppressed density scaling relative to CMC/Newtonian expectations, a null field-binary control, GC binaries being quieter than isolated (consistent with the CMC prediction), and the failure of observational selection to remove the predicted high-acceleration population. A successful standard-dynamics explanation must reproduce all of these signatures simultaneously.

The pulsar signal satisfies three independent criteria consistent with TEP: (i) *Spatial Resolution:* The spin-down anomaly is concentrated in cluster cores (−0.30 dex for inner binaries, p = 0.074) and absent in the outskirts (−0.14 dex, p = 0.41), directly tracking the local temporal-field environment. (ii) *Environmental Isolation:* The Field Binary Control supports an environmental rather than intrinsic origin—the binary vs isolated difference vanishes in the galactic field (p = 0.78), while the GC/field differential test supports an environmental origin (one-sided p = 0.036). (iii) *Suppressed Density Scaling:* While standard dynamics predicts residuals scaling strongly with density (ensemble slope ≈ 0.75), the observed slope is only 0.39 ± 0.08—a 4.1σ rejection. Leave-one-cluster-out validation confirms this result is stable (3.7% relative instability, STABLE assessment). All 15 clusters with sufficient statistics (N ≥ 4) show positive controlled residuals (+0.009 to +0.695 dex), consistent with an environmental enhancement that saturates rather than scaling with density across the clusters entering the mixed-effects analysis. Two of the remaining 10 lower-N clusters show marginally negative residuals, but these carry large uncertainties and do not alter the primary inference.

2. Theoretical Framework: The Screening Transition

The Temporal Equivalence Principle predicts that gravitational time dilation is enhanced at intermediate astrophysical scales while remaining consistent with precision tests in the screened Solar System regime. This section establishes the theoretical basis for the time-domain probe examined in this work: pulsar spin-down in globular clusters, based on the continuous geometric screening framework established in TEP v0.16 Jakarta. This theoretical foundation is necessary to derive the specific quantitative predictions (Pulsar $\dot{P}$ drift) tested in the subsequent sections.

2.1 The TEP Modification

Notation and Conventions

To ensure consistency with the foundational theory (see Section 1) while adapting for astrophysical phenomenology, the following conventions are adopted:

- Metrics: $g_{\mu\nu}$ denotes the gravitational metric (Einstein frame); $\tilde{g}_{\mu\nu}$ denotes the physical matter metric (Jordan frame) to which clocks and rulers couple.
- Fields: ϕ represents the fundamental scalar time field. Φ represents the standard Newtonian gravitational potential ($\Phi \leq 0$).
- Weak-Field Limit: In the non-relativistic limit appropriate for clusters and halos, a linear mapping $\phi \propto \Phi$ is assumed, absorbing coupling constants into the observable pulsar response coefficient $\kappa_{\rm MSP}$.
- Proper Time (τ): Always refers to the physical time measured by an atomic clock ($\tilde{g}$-frame invariant).

Under the Temporal Equivalence Principle, the local proper time τ is related to coordinate time t by:

$$\frac{d\tau}{dt} = 1 + \frac{\Phi}{c^2} + \kappa_{\rm MSP} \cdot f(\Phi, \nabla\Phi) \tag{1}$$

where Φ is the gravitational potential, and $\kappa_{\rm MSP}$ is the enhancement factor. Standard GR corresponds to $\kappa_{\rm MSP} = 0$. The function f(Φ, ∇Φ) encodes the scale-dependent modification.

2.1.1 The Screening Mechanism: A Self-Contained Derivation

To make the screening physics explicit and evaluable without recourse to external references, the effective screening framework is presented here. The TEP framework employs a scalar-tensor structure with two metrics on a single spacetime manifold:

Axiomatic Foundation

- Two-metric structure: Gravity is described by a Lorentzian metric $g_{\mu\nu}$ (Einstein frame); matter fields and clocks couple to the canonical causal (matter) metric $\tilde{g}_{\mu\nu} = -A_{\rm loc}^2(u_{\rm grav}(\mathbf{x}))U_\mu U_\nu + A_{\rm sp}^2(u_{\rm drift}(t,\mathbf{x}))\Pi_{\mu\nu} + B(\phi)\Pi^\alpha_\mu\Pi^\beta_\nu\nabla_\alpha\phi\nabla_\beta\phi$ ($U_\mu \equiv -\nabla_\mu\bar{\phi}/\sqrt{2\bar{X}}$, $\Pi_{\mu\nu} = g_{\mu\nu} + U_\mu U_\nu$; Paper 0, §2.2, Step 80), written in conformal–disformal shorthand as $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi$, where $A(\phi) = \exp(\beta_A\phi/M_{\rm Pl})$ is the universal conformal factor.
- Temporal Equivalence Principle: All non-gravitational processes evolve according to proper time $d\tau$ defined by $\tilde{g}_{\mu\nu}$. In local freely falling frames, physics reduces to special relativity with invariant c.
- Screening via continuous gradient: Rather than operating via discrete boundary cutoffs or thin-shell transitions, screening manifests as a continuous spatial profile governed by the non-linear superposition of field gradients (Temporal Shear). This suppresses Temporal Shear and the associated clock-rate responses in regions of high curvature (Solar System) while leaving low-curvature astrophysical environments accessible to dynamics.

The Action and Field Equations. For comparison, the canonical scalar baseline in the Einstein frame is:

$$S = \int d^4x\sqrt{-g}\left[\frac{M_{\rm Pl}^2}{2}R - \frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi)\right] + S_{\rm matter}[\psi, \tilde{g}_{\mu\nu}, \phi] \tag{2}$$

where $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi$ (in isotropic shorthand). The operative TEP screening realization used across the corpus replaces this canonical scalar kinetic sector with the selected two-branch $P(X,\phi)$ completion

$P(X,\phi) = -V(\phi) + \frac{2k}{3A_X^2} X|X|^{1/2} + X|X|/\Lambda_X^4$ ($P_{,X} = k\sqrt{\xi} + 2\xi$, $k \simeq 16.03$; Paper 0, §2.2), for which static ellipticity and time-dependent hyperbolicity are certified on the adopted branch; the canonical equations below are retained to display the baseline scalar-tensor sourcing structure. Variation of the baseline action yields the Einstein equations $G_{\mu\nu} = M_{\rm Pl}^{-2}[T_{\mu\nu}(\phi) + T_{\mu\nu}^{(\rm matt)}]$ and the scalar field equation:

$$\Box\phi - V'(\phi) = -\alpha(\phi)T + S_{\rm disf} \tag{3}$$

where $\alpha(\phi) \equiv d\ln A/d\phi = \beta_A/M_{\rm Pl}$, $T \equiv \tilde{g}^{\mu\nu}\tilde{T}_{\mu\nu}$ is the matter-frame trace, and $S_{\rm disf}$ represents disformal source terms. In the non-relativistic limit for slow observers, the proper time increment is $d\tau/dt \approx A(\phi) = \exp(\beta_A\phi/M_{\rm Pl})$.

Observable Response Coefficient

$\kappa_{\rm MSP}$ *is treated as an observable pulsar response coefficient.* It is not identified with the microscopic coupling β_A. Solar-System PPN and MICROSCOPE tests constrain the environmentally screened effective source and composition-dependent couplings rather than directly determining the unsuppressed conformal parameter β_A. In the corpus-wide transfer architecture (Paper 0 §7; Paper 11, Step 62), the unsuppressed $\sim 10^6$ reference scale arises from the volumetric saturation-to-envelope density ratio $\rho_T/\bar{\rho}_{\rm proc}$ at $\rho_T \approx 20\ {\rm g\,cm^{-3}}$ and the unit-transport benchmark $\lambda_0 = 1$ (with bare coupling $|\beta_A| = 1$ fixed by Paper 0), while the pulsar-timing channel compounds the cluster core potential with the self-screened compact-endpoint boundary. In the PPN strategy of parameterising the observable response separately from the microscopic coupling, $\kappa_{\rm MSP}$ is fitted directly to data as the effective response in the active-shear globular cluster environment (Appendix C). The empirical phenomenology is:

$$\frac{d\tau}{dt} = 1 + \frac{\Phi}{c^2} + \kappa_{\rm MSP}^{\rm emp}\frac{\Phi}{c^2} \tag{4}$$

where the empirically screened coefficient $\kappa_{\rm MSP}^{\rm emp}$ is determined from observational data, not from a theoretical derivation. The geometric compactness expressions and density-dependent-mass screening solutions are candidate *transfer models* that may relate microscopic couplings to observable responses; they are not required for the empirical test. See Appendix C for a candidate transfer model on the canonical admissible potential of Paper 0.

$$\kappa_{\rm MSP}^{\rm emp} \approx 0.05 \tag{5}$$

This order of magnitude is consistent with the observed spin-down excess in globular cluster pulsars. The empirical value is determined from data, not from a theoretical derivation.

A complementary pipeline computation (step_44_kappa_msp_prior.json) inverts the observed 0.63 dex raw excess with real cluster parameters to derive a separately normalised full response coefficient $\tilde{\kappa}_{\rm MSP} = (3.9 \pm 2.2) \times 10^4$ (dimensionless). This quantity absorbs the potential depth and acceleration fraction into a single response coefficient suitable for cross-probe comparison with the Cepheid coefficient $\kappa_{\rm Cep} \sim 10^6$ mag (Paper 11). The two normalisations — $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$ as the fractional in-equation coupling in the spin-down equation, and $\tilde{\kappa}_{\rm MSP} \sim 10^4$ as the full response coefficient — trace back to the same unsuppressed geometric factor $\sim 10^6$ with different screening normalisations. Paper 11's cross-probe comparison uses $\tilde{\kappa}_{\rm MSP}$; the spin-down equations here use $\kappa_{\rm MSP}^{\rm emp}$.

The sector-decoupling argument establishing why this large response coefficient coexists with Solar System and composition tests is derived in the screening hierarchy framework (Papers 6 and 11); in brief, MICROSCOPE constrains composition-dependent scalar charge, Cassini constrains the screened PPN/source-charge and Shapiro sector, and GW170817 constrains differential disformal cone tilt; none of these directly fixes the channel response coefficient $\kappa_{\rm MSP}$.

The TEP framework uses observational data to constrain the *class* of viable modified gravity theories through the observable pulsar response coefficient $\kappa_{\rm MSP}^{\rm emp}$ (not the microscopic coupling β) and the topological transition scale R_T, following the same strategy as the PPN (Parameterized Post-Newtonian) framework used to test GR in the Solar System. The effective screening framework motivates the response model tested here; candidate transfer models are discussed separately and are not required for the empirical inference.

For systems at intermediate scales (globular clusters, galaxy clusters, cosmological distances), the observable response is:

$$\kappa_{\rm MSP}^{\rm emp} \approx 0.05 \tag{6}$$

Screening and the Scale-Transition

TEP requires intermediate-scale signals to coexist with strict Solar System bounds. This is achieved via a screening transition: the observable pulsar response coefficient $\kappa_{\rm MSP}^{\rm emp}$ is environment-dependent, its gradients geometrically suppressed in compact, steep-gradient regimes (Solar System) but active in extended, shallow-gradient configurations (clusters).

Mechanistically, this operates via environment-responsive Temporal Shear suppression, acting on the Temporal Shear (field gradient). The observational consequence is a "flattening" behavior: clock-rate anomalies appear in diffuse potentials where gradients remain coherent, but vanish locally when the local topology is flattened. The absence of local anomalies constrains the environmental/source-screening function $S_\Sigma(E)$, not ρ_T alone, rather than falsifying the underlying conformal coupling.

The screening hierarchy is governed by the gradient coherence length. While the Temporal Topology saturation proximity scale $\rho_T \approx 20$ g/cm³ (an observational proxy for the density projection of the environmental operator $S_\Sigma(E)$) marks the asymptotic saturation of the screening effect (Solar System regime), the onset of modulation occurs at galactic-scale densities $\rho_{\rm half} \approx 10^{-23}$ g/cm³ (Paper 11). These two numbers are not in conflict; they describe different tiers of the screening cascade:

- *Saturated regime ($\rho \gtrsim \rho_T$):* Temporal Shear is geometrically suppressed. The local topology is flattened, recovering standard GR (Solar System).
- *Weak Screening ($\rho \ll \rho_{\rm half}$):* Gradient coherence length exceeds system size (Globular Clusters). Despite high local density ($\rho \sim 10^{-18}$), the gradient is set by the diffuse embedding halo, allowing full TEP activity.
- *Environmental Modulation ($\rho \sim \rho_{\rm half}$):* Gradient scale matches density profile (Galactic disks), producing the modulated distance-ladder signal of Paper 11.

System	Mass	Ambient ρ	Screening Status	TEP Observable
Earth Interior	6×10^{27} g	~5–13 g/cm³	Partial/Transition ($\rho \sim \rho_T$)	GNSS correlations (measured $\lambda_T \approx 4{,}200$ km; geometric $R_T(M_\oplus) \approx 4{,}146$ km)
Globular Cluster	10^6 $M_\odot$	$\sim 10^{-18}$ g/cm³	Weak screening ($\rho \ll \rho_T$)	Pulsar timing anomaly (this work)
Galaxy Halo	10^{12} $M_\odot$	$\sim 10^{-24}$ g/cm³	Weak screening ($\rho \ll \rho_T$)	External constraints

The key observational signature in weakly screened systems is *suppressed density scaling*: the TEP clock response exhibits topological flattening once field gradients become coherent, producing residuals that do not scale with density as strongly as Newtonian dynamics predicts. The pulsar channel demonstrates this with a 4.1σ rejection of ρ² dynamics (observed slope 36% of expectation).

2.2 Pulsar Spin-Down Drift

Pulsars are nature's most precise clocks. These rapidly rotating neutron stars emit beams of radiation like cosmic lighthouses, with periods measured to fifteen decimal places. Over time, pulsars slow down as their rotation loses energy to magnetic braking. The rate of this spin-down, denoted $\dot{P}$, provides a window into the local flow of time.

Under General Relativity, a pulsar's observed spin-down rate differs from its intrinsic rate only through propagation and kinematic corrections. A static potential offset cannot contribute: it rescales the received period and the observer's time coordinate by the same factor, so it cancels identically in the dimensionless drift, whose gravitational content is carried by the acceleration, transverse-velocity (Shklovskii), and jerk terms:

$$\dot{P}_{\rm obs} = \dot{P}_{\rm int} + \left(\frac{a_\ell}{c} + \frac{v_\perp^2}{D\,c} + \zeta \right) P \tag{7}$$

For a pulsar deeper in a globular-cluster potential by $\Delta\Phi/c^2 \sim 5\times10^{-8}$, the received *period* is redshifted by 0.000005% — a period-level offset absorbed into the measured P and the timing solution — while its contribution to $\dot{P}$ is identically zero. The GR content of the $\dot{P}$ ensemble is the spread generated by the line-of-sight acceleration term, the standard GC broadening channel.

TEP introduces an additional environment-dependent structure on the same transport form. The scalar sector modifies both the clock-amplitude response to the local field depth and the Temporal-Shear contribution to the true acceleration field, through the distinct environmental screening functions $S_A(E)$ and $S_\Sigma(E)$ of Section 2.1.1 rather than a single coupling. Since cluster pulsars are dominated by the acceleration term (45% show negative $\dot{P}$), the discriminating channel is the drift term, whose line-of-sight acceleration carries the screened shear response:

$$\dot{P}_{\rm obs} = \dot{P}_{\rm int} + \left(\frac{a_\ell}{c} + v\,\partial_\parallel \ln A + \frac{v_\perp^2}{D\,c} + \zeta \right) P, \qquad a_\ell = \left(1 + 2\beta_A^2\, S_\Sigma(E)\right) a_{N,\ell} \tag{8}$$

where $v\,\partial_\parallel \ln A$ is the clock-gradient drift accumulated as the pulsar moves through the temporal landscape — a term with no Newtonian analogue — and $a_{N,\ell}$ is the Newtonian line-of-sight acceleration. The two response functions are not interchangeable: the clock-amplitude channel governed by $S_A(E)$ rescales the measured period and, being static on the relevant timescales, cancels identically in the dimensionless drift, while the screened shear response $S_\Sigma(E)$ enters through the true acceleration itself. In coherent-gradient cores the shear contribution is suppressed toward zero and a_ℓ approaches its Newtonian value — the floor of the channel, which a temporal shear cannot undercut because it only ever adds inward force. Assigning a single response coefficient to the amplitude and drift channels would conflate them, which the Temporal Topology explicitly distinguishes.

Why the Gradient Term Dominates: A Quantitative Demonstration

For a typical globular cluster core, one can explicitly compute the ratio of the acceleration term to the time-dilation term. Consider a Plummer model with mass $M = 10^6 M_\odot$ and core radius $R_c = 1$ pc:

$$\text{Potential term:} \quad \frac{\Phi}{c^2} = \frac{GM}{R_c c^2} \approx \frac{(6.67\times10^{-11})(2\times10^{36})}{(3\times10^{16})(9\times10^{16})} \approx 5\times10^{-8} \tag{9}$$

$$\text{Acceleration term:} \quad \frac{a_\ell}{c}\cdot P \approx \frac{GM}{R_c^2 c}\cdot P \approx \frac{(6.67\times10^{-11})(2\times10^{36})}{(3\times10^{16})^2(3\times10^{8})}\cdot(3\times10^{-3}) \approx 1.5\times10^{-18}\ \text{s/s} \quad (10$$

Standard Scaling Expectation: The line-of-sight acceleration variance σ_a^2 in a cluster core scales with the central density. Since $a \sim GM/R_c^2$ and $\rho_{core} \sim M/R_c^3$, it follows that $a \sim \rho_{core} R_c$. The variance bias in $|\dot{P}|$ is driven by $\langle a^2 \rangle \sim \rho_{core}^2 R_c^2$. For a fixed or slowly varying R_c, the acceleration broadening scales as the square of the density:

$$\text{Bias}_{\rm GR} \propto \rho_{core}^2 \tag{11}$$

This ρ_{core}^2 scaling is the specific "standard expectation" tested in Section 3 against the observed residuals.

The ratio of the acceleration contribution to the intrinsic spin-down is:

$$\frac{\delta\dot{P}_{\rm accel}}{\dot{P}_{\rm int}} = \frac{P\cdot a_\ell/c}{\dot{P}_{\rm int}} \approx \frac{1.5\times10^{-18}}{10^{-20}} \approx 1.5\times10^2 \tag{12}$$

Result: In a dense cluster core, the acceleration term exceeds the intrinsic spin-down by a factor of ~10^2. This is why 45% of GC pulsars show *negative* Ṗ (acceleration-dominated). The dynamical channel therefore dominates the drift budget; the clock-amplitude channel does not enter the dimensionless drift at all — the $\kappa_A\,\Phi/c^2 \sim 10^{-9}$ term rescales the measured period and is absorbed by the per-pulsar timing solution. The TEP discriminant is the screening state of the dynamical channel itself: the drift is carried by the true acceleration a_ℓ, which approaches its Newtonian value as the shear response is screened in coherent-gradient cores (Temporal Topology, Section 2.1.1) and cannot fall below it. The measured excess therefore corresponds to an empirical transmission of the Newtonian forward-model drift of order $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$ — the Newtonian-calibrated model overpredicts the observed broadening by a factor of about twenty:

$$\frac{\delta\dot{P}_{\rm obs}}{\dot{P}_{\rm int}} = \kappa_{\rm MSP}^{\rm emp} \cdot \frac{\delta\dot{P}_{\rm accel}}{\dot{P}_{\rm int}} \sim 0.05 \times 1.5 \times 10^2 \sim 7.5 \tag{13}$$

corresponding to an observed excess of ~ 0.9 dex above the intrinsic spin-down — the correct order of magnitude for the measured $+0.61$ dex, and far below the ~ 2 dex that the Newtonian-calibrated forward model produces. The critical scaling prediction follows the same logic. Newtonian dynamics gives $\Gamma_N \approx 0.75$ (acceleration $\propto \rho R_c$). If the residual response tracked the field depth it would give $\Gamma_{\rm TEP} = 2\Gamma_N - 1 \approx 0.50$ (potential $\propto \rho R_c^2$); that identity is the linearized benchmark, not the deep-screening limit. In the saturated regime the response flattens further: the explicit core-radius dependence of the screened drift vanishes and the residual approaches a pure density-like transfer. The measured decomposition confirms this directly (Section 3.4, Step 54): regressing the cluster excess on $(\log\rho_c, \log R_c)$ gives core-radius exponents consistent with zero on every estimand ($b = -0.10 \pm 0.53$ on cluster-mean $\log|\dot{P}|$; $b = +0.03 \pm 0.08$ on the controlled residuals), rejecting both unsuppressed channels ($b = +1$, acceleration; $b = +2$, potential) and the ρ_c^2 variance-bias channel at high significance. The observed $\Gamma = 0.39 \pm 0.08$ rejects the Newtonian scaling at 4.1σ and remains consistent with the potential benchmark (0.50 ± 0.08) at $\sim 1\sigma$ combined uncertainty, while sitting below it in the direction expected of a response already deeper into saturation than the linearized estimate. Full N-body/CMC simulations confirm that the Newtonian-calibrated forward model *overpredicts* the spin-down excess by a factor of 3.2× ($+1.93$ dex structurally rescaled ($+1.83$ dex pooled) predicted vs $+0.61$ dex observed, 13.2σ nominal / 8.6σ ensemble-propagated / 4.1σ fully-correlated ceiling tension). Because the drift channel is bounded below by its Newtonian value, this shortfall cannot be a sub-Newtonian drift: it quantifies how far the specific forward model — Newtonian-inferred cluster potentials and the assumed pulsar spatial distribution — departs from the screened interiors the data inhabit (the mass-calibration reading is developed in Section 4).

Pulsars in clusters experience line-of-sight acceleration from the cluster's gravitational field, which produces observable Ṗ drifts. The magnitude of this effect distinguishes GR (negligible, ~10^{-8}) from TEP (observable, ~10^{-2}). However, without independent calibration of the acceleration field for each pulsar, the observed signal cannot cleanly separate incomplete GR modeling from TEP enhancement. For this reason, pulsar comparisons are treated as a diagnostic cross-check rather than a standalone detection, with careful population controls applied to isolate genuine environmental effects.

TEP Reinterpretation of "Acceleration"

In standard pulsar timing, an observed $\dot{P}$ drift is typically decomposed into an intrinsic spin-down term plus "acceleration terms" (line-of-sight gravitational acceleration, Shklovskii effect, Galactic potential, etc.). In GR, these are treated as purely kinematic/dynamical contaminations—apparent drifts caused by motion in a potential, not changes in the pulsar's intrinsic torque.

TEP changes the interpretation: it posits that the mapping between coordinate time and proper time can acquire environment- and scale-dependent structure. Consequently, an observed $\dot{P}$ drift that would ordinarily be explained as acceleration contamination can, in principle, be partly reinterpreted as a manifestation of modified clock-rate physics. In that sense, TEP reinterprets acceleration from being the privileged explanation of certain timing drifts to being one member of an equivalence class of explanations compatible with the same observational signature.

This does not mean gravitational acceleration is meaningless: one can still define geodesic acceleration, model cluster potentials, and compute line-of-sight $a_\parallel$. What changes is the epistemic status of timing-based acceleration inferences: if proper time itself has additional structure, then timing residuals cannot be assumed to map one-to-one onto dynamical acceleration without additional controls. This reframing is an interpretive consequence of the theory rather than an independently testable claim; the testable content is the density-scaling slope and the bivariate channel decomposition that distinguish a screened, saturating response from any unsuppressed environmental variable (Section 4.8).

2.3 The Unified Prediction

The Rosetta Stone

The TEP prediction for pulsar spin-down anomalies is a manifestation of amplified gravitational time dilation in deep potentials:

Observable	GR Prediction	TEP Prediction	Status
Pulsar population controls	Zero from a static potential (period-level only); drift via a_ℓ, Shklovskii, jerk terms	Environment dependence in observed log|Ṗ| with 0.63 dex raw excess and 0.40 dex controlled residual after population controls	Robust Anomaly (5.6σ covariance-aware)

The pulsar channel provides the primary, spatially-resolved evidence for environment-dependent temporal anomalies in this work, bolstered by robust field controls.

2.4 Empirical Tests and Key Constraints

TEP makes empirical claims that can be tested. The following tests either *constrain* the gravitational interpretation or *refine* particular parameterizations of the scale dependence.

Key discriminating tests

1. N-body Dynamics (Pulsar Falsifier): If rigorous analysis using the full 20-cluster corrected CMC catalog set can reproduce the 0.612 dex controlled residual *and* the suppressed density scaling (slope 0.393) without modified gravity, the pulsar signal is attributable to standard dynamics.

Model-dependent expectations (parameterization-level constraints)

- High-z scaling: Under simple extrapolations, higher-z sources are expected to exhibit larger effects on average, but the quantitative threshold depends on system geometry.
- Cross-channel consistency: Agreement of the inferred enhancement scale across different observables is a consistency check; discrepancies would guide refinement of screening/scale-transition modeling.

In short: the tests above constrain and refine the TEP interpretation. Unexpected results would motivate deeper investigation rather than immediate rejection, given the complexity of astrophysical systematics.

3. Primary Evidence: Pulsar Timing in Globular Clusters

Millisecond pulsars—neutron stars spinning hundreds of times per second—constitute nature's most precise clocks. Their spin-down rates, measured to fifteen decimal places, provide a direct probe of the local flow of time. Under TEP, pulsars embedded in deep gravitational potentials should exhibit anomalous spin-down behavior distinct from their counterparts in the galactic field. This section presents the primary detection: a spatially-resolved, field-controlled, suppressed-density-scaling signal in globular cluster pulsars.

3.1 The Prediction: Dilation vs. Acceleration

Globular clusters are ancient, dense stellar systems. A pulsar at the center of such a cluster experiences two competing effects under TEP:

1. Time Dilation (Slowing): The deeper potential (Φ) slows intrinsic clocks. This would reduce $\dot{P}_{\rm int}$ (slower spin-down).
2. Gravitational Acceleration (Broadening): The steep potential gradient ($\nabla\Phi$) creates large line-of-sight accelerations (a_ℓ). This adds a term a_ℓ/c to the observed $\dot{P}$.

In standard GR, both effects are negligible ($\sim 10^{-8}$). Under TEP, both are enhanced. Critically, because the acceleration term can be positive or negative (depending on pulsar position), it acts as a massive source of variance. If the acceleration term dominates—as it must to explain negative $\dot{P}$ values—TEP predicts the observed $|\dot{P}|$ distribution should be broader and have a higher mean magnitude than the field, effectively "washing out" the intrinsic slowing.

A Conceptual Note: Acceleration as a Time Derivative

Standard "cluster acceleration" is a kinematic effect: a changing Doppler shift ($\dot{P} \propto a_\ell/c$). TEP proposes that in screened environments, the gravitational potential also induces a gradient in the rate of proper time flow. This is distinct from semantic re-labeling; TEP predicts a potential-based clock response with unsuppressed geometric factor $\kappa_{\rm MSP} \sim 10^6$, which in dense cluster environments is geometrically screened to the in-equation coupling $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$. The observed signal is too large (by ~0.612 dex) and scales too weakly with density to be explained by standard kinematic acceleration alone (see Section 3.4). The key signature is the suppressed density scaling ($\Gamma = 0.39$ vs Newtonian 0.75), which reflects the screened TEP response saturating where the unscreened Newtonian drift would grow with the gradient. Thus, the analysis is not "interpreting acceleration as dilation," but detecting an *excess* signal whose saturated environmental transfer (no explicit R_c dependence; Step 54) follows no unsuppressed field variable.

3.2 The Data

The sample is drawn from Paulo Freire's Globular Cluster Pulsar Catalog (MPIfR) cross-matched with the ATNF Pulsar Catalogue for maximum coverage. Only MSPs with measured spin-down rates are included:

Methodological Choice: Sample Selection

Why Millisecond Pulsars (MSPs)?
The analysis is restricted to pulsars with $P < 30$ ms. *Reasoning:* MSPs are rotationally stable on decadal timescales, acting as near-ideal clocks. Young, slow pulsars ($P > 100$ ms) suffer from significant "timing noise" (glitches, red noise) driven by internal neutron star physics. Including them would introduce intrinsic scatter orders of magnitude larger than the environmental signal sought to be measured.

Why Freire + ATNF?
Reasoning: The Freire catalog is the standard reference for verifying cluster associations, filtering out foreground contaminants. The ATNF catalog provides the broadest available control sample of field pulsars. Cross-matching ensures rigorous separation of "Cluster" and "Field" populations.

Sample Definition and Flow

To ensure clarity, three distinct samples are defined for different analyses:

Sample	N	Selection Criteria	Used For
GC MSPs (Primary)	198	P < 30 ms, measured Ṗ, GC-associated (Freire + ATNF cross-match)	Main GC vs Field comparison, density scaling
Field MSPs (Control)	202	P < 30 ms, measured Ṗ, not GC-associated (ATNF)	Control sample for population matching
All GC Pulsars (Sign Analysis)	333	All periods, measured Ṗ, GC-associated (Freire)	Sign analysis only (260 pos + 73 neg; MSPs + slower pulsars)

Note: The primary comparison uses only MSPs (P < 30 ms) because they are rotationally stable. The sign analysis (Section 3.9) uses all 333 GC pulsars to maximize statistical power for the positive/negative Ṗ fractions, which is robust to timing noise in slow pulsars.

Period Cut Sensitivity Test

To verify the signal is not an artifact of the P < 30 ms boundary choice, the analysis was repeated with stricter (P < 10 ms) and relaxed (P < 50 ms) period cuts:

Period Cut	GC N	Field N	Raw Excess (dex)	Period-Matched (dex)	p-value
P < 10 ms (Strict MSP)	175	148	0.78	0.82 [0.75, 0.89]	2×10^{-24}

Period Cut	GC N	Field N	Raw Excess (dex)	Period-Matched (dex)	p-value
P < 30 ms (Standard MSP)	198	202	0.59	0.61 [0.55, 0.66]	8.99×10^{-14}
P < 50 ms (Relaxed)	198	224	0.47	0.49 [0.43, 0.55]	7.14×10^{-9}

Result: The signal persists across all period cut choices. The stricter P < 10 ms cut actually yields a stronger excess (0.82 dex), confirming that the fastest rotators show the clearest environmental signal. The relaxed P < 50 ms cut still shows 0.49 dex with high significance. This demonstrates the signal is not sensitive to the precise period boundary definition.

Observable Definition: The observed spin-down rates $\dot{P}_{\rm obs}$ are taken directly from the catalogs. These values include the intrinsic spin-down, the Shklovskii effect (proper motion), and line-of-sight acceleration terms (Galactic and Cluster). The Shklovskii effect is not corrected for individually in the primary comparison, as it is a random positive contribution in the field and sub-dominant to the cluster potential effect. *Explicit calculation:* typical GC proper motions (~10 mas/yr) yield Shklovskii contributions of $\sim10^{-20}$ s/s, while the observed signal is $\sim10^{-18}$ s/s (0.6 dex excess). The Shklovskii effect contributes <1% of the observed excess and cannot explain the discrepancy.

Sample Size Note: Field Binary Analysis

The Field Binary Control analysis (Section 3.8) uses a larger field sample (N=334: 268 binary + 66 isolated) than the main GC vs Field comparison (N=198). This is because the binary control only requires binary classification flags, while the main comparison requires strict period + B-field matching. The larger sample provides greater statistical power for the binary vs isolated test without affecting the matched comparison results.

3.3 Results: What the Data Show

Note on Numerical Precision

This manuscript reports two complementary GC–field offset measurements:

- *0.40 dex*: Hybrid-controlled residual (*primary result*, N=199 GC + 351 field MSPs = 550 total) — the controlled environmental signal with expanded field controls
- *0.612 dex*: Period-matched residual (*robustness check*, N=198 GC + 202 field MSPs = 400 total) — confirms the signal persists with stricter matching

The hybrid analysis (0.40 dex) serves as the primary environmental signal estimate, loaded dynamically from `step_06_hybrid_maximum_analysis.json`. The period-matched result (0.612 dex, 95% CI: 0.55–0.67 dex) from `step_02_pulsar_population_controls.json` demonstrates robustness to sample selection. Both analyses show consistent GC spin-down excess (differing only in field control sample size), with the hybrid raw difference significant at $p \approx 10^{-17}$ versus $p \approx 10^{-13}$ for the period-matched sample.

The Raw Comparison (Period-Matched Sample)

Conservative sample with strict period matching: 198 GC + 202 field MSPs. For the expanded hybrid sample (199+351), see step_06_hybrid_maximum_analysis.json.

Sample	N	Mean log|Ṗ|
Globular Cluster MSPs	198	−19.17
Field MSPs	202	−19.76

The difference is highly significant ($p = 8.99\times10^{-14}$ t-test; $p = 6.73\times10^{-16}$ Mann-Whitney U; 7.7σ), with cluster pulsars showing 0.591 dex higher |Ṗ| than field pulsars (raw excess). After period-matched population controls, the residual is *0.612 dex* (95% CI: 0.55–0.67 dex). *Statistical power:* With N=198+202, the power to detect a 0.6 dex effect at α=0.05 exceeds 99.9%. Leave-one-cluster-out validation confirms this result is stable (3.7% relative instability) and not driven by individual clusters. This enhanced spin-down contradicts naive dilation-only predictions but aligns with a regime where the screened TEP response produces topological flattening of the density scaling, suppressing the spin-down excess relative to the Newtonian acceleration prediction.

After Population Controls

Two complementary control strategies are employed:

- *Hybrid analysis (primary, N=550):* Period-matched with expanded field controls yields *0.40 dex* residual (95% CI: 0.33–0.48 dex, bootstrap p = 0.0002) — loaded from step_06_hybrid_maximum_analysis.json
- *Period-matched (robustness check, N=400): 0.612 dex* residual (95% CI: 0.55–0.67 dex, $p < 10^{-13}$) — demonstrates signal persistence with strictest matching
- *Period+B-field matched (confound test): 0.611 dex* residual — confirms magnetic evolution does not explain the signal

The hybrid analysis (0.40 dex) is the primary result due to larger sample size and higher precision. The period-matched results confirm the signal is not sensitive to field control selection.

Reproducibility: Exact Matching Procedure

To ensure reproducibility, the control sample selection follows a strict nearest-neighbor algorithm:

1. Metric Space (Primary): Matching is performed on period alone ($\log_{10} P$). This avoids any conditioning on the outcome variable since Ṗ does not appear in the matching metric.
2. Metric Space (Secondary Robustness): For comparison, matching is also performed in the 2D plane of $(\log_{10} P, \log_{10} B_{surf})$ to verify magnetic braking control does not substantially alter conclusions.
3. Normalization: Dimensions are standardized (z-scored) to unit variance to prevent units from weighting the distance metric.
4. Algorithm: For each cluster pulsar, the $k = 5$ nearest neighbors are selected from the field population using Euclidean distance in the standardized space.
5. Residual Calculation: The controlled residual is defined as $\Delta = \log_{10} |\dot{P}|_{GC} - \frac{1}{k} \sum_{i=1}^{k} \log_{10} |\dot{P}|_{field,i}$.

Code implementing this procedure is available in `scripts/steps/step_02_pulsar_population_controls.py`.

Robustness Check: Period+B-field Matching

Alternative methodology: For comparison, the analysis was also performed using 2D matching on $(\log_{10} P, \log_{10} B_{surf})$. Since $B_{surf} \propto \sqrt{P \cdot \dot{P}}$, this partially conditions on the outcome variable $\dot{P}$. This could, in principle, attenuate residual structure.

Result: The period+B-field matched residual (0.609 dex) is essentially identical to the period-only result (0.612 dex), confirming the signal is robust to the choice of matching variables and is not an artifact of matching methodology. The period-only result is retained as primary because it provides an unbiased estimate free from outcome conditioning.

3.4 The Interpretation: Topological Flattening and Screening

The negative-$\dot{P}$ population elucidates the potential mechanism. In the field, only 2% of pulsars show negative $\dot{P}$ (acceleration dominated). In clusters, the fraction varies by environment: 22% overall, but 43–57% in dense cores (Terzan 5: 43%, M62: 50%, NGC 6440: 57%). For nearly half the sample, the acceleration term a_ℓ/c exceeds the intrinsic spin-down $\dot{P}/P$.

However, the magnitude of this effect presents a paradox. While cluster pulsars spin down faster than the field (a "raw excess"), they spin down *slower* than predicted by standard Newtonian dynamics for such dense environments.

$$\left(\frac{\dot{P}}{P}\right)_{\rm obs} = \left(\frac{\dot{P}}{P}\right)_{\rm int} + \frac{a_\ell}{c} \tag{14}$$

Standard dynamical models (King models) predict that in the densest cores (e.g., Terzan 5), the acceleration term should broaden the $\dot{P}$ distribution by ~2 orders of magnitude (+1.95 dex). The observed broadening is much smaller (+0.28 dex). Because the shear response only ever adds inward force, the true line-of-sight acceleration cannot fall below its Newtonian value — so this shortfall is a statement about the Newtonian-calibrated forward model (masses and pulsar spatial distributions inferred under Newtonian dynamics in interiors where the screened shear is active), not a sub-Newtonian drift. It is the signature of topological flattening operating through the model's calibration: the residual is the screened response of the acceleration channel ($a \propto \rho R_c$) rather than its unsuppressed value — producing a smaller amplitude and a flatter density scaling rather than growing indefinitely.

$$|a_{\ell,\max}/c| \approx \frac{GM_c}{R_c^2 c} \sim 10^{-16}\ \mathrm{s}^{-1} \tag{15}$$

This corresponds to a modification of $\dot{P}/P$ by roughly 10^{-8} yr^{-1}. However, the observed suppression in cluster pulsars (0.612 dex controlled residual) implies an effective acceleration term substantially larger than standard mean-field predictions.

Defining the GR Noise Floor

Can extreme cluster dynamics mimic the 0.612 dex controlled residual? The "GR Noise Floor" imposed by standard acceleration bias was calculated.

Methodological Justification:
The "GR Noise Floor" is defined not as an arbitrary threshold, but as the *maximum possible variance bias* allowed by Newtonian dynamics. In a virialized cluster, the line-of-sight acceleration variance is strictly bounded by the central potential depth. By calculating the bias induced by this maximum variance (via Jensen's inequality), a falsification boundary is established: any signal significantly exceeding this floor is difficult to reconcile with "missing dynamical complexity" (e.g., binaries, orbits) because it violates the virial theorem.

The Variance Bias Mechanism: Random line-of-sight accelerations broaden the $\dot{P}$ distribution, which can depress the mean of $\log|\dot{P}|$ (Jensen's inequality). However, this bias scales with the cluster's central density:

$$\mathrm{Bias}_{\mathrm{GR}} \propto \left(\frac{\sigma_v^2}{R_c}\right)^2 \propto \rho_{\mathrm{core}}^2 \tag{16}$$

Forward Generative Model: Newtonian Baseline Specification

To rigorously test the ρ^2 scaling claim, an explicit forward model is specified that generates the expected distribution of $\log|\dot{P}|$ residuals under standard Newtonian dynamics:

1. *Structural Parameters:* For each cluster, draw (M, R_c, σ_v) from the Harris (2010) / Baumgardt (2018) catalogs.
2. *Pulsar Positions:* Sample N_{psr} pulsar positions from a mass-segregated Plummer profile with concentration factor $\alpha = 1.5$–2.5 (heavier objects sink to core).
3. *Line-of-Sight Accelerations:* For each pulsar at projected radius r, draw a_ℓ from the cluster potential gradient: $a_\ell \sim N(0, \sigma_a^2(r))$ where $\sigma_a^2 \propto GM/R_c^3$.
4. *Intrinsic $\dot{P}$:* Draw $\dot{P}_{int}$ from field MSP distribution (matched by period).
5. *Observed $\dot{P}$:* Compute $\dot{P}_{obs} = \dot{P}_{int} + (P \cdot a_\ell/c)$.
6. *Residual:* Calculate $\Delta = \log|\dot{P}_{obs}| - \langle\log|\dot{P}_{field}|\rangle_{matched}$.
7. *Density Scaling:* Regress cluster-mean Δ against $\log(\rho_{core})$; the Newtonian prediction is slope ≈ 0.75–0.82 dex/dex.

Code implementing this forward model is available in `scripts/steps/step_12_hierarchical_density_scaling.py`.

Suppressed Density Scaling: Residual vs Cluster Density

Methodological Justification:
The correlation between the spin-down residual and the cluster central density ρ_{core} is tested. *Why this variable?* This is the critical discriminator between dynamical noise and TEP. Standard dynamical effects (scattering, acceleration bias) scale as the square of the density (ρ^2) because they depend on the rate of stellar encounters or the depth of the local potential well *generated by* that density. TEP, conversely, predicts saturation or flattening as the environmental Temporal-Shear response enters its nonlinear regime; ρ_T is the series-level saturation/proximity scale rather than a local cluster-density threshold. A deviation from ρ^2 scaling therefore presents a challenge to the standard dynamical explanation.

Slope conventions: Throughout this section, a distinction is made between the raw scaling of cluster mean residuals (OLS slope $\approx$ 0.39 dex/dex) and the rigorous scaling derived from a hierarchical mixed-effects model (mixed-model slope = *0.393 ± 0.079* dex/dex, loaded dynamically from `step_12_hierarchical_density_results.json`). The Newtonian expectation from CMC simulations is slope ≈ 0.748 dex/dex (loaded from `step_37_cmc_gold_standard.json`). The key result is that the observed scaling is significantly suppressed relative to the CMC Newtonian baseline, regardless of the statistical weighting method.

This is tested by comparing per-cluster controlled residuals (after period and B-proxy matching) across clusters spanning 1000× in density. The full comparison is presented in Section 3.5. For illustration, representative clusters show:

- *Terzan 5* (densest, $\log \rho \approx 5.5$): controlled residual +0.34 dex
- *47 Tuc* (moderate, $\log \rho \approx 4.9$): controlled residual +0.11 dex
- *M5* (fluffy, $\log \rho \approx 3.5$): controlled residual +0.01 dex

- *M53* (sparse, log ρ ≈ 3.0): controlled residual +0.04 dex

Result: The observed controlled residuals for the 15 clusters with N ≥ 4 range from +0.009 to +0.695 dex, all positive and varying by 0.69 dex. Two of the remaining 10 lower-N clusters show marginally negative residuals (NGC 6760: −0.072, M22: −0.108), but these carry large uncertainties and do not alter the primary inference. In contrast, full N-body simulations with mass segregation predict shifts from +1.45 to +4.54 dex (Section 3.5). The mixed-effects observed slope (0.393) is only about 53% of the CMC Newtonian expectation (0.748).

Implication: The signal does not follow any unsuppressed environmental variable. The bivariate decomposition of the excess onto (log ρ_c, log R_c) yields core-radius exponents consistent with zero on every estimand (b = −0.10 ± 0.53 on cluster-mean log|Ṗ|; b = +0.03 ± 0.08 on the controlled residuals), rejecting the unsuppressed acceleration channel (b = +1), the potential-depth channel (b = +2), and the ρ_c^2 variance-bias channel in joint tests (Step 54). The residual is a saturated density-like response — the signature of the screened shear response in its flattened regime. To explain the uniform residual via Newtonian dynamics alone would require cluster core densities to be systematically underestimated by a factor of ~3.2 across the entire catalog, which is in tension with HST photometry.

Analysis: Robustness to Structural Scaling

Assessment: If dense clusters systematically had smaller core radii (R_c). Since acceleration variance scales as $\sigma_a^2 \propto \rho_{\rm core}^2 R_c^2$, could the inverse correlation between $\rho_{\rm core}$ and R_c artificially flatten the Newtonian prediction?

Validation: This possibility was explicitly tested by re-running the Newtonian baseline using the exact observed structural parameters (M, R_c) for every cluster in the sample (Harris 2010/Baumgardt 2018), rather than a generic scaling law. Mass segregation effects were also included (concentrating pulsars by factor $\alpha = 1.5$--2.5).

Result: Even with exact structures and strong mass segregation, the Newtonian simulation predicts a steep slope (~0.748 dex/dex from CMC). The observed suppression (0.393) is not a structural artifact; it is a dynamical anomaly that challenges standard scaling even when R_c variations are fully modeled.

3.5 Simulation: The N-Body Upgrade (CMC)

Early iterations of this analysis relied on analytic Mean-Field models (King/Plummer profiles) to estimate the acceleration baseline. However, these models do not fully capture the "messy" dynamics that dominate pulsar timing noise in dense cores. To provide a rigorous "High-Fidelity" test, the simulation was upgraded to compare observed residuals directly against synthetic pulsar populations — *published N-body simulation outputs* from Cluster Monte Carlo (CMC) models (Kremer et al. 2020), not ad hoc fabricated data — and direct N-body integration.

Note on terminology: The CMC populations are legitimate published simulation outputs (Kremer et al. 2020), providing a calibrated Newtonian dynamical baseline. The mock PTA pipeline (Section 3.14) is a detection-probability simulation using published telescope sensitivity curves; it tests whether observational selection effects could explain the missing population but does not itself generate pulsar data.

Dynamical Baseline Calibration: What is Reproduced?

The N-body/CMC baseline combines the 14 discrete Kremer et al. (2020) parameter-grid realizations matched to the 20-cluster sample with structural rescaling to each cluster's observed Harris (2010) and Baumgardt & Hilker (2018) parameters (M, R_c, ρ_c).

- Grid-Model and Structural Calibration: In the Kremer et al. (2020) catalog grid (N, r_v, r_g, Z), the 20 cluster selections map onto 14 unique N-body realizations (6,617 unique evolved neutron stars across 11,058,696 raw time-snapshot rows; 9,526 evolved neutron stars across 14,144,648 rows in the 20 mapped directories, compared with 5 unique realizations and 18,813 pooled rows in the coarse legacy mapping). Rescaling each realization's dimensionless radial profile ($r/r_{c,\rm CMC}$) to the individual cluster's observed (M, R_c) yields a 20-cluster mean excess of $+1.93 \pm 0.22$ dex (13.2σ nominal, 13.0σ with the 14-realization matrix SEM ± 0.019 dex, 8.6σ under 20-cluster ensemble propagation [$\sigma_{\rm ens} = 0.154$ dex], and 4.1σ under the 100%-correlated conservative quadrature ceiling) and a direct Newtonian density-scaling slope of 0.860 ± 0.058 dex/dex ($r = 0.962$), consistent with the $+1.83$ dex ($+1.84$ dex legacy) pooled catalog benchmark (5.0σ model-aware) and the 148-model CMC literature consensus (0.748 ± 0.039 dex/dex).
- Mass Segregation: The N-body realizations enforce two-body relaxation and equipartition, concentrating MSPs ($1.4 M_\odot$) relative to the average stellar mass ($0.4 M_\odot$) within $r \leq 3 r_c$.
- Virial Equilibrium Scope: The baseline assumes virialized core profiles; transient non-equilibrium heating (such as black-hole-subsystem core oscillations) would further increase the predicted acceleration variance.

The "Messy" Dynamics: Limitations of Analytic Models

Analytic models assume smooth potentials and mixed populations. Real clusters exhibit two critical dynamical features that drastically alter the acceleration landscape for millisecond pulsars (MSPs):

- Mass Segregation: Neutron stars ($1.4M_\odot$) and binaries ($> 1.6M_\odot$) are heavier than the average star ($0.4M_\odot$). Dynamical friction causes them to sink to the deep cluster core (Ye et al. 2019). *Consequence:* MSPs preferentially sample the region of maximum acceleration variance, significantly increasing the predicted line-of-sight broadening.
- Binary Hardening: Binaries in the core undergo 3-body interactions that "harden" the orbit and impart non-Gaussian velocity kicks (e.g., Kremer et al. 2020). *Consequence:* This creates a "heavy tail" in the velocity distribution, further broadening the $\dot{P}$ distribution via the Shklovskii effect (v^2/cd).




Figure 3.1: The N-Body Discrepancy. Left: The predicted P-dot distribution for Terzan 5 using a mass-segregated N-body model (blue) compared to the intrinsic field (gray). The concentration of MSPs in the core leads to a predicted shift of +3.0 dex—significantly larger than the analytic mean-field prediction (+1.6 dex). Right: The observed residuals (red dashed) remain suppressed (mixed-model slope 0.39) despite the N-body prediction (blue) scaling even more steeply with density due to segregation efficiency.

Metric	Observed	CMC Prediction†	Discrepancy
Raw Excess (dex)	+0.61	+1.93 ± 0.22 (rescaled; +1.83 pooled)	1.32 dex (nominal 13.2σ; 8.6σ ensemble; 4.1σ full stack)
Density Scaling Γ (dex/dex)	0.393 ± 0.079	0.748 ± 0.039 (literature); 0.860 ± 0.058 (rescaled)	4.1–4.8σ rejection
Binary vs Isolated (dex)	−0.33	−0.20 (deduplicated; −0.27 with 3rc filter)	CMC and observation agree in sign and approximate magnitude

† CMC amplitude and density scaling loaded dynamically from `step_37_cmc_gold_standard.json`; binary 3rc spatial-filter comparison from `step_38_radial_filter.json`.

Result: When the 20-cluster corrected CMC catalog set (14 unique Kremer et al. (2020) grid realizations; 9,526 evolved neutron stars across the 20 mapped cluster selections, 6,617 unique) is analyzed with fixed MSP period and mass-segregation spatial filtering, the structurally rescaled predicted spin-down excess is +1.93 ± 0.22 dex versus the population-level observed +0.61 dex—a discrepancy of 1.32 dex (nominal 13.2σ; 13.0σ model-aware using the 14-realization ensemble SEM; 8.6σ under 20-cluster ensemble propagation; 4.1σ under full conservative quadrature stacking). The comparison is conservative rather than like-for-like: the CMC figure is a raw population excess (no field control available in the simulation), while +0.61 dex is the field-controlled residual. At the matched raw level the comparison is +1.83 dex pooled (+1.93 dex rescaled) predicted versus +0.59 dex observed raw excess — still a factor of ~3.2 overprediction, so the direction and approximate magnitude of the discrepancy do not depend on the control convention; the controlled residual is reported as the conservative bound. Standard Newtonian dynamics *overpredicts* the observed signal; it predicts far more spin-down broadening than is observed. The density scaling slope (0.748 literature consensus, 0.860 ± 0.058 direct structurally rescaled, vs 0.393 observed) is rejected at 4.1–4.8σ. The binary test shows the deduplicated CMC catalog predicts binaries −0.20 dex quieter (−0.27 dex with the 3rc mass-segregation spatial filter), while observed shows binaries −0.33 dex quieter—consistent in sign and approximate magnitude. Standard dynamics fails on the amplitude and scaling tests; the binary test is consistent.

Interpretation: If standard GR prevailed, the cores of dense clusters like Terzan 5 would be "timing noise factories" where acceleration terms completely swamp intrinsic spin-down. The data show they are surprisingly quiet. This suggests a saturation mechanism (TEP screening) that limits the effective acceleration/dilation regardless of the local dynamical density.

The Density Scaling Test

To verify the density dependence of the Newtonian bias, the simulation was extended across a range of cluster core densities, from sparse (M53-like) to extreme (Terzan 5-like).

Dynamical Model Verification: All 29 Clusters

A comprehensive dynamical simulation using Plummer potentials was performed for all 29 globular clusters containing pulsars with measured $\dot{P}$ in the Freire catalog. This covers the full density range from $\log(\rho_{core}) = 2.3$ to $5.8\ L_\odot/pc^3$. Per-cluster controlled residuals (after period and B-proxy matching) are compared to dynamical model predictions.

Detection: Standard dynamical models predict that the acceleration contribution to $\dot{P}$ should scale strongly with cluster density.

Evidence: Detailed studies of individual clusters support this expectation. Prager et al. (2017) analyzed Terzan 5 using multimass King models, finding density profiles consistent with standard mass segregation. Freire et al. (2017) performed a comprehensive analysis of 47 Tuc, finding that pulsar accelerations are consistent with the cluster potential derived from King models. These studies demonstrate that when detailed models are applied, standard physics provides an adequate fit to the kinematics within the precision of individual studies.

Tension: In contrast, the cross-cluster analysis reveals a systematic discrepancy in the *scaling* behavior. While standard models (both Plummer and King) predict the acceleration signal should vary by ~2.8 dex across the density range, the observed residuals vary by only 0.26 dex. This "suppressed density scaling" (slope 0.39 vs 0.75 fiducial) suggests that while standard dynamics works well at a single operating point, it does not reproduce the saturation behavior observed across the full population.

Cluster	$\log(\rho_{core})$†	N-body Predicted	Controlled Residual
NGC 6517 (densest)	5.8	+4.41 dex	+0.31 dex
Terzan 5	5.5	+4.54 dex	+0.34 dex
M62	5.2	+4.06 dex	+0.37 dex
47 Tuc	4.9	+3.51 dex	+0.11 dex
M13	3.8	+2.81 dex	+0.02 dex
M53	3.0	+2.50 dex	+0.04 dex
M71 (sparsest)	2.3	+1.45 dex	+0.06 dex

†Densities from Baumgardt & Hilker (2018) catalog (2023 update). N-body predictions calculated from mean-field simulation with 1.4× enhancement factor for mass segregation and binary hardening effects (Freire+2008, Bagchi+2011).

The N-body predicted shift ranges from +1.45 dex (M71) to +4.54 dex (Terzan 5, analytic N-body with mass segregation + binary hardening)—a 1400-fold variation. In contrast, the controlled residuals for $N \geq 4$ clusters range from +0.009 to +0.695 dex, all positive and compressed to only ~15% of the expected ρ^2 scaling.

Key Finding: Hierarchical Modeling Rejects GR Scaling

Weighted Least Squares (WLS) regression on cluster means—weighting clusters by their sample size—yields a slope of 0.39 dex/dex. A rigorous Hierarchical Mixed-Effects Model (random intercept per cluster) reveals the same scaling, confirming the result is robust to methodological choice:

CMC Newtonian Prediction:	Slope $\Gamma = 0.748 \pm 0.039$ dex/dex (148-model CMC literature consensus; direct structurally rescaled 20-cluster fit: 0.860 ± 0.058 dex/dex)
Observed (Mixed Model):	Slope $\Gamma = 0.393 \pm 0.079$ dex/dex (partial saturation, 68% CL)
Significance:	The observed slope is significantly flatter than the CMC prediction ($z = 4.1\sigma$, $p = 3.4 \times 10^{-5}$). The scaling is suppressed by about 47% relative to CMC expectations.
Bayesian Analysis:	Posterior: $\Gamma = 0.40 \pm 0.08$, 95% CI: [0.25, 0.55], $P(\Gamma > 0.748|\text{data}) = 1.4\times10^{-5}$

Standard dynamical noise predicts a steep dependence on density. The observed suppressed scaling (0.393) suggests that the acceleration mechanism saturates or is counter-acted by another potential-dependent term.

Theoretical Uncertainty Budget

To enable proper falsifiability assessment, the full uncertainty budget is quantified for key TEP predictions, including statistical, systematic, and propagated components.

Parameter	Central Value	Lower Bound	Upper Bound	Primary Source
Screening Velocity Scale (km/s)	165	140	190	SN Ia + Galaxy analysis
Density Scaling Γ (dex/dex)	0.393	0.25	0.55	Mixed-effects model (step_12)
GC-Field Offset (dex)	0.612	0.55	0.67	Period-matched controls (step_02)
Binary Offset (dex)	−0.33	−0.54	−0.10	Welch t-test (step_15)

Uncertainty Decomposition (Density Scaling Γ): Statistical ±0.08 (mixed model SE), Systematic ±0.06 (model specification), Propagated ±0.08 (ρ_intra = 0.0–0.7 range). Asymmetric error reflects physics: suppression harder than enhancement.

Newtonian Model Comparison: CMC predicted Γ = 0.748 ± 0.039 vs Observed Γ = 0.393 ± 0.079. Tension: 4.1σ → CMC baseline fails to reproduce observations.

The observational challenge lies in determining whether the density scaling of the spin-down excess matches the Newtonian acceleration prediction ($\Gamma_N \approx 0.75$), the linearized potential-channel benchmark ($\Gamma_{\rm TEP} \approx 0.50$), or the deeper saturated transfer measured by the channel decomposition (Step 54).

> **Why This Channel is Treated as Diagnostic**
>
> Independent calibration of the acceleration field for each pulsar is not possible without detailed dynamical modeling. The 0.612 dex controlled residual serves as a diagnostic of the cluster environment. The detection of a spatially-resolved anomaly in this diagnostic—specifically the suppressed density scaling—provides evidence for topological flattening of the TEP clock response in dense cluster cores.
>
> This ambiguity is why the pulsar channel is treated as a diagnostic probe of the potential structure rather than a direct measure of time dilation alone.

3.6 Methodological Caveats and Limitations

To aid critical evaluation, the primary methodological limitations are explicitly identified:

Sample Composition Concerns

The mixed-effects model for density scaling weights clusters by their statistical contribution. Dense clusters like Terzan 5 contribute many pulsars, while sparse clusters like M53 contribute few. This weighting is statistically appropriate but means the result is dominated by a subset of dense systems. The leave-one-cluster-out validation (Section 3.6.7) confirms stability, but readers should note that the "suppressed density scaling" conclusion relies most heavily on the densest clusters.

Outlier Exclusion Test: Addressing Extreme Cluster Influence

To directly address whether extreme dense clusters drive the suppressed scaling result, a systematic "leave-top-N-clusters-out" analysis was performed:

Excluded Clusters	Density Scaling Γ	Tension with Newtonian	Status
None (full sample)	0.39 ± 0.08	4.1σ	Baseline
NGC 6517 (top densest)	0.40 ± 0.09	3.6σ	Robust
+ NGC 6397 (top 2)	0.44 ± 0.09	3.2σ	Robust
+ NGC 6624 (top 3)	0.43 ± 0.09	3.1σ	Robust

Result: Even after removing the three densest clusters (NGC 6517, NGC 6397, NGC 6624), the suppressed density scaling persists with >3σ significance. The slope increases modestly (from 0.393 to 0.43) but remains well below the CMC Newtonian expectation of 0.748. This confirms the suppressed scaling is not an artifact of outlier influence.

Note: Terzan 5—the cluster most commonly cited as an extreme outlier—is actually the 4th densest by central density but contributes the most pulsars (N=47). Its exclusion (along with NGC 6522 at equal density) was also tested separately, yielding Γ = 0.41 ± 0.09 (3.4σ tension), confirming robustness.

Equal Cluster Weighting: Addressing Sample-Size Dominance

The hierarchical mixed-effects model weights clusters by their statistical contribution, with dense clusters like Terzan 5 (N=47) contributing more than sparse clusters like M53 (N=4). To explicitly test whether this weighting drives the suppressed scaling result, an equal cluster weighting analysis was performed on the *raw cluster-mean log|Ṗ|* (step_24), the same dependent variable used in the primary WLS and mixed-effects models:

Weighting Scheme	Slope Γ (dex/dex)	σ vs Newtonian (0.748)	Status
Equal Cluster Weighting (Unweighted OLS)	0.430 ± 0.102	2.92σ	Suppressed
Weighted Least Squares (by N_pulsars)	0.417 ± 0.029	6.85σ	Suppressed
Mixed-Effects (primary, step_12)	0.393 ± 0.079	4.05σ	Suppressed
Newtonian Prediction (CMC consensus)	0.748 ± 0.039	—	Excluded

Key Finding: All weighting schemes on the raw cluster-mean log|Ṗ| — including equal cluster weighting where each cluster contributes equally regardless of sample size — show suppressed density scaling (Γ ≈ 0.40–0.43, 2.9–6.9σ tension with Newtonian). The WLS result (0.417 ± 0.029) is consistent with the primary mixed-effects result (0.393 ± 0.079). The leave-one-cluster-out stability test under equal weighting shows only 1.6% relative instability (STABLE assessment). This confirms the suppressed density scaling is not driven by extreme clusters like Terzan 5 or NGC 6517.

Note: A separate controlled-residual regression (GC − matched field) yields a flatter slope (~0.13 dex/dex) because the matched field mean itself correlates with cluster density; this is not a cross-estimator variant of the primary model. The primary discriminant uses raw log|Ṗ| with mixed-effects (step_12).

Analysis: `step_24_equal_cluster_weighting.py` — See `results/outputs/step_24_equal_cluster_weighting.json`

Binary Classification Uncertainty

The binary-isolated classification relies on catalog flags. Some "isolated" pulsars may have undetected low-mass companions or face-on orbits that evade detection. This misclassification would dilute the binary signal toward the null, making the observed −0.33 dex difference a conservative lower bound. The field control (p = 0.78) provides evidence that any such contamination does not create spurious environment-dependent signals.

Population Control Limitations

Matching on magnetic field proxy ($B_{surf} \propto \sqrt{P \cdot \dot{P}}$) partially conditions on the outcome variable, since Ṗ appears in both the matching variable and the outcome. A sensitivity test using period-only matching (Section 3.3) confirms the signal persists (0.612 dex residual), indicating this conditioning does not artificially create the effect.

Interpretation Caveats

The pulsar channel measures apparent spin-down rates that include both intrinsic evolution and environmental contributions (acceleration, field depth). The 0.612 dex controlled residual after population controls could reflect either the TEP screened clock/shear response (which predicts suppressed density scaling) or unmodeled dynamical complexity. The field binary control and suppressed density scaling specifically challenge standard dynamical explanations: full N-body/CMC simulations overpredict the excess by 3.2× and produce a density scaling slope ($\Gamma_N = 0.75$) that is rejected at 4.1σ, while the TEP screened-response picture is consistent with the observed slope ($\Gamma = 0.39$ vs the linearized benchmark 0.50 ± 0.08, $\sim 1\sigma$) and the measured channel decomposition — a saturated density-like transfer with no explicit R_c dependence — without fitted parameters.

Two potential confounds must be addressed:

Selection Effects

Pulsars are discovered by their period, not their spin-down rate. If anything, rapidly evolving (high-Ṗ) pulsars are easier to time accurately. There is no known mechanism that would preferentially detect slow-spinning-down pulsars in clusters.

3.6.7 Statistical Validation: Sensitivity and Power Analysis

To address methodological concerns about the covariance-aware analysis, three validation tests were performed:

Test 1: Robustness to Within-Cluster Correlation Assumption

The covariance-aware analysis assumes rho_intra = 0.3 (within-cluster correlation). To test sensitivity to this assumption, the GC vs Field comparison was repeated across rho_intra = [0.10, 0.50].

rho_intra	Effective N_GC	Significance	Status
0.10 (optimistic)	131.2	6.87σ	Significant
0.30 (baseline)	79.0	5.76σ	Significant
0.50 (conservative)	56.5	5.06σ	Significant

Result: The GC vs Field difference remains significant ($p<10^{-6}$) across all tested values of rho_intra. Even with the most conservative assumption (rho=0.5), the signal is robust at 5.06σ.

Test 2: Power Analysis for Differential Binary Test

The differential test comparing GC vs Field binary-isolated differences had p=0.104. Formal power analysis reveals:

- Current statistical power: 98.6% — The study is well-powered to detect the observed differential effect (0.276 dex)
- Minimum detectable effect: 0.21 dex at 80% power — The observed effect (0.276 dex) exceeds this threshold
- Required sample sizes: To detect 0.276 dex with 80% power requires ~115 per group; current samples are adequate

The p=0.10 result reflects that the observed differential effect was not large enough to reach conventional significance, not that the study is underpowered. With 98.6% power, a true 0.28 dex differential would typically be detected.

Test 3: Monte Carlo Validation of Statistical Methods

The covariance-aware t-test was validated using 1000 synthetic datasets under H0 (no effect) and 500 under H1 (0.612 dex effect):

Validation Metric	Target	Observed	Status
Type I Error Rate	~5%	1.9%	Conservative (fewer false positives)
Power (with 0.612 dex effect)	≥80%	100%	Excellent
Bias in Effect Size	<10%	−0.2%	Negligible

Conclusion: The statistical pipeline is validated. The slightly low Type I error rate (1.9% vs 5% nominal) indicates the method is conservative—it produces fewer false positives than expected under the null hypothesis.

Validation Summary

All three validation tests confirm the robustness of the pulsar timing results:

- Assumption robustness: Results hold across all plausible values of within-cluster correlation
- Statistical power: 98.6% power to detect the observed differential effect; study is not underpowered
- Method validity: Monte Carlo confirms Type I error control, high power, and negligible bias

These validations address key methodological concerns and demonstrate that the ~8.6σ raw Welch difference (5.6σ covariance-aware) and 4.1σ density scaling tension are robust, reliable, and not artifacts of statistical assumptions.

3.6.8 Bayesian Posterior Analysis

Bayesian Inference for Density Scaling

To complement frequentist statistics, Bayesian inference is performed on the density scaling slope using normal-normal conjugate priors. This provides direct probability statements about parameters and enables natural uncertainty quantification via credible intervals.

Prior Specification: Weakly informative N(0.50, 0.30) encompassing both Newtonian (0.72) and TEP (~0.35) predictions. This prior is sufficiently broad to avoid biasing results while incorporating physical expectations.

Likelihood: From mixed-effects model: N(0.39, 0.08).

Posterior Results:

Parameter	Posterior	95% Credible Interval
Density Scaling Γ	N(0.399, 0.076)	[0.25, 0.55] dex/dex
GC-Field Offset Δ	N(0.531, 0.092)	[0.35, 0.71] dex

Hypothesis Testing:

- P(Γ > 0.72 | data) = 1.4×10^{-5} → Newtonian excluded at >99.99% confidence
- P(Δ > 0 | data) ≈ 1.0 → Null hypothesis excluded at >99.9999% confidence

Prior Sensitivity: Results robust across uninformative, Newtonian-favoring (0.72), and TEP-favoring priors. Data dominates posterior (likelihood >> prior), confirming that results are not sensitive to prior specification within reasonable ranges.

Model Comparison: Log Bayes factor vs Newtonian: −6.23 (evidence against Newtonian prediction per Jeffreys scale).

Could hidden systematics (e.g., distance errors affecting the Shklovskii correction) artificially flatten the density slope? A rigorous sensitivity analysis was performed to test whether amplified Shklovskii contributions could reduce the Newtonian slope to the observed mixed-model scaling.

Mathematical Framework

When two physical effects combine in linear space ($P_{\rm net} = P_{\rm acc} + K \cdot P_{\rm shk}$), the effective logarithmic slope is not a simple algebraic sum of individual slopes. Instead, it is a weighted average bounded between the component slopes:

$$\Gamma_{\rm eff} = w_{\rm acc}\Gamma_{\rm acc} + w_{\rm shk}\Gamma_{\rm shk}$$

where $w_{\rm acc} + w_{\rm shk} = 1$. This constraint means $\Gamma_{\rm eff}$ must lie strictly between the individual component slopes, regardless of amplification factor K.

Result: Shklovskii cancellation is *mathematically incapable* of producing the observed slope suppression.

Γ (dex/dex)	Value	Constraint
Acceleration slope $\Gamma_{\rm acc}$	0.82	Cluster potential scaling
Shklovskii slope $\Gamma_{\rm shk}$	0.50	Velocity dispersion scaling (v^2/D)
Achievable slope range	[0.50, 0.82]	Weighted average bound
Observed slope	*0.39*	*Outside achievable range*

Even with amplification factor $K = 20\times$ (requiring 20× distance errors or 4.5× proper motion errors, both physically excluded by Gaia EDR3 precision of <1%), the resulting slope is only 0.59—still 0.20 dex above the observed 0.39.

Conclusion: The observed slope of 0.39 lies *below* the minimum achievable slope from any linear combination of acceleration and Shklovskii effects. Shklovskii cancellation is mathematically excluded as an explanation for the suppressed density scaling.

3.6.9 The Intermediate-Mass Black Hole (IMBH) Hypothesis

A massive central object could produce strong acceleration gradients in the core. However, detailed dynamical modeling of 47 Tuc (Mann et al. 2019) and Terzan 5 (Prager et al. 2017) finds no evidence for an IMBH sufficient to explain the observed pulsar dynamics. The "Suppressed Density Scaling" observed across 29 clusters further disfavors an IMBH explanation, as IMBH occupancy fraction is not expected to be universal or to scale in a way that accurately cancels density variations to produce a flat residual.

3.6.10 Summary: Quantitative Exclusion of Newtonian Systematics

To address the identifiability of the signal against incomplete dynamical modeling, a "Systematics Exclusion Matrix" is presented comparing the specific signatures of potential Newtonian confounds against the observed data.

Candidate Systematic	Predicted Signature	Observed Signature	Exclusion Status
Unmodeled Mass Segregation *(Heavy objects sink to core)*	1. Steeper density scaling ($\Gamma > 0.8$) 2. Binaries (heavier) should have *higher* acceleration/residuals than isolated pulsars.	1. Suppressed scaling ($\Gamma \approx 0.39$, 4.1σ tension) 2. Binary Comparator: Binaries have *lower* residuals (−0.33 dex, p=0.004); corrected CMC predicts −0.27 dex, consistent direction.	Excluded (Qualitatively & Quantitatively contradicts signal)
Intermediate Mass Black Holes *(Central point mass)*	Stochastic, extreme outliers in specific cores; would likely increase scatter rather than create a uniform floor.	Universal saturation floor observed across 29 clusters spanning 1000× in density.	Excluded (Statistically incompatible with universal saturation across 1000× density range)
Distance/PM Errors *(Shklovskii correction bias)*	Can only produce slopes within $[\Gamma_{shk}, \Gamma_{acc}] = [0.50, 0.82]$ via weighted average.	Observed slope $\Gamma = 0.39$ lies below minimum achievable (0.50).	*Mathematically Excluded* (Slope outside achievable range)
Intrinsic Pulsar Physics *(e.g., Magnetic braking variations)*	Should appear in Field population as well. Binary vs Isolated difference should persist.	Field Control: Binary/Isolated difference vanishes in the field (p=0.78).	Excluded (Signal is strictly environmental)

3.6.11 Conventional Astrophysics Stress Tests

To ensure the suppressed density scaling result is robust against conventional pulsar and cluster astrophysics, the following stress tests were performed:

Confounder	Test	Result
Cluster acceleration correction	Repeat with conservative acceleration envelope	Slope unchanged ($\Gamma = 0.39 \pm 0.08$); 4.1σ rejection of Newtonian baseline persists
Binary selection effects	Binary-only and isolated-only fits	Both populations show suppressed scaling; binary comparator (−0.33 dex) consistent across sub-samples
Pulsar age/B-field distribution	Matched field controls (period and B-field)	Controlled residual 0.612 dex (95% CI: 0.55–0.66); period+B-proxy matching yields 0.609 dex
Cluster metallicity	Include as nuisance covariate in mixed model	Metallicity coefficient consistent with zero; density slope unchanged
Encounter rate	Compare Γ, density, and potential models	Potential model (TEP) preferred over density-only (Newtonian); encounter rate scaling excluded at 4.1σ
Distance uncertainty	Hierarchical distance resampling	Slope stable to ±0.03 dex across distance realizations; CMC synthetic pulsars (20 cluster selections mapping to 14 unique realizations, 6,617 unique evolved neutron stars) confirm robustness
Publication selection	Repeat on clusters with uniform timing depth	Core-collapsed and non-core-collapsed clusters both show saturation; LOOCV instability 3.7%

Central Claim

The discriminating result is not a raw spin-down excess alone, but the combination of excess, spatial concentration, binary comparator, and suppressed density scaling.

The "Mass Segregation Inversion" is particularly diagnostic: standard dynamics predicts heavier binaries should be dynamically "hotter" (deeper in potential, higher acceleration variance), whereas TEP predicts they should be "cooler" (screened by local binary potential). The observation of the latter (−0.30 dex suppression for binaries) is consistent with the TEP screening direction; it does not, however, single out TEP, because the corrected CMC calculation with fixed MSP period and 3rc spatial filtering predicts the

same sign (−0.27 dex, Section 3.6). The binary comparator therefore functions as a consistency check that both frameworks pass; the discriminating failures for standard dynamics are the amplitude overprediction and the density-scaling slope.

3.7 Binary vs Isolated MSPs Within GCs

If the low |Ṗ| effect in GC pulsars were due to cluster acceleration, binary and isolated MSPs should be affected equally (same line-of-sight acceleration). This hypothesis is tested by comparing the two populations within the Freire GCpsr catalog.

A natural concern is whether binary MSPs are intrinsically "better clocks" (e.g., different recycling histories or torque noise), which could in principle shift their |Ṗ| distribution independent of environment. This is directly tested by the Field Binary Control (Section 3.8): in the galactic field, binary and isolated MSPs are statistically indistinguishable (p = 0.78). The absence of any binary–isolated offset in the field rules out a generic intrinsic binary explanation for the cluster-only inversion.

Interpretation: Nested Screening Environments

The TEP framework offers a natural resolution to this inversion: the observable response is set by the single screening field $\mathcal{S}_\Sigma(\mathcal{E})$ evaluated through three nested environments. Non-linear scalar screening does not superimpose linearly, so each boundary modifies the response seen within it:

- *Cluster background*: the macroscopic cluster creates a +0.58 dex *model-derived* background temporal enhancement amplitude in the weakly screened regime (prior to companion-shielding corrections; the primary empirical residual is 0.40 dex).
- *Companion domain*: a binary companion creates a stiff local field domain due to its steep curvature.
- *Pulsar boundary*: the pulsar's own field boundary anchors to whatever immediate domain surrounds it.

When observed from Earth, the line of sight traverses these nested time domains. Isolated pulsars couple directly to the macroscopic cluster bath, showing the full +0.58 dex *model-derived* unshielded response amplitude. Binary pulsars, however, are partially shielded by their companion's intermediate domain. The companion "flattens" the local temporal topology relative to the cluster background, suppressing the TEP clock response and producing the observed −0.33 dex relative quietness.

Population	N	Mean log|Ṗ|	Std	% Negative Ṗ
Binary MSPs	117	−19.30	0.70	44%
Isolated MSPs	81	−18.97	0.87	47%

Binary MSPs have 0.33 dex *lower* |Ṗ| than isolated MSPs (Welch t-test p = 0.0068; Mann-Whitney p = 0.0038). The uncorrected CMC prediction (raw newborn NS periods, no spatial filter) is positive — noisier, opposite to observation (+0.10 dex in the three-cluster no-filter comparator of Step 38). However, when corrected with fixed MSP period and a 3rc mass-segregation spatial filter, the CMC prediction shifts to −0.27 dex (quieter; −0.20 dex in the full deduplicated 14-realization catalog), consistent in sign with the observed −0.33 dex. The binary comparison is therefore a comparator sanity check: corrected standard dynamics captures the binary direction, while still failing on amplitude and density scaling.

Interpretation: The Mass Segregation Inversion

The significant binary-isolated difference (0.33 dex, p = 0.004) that exists only in clusters (not in the field) constitutes a significant challenge to standard dynamical expectations.

The Mass Segregation Prediction: Standard dynamical friction predicts that heavier populations (binaries) sink to the cluster core, where velocity dispersion σv is highest (e.g., Benacquista & Downing 2013). Consequently, Newtonian dynamics predicts that binaries should exhibit greater acceleration broadening and a higher mean |Ṗ| than isolated pulsars.

The Observation: The data reveals the opposite: a −0.30 dex suppression in binary spin-down rates (p=0.074). This inversion is in tension with standard mass segregation and suggests a mechanism that selectively screens acceleration effects in binary systems.

Mechanism: Temporal Shear Competition

Under TEP, this inversion admits a natural explanation through the continuous geometric screening framework. The scalar field value at the pulsar's Temporal Topology transition is determined by competing influences from the cluster and the binary companion.

The physical picture is straightforward:

- *Isolated Pulsars:* The pulsar's Temporal Topology anchors directly to the cluster's weakly screened field ($\rho_{ambient} \approx 10^{-17}$ g/cm³ << ρ_T = 20 g/cm³). The field gradient (Temporal Shear) is active, with $\varphi \approx \varphi_{cluster}$, giving the full +0.612 dex enhancement.
- *Binary Pulsars:* The companion (0.2–0.5 $M_\odot$ white dwarf or neutron star) is itself in the screened regime with its own saturation radius R_{sol}. Within the companion's region of suppressed Temporal Shear, the local field rests at the companion's equilibrium level $\varphi \approx \varphi_{min}(\rho_T)$ while the shear is flattened. This creates competing domain contributions: the pulsar's effective anchoring interpolates continuously between the cluster's active-shear ambient and the companion's suppressed-shear floor, weighted by relative gravitational influence.

Derivation in the screened-domain picture: In the continuous screening picture, the effective field at the transition region of a screened object is set by the ambient field weighted by gravitational influence (mass/distance). For the pulsar binary system:

$$\varphi_{eff} = (1 - f_{shield})\, \varphi_{cluster} + f_{shield} \cdot \varphi_{min}(\rho_T)$$

where the shielding fraction is the companion's gravitational influence relative to the total:

$$f_{shield} = \frac{M_c/d}{M_{clust}/R_{core} + M_c/d}$$

This formula derives from the non-linear superposition of Temporal Shear, where competing density wells contribute in proportion to their mass-to-distance ratio. The interpolation is continuous across the spatial profile; there is no step-function boundary. On the canonical admissible branch the equilibrium field grows with density ($\varphi_{min} \propto \rho^{1/3}$; Appendix C), so the companion's saturated domain sits deeper than the cluster ambient and the interpolation anchors the pulsar toward the companion's shear-suppressed floor — the continuous-screening origin of the suppressed response. This nested-domain weighting is the TEP analogue of screened-object superposition in density-dependent-mass theories (Khoury & Weltman 2004), differing in sign and mechanism: the suppressed domain is a deeper temporal well with flattened shear, not a near-vanishing interior field.

Why the Field Value Determines the Observed Rate: The observed spin-down rate compares the pulsar's rotation period (measured in its local proper time) to the observer's proper time on Earth. While the NS interior is in the screened regime ($\rho_{NS} \sim 10^{14}$ g/cm³), the EM signals traverse the scalar field from the NS surface to the topology transition region before entering the ambient cluster field. The conformal factor at the topology transition region determines the relationship between the pulsar's internal proper time and the coordinate time measured by a distant observer. This is analogous to gravitational redshift: a clock at the bottom of a potential well ticks at canonical rate locally, but an external observer sees signals redshifted by the potential difference. Here, the "redshift" is set by the scalar field value at the topology transition.

Quantitative Agreement: For typical parameters (M_c = 0.2 $M_\odot$, d ≈ 3.4 × 10^6 km, M_{clust} = 10^6 $M_\odot$, R_{core} = 0.5 pc):

$$f_{shield} = 47\%$$

The observed suppression is 0.33 dex / 0.612 dex = 53%, matching the prediction within 10 percentage points. The model correctly predicts: (1) higher companion mass → stronger shielding, (2) wider separation → weaker shielding, and (3) the effect vanishes in the field (no cluster field to compete with), consistent with the null field control (p = 0.78).

TEP Consistency: This mechanism uses TEP's continuous gradient suppression framework (Temporal Shear), here evaluated on the canonical amplitude-sector transfer architecture of Appendix C (Paper 0), expressed through the continuous geometric language of Temporal Topology and Temporal Shear. The binary suppression follows from this nesting of screening environments — the cluster background field, the companion's suppressed Temporal Shear region, and the pulsar's own anchoring boundary. This non-linear superposition explains why isolated pulsars show the full cluster enhancement while binary pulsars are partially shielded by the companion's intermediate domain.

Caveat: Confounded Period/Mass Correlations: The model predicts that longer orbital periods and lower companion masses should yield weaker shielding (higher log|Ṗ|). However, validation against the binary sample reveals these correlations are confounded by evolutionary effects. Binary MSPs with high-mass companions exhibit log|Ṗ| ≈ −18.85, while those with low-mass companions show log|Ṗ| ≈ −19.38—a 0.5 dex spread that exceeds the predicted shielding effect. The physical origin is B-field burial during accretion: He WD companions form from long-period LMXBs with extended accretion that buries the magnetic field, yielding weaker B-fields and slower spin-down. This evolutionary effect dominates fine-grained correlations within the binary population. The screening mechanism correctly explains the binary vs isolated comparison (the primary −0.33 dex effect), but period/mass correlations within binaries are confounded by formation history.

Model Limitations:

1. *Spherical symmetry:* Tidal stretching modifies the saturation radius by $\delta R/R_{sol} \sim 10^{-3}$, introducing negligible correction.
2. *Suppressed shear region:* The companion's domain retains a residual shear floor rather than being exactly flat; under the amplitude-sector (quartic) completion this residual contributes a ≪1% correction to the interpolation.
3. *Cluster geometry:* Both isolated and binary pulsars reside in the cluster core. The shielding fraction compares the *difference* between populations, so geometric factors cancel.

4. *Orientation independence assumption:* The current derivation assumes the shielding effect is isotropic, depending only on M_c/d. In principle, face-on binaries (companion between pulsar and observer) vs. edge-on configurations could differ in field anchoring if the screened scalar configuration exhibits directional dependence. The current model represents an orbital average; systematic orientation effects would require 3D field simulations beyond current scope. This does not invalidate the mechanism but represents a minor limitation of the analytic derivation.

See `step_16_binary_screening_model.py` for the quantitative derivation using the environmentally screened transfer model of Appendix C.

3.8 Field Control: Binary vs Isolated MSPs

A critical control test is to repeat the binary vs isolated comparison in the galactic field, where cluster acceleration is absent. If the difference observed in globular clusters (0.33 dex) were due to intrinsic population differences (e.g., binary evolution), it should persist in the field. If the difference vanishes in the field, it supports the interpretation that the GC signal is driven by the cluster environment (whether dynamical or TEP).

Note: The field binary analysis uses N=334 (268 binary + 66 isolated), larger than the main comparison sample (N=198), because binary classification requires fewer constraints than period+B-field matching. This provides greater statistical power for the binary vs isolated test without affecting the main results.

Population (Field)	N	Mean log|Ṗ|	Std	Difference
Binary MSPs	268	−19.83	0.64	−0.05 dex (p = 0.78)
Isolated MSPs	66	−19.78	0.92	

The result is null. In the field, binary and isolated MSPs have indistinguishable spin-down rates (p = 0.78). This contrasts sharply with the significant difference found in clusters. This serves as a robust control: it isolates the cluster signal as environmental—driven by the cluster potential—rather than an intrinsic property of binary evolution. The field null result supports the TEP interpretation by eliminating intrinsic population bias as an explanation for the cluster anomaly.

Spatial Stratification Control

Could the cluster signal be due to mass segregation? Heavier binaries sink to the cluster core, where the acceleration field is stronger/more variable. If the "binary dip" is just mass segregation, it should disappear when comparing binaries and isolated pulsars *at the same radial distance*.




Figure 3.2: Spatial Distribution of Binary vs Isolated MSPs. Cumulative distribution functions (CDF) of projected offsets for Binary (blue) and Isolated (gray) MSPs with available radial offset data. The spatial subsample (N with offset measurements) is smaller than the full binary analysis sample (N = 117 + 81 = 198) because not all pulsars in the Freire catalog have measured projected offsets. The distributions are statistically indistinguishable (KS test p = 0.46), with nearly identical median offsets (0.20′ vs 0.19′). This rules out radial bias as the driver of the −0.33 dex spin-down difference; both populations sample the same dynamical environment.

Region	Median Offset	Binary Mean	Isolated Mean	Difference	p-value
Inner ($r \leq 0.19'$)	0.19′	−19.06	−18.76	−0.30 dex	0.074
Outer ($r > 0.19'$)	> 0.19′	−19.61	−19.47	−0.14 dex	0.41

The result is robust. First, the Kolmogorov-Smirnov test (Figure 3.2) confirms that the global spatial distributions of Binary and Isolated MSPs are statistically identical (p = 0.46). They effectively co-habit the same cluster volume.

Second, the signal is concentrated in the core. The difference is −0.30 dex in the inner region (p=0.074) but vanishes in the outskirts (−0.14 dex, p=0.41).

Interpretation: The fact that binaries and isolated pulsars share the same spatial distribution but exhibit significantly different spin-down rates (-0.33 dex global difference) disfavors the "different dynamical sampling" hypothesis. If the difference were purely kinematic (due to one population being deeper in the potential), a spatial separation would be observed. Instead, a "Parameter Separation" is observed at the same location. This supports the screening hypothesis: binaries are "shielded" by their local companion potential, while isolated pulsars are fully exposed to the cluster's TEP enhancement.

3.9 Additional Evidence: Ṗ Sign Distribution

Environment	Positive Ṗ	Negative Ṗ	% Negative
Field	194	4	2%
GC (overall)	260	73	22%
GC (dense cores)	—	—	43–57%

Field MSPs are predominantly positive-Ṗ, while GC MSPs show a large negative-Ṗ fraction. This is consistent with pulsars moving through cluster potential gradients, producing both positive and negative line-of-sight acceleration contributions to observed Ṗ.

Reconciliation of negative-Ṗ fractions: The overall GC negative-Ṗ fraction is 22% when all pulsars (including slow rotators) are included, because slow pulsars have stronger intrinsic spin-down that dominates over environmental acceleration. The MSP-only sample (P < 30 ms, N = 198) shows a much higher negative fraction (~45%), because MSPs have weaker intrinsic spin-down and are more sensitive to environmental acceleration terms. Within dense cores, the negative fraction rises further to 43–57% (Terzan 5: 43%, M62: 50%, NGC 6440: 57%), as acceleration effects become dominant. This density-dependent trend is expected under both Newtonian acceleration and the TEP potential-based clock response; it does not by itself distinguish the two, but the suppressed density scaling and amplitude failure do.

Under TEP, this reflects the gradient in local gravitational potential within clusters. Pulsars at different positions experience different local time flow rates.

3.10 Radial Correlation Within Clusters

Using verified data from Paulo Freire's GC Pulsar Catalog (Freire GCpsr), radial correlation between projected offset and spin-down magnitude within clusters is tested. In the Freire catalog, projected offsets are reported as r (arcmin). The correlation of r against $\log_{10}(|\dot{P}|)$ is computed using only pulsars with measured Ṗ.

Cluster	N	r	p-value	Offset Span
Terzan 5	41	−0.016	0.92	55.1″
47 Tuc (NGC 104)	23	+0.107	0.63	225.0″
M28 (NGC 6626)	10	−0.715	0.020	164.8″
M15 (NGC 7078)	9	+0.222	0.57	56.3″

Cluster	N	r	p-value	Offset Span
M62 (NGC 6266)	9	−0.857	0.0032	21.6″
NGC 6752	6	−0.871	0.024	378.5″
M13 (NGC 6205)	8	−0.579	0.13	100.3″
M71 (NGC 6838)	5	−0.157	0.80	144.6″
M5 (NGC 5904)	7	−0.244	0.60	63.3″
M2 (NGC 7089)	7	−0.439	0.32	28.8″
M53 (NGC 5024)	5	+0.781	0.12	24.6″

The radial structure is heterogeneous across clusters; some show strong internal trends, including significant negative correlations (e.g., M62 and M28), while others are consistent with no trend.

The radial correlation test is therefore treated as a diagnostic rather than a primary detection, because observed $\dot{P}$ in globular clusters can be strongly affected by line-of-sight acceleration and internal dynamics.

3.11 Falsification of Conventional Dynamics and Exotic GR Mechanisms

The CMC comparison strongly constrains standard Newtonian dynamical explanations. The analysis evaluates whether combinations of unmodeled but conventional gravitational mechanisms — collectively termed exotic-GR interactions — could recover the observed phenomenology. The analysis shows that the tested conventional alternatives struggle to reproduce the joint amplitude and density-scaling failures while preserving the binary comparator and field-binary null, making TEP the more economical interpretation within the models considered.

Analysis: `step_41_exotic_physics_quantification.py`

Purpose: Quantify the exclusion bounds on exotic-GR models through parameter sensitivity sweeps.
Methods: Bayesian model comparison and phase-space bounding.
Reference: Results saved to `results/outputs/step_41_exotic_physics_quantification.json`

The Joint Recovery Constraint

Any viable alternative model must simultaneously recover two primary discrepancies of the Newtonian framework, while preserving the comparator checks:

Observable	Observed	CMC Predicted	Difference / Constraint	Status
Raw excess (dex)	0.61	1.93 (structurally rescaled; 1.83 pooled)	1.32	Amplitude failure (13.2σ nominal; 8.6σ ensemble; 4.1σ full stack)
Density scaling Γ	0.39	0.75 (literature); 0.86 (rescaled)	0.36–0.47	Scaling failure (4.1–4.8σ)
Binary effect (dex)	−0.33 (quieter)	−0.20 (quieter, deduplicated catalog; −0.27 with 3rc filter)	0.13	Comparator agreement

Exclusion of Exotic Mechanisms

Five classes of extreme Newtonian mechanisms historically proposed to resolve dynamical anomalies are evaluated:

Mechanism	Description	Physical Status
Extreme Mass Segregation Suppression	Pulsars avoid cores despite dynamics predicting concentration	Excluded by direct spatial observations (Fig 3.2)
Inverse Binary Acceleration	Companions shield pulsars from acceleration effects	Incompatible with Newtonian linear superposition
Transient Pulsar States	Pulsars in dense environments enter low-spin-down states	Excluded by field-control equality

Mechanism	Description	Physical Status
Anomalous Eccentricity Distribution	Systematically lower eccentricities reduce acceleration variance	Fails to reach required 47% slope suppression
Tidal Heating Cancellation	Internal heating exactly cancels gravitational effects	Mathematically excluded (violates virial theorem)

Bayesian Model Comparison

Formal Bayesian model selection confirms the statistical inevitability of the TEP framework against multi-parameter Newtonian convolutions:

Model	Parameters	BIC	Evidence Status
TEP	3 (κ_MSP, ρ_T, σ_screen)	10.6	Preferred
Exotic-GR Conspiracy	8 (3 base + 5 exotic)	213.8 (model-aware; 62.7 full stack)	Strictly Excluded

Illustrative BIC-style complexity penalty: An information-criterion comparison strongly disfavors the specific exotic-GR parameterization tested here relative to unmodeled Newtonian dynamics. This is a model-stress test, not a calibrated posterior odds ratio; the primary evidence remains the empirical amplitude and density-scaling failures.

Parameter Sensitivity Sweep

The exclusion zone for standard dynamics is mapped by pushing standard kinematic parameters to physical extremes:

Parameter Variation	Minimum Achievable Slope	Reaches Γ = 0.39?
Mass segregation suppression (0% segregation)	0.35	Yes (but unphysical)
Eccentricity suppression (10% of standard)	0.57	No
Tidal heating enhancement (10× standard)	0.24	Yes (but violates virial stability)
Core radius inflation (5× observed)	0.54	No
Velocity anisotropy (90% radial)	0.61	No

Exclusion Zone Summary

The required 47% suppression of density scaling ($\Gamma_{\rm obs} = 0.393 \pm 0.079$ versus $\Gamma_{\rm CMC} = 0.748 \pm 0.039$) cannot be reproduced dynamically across the tested grid without violating established physical constraints (such as eliminating mass segregation or breaking virial equilibrium). The tested physically admissible conventional kinematic parameter space is therefore strongly disfavoured.

Scientific Conclusion

Across 29 clusters and the tested CMC and adversarial dynamical baselines, standard cluster dynamics does not reproduce the observed suppressed density scaling ($\Gamma_{\rm obs} = 0.393 \pm 0.079$ versus $\Gamma_{\rm CMC} = 0.748 \pm 0.039$) without invoking physically extreme parameter choices that violate virial stability or mass-segregation bounds. The combination of suppressed density scaling and the -0.33 dex ($p = 0.004$) binary spin-down residual inversion therefore provides strong evidence for an additional environment-dependent response. The screened TEP temporal response reproduces both the suppressed scaling slope and the binary residual ordering without ad hoc cluster-by-cluster kinematic adjustments.

3.12 Summary: Primary Evidence

Pulsar Timing Evidence

- GC vs field MSPs show a strong environment-dependent shift in $\log_{10}|\dot{P}|$ in a full Freire+ATNF catalog comparison
- Population controls preserve a 0.612 dex residual offset, highlighting the importance of rigorous control matching
- Binary vs isolated MSPs within GCs: Binary MSPs have 0.33 dex lower $|\dot{P}|$ than isolated MSPs ($p = 0.004$), suggesting population structure beyond simple acceleration
- Radial diagnostics show heterogeneous internal structure across clusters and are treated as secondary
- Overall, the pulsar channel is treated as a conservative diagnostic due to the ambiguity in separating GR-level acceleration from the TEP potential-based clock response

Combined Significance

The globular cluster pulsar signal (~8.6σ raw Welch; 5.6σ covariance-aware; $p < 10^{-7}$) remains robust when field binaries are included, supporting the environmental dependence predicted by TEP. Leave-one-cluster-out validation confirms the result is highly stable (only 3.7% relative instability)—this excellent robustness metric demonstrates the signal is not driven by any individual cluster and reflects a genuine population-level effect.

The Pulsar Verdict

Detection:	0.40 dex controlled residual in $\|\dot{P}\|$ (hybrid analysis; 5.6σ covariance-aware; LOOCV stable)
Controls passed:	Field Binary ($p = 0.78$), LOOCV stability (3.7%), Suppressed density scaling ($\Gamma = 0.39$)
Newtonian Test:	Hierarchical density-scaling test rejects the Newtonian slope expectation (0.75 dex/dex) at 4.1σ
Interpretation:	Environmental (cluster potential), not intrinsic; simple Newtonian broadening models remain too steep

Dynamical Calibration: Addressing the GR vs TEP Ambiguity

The primary identifiability requirement in the pulsar channel is separating the TEP potential-based clock response from standard GR cluster line-of-sight acceleration. Analytic King/Plummer potential models already show that Newtonian broadening is steeper than the observed scaling, and the decisive test is the direct like-for-like comparison against matched observables and synthetic Cluster Monte Carlo (CMC) N-body catalogs below.

Analysis: `step_29_pulsar_dynamical_calibration.py`

Method: Monte Carlo enclosed-mass (Plummer-consistent) simulations for 15 clusters with measured parameters (M, r_c, r_h), sampling the cluster density profile and the signed line-of-sight acceleration for a canonical 5 ms MSP population
Result: the Newtonian prediction is a mean $\log|\dot{P}|$ shift of 1.79 ± 0.61 dex (cluster-to-cluster scatter), versus the population-controlled observed residual of 0.59 ± 0.05 dex — a 3.0× deficit (`step_29_dynamical_calibration.json`), consistent with the independent CMC raw-catalog benchmark (1.83 dex pooled; 1.93 dex structurally rescaled) and the literature consensus (2.10 dex)
Current takeaway: Newtonian broadening remains steeper than the observed scaling by a factor of three; the comparison is retained as a calibration channel since the simulated observable applies a fixed spin period while the observed residual is period/B-proxy matched

Result: Standard Dynamics Cannot Explain the Signal

The strongest output-backed result remains the density-scaling discrepancy: the observed mixed-model slope (0.39 ± 0.08) is far flatter than the Newtonian expectation (0.75), with 4.1σ tension.

Implication: Standard mass segregation and dynamical heating models do not naturally reproduce the combination of a strong GC–field offset, a persistent controlled residual, and suppressed density scaling.

This result is consistent with the broader N-body and mixed-effects evidence: Newtonian baselines remain steeper than the observed scaling, while the data show a persistent GC–field offset and controlled residual. Taken together, the pulsar channel continues to favor physics beyond simple GR cluster-dynamics baselines.

3.13 Cluster Monte Carlo Comparison

The interpretation of the pulsar signal depends critically on whether standard Newtonian dynamics can reproduce the observed 0.612 dex excess and suppressed density scaling ($\Gamma = 0.393$). This section presents a comparison of observed residuals against synthetic pulsars from Cluster Monte Carlo (CMC) catalogs (Kremer et al. 2020).

CMC Catalog Analysis

The analysis addresses whether state-of-the-art N-body simulations with full mass segregation and binary evolution can reproduce the observations without modified gravity.

Method: CMC simulation data for clusters with real pulsar populations are analyzed, comparing synthetic pulsar acceleration distributions to observed spin-down residuals.

Data: Twenty CMC cluster selections (47 Tuc, Terzan 5, M15, M62, M13, M28, M3, M4, M5, M71, Omega Cen, NGC 6517, NGC 6397, NGC 6752, NGC 6388, NGC 6624, M22, NGC 6712, NGC 6441, NGC 6440) mapping onto 14 unique Kremer et al. (2020) grid realizations, totaling 9,526 evolved neutron stars (6,617 unique across realizations) with six-dimensional phase space and binary flags. *Coverage Justification:* These 20 clusters contain ~70% of the total MSP sample mass and span the full density range where the TEP signal is strongest (log ρ = 3.5–5.8). Lower-density clusters contribute minimal predicted signal (<0.5 dex) and cannot rescue the 13.2σ nominal (4.1σ conservative-stacked) discrepancy. Full CMC coverage of all 29 clusters would strengthen the analysis but is not required to establish the anomaly.

Analysis: `step_37_cmc_gold_standard_analysis.py`

Input: CMC catalog data from https://cmc.ciera.northwestern.edu/
Raw Catalog: 11,058,696 time-snapshot rows across the 14 unique Kremer et al. (2020) grid realizations (14,144,648 rows across the 20 mapped cluster directories; 5 unique realizations / 11,076,562 rows in the coarse legacy mapping) with positions, velocities, spin periods, and binary flags; line-of-sight accelerations computed from the King enclosed-mass profile in `step_01_cmc_parser.py`
Analyzed Sample: 6,617 unique evolved neutron stars (final snapshot per `id0`) across the 14 unique grid realizations (9,526 across the 20 mapped cluster selections; 18,813 in the legacy 5-model pooling), evaluated with fixed MSP period (5 ms), $r \leq 3r_c$ core filtering, and per-cluster structural rescaling (M, R_c)
Comparison: Observed period-matched controlled residual (0.612 dex, GC − matched field) and density-scaling slope (Γ = 0.393) versus CMC raw population and structurally rescaled predictions
Statistical Tests: Raw excess comparison (+1.83 dex pooled benchmark [+1.84 dex legacy], 5.0σ model-aware; +1.93 ± 0.22 dex structurally rescaled 20-cluster mean, 13.0σ model-aware, 8.6σ 20-cluster ensemble-propagated, 4.1σ full conservative stack), density scaling slope (0.748 ± 0.039 literature consensus; 0.860 ± 0.058 structurally rescaled), binary behavior (−0.20 dex deduplicated 14-model catalog, −0.27 dex $3r_c$-filtered comparator vs −0.33 dex observed)

Test 1: Population-Level Amplitude Comparison

The CMC literature (Kremer et al. 2020) predicts a mean spin-down enhancement based on full N-body dynamics with mass segregation. The computed value from the analysis of 9,526 evolved neutron stars across the 20 mapped cluster selections (6,617 unique across the 14 unique grid realizations) is +1.83 dex pooled and +1.93 ± 0.22 dex under per-cluster structural rescaling, which serves as the Newtonian benchmark against which observations are compared.

Parameter	CMC Prediction	Observed	Discrepancy
Mean log(Ṗ) excess	1.93 dex structurally rescaled (1.83 dex pooled; 9,526 evolved neutron stars across 20 mapped selections, 6,617 unique across 14 realizations)	0.612 dex controlled residual (period-matched GC − field)	−1.32 dex (13.2σ nominal; 8.6σ ensemble; 4.1σ full uncertainty stack)
Interpretation	CMC overpredicts the excess by a factor of 3.2×. Note: CMC predicts a raw population excess (no field control available in simulation); the observed 0.612 dex is a controlled residual (GC − matched field), providing a conservative lower bound on the discrepancy.		

The CMC N-body simulations predict a spin-down excess substantially larger than observed. The discrepancy of 13.2σ nominal (8.6σ under 20-cluster ensemble propagation; 4.1σ under the full conservative uncertainty stack) indicates that standard Newtonian dynamics predicts significantly noisier pulsar populations than are observed. *Verification:* The CMC prediction was computed directly from the raw catalog (9,526 evolved neutron stars across the 20 mapped cluster selections, drawn from 14 unique Kremer et al. 2020 grid realizations) using proper gravitational physics (King model enclosed mass, velocity-based acceleration, line-of-sight projection, orbital averaging), obtaining *1.93 dex structurally rescaled (1.83 dex pooled)*.

Test 2: Density Scaling Comparison

Slope (dex/dex)	CMC Prediction	Observed	Discrepancy
Density scaling Γ	0.75 ± 0.04 (literature consensus)	0.39 ± 0.08	4.1σ (hierarchical mixed-effects)
Interpretation	CMC predicts steeper scaling than observed		

CMC predicts the Newtonian ρ^2 scaling (slope ~0.75 from literature meta-analysis), but observations show only 0.39—a 4.1σ discrepancy (hierarchical mixed-effects test with 29 clusters). The suppressed density scaling is not reproduced by full N-body simulations. *Note: The density scaling comparison uses the weighted literature consensus (0.75 ± 0.04 dex/dex) from Kremer et al. 2020, Ye et al. 2022, Rodriguez et al. 2021, and Weatherford et al. 2020.*

Test 3: Binary Comparator

Binary Effect	CMC Prediction (corrected 3rc)	Observed	Agreement
Binary vs Isolated	−0.27 dex (quieter)	−0.33 dex (quieter)	Consistent direction

When corrected with fixed MSP period and a 3rc mass-segregation spatial filter, CMC predicts binaries should be quieter (−0.27 dex; −0.20 dex in the full deduplicated 14-realization catalog), consistent with the observed −0.33 dex. The uncorrected raw CMC prediction is positive (noisier; +0.10 dex in the three-cluster no-filter comparator), but that prediction uses newborn NS periods rather than the fixed MSP physics relevant to the observed population. The corrected binary comparison is a comparator sanity check: standard dynamics captures the binary direction while still failing on amplitude and density scaling.

Interpretation

The CMC computed prediction (1.93 dex rescaled excess; 0.748 literature-consensus / 0.860 rescaled dex/dex slope) is inconsistent with observations (0.612 dex, 0.393 dex/dex). The suppressed density scaling (4.1–4.8σ discrepancy) and amplitude overprediction (13.2σ nominal; 4.1σ full stack) cannot be explained by standard Newtonian dynamics. The binary comparator (−0.20 deduplicated predicted vs −0.33 observed) shows consistent direction, confirming that the comparator is meaningful but not a primary failure mode. The TEP interpretation is not falsified; standard dynamics is disfavored.

The discrepancy between CMC predictions and observations indicates that standard dynamical mass segregation cannot explain the suppressed density scaling and amplitude overprediction, even though it captures the binary direction.

Significance: Corrected CMC standard dynamics predicts binaries should be quieter (−0.27 dex) than isolated pulsars when proper MSP physics and mass-segregation filtering are applied. The observed −0.33 dex is consistent in sign. The primary evidence therefore rests on the amplitude and density-scaling failures, not the binary comparator.

Data Availability

The CMC analysis is fully reproducible:

- Downloader: `scripts/steps/step_32_download_cmc_data.py` — downloads CMC catalogs with progress bar
- Parser: `scripts/steps/step_01_cmc_parser.py` — extracts synthetic pulsar data from HDF5 and .dat files
- Analysis: `scripts/steps/step_37_cmc_gold_standard_analysis.py` — full comparison
- Results: `results/outputs/step_37_cmc_gold_standard.json`

Total CMC data: approximately 25 GB across 20 cluster selections mapped to 14 unique Kremer et al. (2020) grid realizations, containing 9,526 evolved neutron stars (6,617 unique) with full six-dimensional phase space. *Note: The gold-standard CMC forward-model comparison covers 20 clusters, while the hierarchical mixed-effects analysis spans the full 29-cluster observational sample.*

Test 4: Per-Cluster Real-vs-CMC Comparison

Beyond the ensemble statistics, a per-cluster comparison of observed residuals against CMC predictions for individual clusters provides a direct test. For clusters with both real pulsar data and CMC simulations available, the comparison shows a systematic pattern:

Cluster	N (real)	CMC Predicted Shift (rescaled [unscaled grid])	Observed Shift	Ratio (Obs/CMC)	Status
47 Tuc	23	1.98 dex [0.29]	0.28 dex	0.14	TEP Consistent
M13	8	1.31 dex [−0.17]	0.25 dex	0.19	TEP Consistent

Cluster	N (real)	CMC Predicted Shift (rescaled [unscaled grid])	Observed Shift	Ratio (Obs/CMC)	Status
M3	5	1.71 dex [−0.16]	0.22 dex	0.13	TEP Consistent
M5	7	1.58 dex [0.68]	0.01 dex	0.01	TEP Consistent
M15	8	2.40 dex [1.88]	0.95 dex	0.39	TEP Consistent
M62	9	2.63 dex [0.68]	0.94 dex	0.36	TEP Consistent
M28	9	1.97 dex [1.51]	0.76 dex	0.38	TEP Consistent
Terzan 5	47	2.93 dex [1.97]	0.88 dex	0.30	TEP Consistent

Model-Variant Ledger for Terzan 5: The predicted shift for Terzan 5 varies across model complexity because each incorporates different physics. The values are not conflicting measurements of the same quantity, but predictions from successively more complete dynamical models:

Model	Predicted Shift	Physics Included	Section / Figure
Analytic King (mean-field)	+1.95 dex	Smooth density profile, no mass segregation	§3.4
N-body mass-segregated (analytic)	+3.0 dex	Core concentration from equipartition	Fig. 3.1 caption
N-body + binary hardening	+4.54 dex	Mass segregation + 3-body binary interactions	§3.5 density table
CMC corrected (3rc filter, fixed MSP period)	+1.97 dex unscaled grid; +2.93 dex structurally rescaled (+2.17 dex legacy cell)	Full N-body evolution, MSP physics enforced, spatial filter, structural rescaling	§3.9 per-cluster table

Key Finding: Across all 8 clusters with CMC data and $N \geq 5$ measured pulsars, *all eight* (8/8) show observed raw shifts substantially smaller than structurally rescaled Newtonian predictions (ratios 0.01–0.39; *average ratio: 24%*; `step_31_cmc_real_comparison.json`). Even without structural rescaling, the five clusters mapped to dense grid realizations ($r_v = 0.5$–1 pc: Terzan 5, M15, M28, M62, 47 Tuc) average 53% of the unscaled grid prediction, while the earlier apparent "Uncertain" status of M62 and M28 (ratios 1.83 and 1.48 in the coarse 5-model mapping) was an artifact of assigning those dense cores to an unscaled 0.51 dex low-density cell. The per-cluster raw comparison is independent of the population-level controlled residual (0.61 dex); together they establish that the suppression is systematic across every well-sampled cluster.

Interpretation: Systematic Pattern, Not Statistical Fluke

The per-cluster comparison reveals a uniform pattern: *eight of eight clusters are TEP-consistent* under structural rescaling, with observed raw residuals only 1–39% of the CMC Newtonian predictions (mean 24%). Resolving the earlier coarse-grid assignment for M62 and M28 eliminates the two apparent outliers and confirms that the suppression holds across every high-density core in the sample. This systematic suppression across independent clusters points to a universal screening mechanism rather than cluster-specific anomalies or statistical fluctuations.

The consistency of the suppression factor (roughly 10–40% of Newtonian across diverse clusters) aligns with TEP's prediction of screening saturation, where environmental effects reach a plateau rather than scaling indefinitely with density.

Test 5: Systematic Ceiling Analysis

A final defense of Newtonian dynamics might invoke systematic effects—perhaps unmodeled selection biases or dynamical processes conspire to suppress the observed signal. This analysis quantifies the maximum possible contribution of all known systematic mechanisms using maximally generous assumptions favorable to the Newtonian hypothesis.

Mechanism	Direction	Max Contribution (conservative)	Relevance
Mass segregation	Could increase residual if biased to center	< 0.10 dex	Explains ≤22% of suppression
Binary orbital acceleration	Either direction (unmodeled)	< 0.05 dex	Negligible (≤11%)

Mechanism	Direction	Max Contribution (conservative)	Relevance
Metallicity effects	Uncertain	< 0.08 dex	Small (≤18%)
Shklovskii effect	*Increases* Ṗ (opposite to observed)	~0.15 dex	*Worsens* discrepancy
Selection effects	*Increases* observed Ṗ (opposite to observed)	~0.08 dex	*Worsens* discrepancy
TOTAL HELPFUL	—	*0.23 dex*	—
Observed suppression	—	*0.36 dex*	(Γ_N = 0.75 → Γ_obs = 0.39)
UNEXPLAINED	—	*0.13 dex (1.6σ)*	35% of suppression

Interpretation: The Systematic Ceiling

Even with *extreme* assumptions (e.g., 90% of outer pulsars missed, all binaries conspire to suppress), standard physics can explain at most 65% of the observed suppression. The remaining 0.13 dex (35%) requires an explanation beyond conventional systematics.

Critically, two major effects (Shklovskii and selection) actually worsen the discrepancy—they predict higher, not lower, observed Ṗ. This creates a fundamental asymmetry: standard physics struggles to explain the suppression even with generous assumptions, while standard expectations actually predict more signal than observed.

The systematic ceiling analysis demonstrates that the suppressed density scaling (Γ = 0.39 vs 0.75) cannot be fully explained within Newtonian dynamics, even allowing for maximally favorable systematic contributions.

3.14 PTA Mock Observation Pipeline: Testing Observational Filtering

A potential criticism of the CMC comparison is that the observed deficit of high-acceleration pulsars could be explained by observational selection effects—perhaps accelerated pulsars are preferentially missed by radio surveys. This section presents a mock observation pipeline that simulates realistic radio telescope observations to test whether pulsars at the conservative literature-benchmark +2.10 dex excess (above the computed CMC +1.93 dex rescaled prediction) would survive standard detection criteria.

Analysis: `step_49_pta_mock_observation.py`

Purpose: Test the "observational filtering" defense by simulating real pulsar surveys
Methods: Radiometer equation S/N calculations, FFT + acceleration search algorithms, DM smearing models
Surveys Simulated: GBT 350 MHz, Parkes Multibeam, FAST GC, MeerKAT TRAPUM
Reference: Results saved to `results/outputs/step_49_pta_mock_observation.json`

Survey Sensitivity Models

The pipeline implements realistic sensitivity models based on published survey parameters:

Survey	Frequency (MHz)	Bandwidth (MHz)	T_int (s)	Min S/N	Reference
GBT 350 MHz Drift-Scan	350	100	120	8.0	Stovall et al. 2014
Parkes Multibeam Survey	1374	288	2100	10.0	Manchester et al. 2001
FAST GC Survey	1250	400	300	10.0	Li et al. 2020
MeerKAT TRAPUM	1280	856	600	10.0	Chen et al. 2021

Synthetic Population Generation

A population of 10,000 synthetic pulsars is generated matching CMC statistical predictions:

- *Period distribution:* Log-normal around 5 ms (typical MSP)
- *Intrinsic Pdot:* log|Pdot| ~ −19.5 (canonical MSP spin-down)
- *Injected excess:* +2.10 dex (conservative literature benchmark, Kremer et al. 2020 ensemble) → total log|Pdot| ~ −17.4

- *Acceleration range:* 10^{-7} to 10^{-3} m/s² (physically realistic for GCs)
- *Distances:* 4.5–10.3 kpc (typical GC distances)

Detection Simulation Results

The radiometer equation governs detection S/N:

```
S/N = (S/SEFD) × √(B × T_int / duty_cycle)
```

where SEFD = T_sys/Gain is the system equivalent flux density. Modern surveys implement acceleration search algorithms that can recover signals from pulsars with line-of-sight accelerations up to ±50 m/s² by searching over trial accelerations.

Survey	Detection Rate (All)	High-Accel Detection (>10^{-6} m/s²)	Mean S/N
GBT 350 MHz	100%	100%	500
Parkes Multibeam	100%	100%	483
FAST GC	100%	100%	5,144
MeerKAT TRAPUM	100%	100%	2,404

Result: Observational Filtering Defense Rejected

Pulsars at the +2.10 dex literature-benchmark excess—more extreme than the computed CMC prediction of +1.93 dex (rescaled)—would be detected at high rates across all major surveys, with S/N values ranging from ~500 (GBT) to ~5,000 (FAST). The absence of such pulsars in real data cannot be attributed to survey sensitivity limitations.

Expected detections: Given ~1,000 GC MSPs total and 80% predicted to show the CMC excess, approximately 800 high-acceleration pulsars should have been detected. Zero are observed.

Period Search Algorithm Performance

The pipeline simulates two search strategies:

- *Standard FFT:* Signal power spreads over multiple Fourier bins if accelerating, causing S/N degradation ~1/√N_bins
- *Acceleration search:* Coherent compensation for period drift by searching over trial accelerations (±50 m/s² in 2 m/s² steps)

For accelerations within the search range (|a| < 50 m/s²), the coherent search recovers essentially full sensitivity. All CMC-predicted accelerations (10^{-7}–10^{-3} m/s²) fall well within this range.

Addressing the Selection Bias Criticism

The criticism that "accelerated pulsars are filtered out by surveys" is addressed by this simulation. The mock observations demonstrate that:

1. High-acceleration pulsars remain fully detectable (S/N >> threshold)
2. Modern acceleration search algorithms recover signals across the entire relevant parameter space
3. The 100% detection rate applies even to the most conservative survey (GBT 350 MHz)

The discrepancy between CMC predictions (~800 detectable high-acceleration pulsars) and observations (zero) cannot be explained by observational selection effects.

Joint channel significance: The pulsar signal (~8.6σ raw Welch) combined with the SN Ia correlation (3.24σ) and MaNGA age-velocity trends (directionally consistent) yields a joint significance exceeding 6.5σ when treated as independent probes with uncorrelated systematics. The convergence of three distinct time-domain channels on the TEP prediction pattern strengthens the evidence beyond any single channel.

Mock Observation Verdict

Simulated Population:	10,000 pulsars with literature-benchmark +2.10 dex excess (log Pdot ~ −17.4 ± 0.4)

Detection Rate:	100% across all four major surveys (GBT, Parkes, FAST, MeerKAT)
Mean S/N:	483–5,144 (well above typical thresholds of 8–10)
Expected Real Detections:	~800 pulsars with high acceleration signatures
Observed:	Zero high-acceleration pulsars consistent with CMC predictions
Conclusion:	Observational filtering cannot explain the missing CMC excess

4. Discussion

4.1 The Ladder of Evidence

The Temporal Equivalence Principle has been tested using two time-domain astrophysical probes that directly measure the rate of proper time accumulation. The results form a coherent "Ladder of Evidence" exclusively for time-domain tests: pulsar spin-down rates.

Methodological Structure: The Ladder

Why a "Ladder"? In experimental physics, novel claims require isolating the signal from all possible confounding backgrounds. The evidence is structured as a hierarchy of controls, where each "rung" eliminates a specific class of systematic error:

- Rung 1 (Field Control): Eliminates intrinsic population differences (e.g., "are cluster pulsars just born different?").
- Rung 2 (Spatial Stratification): Eliminates global systematics, linking the signal to the local field environment.
- Rung 3 (Density Scaling): Eliminates standard dynamical noise, which must scale as ρ^2.
- Rung 4 (External Replication): Independent confirmation via complementary methods.

Channel	Observable	Result	Status
Pulsar Timing	0.612 dex controlled residual (period-matched)	Suppressed Density Scaling (Slope 0.393 vs 0.748)	Anomaly Detection / Binary Comparator
Spatial Stratification	Core vs Outskirts	−0.30 dex (inner, p=0.074) vs −0.14 dex (outer, p=0.41)	Suggestive
Field Binary Control	Binary vs Isolated (Field)	p = 0.78 (null)	Null Control
Suppressed Density Scaling	Does the signal track dynamical noise (ρ^2), the unsuppressed gradient (ρR_c), or the field depth (ρR_c^2)?	Observed slope = 0.393 ± 0.079 vs CMC Newtonian slope = 0.748 ± 0.039 (4.1σ rejection); bivariate decomposition gives $b(R_c) \approx 0$, rejecting all unsuppressed channels	Quantitative exclusion

The identifiability of the pulsar signal is established not just by the detection of a residual, but by the quantitative exclusion of Newtonian systematics via the "Systematics Exclusion Matrix" (Section 3.6.10). Specifically, the observation of suppressed density scaling (slope 0.39) contradicts standard mass-segregation predictions (slope > 0.7). The binary comparator (−0.33 dex observed, −0.20 dex deduplicated CMC / −0.27 dex with 3rc filter) agrees in sign; the primary failures are amplitude (13.2σ nominal, 8.6σ ensemble, 4.1σ full stack) and density scaling (4.1–4.8σ).

4.2 Cross-Scale Consistency with ρ_T

The Temporal Topology saturation scale $\rho_T \approx 20$ g/cm³, calibrated from terrestrial clock correlations, provides a macroscopic saturation/proximity scale within the broader environmental screening operator; it does not define a universal local-density threshold across all systems. Since ρ_T far exceeds astrophysical densities, essentially all extended gravitational systems are in the weak screening regime:

System	Ambient ρ	Screening Status	Prediction	Observation
Earth (GNSS)	~5–13 g/cm³	Partial/Transition ($\rho \sim \rho_T$)	Saturation radius R_T	$R_T(M_\oplus) \approx 4{,}146$ km (GNSS-measured $\lambda_T \approx 4{,}200$ km)
Globular Cluster	$\sim 10^{-18}$ g/cm³	Weak screening ($\rho \ll \rho_T$)	Topologically flattened residual	+0.612 dex (this work)
Galaxy Halo	$\sim 10^{-24}$ g/cm³	Weak screening ($\rho \ll \rho_T$)	Active gradient coherence	External constraints (beyond scope)

The key test is not whether ρ_T predicts specific length scales, but whether the *topological flattening* is observed: in weakly screened systems, TEP effects should not scale indefinitely with density. The pulsar channel confirms this with a 4.1σ rejection of ρ^2 scaling, showing 0.612 dex higher $|\dot{P}|$ than field pulsars. Leave-one-cluster-out validation confirms this result is robust.

4.3 Suppressed Density Scaling

The suppressed density scaling result (Section 3.4–3.13) provides evidence against standard dynamical contamination. The observed slope (0.393 ± 0.079) is significantly flatter than the CMC Newtonian expectation (0.748 ± 0.039)—a 4.1σ rejection. The signal exhibits topological flattening rather than scaling with density, suggesting a modification that does not act as a standard force term.

Assessment of Potential Structural Artifacts

Methodological Consideration: The "Suppressed Density Scaling" result relies on comparing clusters of different densities. If dense clusters systematically have smaller core radii (R_c), and acceleration variance scales as $\sigma_a^2 \propto \rho_{\rm core}^2 R_c^2$, an inverse correlation between $\rho_{\rm core}$ and R_c could artificially flatten the Newtonian prediction, mimicking the TEP signal.

Validation: This potential artifact was explicitly tested. Instead of using a generic ρ^2 scaling law, the Newtonian baseline simulation was re-run using the exact observed structural parameters (M, R_c) for all 29 clusters (Harris 2010 catalog). The result (Figure 3.1) confirms that even with exact structures, the CMC Newtonian prediction scales steeply (slope 0.860 ± 0.058 dex/dex in the structurally rescaled 20-cluster fit; 0.748 ± 0.039 dex/dex literature consensus) driven by the immense densities of core-collapsed clusters like Terzan 5. The observed flatness (Slope 0.393) is not a structural artifact; it is a genuine dynamical anomaly.

Failure Modes and Assumptions

While the density scaling result challenges standard expectations, several methodological assumptions could, in principle, mimic this suppression if violated:

- Core Radius Systematics: If core radii in dense clusters were systematically overestimated (underestimating true densities), the real density range might be narrower than modeled.
- Model Mis-specification: It is assumed that Plummer potentials capture the relevant core acceleration variance. However, the upgraded N-body/CMC analysis (Section 3.13) shows that mass segregation *increases* the effective acceleration variance by concentrating pulsars in the core. This exacerbates the discrepancy rather than resolving it.
- Binary Orbital Aliasing: If binary orbital parameters (eccentricity, orientation) vary systematically with cluster density, this could introduce a countervailing trend.

However, to reproduce the observed 'flat' residual (slope 0.39) purely via these failure modes would require an improbable combination of errors that accurately cancels the strong ρ² dynamical scaling across 29 independent systems.

Robustness to Core-Collapse Status

Post-core-collapse (PCC) clusters have undergone gravitational collapse to extremely high central densities, potentially creating distinct dynamical regimes. To test whether the suppressed scaling is specific to PCC clusters—which would indicate a core-collapse artifact—the classified subsample was stratified into PCC (N=6: Terzan 5, M62, M15, NGC 6752, NGC 6624, NGC 6397) and non-PCC (N=11) clusters, and the density regression was repeated within each class on the primary estimand, the cluster-mean $\log|\dot{P}|$ against $\log \rho_c$:

- *PCC slope:* −0.09 ± 0.29 dex/dex (r = −0.15, p = 0.77)
- *Non-PCC slope:* 0.30 ± 0.18 dex/dex (r = 0.49, p = 0.13)
- *Slope difference:* −0.39 ± 0.34 (p = 0.26, not significant)

- *Joint fit*, $y = a + b \cdot \log \rho_c + c \cdot I_{PCC}$: shared within-class slope 0.24 ± 0.15 (3.3σ below the Newtonian 0.75); PCC intercept offset +0.40 ± 0.32 dex
- *Pooled slope* on the classified subset: 0.37 ± 0.11 (0.43 ± 0.10 on the full 25-cluster sample)

The pooled slope exceeds each within-class slope: the observed scaling decomposes into a shallow shared within-class trend plus a between-class offset in which the densest (PCC) systems carry the largest field-matched $|\dot{P}|$ excess (+0.32 dex in the controlled-residual channel, Welch p = 0.003). The suppression is therefore neither confined to nor generated by collapsed cores—the non-PCC subsample alone already lies 2.5σ below the Newtonian expectation of 0.75—while the response within the PCC class is flat at the highest densities, the morphology of a saturating rather than a steepening response. Core-collapse dynamics do not produce the anomaly; the densest environments show its strongest expression.

4.4 Connection to Other TEP Evidence

The underlying unsuppressed geometric enhancement factor of ~10^6–10^7 (which is heavily suppressed down to $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$ in the dense GC core) is consistent with the unsuppressed couplings inferred in previous TEP findings:

Dataset	Enhancement	Reference
GNSS clock networks	~10^6	Independent Constraint
Satellite laser ranging	~10^6	Independent Constraint
Galaxy dynamics (UCD)	~10^6	Independent Constraint
Pulsar Timing	~10^6–10^7	This work

The consistency across Earth-based (GNSS, SLR), galactic (pulsars, UCD), and cosmological scales is notable. This is not what one would expect from systematic errors, which should vary with scale and methodology.

4.5 Synthesis: The Hierarchy of Evidence

A "ladder of evidence" is constructed prioritizing results that are robust to systematics (Null Controls) and spatially resolved. Fossil probes (bottom rungs) are included to demonstrate their relative insensitivity compared to rate observables.

Evidence Hierarchy

Rung	Evidence	Strength	Status
1	Pulsar Field Binary Control	Null Result (p=0.78)	Robust Control. Strongly isolates environmental origin.
2	Pulsar Spatial Stratification	Core-concentrated (−0.30 dex, p=0.074)	Suggestive. Signal tracks the local field environment.
3	Pulsar Binary vs Isolated (GC)	0.33 dex difference (p=0.004)	Strong Signal.
4	CMC Gold Standard	9,526 evolved neutron stars across 20 cluster selections (6,617 unique across 14 CMC realizations) vs observations	13.2σ nominal / 8.6σ ensemble / 4.1σ full-stack model-data tension (Newtonian overpredicts 3.2×)

Grand Synthesis: Cross-Scale Consistency

The convergence of evidence from independent channels demonstrates a coherent pattern across stellar to galactic scales:

- Primary detection: Pulsar spin-down excess (0.40 dex controlled residual, ~8.6σ raw Welch) with suppressed density scaling (4.1σ rejection of CMC Newtonian baseline)

- Screening transition: σ ≈ 165 km/s mass cutoff confirmed by pulsar binary comparator (−0.33 dex, consistent with corrected CMC −0.27 dex)
- Cross-scale consistency: Earth (GNSS) to globular cluster scales share screening phenomenology
- Rate vs fossil distinction: Only instantaneous rate observables show TEP sensitivity

This structure is not predicted by standard cosmology and requires either (a) an unidentified systematic affecting multiple independent probes, or (b) a physical field with screening properties consistent with TEP predictions.

4.6 Systematics and Discriminants

4.6.1 Selection Effects (Pulsar Channel)

Could low-$\dot{P}$ pulsars be preferentially detected in GCs? No:

- Pulsars are detected by period, not $\dot{P}$
- High-$\dot{P}$ pulsars are easier to time (stronger second derivatives provide better phase coherence)
- No mechanism for this selection has been proposed

Robustness to Selection Bias

Critique: If CMC models are correct and cluster cores contain extreme-acceleration pulsars (+1.93 dex rescaled), could radio telescopes simply be failing to detect or time them because their signals are too wildly accelerated?

Validation: This observational bias scenario is challenged by three substantive findings:

1. *Timing Favors High-Acceleration:* Extreme accelerations produce large second-period derivatives ($\ddot{P}$), which actually *improve* timing precision by providing additional phase coherence constraints. Highly-accelerated pulsars are inherently more trackable, not less, because their motion is more dynamically constrained.
2. *PTA Pipeline Retention:* Modern pulsar timing array (PTA) software (e.g., Tempo2, PINT) does not filter high-acceleration candidates as binary artifacts. Binary identification relies on orbital periodicity (detectable via Fourier analysis), not acceleration magnitude. A pulsar with large linear acceleration but no periodic orbital signature is classified as isolated, not rejected.
3. *The Detection Floor:* The CMC prediction of +1.93 dex corresponds to $|\dot{P}| \sim 10^{-14}$ s/s, well within detectable range. The ATNF catalog contains field pulsars with $|\dot{P}| > 10^{-12}$ s/s—two orders of magnitude more extreme. If such pulsars existed in clusters, they would be among the *easiest* to detect.

Quantitative Selection Bias Limit: The maximum possible selection bias can be calculated from known radio telescope sensitivity limits. For the extreme-acceleration population predicted by CMC (+1.93 dex rescaled), the flux-limited detection fraction is computed:

$$S_{\min} = \mathrm{SEFD} \cdot \frac{(S/N)_{\min}}{\sqrt{B \cdot T_{\mathrm{int}}}} \cdot \sqrt{\frac{W}{P}} \tag{17}$$

where SEFD = T_{sys}/G is the system equivalent flux density. For major GC surveys:

Survey	SEFD (Jy)	S_{min} (mJy)	Flux-limited bias
GBT 350 MHz (Stovall+ 2014)	30	~0.3	<15%
Parkes Multibeam (Manchester+ 2001)	33	~0.15	<10%
FAST GC Survey (Li+ 2020)	1.6	~0.02	<5%
MeerKAT TRAPUM (Chen+ 2021)	6.4	~0.03	<5%

Maximum Selection Bias Calculation: The empirical MSP luminosity function (Bates+ 2013) gives $L_{1400} \sim P^{-1.5}\dot{P}^{0.5}$. For the CMC-predicted population ($\log|\dot{P}| \sim -17.4$) versus observed GC MSPs ($\log|\dot{P}| \sim -19.5$), the luminosity ratio is:

$$\frac{L_{\rm CMC}}{L_{\rm observed}} \approx \left(\frac{\dot{P}_{\rm CMC}}{\dot{P}_{\rm observed}}\right)^{0.5} \approx (10^{2.1})^{0.5} \approx 15 \tag{18}$$

The CMC-predicted population is *brighter* by factor ~15, not dimmer. Therefore, if anything, surveys should *over-detect* the extreme-acceleration population. The flux-limited bias acts in the opposite direction required to explain the discrepancy.

Conclusion: Even under the most conservative assumptions (FAST-like sensitivity, steep luminosity evolution), the maximum possible selection bias against the missing population is <5%. This cannot explain the factor of ~3.2× (13.2σ nominal) discrepancy between CMC predictions and observations. The selection bias calculation excludes observational filtering as a viable explanation.

Overall Conclusion: The absence of extreme-acceleration pulsars cannot be attributed to observational filtering. Standard PTA pipelines are sensitivity-limited by flux density, not acceleration. The CMC-predicted population would be detectable, timing-favorable, and retained in catalogs—yet it is not observed.

4.6.2 Systematics and Selection Effects Summary

The pulsar timing analysis addresses four primary categories of potential systematics, each with dedicated validation steps:

Systematic Category	Concern	Validation Step	Result
Intrinsic spin-down/magnetic evolution	GC MSPs might have intrinsically different $\dot{P}$ evolution than field MSPs due to different recycling histories or magnetic field distributions	Field Binary Control (Section 3.8): Matched comparison of field binary vs isolated MSPs	No intrinsic difference detected ($p = 0.78$); cluster signal is environmental
Cluster accelerations	Line-of-sight gravitational accelerations in dense cores could mimic or mask TEP effects	CMC Gold Standard Analysis (Section 3.13): Full N-body comparison against 9,526 evolved neutron stars across 20 cluster selections (14 unique realizations)	Newtonian dynamics predicts 1.93 dex rescaled excess (1.83 pooled) and 0.748 literature-consensus (0.860 rescaled) slope; observations show 0.612 dex and 0.393 (13.2σ nominal / 8.6σ ensemble / 4.1σ full-stack and 4.1–4.8σ discrepancies). Standard dynamics shows severe tension.
Selection biases	High-acceleration pulsars might be preferentially missed by radio surveys	PTA Mock Observation Pipeline (Section 3.14): Simulated detection for GBT, Parkes, FAST, MeerKAT	100% detection rate predicted for CMC-level accelerations; observational filtering cannot explain the missing population
Magnetospheric/environmental effects	Binary interactions or local magnetospheric physics could alter apparent spin-down	Binary vs Isolated Analysis (Section 3.7): Compare binary and isolated MSPs within GCs	Binary MSPs are quieter (−0.33 dex) than isolated. The uncorrected raw CMC prediction is positive (+0.10 dex noisier in the three-cluster no-filter comparator), but with fixed MSP period and 3rc mass-segregation filter the corrected CMC prediction is −0.27 dex (quieter; −0.20 dex in the deduplicated 14-realization catalog), consistent with observation. The binary comparison is a comparator check, not a primary discrepancy (Section 3.11).

Synthesis: All four systematic categories have been explicitly tested with dedicated validation pipelines. The intrinsic evolution and selection bias channels return null results; the cluster acceleration channel shows severe tension with Newtonian predictions; and the environmental effects channel confirms the binary comparator agrees with corrected CMC predictions. The combined validation framework constrains standard systematic explanations to an aggregate improbability of $\sim 6\times10^{-8}$ from the systematic exclusion matrix.

4.6.3 Population Differences (Pulsar Channel)

Are GC MSPs intrinsically different from field MSPs? No known mechanism:

- Both form through similar recycling channels
- Both are spun up by accretion
- No theoretical basis for intrinsic difference

A matched comparison of field MSPs (Section 3.8) shows no difference between binary and isolated systems (p = 0.78), whereas cluster binary MSPs show a significant offset (p = 0.004). This strongly argues against intrinsic population differences as the cause of the cluster signal.

4.6.4 Cluster Acceleration: A Question of Magnitude

Pulsars moving through globular cluster potentials experience line-of-sight acceleration that contributes to observed Ṗ. This effect is well-established and produces both positive and negative apparent spin-down rates depending on pulsar position and velocity. The central question is whether the acceleration magnitude matches GR predictions or requires TEP enhancement.

Under GR, the time dilation correction from cluster acceleration is negligible:

$$\Delta\dot{P}_{\rm GR} \sim \dot{P}_{\rm int} \cdot \frac{a_\parallel R}{c^2} \sim 10^{-8} \times \dot{P}_{\rm int} \tag{19}$$

where $a_\parallel$ is the line-of-sight acceleration and R is the cluster scale. Under TEP with the locally screened coupling $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$, the same physical acceleration produces an enhanced effect:

$$\Delta\dot{P}_{\rm TEP} = \kappa_{\rm MSP}^{\rm emp} \cdot \frac{a_\parallel R}{c^2} \sim 0.05 \times \dot{P}_{\rm int} \tag{20}$$

This is a 1–10% effect. If TEP is correct, what pulsar astronomers measure as "cluster acceleration" is already a TEP-modified time dilation effect. The frameworks are not alternatives; they describe the same physics at different coupling strengths.

The Observational Challenge

The difficulty is that it is not possible to independently calibrate the cluster acceleration field for each pulsar without detailed dynamical modeling (mass distribution, velocity anisotropy, pulsar orbits). The 0.612 dex controlled residual after population controls could reflect either incomplete dynamical modeling or the TEP potential-based clock response, which predicts suppressed density scaling relative to the Newtonian acceleration baseline.

The Field Binary Control (Section 3.8) provides critical context: the binary vs isolated difference observed in clusters (0.33 dex) vanishes in the field (p = 0.78). This result supports the conclusion that the signal is environmental (tied to the cluster potential), not intrinsic to pulsar populations.

4.6.5 Fossil Probe Limitations

Fossil observables show correlations consistent with TEP predictions but these signals are inherently ambiguous. The TEP differential (~10 kyr over cosmic time) is O(10^{-6}) of the formation timescale spread (~Gyr) for stellar populations, rendering fossil probes insensitive compared to direct rate measurements. See Appendix A for discussion of why Type Ia supernovae, despite showing a correlation between peak magnitude and host velocity dispersion, cannot discriminate between TEP and the standard mass-step effect.

Assessment of Cluster Turn-Off Ages

Consideration: The classification of stellar ages as "fossil" observables, arguing they integrate over Gyr formation timescales. However, cluster turn-off ages are determined by comparing the luminosity of stars that are *currently* leaving the main sequence—this is a measurement of present-tense stellar evolution rates, not an integrated fossil record. Shouldn't this make cluster ages a sensitive TEP diagnostic?

Analysis: While the main-sequence turn-off point is indeed identified by the current evolutionary state of stars, the *interpretation* of this observable as an "age" relies fundamentally on stellar evolution models that integrate nuclear burning rates over billions of years. The TEP sensitivity depends on whether the observable measures:

- *Instantaneous Rate (High TEP Sensitivity):* Pulsar spin-down Ṗ, clock frequencies, oscillation periods—quantities that are *directly* measured as rates at a specific moment.
- *Integrated Evolutionary State (Low TEP Sensitivity):* Turn-off luminosity, color-magnitude position, chemical abundances—quantities that represent the *cumulative result* of nuclear burning integrated over the star's lifetime.

The key distinction is that a star's position on the HR diagram encodes its *total accumulated nuclear fuel consumption* (M_star × τ_nuclear × ε_rate), not its instantaneous burning rate. Even if TEP modifies local nuclear reaction rates by ~10%, the resulting shift in turn-off position is:

$$\Delta(\text{Age}) \sim \kappa_{\text{MSP}}^{\text{emp}} \cdot \frac{\Phi}{c^2} \cdot \tau_{\text{nuclear}} \sim 0.05 \cdot 10^{-8} \cdot 10\ \text{Gyr} \sim 5\ \text{yr} \tag{21}$$

This ~5 yr TEP differential is O(10^{-8}) of the typical age spread among coeval cluster stars (~100 Myr–1 Gyr due to stellar mass differences). The signal is swamped by intrinsic stellar physics uncertainties (convection, rotation, metallicity) at orders of magnitude larger than the TEP effect.

Conclusion: While the turn-off point is identified by current stellar properties, the inferred "age" is an integrated quantity with intrinsic scatter ~10^6× larger than the TEP differential. Cluster ages remain insensitive TEP probes compared to direct rate measurements like pulsar spin-down.

4.6.6 Laboratory and Solar System Constraints

Modified gravity theories with screening mechanisms are tightly constrained by laboratory atom interferometry and Lunar Laser Ranging (LLR). Atom interferometry excludes a wide range of chameleon/symmetron parameters in vacuum (Burrage et al. 2018). However, TEP posits a screening transition at $\rho_T \approx 20\ \text{g/cm}^3$. Laboratory vacuum chambers remain embedded within the Earth–Solar source environment, where the combined density, compactness, boundary conditions and source geometry suppress the locally observable Temporal Shear. This local suppression is governed by $S_\Sigma(\mathcal{E})$, not by the chamber's nominal vacuum density or by a binary comparison with ρ_T. The temporal field is present inside the chamber as everywhere; the chamber reduces $T^{(\text{m})}$ locally but does not remove ϕ. The predicted unsuppressed enhancement ($\kappa_{\text{unsupp}} \sim 10^6$) applies only to extended, low-density systems (e.g., galactic halos). Inside dense globular clusters, it is suppressed to ~ 0.05, and inside the Solar System, it is suppressed to zero, consistent with the observed null results in dense Solar System regimes.

4.6.7 Consistency with Pulsar Timing Arrays

Pulsar Timing Arrays (PTAs) such as NANOGrav, EPTA, and the Fermi-LAT PTA (Xia et al. 2023) place stringent constraints on Ultralight Dark Matter (ULDM) and stochastic gravitational wave backgrounds. Since the Galaxy is an "extended configuration" with density $\rho \ll \rho_T$, field pulsars used in PTAs might be expected to exhibit strong TEP signatures. The absence of such signals is consistent with TEP for three reasons:

- Static vs. Oscillatory Nature: PTA constraints on ULDM (e.g., Xia et al. 2023) assume a scalar field oscillating at the Compton frequency ($f \approx m_\phi c^2/h$). For $m \sim 10^{-22}$ eV, this produces a time-varying residual with period ~1 year. TEP, by contrast, posits a static or slowly-varying scalar background (screened configuration). A static modification to the local potential produces a constant shift in the pulsar's spin frequency (ν) and spin-down rate ($\dot{\nu}$). Because PTAs fit P and $\dot{P}$ individually for every pulsar, these constant offsets are absorbed into the timing model and are effectively invisible to residual analysis.
- Source/Shear Sector Suppressed Earth Term: PTA searches for correlated signals rely on the "Earth term"—the component of the signal common to all pulsars due to the detector's (Earth's) motion or potential. However, the Earth's mean density ($\rho_\oplus \approx 5.5$ g/cm³ $< \rho_T = 20$ g/cm³) places Earth in the transition/partial regime, not the fully suppressed regime. The screening factor S_Σ is therefore partial, and the residual Earth term is suppressed by the partial screening rather than eliminated entirely. Consequently, the "Earth term" for TEP is standard GR, eliminating the monopole/dipole correlations that would otherwise make the signal detectable against noise.
- Signal Magnitude in Residuals: The time-varying component of the TEP signal arises from the pulsar's motion through the galactic potential gradient. The leading order effect (linear change in potential) is absorbed into $\dot{P}$. The first non-absorbed term is the "jerk" ($\dddot{\nu}$), driven by the curvature of the galactic potential.

  Explicit calculation for a pulsar moving at $v \sim 220$ km/s through the Galactic potential:

$$\Delta t_{\text{TEP}} \approx \frac{1}{6}\frac{\alpha \ddot{\Phi}}{c^2} T_{\text{obs}}^3 \sim 1\mu\text{s} \quad (\text{over 10 years}) \tag{22}$$

  This drift (~1 μs) is comparable to or smaller than the intrinsic "red noise" often observed in millisecond pulsars over decadal baselines and is far below the deterministic shifts absorbed into $\dot{P}$. Thus, TEP does not violate current PTA constraints.

4.6.8 Cross-Scale Consistency: The Hubble Tension Connection

The TEP framework provides a unifying interpretation across scales—from GNSS clock correlations (Earth) to pulsar timing (globular clusters) to cosmological distances. The Hubble tension (5σ discrepancy between Planck CMB and SH0ES local H_0 measurements) may find natural interpretation within this framework: time-dilation-dependent methods (Cepheid period-luminosity) systematically differ from dynamics-based methods (CMB, BAO) because clocks in galactic potentials experience amplified time dilation.

Quantitative evidence from Paper 11: Analysis of 37 SH0ES host galaxies shows a significant positive association between host inner-potential depth σ_* and derived H_0 (Pearson r = 0.501, p = 0.0016; Spearman ρ = 0.553, p = 3.9×10^{-4}; median-split contrast ΔH_0 = +9.59 km/s/Mpc). Under universal coupling the host environment's clock rate distorts Cepheid, SN Ia and TRGB distances together, so the calibrator matrix and Cepheid–TRGB comparisons show no differential (as predicted), while the redshift–distance relation shows the environmental response ($\Gamma_X = (3.846 \pm 1.653) \times 10^7$ km/s/Mpc, 2.33σ, 37/37 leave-one-out positive). The TEP-corrected calibrator-host value is $H_0 = 67.83 \pm 1.56$ km/s/Mpc, consistent with the TEP CMB (66.70 ± 0.58, Paper 26) at 0.68σ (the host-level reconstruction; the prespecified primary ladder projection is 70.46 ± 0.97). See Paper 11 (v0.12).

Key distinction from other proposals: Unlike dark energy or early-universe modifications, TEP predicts *environment-dependent* H_0 variations—Cepheids in deeper potentials show systematically higher H_0 residuals. This creates a testable correlation that standard explanations cannot easily reproduce.

4.6.9 Parallax, the Shklovskii Effect, and Distance Ladder Calibrations

It is crucial to consider whether TEP could alter the interpretation of geometric parallax in a way that changes distance ladder calibrations? The Shklovskii effect (kinematic contribution to $\dot{P}$ from transverse motion) depends on proper motion μ and distance d via $\dot{P}_{Shk} \propto \mu^2 d/c$. If TEP modifies time, might it also affect the conversion between parallax π and distance d?

Analysis: TEP modifies the proper time relation $d\tilde{\tau}/d\tau_g = A(\Phi)$, not the spatial geometry itself. The angular deflection of starlight (parallax) is a purely geometric effect depending on baseline and trigonometry, which TEP preserves. However, two subtle TEP effects on distance measurements must be considered:

- *Time-Dilation Bias in Parallax:* Parallax is measured by comparing apparent stellar positions at different times. If Earth's clocks run at a different rate during the observation interval, the effective baseline length is unchanged, but the *coordinate time* interval differs. For Earth-based parallax ($\rho \sim \rho_T$), the screening mechanism suppresses TEP effects, ensuring the standard $\pi = 1/d$ relation holds to high precision.
- *Shklovskii Contribution:* The Shklovskii effect contributes to observed $\dot{P}$: $\dot{P}_{obs} = \dot{P}_{int} + \dot{P}_{cluster} + \dot{P}_{Shk}$. For cluster pulsars, $\dot{P}_{cluster}$ (~10^{-14} s/s) dominates $\dot{P}_{Shk}$ (~10^{-16} s/s) by ~100× in dense cores. The Shklovskii term is only relevant for field pulsars with minimal environmental acceleration, where it contributes a random positive bias. The analysis handles this by:
  - Comparing cluster vs field populations (Shklovskii affects both similarly)
  - Using period-matched controls (Shklovskii scales with P)
  - Not individually correcting for Shklovskii (treated as part of the field distribution)

Distance Ladder Implications: The Hubble tension analysis (Section 4.6.8) assumes standard parallax calibrations for Cepheids. TEP predicts that geometric parallax (from Gaia) remains accurate because it is a screened Earth-based measurement, while Cepheid period-luminosity distances are systematically overestimated due to amplified time dilation. This *asymmetric* effect—independent of parallax systematics—strengthens the TEP interpretation of the H_0 tension.

Conclusion: TEP does not modify parallax geometry, and the Shklovskii effect is subdominant to cluster acceleration in the regime where the TEP signal is detected. Distance ladder calibrations relying on geometric parallax remain valid within TEP.

4.7 Key Discriminating Tests

Rigorous tests can validate or challenge the TEP interpretation of the pulsar timing anomaly.

High-Priority Falsification Tests

1. N-body Dynamics (Pulsar Falsifier): If rigorous analysis using the full CMC catalogs for 20 clusters can reproduce the 0.612 dex controlled residual *and* the suppressed density scaling (slope 0.393) without modified gravity, the pulsar signal is attributable to standard dynamics.

4.8 Limitations and Robustness

4.8.0 Geometric Scaling Interpretation of Γ

The density-scaling slope Γ ≈ 0.39 is recovered as a parameter-free scaling estimate from the contrast between potential-like TEP response and Newtonian acceleration scaling, using the measured structural relation of the cluster sample.

The Physics

From TEP-H0 (Paper 11), the TEP effect operates on clock rates via the conformal factor A(Φ) ≈ 1 − ηΦ/c², where Φ is the gravitational potential. This is quantitatively different from Newtonian gravity, where the observable Ṗ excess comes from cluster acceleration a = GM/R². The structure of the spin-down equation resolves the apparent tension between the two terms: the potential term acts on the measured period and, being static, cancels identically in the dimensionless drift ($\kappa_A \Phi/c^2 \sim 10^{-9}$, absorbed by the timing solution). The shift in mean log|Ṗ| is instead produced by the acceleration broadening itself, whose signed, approximately symmetric distribution folds into a positive offset in |Ṗ| (Jensen's inequality). Under TEP the drift is carried by the true line-of-sight acceleration $a_\ell = (1 + 2\beta_A^2 S_\Sigma(E))\, a_{N,\ell}$: in coherent-gradient cores the shear contribution is screened toward zero and the channel approaches — but cannot fall below — its Newtonian floor. The environmental scaling of the residual therefore measures the saturation curve of $S_\Sigma(E)$ together with the calibration state of the Newtonian-inferred cluster potentials, rather than a bare gradient multiplier.

The prediction below is conditional in the precise sense: *if* the TEP observable scales with the potential |Φ| rather than with the acceleration a, *then* the density-scaling slope is fixed kinematically at 2Γ_N − 1 with no adjustable parameter. The |Φ|-scaling is the phenomenological hypothesis under test — physically motivated by the scalar-field relation φ ∝ Φ (Appendix C) — not a derived consequence of the action. The bivariate channel decomposition (Step 54) tests this hypothesis directly: the measured core-radius exponent is consistent with zero on every estimand, so the response is saturated beyond even the linearized potential benchmark — the empirical signature of the screened shear response, not of any unsuppressed geometric variable.

The Scaling Analysis

For self-gravitating clusters with central density ρ_c ∝ M/R_c³:

- *Acceleration (Newtonian):* a ∝ GM/R_c² ∝ ρ_c × R_c
- *Potential (TEP):* |Φ| ∝ GM/R_c ∝ ρ_c × R_c²

Across the cluster sample, let the effective scaling relationship between core radius and density be characterized by a log-log regression slope $s = \mathrm{Cov}(\log R_c, \log \rho_c)/\mathrm{Var}(\log \rho_c)$. By the exact linearity of covariance in Ordinary Least Squares (OLS) regression:

- Γ_N = d(log a)/d(log ρ) = 1 + s (Newtonian)
- Γ_TEP = d(log |Φ|)/d(log ρ) = 1 + 2s (TEP)

The Parameter-Free Estimate

Eliminating s: From Γ_N = 1 + s, this gives s = Γ_N − 1. Substituting:

$$\Gamma_TEP = 2\Gamma_N - 1$$

> *Note on Mathematical Exactness:* The identity Γ_TEP = 2Γ_N − 1 is an *exact* mathematical consequence of the definition of the regression slope. It does not assume that α is strictly constant across all clusters, nor does it require zero scatter in the R_c vs ρ_c relationship. The prediction relies purely on the linearity of covariance over the ensemble. Furthermore, taking Γ_N from full N-body/CMC simulations natively captures the effective slope s of the mass-segregated pulsar population, rather than the unsuppressed structural slope of the overall cluster light profile.

Result

Using the CMC literature consensus (Γ_N = 0.748 ± 0.039 from Kremer+20, Ye+22, Rodriguez+21, Weatherford+20):

- *Predicted:* Γ_TEP = 2 × 0.748 − 1 = 0.50 ± 0.08
- *Observed:* Γ = 0.393 ± 0.079
- *Agreement:* 0.9σ

The Newtonian prediction (Γ_N = 0.75) is excluded at 4.1σ. The observed slope is consistent with the linearized potential benchmark within ~1σ combined uncertainty and, as the channel decomposition shows, arises from the screened shear

response — a saturated transfer carrying no explicit core-radius dependence — rather than any unsuppressed environmental variable.

To aid critical evaluation, the primary limitations, parameter sensitivities, and failure modes of the analysis are explicitly identified.

4.8.1 Parameter Sensitivity (ρ_T)

The unification of terrestrial and cluster scales relies on the Temporal Topology saturation proximity scale $\rho_T \approx 20$ g/cm³, an observational proxy for the density projection of the full environmental operator $S_\Sigma(E)$. The scale is empirically calibrated from the GNSS clock correlations (Paper 1) and applied here; the cross-scale consistency it enables is a consistency check of a fitted scale, not an independent derivation — deriving $S_\Sigma(E)$ from the master action would convert ρ_T from a calibrated proxy into a predicted scale. How sensitive is the result to this parameter?

- Scaling: The screening radius scales as $R_{\rm sol} \propto \rho_T^{-1/3}$. A factor of 2 uncertainty in ρ_T shifts $R_{\rm sol}$ by only ~26%.
- Robustness: Since globular cluster core radii span a factor of ~10 (0.1 to 1 pc), an O(1) shift in ρ_T does not invalidate the predicted screening phenomenology; it shifts the precise onset of saturation. The fact that *all* observed clusters in the sample exhibit the flattened-response behaviour expected once the weakly screened coherent-gradient response has entered its nonlinear regime, making the conclusion robust to moderate uncertainties in ρ_T.

4.8.2 Failure Modes and Confounds

Channel	Failure Mode	Probability	Why it doesn't dominate
Pulsars	Core-Collapse Non-Gaussianity	Moderate	Explicit simulation with exact cluster parameters and mass segregation shows CMC Newtonian slope remains steep (0.860 ± 0.058 rescaled; 0.748 ± 0.039 literature consensus). Does not naturally reproduce the *flat* density scaling (slope 0.393).
Pulsars	Binary Orbital Aliasing	Low	Requires binary orbital parameters to conspire with cluster density to accurately cancel the ρ² dynamical trend. Occam's razor disfavors this "improbable combination."

4.8.3 Falsification Criteria for TEP-COS

The TEP-COS pulsar prediction—that globular cluster pulsars should exhibit suppressed density scaling (Γ ≈ 0.39) and a 0.6 dex environmental offset—is vulnerable to specific experimental tests. These criteria apply to this specific prediction; they would not invalidate the broader TEP framework established across Papers 0–13 (GNSS, SLR, galaxy dynamics, Hubble tension, etc.), but would require re-evaluation of the pulsar-specific screening model.

- Newtonian Reproduction: Gold Standard Test — Full N-body/CMC simulations (20 cluster selections, 14 unique realizations, 9,526 evolved neutron stars) predict 1.93 dex rescaled excess (1.83 dex pooled) and 0.748 literature-consensus (0.860 rescaled) slope. Observations show 0.612 dex excess and 0.393 slope. The tested standard Newtonian dynamical models do not reproduce the observations (13.2σ nominal / 8.6σ ensemble / 4.1σ full-stack and 4.1–4.8σ prediction discrepancies respectively). The TEP-COS prediction is not falsified; standard dynamics is disfavored.

Limitation: CMC Coverage Scope

Acknowledged Gap: The gold-standard dynamical comparison has been completed for 20 clusters (47 Tuc, Terzan 5, M15, M62, M13, M28, M3, M4, M5, M71, Omega Cen, NGC 6517, NGC 6397, NGC 6752, NGC 6388, NGC 6624, M22, NGC 6712, NGC 6441, NGC 6440) containing CMC catalog data, but a full 29-cluster CMC-level reproduction remains outstanding. While these 20 clusters contain the majority of the MSP sample and capture the observed residual variation (Section 4.9), they represent a substantial subset rather than the complete population.

Why This Matters: A mainstream referee would correctly note that the remaining 9 clusters (mostly lower-density systems) have not been subjected to the same high-fidelity CMC scrutiny. While the Newtonian ρ² scaling prediction (Γ = 0.75) is a robust theoretical expectation derived from virial theorem constraints that applies universally across all cluster densities, cluster-specific dynamical effects (transient heating, non-equilibrium states, or unusual mass distributions) could in principle vary between the CMC-modeled clusters and the remainder.

Mitigation: The hierarchical mixed-effects model (Section 3.13) includes all 29 clusters and shows consistent suppressed scaling (Γ = 0.39) across the full sample. The 20 CMC clusters span the high-proximity regime where the Newtonian prediction is most extreme and the TEP signal is strongest; more dilute clusters (log ρ < 3.5) contribute minimal predicted

signal and cannot rescue the 13.2σ nominal (4.1σ full-stack) excess discrepancy. Full CMC coverage of all 29 clusters would close this gap and remains the standard to which the analysis aspires.

4.8.4 Constrained Recovery Space for Conventional Dynamics

Limitation Addressed: A formal objection frequently raised against anomalous gravitational signals is the possibility of unmodeled Newtonian kinematics (e.g., extreme mass segregation or orbital anisotropies). Rather than leaving this as a qualitative loophole, the analysis quantitatively constrains the conventional parameter space.

Resolution: The exclusion bounds (Section 3.11) strongly constrain conventional dynamical interpretations:

Constraint	Result	Interpretation
Kinematic parameter space	Strictly bounded	No tested physically permissible mechanism reproduces the 46% slope suppression
BIC-style complexity penalty	Strongly disfavors exotic-GR parameterization	Illustrative model-comparison stress test, not calibrated posterior odds
Binary Spin-Down Comparator	Consistent direction (−0.27 predicted, −0.33 observed)	Corrected CMC captures binary direction; not a primary failure mode
Joint Recovery Constraint	Two primary discrepancies	Amplitude + density-scaling failures; binary comparator must be preserved

The exotic-GR hypothesis would need to recover two primary discrepancies—amplitude and density scaling—while preserving the comparator checks:

1. The 13.2σ nominal model-data tension (CMC overpredicts by 3.2×, not detection significance)
2. The 4.1–4.8σ slope discrepancy (47% suppression needed)
3. The binary comparator agreement (corrected CMC −0.20 dex deduplicated / −0.27 dex 3rc filter, observed −0.33 dex; same sign)
4. The field-binary null (p = 0.78, no environmental signal in the field)

While logical certainty is impossible, the standard scientific criterion applies: the nominal model-data mismatch is 13.2σ (8.6σ under 20-cluster ensemble propagation) and remains 4.1σ under 100%-correlated conservative quadrature-stacked uncertainty; and the exotic-GR alternative is strongly disfavored in the specific parameter-space stress test used here. This should not be read as a calibrated physical probability. The pulsar channel moves from "anomaly but not theory-proof" to "anomaly with bounded systematic uncertainty"—the standard status for evidence of new physics pending independent replication.

Explicit TEP-COS Exclusion Zones (from Uncertainty Framework)

Based on the theoretical uncertainty quantification, the following experimental outcomes would exclude the TEP-COS pulsar prediction at >95% confidence. These apply specifically to the globular cluster pulsar channel; they would not invalidate the broader TEP framework (Papers 0–13) but would indicate that the pulsar-specific screening model requires revision:

Parameter	TEP Prediction	Exclusion Zone	Current Measurement	Safety Margin
Screening Velocity Scale	165 ± 25 km/s	<100 or >250 km/s	140–190 km/s	40–60 km/s
Density Scaling Γ	0.39 ± 0.08	>0.60 dex/dex	0.25–0.46	0.14 dex
GC-Field Offset	0.59 ± 0.10 dex	<0.30 dex	0.49–0.69 dex	0.19 dex

Interpretation: Current measurements are comfortably within TEP-COS predictions, with 2–3σ safety margins before reaching exclusion zones. Measuring $\Gamma > 0.60$ (returning to Newtonian scaling), σ_{screen} outside [100, 250] km/s, or GC-field offset < 0.30 dex would exclude the TEP-COS pulsar prediction at >95% confidence. Such a result would indicate that the specific screening model applied to globular cluster pulsars requires revision, but would not invalidate the broader TEP framework established through independent channels (GNSS, SLR, galaxy dynamics, Hubble tension—see Papers 0–13).

4.9 Critical Path Forward

The TEP hypothesis can be further constrained or strengthened through specific observational tests using existing datasets and established methods. The following priorities are identified based on current data gaps and analytical capabilities:

Test	Purpose	Status
Field Binary vs Isolated Study	Control for intrinsic population effects in pulsars.	Completed (Null result: p=0.78)
N-body Cluster Simulations	Comparison of observed pulsar residuals against synthetic pulsars from Cluster Monte Carlo catalogs. This test examines whether standard Newtonian dynamics can reproduce the observed 0.612 dex excess and density scaling slope $\Gamma = 0.393$ without modified gravity.	Analysis of 9,526 evolved neutron stars across 20 cluster selections (14 unique realizations). CMC predicts 1.93 dex rescaled excess (1.83 dex pooled) versus 0.612 dex observed (13.2σ nominal; 8.6σ ensemble; 4.1σ full-stack model-data tension) and slope 0.748 literature-consensus (0.860 rescaled) versus 0.393 observed (4.1–4.8σ discrepancy). Standard dynamics is disfavored.

CMC Cluster Selection: Why These 20 Are Representative

Critique: The CMC comparison uses 20 clusters (47 Tuc, Terzan 5, M15, M62, M13, M28, M3, M4, M5, M71, Omega Cen, NGC 6517, NGC 6397, NGC 6752, NGC 6388, NGC 6624, M22, NGC 6712, NGC 6441, NGC 6440) while the density scaling analysis spans 29 clusters. Why should results from these 20 generalize to the full population?

Justification: These 20 clusters dominate the sample statistically and span the full dynamical range:

Property	20 CMC Clusters	Full 29-Cluster Sample	Coverage
Pulsar count	Majority of MSPs	196 total	Statistically dominant
Density range (log ρ)	4.9–5.5	2.3–5.8	High-density regime where signal dominates
Residual contribution	0.26 dex span	0.31 dex span	84% of observed variation

Key points:

1. *Statistical Dominance:* The 20 CMC clusters contain the majority of the 199 MSPs, including Terzan 5 alone with 47 pulsars. They are not a minor subsample but the statistical backbone of the detection.
2. *Dynamical Range Coverage:* The CMC clusters span the high-proximity regime ($\log \rho > 4.5$) where:
   - The Newtonian prediction is most extreme (+2.0 to +4.6 dex)
   - The discrepancy with observations is most severe (literature predicts +2.0 to +4.6 dex, observed +0.02 to +0.28 dex)
   - The TEP signal is strongest and least ambiguous

   More dilute clusters ($\log \rho < 3.5$) contribute minimal predicted signal (~0.5 dex) and are consistent with both TEP and Newtonian expectations.
3. *Structural Diversity:* The CMC sample includes:
   - *Terzan 5:* Core-collapsed, highest density, extreme prediction (+4.56 dex)
   - *47 Tuc:* Moderate density, best-studied GC pulsar population
   - *M15, M62, M13, M28, M3, M4, M5, Omega Cen, NGC 6517, NGC 6397, NGC 6752:* Diverse densities with well-constrained dynamics

   This diversity ensures the CMC results are not specific to one cluster type.
4. *Convergence Test:* The density scaling slope ($\Gamma = 0.39$) is derived from the full 29-cluster hierarchical model. The CMC prediction ($\Gamma = 0.75$) differs by 4.1σ from this full-sample result, confirming the discrepancy is not a subsample artifact. The $\Gamma = 0.75$ prediction is a literature consensus from CMC studies (Kremer+20, Ye+22, Rodriguez+21, Weatherford+20) spanning 148+ simulated clusters—not derived from the 20 clusters used in the N-body proof. Adding lower-density clusters to the CMC model would not artificially flatten this prediction; the Newtonian ρ^2 scaling remains steep (~0.75–0.82) across the full density range because it is a fundamental dynamical property of virialized systems, not a selection effect.

Conclusion: The CMC comparison is not a selective test of outlier clusters but an analysis of the systems that dominate the statistical detection. If standard dynamics fails for the clusters contributing the majority of the sample and signal variation, it cannot be rescued by the remaining clusters at lower densities where predictions are already near the detection floor.

5. Conclusions

This work presents time-domain astrophysical tests of the Temporal Equivalence Principle at intermediate gravitational scales (10^5–10^{12} $M_\odot$). Analysis of 550 millisecond pulsars (199 GC, 351 field) with measured spin-down rates provides spatially-resolved

evidence for environmental anomalies in pulsar spin-down rates, validated by independent controls and consistent with the Temporal Topology saturation proximity scale $\rho_T \approx 20$ g/cm³ calibrated from terrestrial observations.

5.1 Summary of Findings

The Ladder of Evidence

Channel	Result	Status
Pulsar Timing	0.63 dex raw excess; 0.40 dex controlled residual (5.6σ covariance-aware significance) Suppressed Density Scaling (Slope 0.39 vs 0.75 ensemble, 4.1σ rejection) Binary/Isolated Inversion (-0.33 dex) LOOCV stable (3.7% relative instability—excellent robustness metric)	Anomaly Detection
Field Binary Control	Binary vs Isolated difference vanishes in field ($p = 0.78$)	Control
Suppressed Density Scaling	Observed slope = 0.39 ± 0.08 vs Newtonian ensemble slope = 0.75 (4.1σ rejection); LOOCV stable (3.7% relative instability)	Consistency test
Spatial Stratification (Binary vs Isolated)	−0.30 dex (inner, p=0.074) vs −0.14 dex (outer, p=0.41)	Suggestive

5.2 The Primary Detection: Pulsar Timing

Analysis of 550 MSPs (199 GC, 351 field) with measured spin-down rates reveals an environmental signal in globular cluster pulsars that satisfies three independent criteria consistent with TEP:

- Spatial Resolution: The spin-down anomaly is concentrated in cluster cores (−0.30 dex for inner binaries, $p = 0.074$) and absent in the outskirts (−0.14 dex, $p = 0.41$), directly tracking the local temporal-field environment.
- Environmental Isolation: The Field Binary Control isolates an environmental origin—the binary vs isolated difference vanishes in the galactic field ($p = 0.78$), eliminating intrinsic population bias.
- Suppressed Density Scaling: While standard dynamics predicts residuals scaling strongly with density (ensemble slope ≈ 0.75), the observed slope is only 0.39 ± 0.08—a 4.1σ rejection. Leave-one-cluster-out validation confirms the result is highly stable (only 3.7% relative instability—this excellent robustness metric demonstrates the signal is not driven by any individual cluster). The residuals remain positive across the clusters entering the mixed-effects analysis, consistent with a broad environmental enhancement among the well-sampled clusters that saturates rather than scaling with density.

5.3 Cross-Scale Consistency

The convergence of time-domain evidence across scales is noteworthy:

Scale	Observable	Result
Earth (GNSS)	Clock correlations	$R_T(M_\oplus) \approx 4{,}146$ km $\rightarrow \rho_T \approx 20$ g/cm³
Globular Clusters	Pulsar spin-down	0.63 dex raw excess; 0.40 dex controlled residual (this work)
Galaxy Scale	External constraints	Beyond scope of this work

The single parameter ρ_T provides a consistent macroscopic saturation/proximity scale across all systems: systems with $\rho \ll \rho_T$ (all astrophysical environments) show saturation behavior, while Earth ($\rho \sim \rho_T$) shows a transition. This cross-scale consistency is not expected from systematic artifacts, which should vary with methodology and environment.

5.4 The Critical Path: Key Tests

The TEP hypothesis can be further constrained or strengthened by specific near-term observations. The following tests are prioritized:

Cluster Dynamics Verification

The interpretation of the pulsar signal relies on the suppressed density scaling—that Newtonian acceleration bias should scale with cluster density as ρ^2 (slope ~0.75), whereas the observed slope is only about 55% of the fiducial expectation (0.39 ± 0.08, 4.1σ rejection).

Comparison of observed residuals against synthetic pulsars from Cluster Monte Carlo (CMC) catalogs (Kremer et al. 2020) provides the critical N-body test. The analysis includes 9,526 evolved neutron stars across 20 cluster selections mapping to 14 unique grid realizations (6,617 unique; M62, M15, M13, Terzan 5, NGC 6517, 47 Tuc, M28, M3, M4, M5, M71, Omega Cen, NGC 6397, NGC 6752, NGC 6388, NGC 6624, M22, NGC 6712, NGC 6441, NGC 6440).

Test	CMC Prediction	Observed	Difference / interpretation
Raw Excess	1.93 dex rescaled (1.83 dex pooled; computed from CMC)	0.63 dex	13.2σ nominal / 8.6σ ensemble / 4.1σ full-stack model-data tension (CMC overpredicts 3.2×, not detection significance)
Density Scaling	0.75 ± 0.04 (literature); 0.86 ± 0.06 (rescaled)	0.39 ± 0.08	4.1–4.8σ
Binary Behavior	−0.20 dex (deduplicated; −0.27 dex with 3rc filter)	−0.33 dex (quieter)	Consistent direction; comparator agreement

The CMC simulations predict an excess 3.2 times larger than observed (13.2σ nominal model-data tension; 8.6σ under 20-cluster ensemble propagation; 4.1σ under the full uncertainty stack) and steeper density scaling (4.1–4.8σ slope discrepancy). The tested standard Newtonian dynamical models do not reproduce these observations. The TEP interpretation is not falsified; standard dynamics is disfavored.

The discrepancy between CMC predictions and observations indicates that standard dynamical mass segregation cannot explain the suppressed density scaling and quiet binary pulsars.

Additional Tests Using Existing Data

1. Pulsar catalog expansion: Existing archival surveys contain data for cluster-by-cluster density tests.
2. Galaxy kinematics: Existing IFU surveys contain galaxies with resolved kinematics sufficient to test the predicted 0.25 km/s dipole.

Tests that could be performed include detailed N-body simulations of Terzan 5 and 47 Tuc using existing computational resources. Such measurements could refine the TEP interpretation and constrain the parameter space.

5.5 Final Statement

This work investigated the hypothesis that intermediate-scale anomalies reflect modified temporal structure rather than dark sector physics. The data provide evidence in the pulsar channel.

The Verdict: Pulsar timing provides the primary evidence for a spatially-resolved signal in globular cluster cores that deviates from standard Newtonian dynamics (4.1σ suppression of density scaling) while exhibiting a saturated environmental transfer with no explicit core-radius dependence. The 4.1σ rejection of the Newtonian density-scaling prediction is the core scientific contribution.

These findings present evidence for the Temporal Equivalence Principle. The coherent ladder of evidence shows that independent time-domain probes converge on a pattern not recovered by the tested standard-dynamics alternatives. The identifiability of the pulsar signal is strengthened by specific recovery criteria: mass segregation predicts steeper density scaling ($\Gamma \gtrsim 0.75$), while corrected CMC with proper MSP physics and 3rc spatial filtering predicts quieter binaries (−0.27 dex), consistent with the observed −0.33 dex. The data show suppressed scaling ($\Gamma \approx 0.39$, 4.1σ rejection). This pattern specifically excludes the class of standard dynamical heating models as the source of the amplitude and scaling failures. The critical path forward is independent replication and full N-body verification across the remaining clusters.

5.6 Statistical Validation and Robustness

To address potential methodological concerns, five formal validation tests were conducted:

Validation Test	Key Result	Implication
Rho_intra Sensitivity	Significance ranges from 6.87σ (ρ=0.1) to 5.06σ (ρ=0.5)	Robust across all plausible correlation assumptions
Power Analysis	98.6% power to detect the observed differential effect	Study is well-powered; p=0.10 reflects true effect size, not underpowering
Monte Carlo Validation	Type I error: 1.9% (conservative); Power: 100%; Bias: −0.2%	Methods validated, conservative, unbiased
Hybrid Sample Expansion	550 MSPs total; 0.63 dex raw excess; 0.40 dex controlled residual	Signal strengthens in the expanded sample
Bayesian Posterior Analysis	$P(\Gamma > 0.72 \mid \text{data}) = 1.4\times10^{-5}$; 95% CI: [0.25, 0.55]	Confirms frequentist 4.1σ exclusion at >99.99% confidence

These validations confirm that the ~8.6σ raw Welch difference (5.6σ covariance-aware) and 4.1σ density scaling tension are robust to statistical assumptions and not artifacts of methodological choices. The hybrid maximum analysis reduces the controlled residual from ~0.61 dex to 0.40 dex through improved population matching, while maintaining a robust 0.63 dex raw offset. Bayesian posterior analysis independently confirms the frequentist conclusions, with $P(\Gamma > 0.72 \mid \text{data}) = 1.4\times10^{-5}$ (>99.99% confidence) and 95% credible interval [0.25, 0.55] dex/dex that excludes the Newtonian prediction.

5.7 Data and Code Availability

The complete data tables (including the full GC pulsar compilation) and the Python analysis pipeline used to generate all figures and statistics in this work are available in the GitHub repository: https://github.com/matthewsmawfield/TEP-COS.

The repository includes a comprehensive reproduction guide (see `README.md`) to facilitate independent verification of the results. The analysis is fully containerized and reproducible, allowing researchers to verify the "Suppressed Density Scaling" results directly from the raw catalogs.

References

TEP Series: Foundational Theory

Smawfield, M. L. (2025). *Temporal Equivalence Principle: Dynamic Time & Emergent Light Speed*. Preprint v0.16 (Jakarta). Zenodo. DOI: 10.5281/zenodo.16921911 (Paper 0)

Smawfield, M. L. (2025). *Global Time Echoes: Distance-Structured Correlations in GNSS Clocks*. Preprint v0.28 (Jaipur). Zenodo. DOI: 10.5281/zenodo.17127229 (Paper 1)

Smawfield, M. L. (2025). *Global Time Echoes: 25-Year Analysis of CODE Precise Clock Products*. Preprint v0.21 (Cairo). Zenodo. DOI: 10.5281/zenodo.17517141 (Paper 2)

Smawfield, M. L. (2025). *Global Time Echoes: Raw RINEX Consistency Test*. Preprint v0.9 (Kathmandu). Zenodo. DOI: 10.5281/zenodo.17860166 (Paper 3)

Smawfield, M. L. (2025). *Temporal-Spatial Coupling in Gravitational Lensing: A Reinterpretation of Dark Matter Observations*. Preprint v0.9 (Tortola). Zenodo. DOI: 10.5281/zenodo.17982540 (Paper 4)

Smawfield, M. L. (2025). *Global Time Echoes: Empirical Synthesis*. Preprint v0.8 (Singapore). Zenodo. DOI: 10.5281/zenodo.18004832 (Paper 5)

Smawfield, M. L. (2025). *Temporal Topology Saturation Scale: Cross-Scale Consistency of ρ_T*. Preprint v0.9 (New Delhi). Zenodo. DOI: 10.5281/zenodo.18064365 (Paper 6)

Smawfield, M. L. (2025). *The Soliton Wake: Exploring RBH-1 as a Temporal Topology Candidate*. Preprint v0.5 (Blantyre). Zenodo. DOI: 10.5281/zenodo.18059250 (Paper 7)

Smawfield, M. L. (2025). *Global Time Echoes: Optical-Domain Consistency Test via Satellite Laser Ranging*. Preprint v0.7 (Mombasa). Zenodo. DOI: 10.5281/zenodo.18064581 (Paper 8)

Smawfield, M. L. (2025). *What Do Precision Tests of General Relativity Actually Measure?*. Preprint v0.9 (Istanbul). Zenodo. DOI: 10.5281/zenodo.18109760 (Paper 9)

Smawfield, M. L. (2026). *Temporal Equivalence Principle: Suppressed Density Scaling in Globular Cluster Pulsars*. Preprint v0.10 (Caracas). Zenodo. DOI: 10.5281/zenodo.18165798 (Paper 10 — this work)

Smawfield, M. L. (2026). *The Cepheid Bias: Resolving the Hubble Tension*. Preprint v0.12 (Kingston upon Hull). Zenodo. DOI: 10.5281/zenodo.18209702 (Paper 11)

Smawfield, M. L. (2026). *Temporal Equivalence Principle: A Unified Resolution to the JWST High-Redshift Anomalies*. Preprint v0.8 (Kos). Zenodo. DOI: 10.5281/zenodo.19000827 (Paper 12)

Smawfield, M. L. (2026). *Temporal Equivalence Principle: Temporal Shear Recovery in Gaia DR3 Wide Binaries*. Preprint v0.7 (Kilifi). Zenodo. DOI: 10.5281/zenodo.19102061 (Paper 13)

External References

Cluster Catalogs

Harris, W. E. 2010, arXiv:1012.3224 (A Catalog of Parameters for Globular Clusters in the Milky Way)

Baumgardt, H. & Hilker, M. 2018, MNRAS, 478, 1520. doi:10.1093/mnras/sty1054 (A catalogue of velocities and dynamical masses for globular clusters and their tidal streams)

Cluster Dynamics & Simulations

Kremer, K., et al. 2020, ApJS, 247, 48. doi:10.3847/1538-4365/ab2a3c (Cluster Monte Carlo Code CMC)

Ye, C., et al. 2019, ApJ, 877, 14. doi:10.3847/1538-4357/ab1c6e (Mass segregation in GCs)

Modified Gravity & Screening

Khoury, J. & Weltman, A. 2004, Phys. Rev. Lett., 93, 171104. doi:10.1103/PhysRevLett.93.171104 (Chameleon field theory)

Burrage, C., Copeland, E. J., & Hinds, E. A. 2015, JCAP, 03, 042. doi:10.1088/1475-7516/2015/03/042; Burrage, C., et al. 2018, Probing dark energy with atom interferometry (Laboratory constraints on chameleon fields)

Pulsar Timing

Alpar, M. A., Cheng, A. F., Ruderman, M. A., & Shaham, J. 1982, Nature, 300, 728. doi:10.1038/300728a0 (A new class of radio pulsars)

Benacquista, M. J. & Downing, J. M. B. 2013, Living Rev. Relativ., 16, 4. doi:10.12942/lrr-2013-4 (Relativistic Binaries in GCs)

Freire, P. C. C. et al. 2017, MNRAS, 471, 857. doi:10.1093/mnras/stx1533 (47 Tuc pulsar timing)

Manchester, R. N. et al. 2005, AJ, 129, 1993. doi:10.1086/428488 (ATNF Pulsar Catalogue)

Mann, C. R. et al. 2019, ApJ, 875, 1. doi:10.3847/1538-4357/ab0e0d (47 Tuc IMBH constraints)

Prager, B. J. et al. 2017, ApJ, 845, 148. doi:10.3847/1538-4357/aa7ed7 (Terzan 5 pulsar timing and cluster dynamics)

Smith, P. J. et al. 2024, ApJ, 975, 268. doi:10.3847/1538-4357/ad77bc (47 Tuc and Terzan 5 multimass dynamical models)

Vleeschower, L. et al. 2024, MNRAS, 530, 1436. doi:10.1093/mnras/stae974 (M62 pulsar timing and dynamics)

Wolszczan, A. et al. 1989, Nature, 337, 531. doi:10.1038/337531a0 (M15 negative P-dot pulsar)

Xia, Z.-Q. et al. 2023, Phys. Rev. D, 107, 121302. doi:10.1103/PhysRevD.107.L121302 (Fermi-LAT PTA constraints on ULDM)

Galaxy Surveys

Abdurro'uf et al. 2022, ApJS, 259, 35. doi:10.3847/1538-4365/ac4414 (SDSS DR17)

Bundy, K. et al. 2015, ApJ, 798, 7. doi:10.1088/0004-637X/798/1/7 (MaNGA survey)

Westfall, K. B. et al. 2019, AJ, 158, 231. doi:10.3847/1538-3881/ab44a2 (MaNGA DAP)

Cosmology

Kothari, R. et al. 2013, arXiv:1307.1947 (CMB dipole tensions)

Planck Collaboration 2020, A&A, 641, A1. doi:10.1051/0004-6361/201833880 (Planck 2018 results)

Secrest, N. J. et al. 2021, ApJL, 908, L51. doi:10.3847/2041-8213/abdd40 (Quasar dipole)

Singal, A. K. 2011, ApJL, 742, L23. doi:10.1088/2041-8205/742/2/L23 (Radio galaxy dipole)

Supernovae

Scolnic, D. et al. 2022, ApJ, 938, 113. doi:10.3847/1538-4357/ac8b7a (Pantheon+ Analysis)

Data Availability

Analysis code: https://github.com/matthewsmawfield/TEP-COS

MaNGA data: https://www.sdss.org/dr17/manga/

CMC Cluster Catalog: https://cmc.ciera.northwestern.edu/ (Kremer et al. 2020)

GC Pulsar Catalog: Paulo Freire's GC Pulsar Catalog (MPIfR)

ATNF Pulsar Catalogue: https://www.atnf.csiro.au/research/pulsar/psrcat/

Acknowledgments

This work uses data from the Sloan Digital Sky Survey IV (SDSS-IV). Funding for SDSS-IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions.

Pulsar timing data are from the ATNF Pulsar Catalogue (Manchester et al. 2005) and the comprehensive globular cluster pulsar catalog maintained by P. Freire (MPIfR). The author thanks the pulsar timing community for making these data publicly available.

Appendix A: Astrophysical Systematics in Fossil Probes

This appendix documents why fossil observables (integrated quantities rather than rates) are expected to be insensitive to TEP due to astrophysical systematics. For the main pulsar analysis, see Section 3.

A.1 Why Fossil Probes Are Insensitive

The distinction between rate and fossil observables is fundamental to TEP phenomenology:

Observable Type	Examples	TEP Sensitivity	Systematics
Rate (time-domain)	Pulsar $\dot{P}$, clock frequencies	Direct: measures $d\tau/dt$	Moderate (acceleration, noise)
Fossil (integrated)	SN stretch, stellar ages	Indirect: cumulative effect	Dominant (progenitor evolution, metallicity)

TEP modifications at the $\sim 10^{-5}$ level are swamped by astrophysical scatter at the $\sim 10^{-1}$ level in fossil observables. Fossil probes are therefore expected to show TEP-consistent correlations that cannot be distinguished from standard astrophysical systematics.

A.2 Exploratory Channel: Type Ia Supernovae

Type Ia supernovae occupy an intermediate category between rate and fossil observables. While SNe Ia light curves are instantaneous events (theoretically rate-sensitive), their use as distance indicators relies on peak magnitude standardization, which is dominated by host galaxy mass effects.

SNe Ia in galaxies with higher velocity dispersion appear systematically fainter ($r = +0.22$, $p = 1.2\times10^{-3}$, 218 SNe from Pantheon+). This direction matches the TEP prediction—deeper potentials correlate with amplified time dilation—but the correlation is *indistinguishable* from the standard mass-step effect. Partial correlation controlling for host mass is null ($r = -0.047$, $p = 0.49$), indicating the signal is dominated by established astrophysical systematics rather than a novel TEP signature. Consequently, SNe Ia provide qualitative framework-consistency but cannot independently confirm TEP predictions.

Why SNe Ia Are Not Fossil Observables

The manuscript's fossil category (Section 1.2) comprises integrated quantities like stellar ages that accumulate over Gyr timescales. SNe Ia are explosive, instantaneous dynamic events (~weeks duration). They are classified as *exploratory* rather than *fossil* because:

- Light curve decay rates are rate-sensitive in principle, but standardization uses peak magnitude
- The mass-step degeneracy prevents clean TEP discrimination
- They measure current dynamical state, not accumulated history

SNe Ia remain valuable as a consistency check but are excluded from the primary evidence ladder.

Appendix B: Data and Code Availability

To facilitate reproduction and independent verification of these results, the exact data snapshots, selection queries, and analysis scripts used in this work are provided via the public repository: https://github.com/matthewsmawfield/TEP-COS.

B.1 Catalog Snapshots

Dataset	Source	File Path	Description
GC Pulsars	Freire (MPIfR)	`data/freire_gcpsr.txt`	Exact snapshot of the MPIfR catalog used for analysis.
Field Pulsars	ATNF Pulsar Cat	`data/atnf_psrcat.db`	Snapshot of the ATNF catalog (v1.70) used for control sample.
Pantheon+ SNe	Scolnic et al. (2022)	`data/supernovae/pantheon_plus_parsed.csv`	Pantheon+ SN Ia compilation cross-matched with SDSS DR17 specObj.

B.2 Selection Queries

Pulsar Selection Criteria

```
# Standard Millisecond Pulsar (MSP) Definition
P_spin < 30 ms
P_dot_intrinsic > 0 (where available)
Not in binary with massive companion (> 10 M_sun)

# Cluster Association
Use Freire catalog "Cluster" field.
Filter out foreground contaminants identified in literature.
```

B.3 Analysis Code

All analysis steps are encapsulated in Python scripts available in the `scripts/` directory. Key reproduction scripts include:

- `scripts/steps/step_02_pulsar_population_controls.py` : Implements the exact matching procedure for pulsar controls.
- `scripts/steps/step_12_hierarchical_density_scaling.py` : Runs the hierarchical mixed-effects model for density scaling.
- `scripts/steps/step_62_sn_ia_stretch_test.py` : SN Ia peak magnitude vs host velocity dispersion correlation (mB-σ test).

Appendix C: Microscopic Transfer Architecture and the κ Bridge

This appendix connects the observable pulsar response coefficient $\kappa_{\rm MSP}$ to the canonical scalar-sector architecture of Paper 0 (v0.16) and the volumetric Temporal-Topology susceptibility bridge established in Paper 11 (Step 62). While the primary empirical test treats $\kappa_{\rm MSP}$ as an observable pulsar response coefficient determined directly from data, the derivation below shows how the same saturation scale $\rho_T \approx 20\ \mathrm{g\,cm^{-3}}$ and universal coupling $\beta_A = -1$ govern both the extragalactic stellar-envelope amplification ($\kappa_{\rm gal} \sim 10^6$ mag) and the screened globular-cluster pulsar response.

The following derives the amplitude-sector transfer model on the admissible branch of the scalar sector, for the clock-amplitude channel of the Temporal-Topology response. As established in Paper 0 (v0.16), the shear-sector screening mechanism is the selected two-branch nonlinear kinetic completion $P(X,\phi) = -V(\phi) + \frac{2k}{3\Lambda_X^2} X|X|^{1/2} + X|X|/\Lambda_X^4$ ($P_{,X} = k\sqrt{\xi} + 2\xi$, $k \simeq 16.03$ fixed by a_0; unit-term control $X - V + X|X|/\Lambda_X^4$) — which pins the slope of the clock-rate landscape where it is already steep rather than flattening the field — while the quartic self-interaction $V(\phi) = \lambda\phi^4/4$ supplies the canonical matter-hosting amplitude-sector completion. What is fixed is the admissible branch itself: on the universal coupling $\beta_A = -1$, the inverse-power chameleon family admits no positive-density equilibrium and predicts the wrong environmental direction, and is excluded by constraint F1. The derivation below therefore uses the canonical matter-hosting potential — the quartic self-interaction retained in Paper 0 as the amplitude-sector completion — rather than the excluded branch.

C.1 Motivation

To satisfy precision tests in dense environments while allowing cosmological dynamics, a density-dependent mass may be adopted. On the canonical branch the operative potential is the matter-hosting factor of the master form of Paper 0, $V(\phi) = \lambda\phi^4/4$, the minimal admissible self-interaction for $\beta_A = -1$. The following derivation shows how such a microscopic framework could, in principle, produce the unsuppressed geometric factor of $\sim 10^6$, which is then heavily attenuated by environmental screening ($\mathcal{S}_{\rm MSP} \approx 0.04$) down to the empirically inferred $\kappa_{\rm MSP}^{\rm emp} \approx 3.8 \times 10^4$ response coefficient (Step 44).

C.2 Effective Potential and Field Equilibrium

For the canonical matter-hosting potential $V(\phi) = \lambda\phi^4/4$ — the weak-field branch of the master potential of Paper 0 — the effective potential including matter coupling is:

$$V_{\rm eff}(\phi;\rho) = \frac{\lambda}{4}\phi^4 + [A(\phi) - 1]\rho \approx \frac{\lambda}{4}\phi^4 + \frac{\beta_A\phi}{M_{\rm Pl}}\rho \tag{23}$$

where $A(\phi) = \exp(\beta_A\phi/M_{\rm Pl})$ and the approximation holds for small $\beta_A\phi/M_{\rm Pl}$. On the universal branch $\beta_A = -1$ the matter-coupling slope is negative, balancing the positive potential slope at a positive equilibrium field — precisely the sign structure the inverse-power family lacks on this branch. The equilibrium field value $\phi_{\rm min}(\rho)$ and effective mass $m_{\rm eff}(\rho)$ are (Paper 0):

$$\phi_{\rm min}(\rho) \approx \left(\frac{\rho}{\lambda M_{\rm Pl}}\right)^{1/3}, \quad m_{\rm eff}(\rho) \approx \sqrt{3}\,\lambda^{1/6}\left(\frac{\rho}{M_{\rm Pl}}\right)^{1/3} \tag{24}$$

Both quantities grow with ambient density and carry the correct signs: the equilibrium field is positive and deepens in denser environments ($\phi_{\rm min} \propto \rho^{1/3}$, consistent with $\phi > 0$ inside temporal wells), while the field is more massive and short-ranged in dense regions (Solar System, compact bodies) and light and dynamical in diffuse ones (clusters, cosmology). The quartic self-coupling is fixed by the corpus operative coupling $\lambda_{\rm Cassini} = 10^5\lambda_{\rm ref} \approx 7.5 \times 10^{-66}$ — the branch consistent with the corrected linear-in-S_Σ Cassini evaluation (Paper 0, step_03; the fiducial reference normalization $\lambda_{\rm ref} \approx 7.5 \times 10^{-71}$ fails that bound by $\sim 250\times$). The terrestrial saturation scale $R_T = \lambda_T$ of Paper 0 is a geometric identification $(3M/4\pi\bar{\rho})^{1/3}$, independent of λ; nothing in this appendix is separately tuned.

C.3 Weak-Field Mapping

The linear relation between the scalar field and Newtonian potential emerges from perturbations around equilibrium in the weak screening regime. Consider a cluster with background density ρ_0 and the associated equilibrium field value $\phi_{\rm min}(\rho_0)$. In the presence of a localized gravitational source, the total matter density becomes $\rho = \rho_0 + \delta\rho$, where $\delta\rho$ is related to the Newtonian potential through the Poisson equation: $\nabla^2\Phi = 4\pi G\delta\rho$.

Expanding the field around its equilibrium value, write $\phi = \phi_{\rm min}(\rho_0) + \delta\phi$. The equation of motion for the perturbation $\delta\phi$ in the static limit is:

$$\nabla^2\delta\phi - m_{\rm eff}^2(\rho_0)\delta\phi = +\frac{\beta_A}{M_{\rm Pl}}\delta\rho \tag{25}$$

In the weak screening regime ($\rho_0 \ll \rho_T$), the mass term is negligible on cluster scales. Substituting $\delta\rho = \nabla^2\Phi/(4\pi G)$:

$$\nabla^2\delta\phi \approx +\frac{\beta_A}{4\pi G M_{\rm Pl}}\nabla^2\Phi \tag{26}$$

Integrating and requiring $\delta\phi \to 0$ as $\Phi \to 0$ at infinity:

$$\delta\phi = +\frac{\beta_A}{4\pi G M_{\rm Pl}}\Phi \tag{27}$$

C.4 Response Coefficient Estimate

For $A(\phi) = \exp(\beta_A\phi/M_{\rm Pl})$, the proper time relation including the perturbation is:

$$\frac{d\tau}{dt} \approx \exp\left(\frac{\beta_A\phi_{\rm min}}{M_{\rm Pl}} + \frac{\beta_A\delta\phi}{M_{\rm Pl}}\right) \approx A(\phi_{\rm min})\left(1 + \frac{\beta_A\delta\phi}{M_{\rm Pl}}\right) \tag{28}$$

Comparing to the phenomenological form and substituting the linear mapping yields a candidate expression for the response coefficient:

$$\kappa_{\rm MSP}^{\rm emp} \approx \frac{\beta_A^2}{M_{\rm Pl}^2}\cdot\left(\frac{c^2}{4\pi G\rho_0 R_c^2}\right)\cdot\mathcal{S}(m_{\rm eff}, R_c) \tag{29}$$

where the unsuppressed geometric factor $c^2/(4\pi G\rho_0 R_c^2) \approx 10^6$–$10^7$ for typical cluster densities ($\rho_0 \sim 10^{-18}$ g/cm³) and core radii ($R_c \sim 1$ pc). On the admissible branch the screening factor $\mathcal{S}(m_{\rm eff}, R_c)$ collects the suppression accumulated across the nested domain hierarchy — the dense endpoints (the pulsar, its companion, the calibrator environment) carry a large $m_{\rm eff}$ and are strongly self-screened, while the diffuse cluster volume retains the lighter, active-shear field — yielding an aggregate attenuation of order $\mathcal{S} \sim 10^{-8}$ at the in-equation level. This strong attenuation squashes the unsuppressed $\sim 10^6$ coupling down to the inferred $\kappa_{\rm MSP}^{\rm emp} \approx 3.8 \times 10^4$ response coefficient — equivalently $\kappa_{\rm MSP}^{\rm emp}/\kappa_{\rm bare} = \mathcal{S}_{\rm MSP} \approx 0.04$ (Step 44). In the corpus-wide transfer architecture (Paper 0 §7; Paper 11, Step 62, `step_62_q_structure_derivation.json`), the unsuppressed $\sim 10^6$ magnitude-sector reference scale $\kappa_{\rm canonical} = 0.96 \times 10^6$ mag is grounded directly in the volumetric Temporal-Topology susceptibility ratio $\rho_T/\bar{\rho}_{\rm proc}$: for a classical Cepheid at the $P_0 = 10$ d pivot ($Q = 0.041$ d), the fundamental-mode period–mean-density relation $\langle\rho\rangle(10\ {\rm d}) = 2.367 \times 10^{-5}\ {\rm g\,cm^{-3}}$ yields $\rho_T/\langle\rho\rangle(10\ {\rm d}) = 8.450 \times 10^5$ at $\rho_T = 20\ {\rm g\,cm^{-3}}$, which multiplied by the Wesenheit Leavitt projector $\Gamma_{\rm PL} = |b_W|/\ln 10 = 1.416$ mag at the unit-transport benchmark $\lambda_0 = 1$ (with bare coupling $|\beta_A| = 1$ fixed by Paper 0) predicts $|\kappa_{P^2}^{\rm pred}(\lambda_0 = 1)| = 1.196 \times 10^6$ mag, matching the direct SH0ES P^2-carrier fit $(1.191 \pm 0.480) \times 10^6$ mag to 0.5% ($\lambda_0 = 0.995 \pm 0.401$, inverting to $\rho_T^{\rm (Cepheid)} = 19.91 \pm 8.02\ {\rm g\,cm^{-3}}$). While the Cepheid and galaxy inference channels (Papers 11–12) sample this unscreened diffuse-envelope amplification directly (with only the order-unity coupled-envelope acoustic weight $\lambda_0 \approx 1$ remaining as the registered microphysical closure target), the globular-cluster pulsar timing observable compounds the diffuse cluster core potential with the self-screened compact-endpoint boundary, attenuating the unsuppressed $\sim 10^6$ scale by $S_{\rm MSP} \approx 0.030$--$0.040$.

The pipeline (step_44_kappa_msp_prior.json) provides the corresponding data-driven normalisation: by inverting the observed 0.63 dex raw excess with real cluster parameters (core radii 0.1–0.5 pc, 64 pulsars across 7 clusters), it derives a full response coefficient $\tilde{\kappa}_{\rm MSP} = (2.9 \pm 4.5) \times 10^4$ with an effective screening factor $S_{\rm MSP} \approx 0.030$ relative to the unsuppressed $\kappa_{\rm canonical} = 0.96 \times 10^6$ scale. The density-dependent screening factor $\mathcal{S} \sim 10^{-8}$ and the pipeline's geometric suppression factor $S_{\rm MSP} \approx 0.030$ operate at different levels of the normalisation hierarchy: $\mathcal{S}$ suppresses the unsuppressed microscopic coupling to the in-equation coefficient $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$, while $S_{\rm MSP}$ suppresses the unsuppressed geometric factor to the full response coefficient $\tilde{\kappa}_{\rm MSP} \sim 10^4$. Paper 11 uses the latter for cross-probe comparison with $\kappa_{\rm Cep}$.

Data Availability & Reproducibility

> This work follows open-science practices. All results are fully reproducible from raw data using the documented pipeline. All numerical results, figures, and statistics are generated by deterministic Python scripts processing real observational data.
>
> *Canonical source:* The canonical source of this manuscript is the HTML component system (`site/components/*.html`). The Markdown and PDF versions are auto-generated build artifacts. Direct edits to derived files are non-canonical and are overwritten on rebuild.

Repository & Code

GitHub Repository: github.com/matthewsmawfield/TEP-COS

The repository contains a deterministic, version-controlled analysis pipeline with analysis steps for pulsar timing data. All steps are orchestrated by `scripts/run_pipeline.py` with comprehensive logging.

Repository Structure

```
TEP-COS/
├── data/                          # Raw observational data
│   ├── supernovae/               # Pantheon+ SN Ia data
│   └── pulsars/                  # Pulsar reference data
├── scripts/
│   ├── steps/                    # Analysis pipeline steps
│   ├── utils/                    # Utility functions (logger.py)
│   └── run_pipeline.py          # Master orchestration script
├── results/
│   ├── outputs/                  # JSON/CSV analytical outputs
│   └── figures/                  # Generated plots (PNG/PDF)
├── logs/                         # Per-step execution logs
├── site/
│   └── components/               # Manuscript HTML sections
├── reproduce_manuscript.py       # Legacy wrapper (deprecated)
├── requirements.txt              # Python dependencies
└── README.md                     # Documentation
```

Data Provenance

Data Source	Provider	Access Method	Size	Location
ATNF Pulsar Catalogue	ATNF	Auto-downloaded	~10 MB	`results/outputs/atnf_psrcat.db`
CMC Cluster Catalogs	Kremer et al. 2020	Auto-downloaded	~25 GB extracted (~4.8 GB compressed)	`data/cmc/` (20 cluster selections, 14 unique realizations, 9,526 evolved neutron stars / 6,617 unique)
Pantheon+ SNe Ia	Scolnic et al.	Auto-downloaded	~2 MB	Via astroquery

Pipeline Architecture

The analysis pipeline comprises 26 deterministic steps organized into logical groups. Each step is a standalone Python script in `scripts/steps/` that produces JSON outputs and detailed logs in `logs/step_*.log` .

Complete Step Inventory & Runtime

Group	Step	Script	Description	Runtime
Section 3: Pulsar Timing Analysis				
Data	3.0 Prep	`step_05_freire_gcpsr_radial_analysis.py`	Radial analysis of GC pulsars (Freire catalog)	~0.5s

Group	Step	Script	Description	Runtime
Data	3.1 Sample	`step_02_pulsar_population_controls.py`	Population controls: period/B-field matching (400 MSPs)	~10.8s
Core	3.2 Maximal	`step_06_hybrid_maximum_analysis.py`	Hybrid maximal sample construction (GC + Field)	~6s
Core	3.3 Density	`step_07_per_cluster_controlled_residuals.py`	Per-cluster controlled residuals vs density	~1s
Core	3.4 Scaling	`step_11_full_density_scaling.py`	Full density scaling simulation	~2s
Core	3.5 Hierarchical	`step_12_hierarchical_density_scaling.py`	Hierarchical mixed-effects density analysis	~1s
Core	3.6 Validation	`step_13_covariance_validation.py`	Covariance-aware statistical validation	~3s
Binary	3.7 GC Binary	`step_15_binary_pulsar_analysis.py`	Binary vs isolated MSPs in GCs	~0.5s
Binary	3.8 Field Binary	`step_20_field_binary_analysis.py`	Field binary control analysis	~0.5s
Binary	3.9 Integrated	`step_21_integrated_binary_control.py`	Integrated binary control test (GC vs Field)	~0.6s
Section 3.6: CMC Catalog Analysis				
CMC	3.10a Download	`download_cmc_data.py`	Download CMC cluster catalogs (20 clusters, ~25 GB extracted)	~12 min
CMC	3.10b Analysis	`step_37_cmc_gold_standard_analysis.py`	Comparison of observed versus CMC synthetic pulsars (20 cluster selections, 14 unique realizations, 9,526 evolved neutron stars)	~66.5s
CMC	3.11 Exotic	`step_41_exotic_physics_quantification.py`	Evaluate exotic-GR mechanism burden	~16.7s
CMC	3.12 PTA Mock	`step_49_pta_mock_observation.py`	Mock radio observations testing observational filtering defense	~2s
Section 4: Sensitivity & Validation				
Valid	4.1 Shklovskii	`step_23_shklovskii_sensitivity.py`	Shklovskii correction sensitivity analysis	~1s
Valid	4.2 Rho Sensitivity	`step_25_rho_sensitivity.py`	Rho_intra sensitivity analysis	~0.7s
Valid	4.3 Power	`step_26_power_analysis.py`	Statistical power analysis	~9.9s
Valid	4.4 Monte Carlo	`step_27_monte_carlo_validation.py`	Monte Carlo validation (Type I, Power, Bias)	~8.7s
Appendix (Brief Supplementary Notes & Archived Tests)				
App	A.1 SN Ia	`step_62_sn_ia_stretch_test.py`	SN Ia mB-σ correlation (exploratory only; indistinguishable from mass-step effect)	~1s
App	A.2 MaNGA (Archived)	`step_59_manga_spatially_resolved.py`	MaNGA age gradients (archived; fossil probe cannot distinguish TEP)	~300s
App	A.3 MaNGA (Archived)	`step_60_manga_test_e_age_discrepancy.py`	LW vs MW age discrepancy (archived; fossil probe cannot distinguish TEP)	~300s
Fig	7.6 Galaxy Fig (Archived)	`step_59_manga_spatially_resolved.py`	MaNGA age gradient figures (archived)	~1s

Total Runtime Summary

Runtimes measured on Apple M4 Pro with `--parallel` flag. First runs include CMC data processing and may take 2–3× longer.

Component	Steps	Runtime
Pulsar Timing (Section 3)	10	~22s
CMC N-Body Analysis (3.10)	4	~85s
Sensitivity & Validation	4	~21s
Figure Generation	4	~3s
Appendix	1	~1s
Total	**23**	**~148s (~2.5 min)**

Reproduction Instructions

Quick Start (Full Reproduction)

```
# 1. Clone repository
git clone https://github.com/matthewsmawfield/TEP-COS.git
cd TEP-COS

# 2. Install dependencies
pip install -r requirements.txt

# 3. Run full pipeline (generates all results & figures)
python scripts/run_pipeline.py

# 4. Results will be in:
#    - results/outputs/   (JSON/CSV data)
#    - results/figures/   (PNG/PDF plots)
#    - logs/              (Detailed execution logs)
```

Command-Line Options

The pipeline supports selective execution for faster testing:

```
# Fast mode: core analysis only (skips long validations)
python scripts/run_pipeline.py --only-core

# Skip validation steps
python scripts/run_pipeline.py --skip-validation

# Skip figure generation
python scripts/run_pipeline.py --skip-figures

# Legacy wrapper (deprecated, calls run_pipeline.py)
python reproduce_manuscript.py
```

System Requirements

Component	Minimum	Recommended	Tested On
CPU	4 cores	8+ cores	Apple M4 Pro (14-core)
RAM	8 GB	16 GB	24 GB (M4 Pro)
Storage	5 GB	15 GB (includes 4.8 GB CMC data)	NVMe SSD
Runtime	~10 min	~6 min	~5.7 min (M4 Pro)

Key Analysis Outputs

- `results/outputs/step_02_pulsar_population_controls.csv` — Base pulsar dataset (400 MSPs with 198 GC, 202 field)

- `results/outputs/step_06_hybrid_maximum_analysis.json` — Expanded hybrid pulsar summary (550 MSPs; 0.40 dex hybrid-controlled residual)
- `results/outputs/step_15_binary_pulsar_analysis.json` — Binary vs isolated analysis results
- `results/outputs/step_12_hierarchical_density_results.json` — Mixed-effects density-scaling results
- `results/outputs/step_13_covariance_validation.json` — Covariance-aware and LOOCV validation
- `results/outputs/step_21_integrated_binary_control.json` — Integrated differential test
- `results/outputs/step_37_cmc_gold_standard.json` — CMC catalog comparison (20 cluster selections, 14 unique realizations, 9,526 evolved neutron stars / 6,617 unique; 13.2σ nominal amplitude overprediction)
- `results/outputs/step_41_exotic_physics_quantification.json` — Exotic-GR mechanism evaluation
- `results/outputs/step_49_pta_mock_observation.json` — PTA mock observation results (observational filtering defense)

Log Files

Each step produces detailed logs:

- `logs/pipeline_master.log` — Master pipeline execution log
- `logs/step_*.log` — Individual step logs (26 files)

Software Dependencies

Package	Version	Purpose
Python	3.10+	Language runtime
NumPy	1.24+	Numerical computing
SciPy	1.10+	Statistical functions
Pandas	2.0+	Data manipulation
Matplotlib	3.7+	Visualization
Astropy	5.0+	Astronomical calculations
Astroquery	0.4.6+	Data queries (SDSS, etc.)
Joblib	1.3.0+	Parallel processing

All dependencies are specified in `requirements.txt`.

Unified TEP Parameter Dictionary

The TEP corpus uses related but distinct symbols across its papers. This dictionary maps every parameter, its definition, the paper where it is primary, and its fiducial or fitted value.

Symbol	Definition	Primary Paper	Fiducial / Fitted Value
β_A	Microscopic conformal coupling: $A(\phi) = \exp(\beta_A \phi / M_{\rm Pl})$	TEP (Paper 0)	-1.0 (locked lab-scale convention; not varied in pulsar channel)
ρ_T	Temporal Topology saturation scale (series-level)	TEP-UCD (Paper 6)	≈ 20 g/cm³
$R_T(M)$	Temporal Topology saturation radius as function of mass	TEP-UCD (Paper 6)	$R_T \propto M^{1/3}$
$\kappa_{\rm MSP}^{\rm emp}$	In-equation acceleration coupling in the pulsar spin-down equation; screened by the density-dependent factor $\mathcal{S} \sim 10^{-8}$ from the unsuppressed $\sim 10^6$ geometric factor	TEP-COS (Paper 10)	≈ 0.05 (dimensionless)
$\tilde{\kappa}_{\rm MSP}$	Full empirical response coefficient from pipeline inversion of the 0.63 dex raw excess with real cluster parameters; suitable for cross-probe comparison with $\kappa_{\rm Cep}$	TEP-COS (Paper 10)	$(2.9 \pm 4.5) \times 10^4$ (dimensionless)
$S_{\rm MSP}$	Effective screening factor: ratio of $\tilde{\kappa}_{\rm MSP}$ to the unsuppressed geometric factor $\sim 10^6$	TEP-COS (Paper 10)	≈ 0.030

Symbol	Definition	Primary Paper	Fiducial / Fitted Value
$\mathcal{S}(m_{\rm eff}, R_c)$	Density-dependent screening factor suppressing the unsuppressed microscopic coupling to the in-equation coefficient	TEP-COS (Paper 10)	$\sim 10^{-8}$ aggregate across the nested domain hierarchy
Γ	Density-scaling slope of the spin-down excess: $d(\log \lvert\dot{P}\rvert)/d(\log \rho)$	TEP-COS (Paper 10)	Observed: 0.39 ± 0.08; Newtonian: 0.75; TEP prediction: 0.50
$\Gamma_{\rm TEP} = 2\Gamma_N - 1$	Parameter-free TEP prediction for the density-scaling slope, derived from potential-based ($\lvert\Phi\rvert \propto \rho R_c^2$) vs acceleration-based ($a \propto \rho R_c$) scaling	TEP-COS (Paper 10)	0.50 ± 0.08 (matches observation at 0.9σ)
$\kappa_{\rm canonical}$	Prespecified theory benchmark (Paper 0 §7); unscreened reference used in the $S_{\rm MSP}$ ratio	TEP (Paper 0 §7)	0.96×10^6 mag
$\alpha_{\rm clock}$	Shared underlying clock-response scale; unifies $\kappa_{\rm Cep}$ and $\tilde{\kappa}_{\rm MSP}$ via channel-specific transfer factors	TEP-H0 (Paper 11)	Derived; not independently fitted
$T_{\rm GC}, T_{\rm disk}$	Channel-specific environmental transfer factors: $T_{\rm GC} \sim 10^{-2}$–10^{-1} (dense clusters), $T_{\rm disk} \sim 1$ (galactic disks)	TEP-H0 (Paper 11)	Derived from cluster/disk density contrast
$S(\rho)$	Continuous shear-suppression factor: $S(\rho) = [1 + (\rho/\rho_{\rm half})^2]^{-1}$	TEP-H0 (Paper 11) / TEP-C0 (Paper 26)	$\rho_{\rm half} \approx 0.5\, M_\odot/{\rm pc}^3$
$\mathcal{S}_\Sigma(\mathcal{E})$	Unified covariant screening operator; reduces to $S(\rho)$ or gradient-dependent $f(g)$ in appropriate limits	TEP-C0 (Paper 26)	Environment-dependent

Note: The two pulsar normalisations — $\kappa_{\rm MSP}^{\rm emp} \approx 0.05$ (in-equation coupling) and $\tilde{\kappa}_{\rm MSP} \sim 10^4$ (full response coefficient) — trace back to the same unsuppressed geometric factor $\sim 10^6$ but absorb different combinations of the potential and acceleration normalisation. The cross-probe comparison with the Cepheid coefficient $\kappa_{\rm Cep}$ uses $\tilde{\kappa}_{\rm MSP}$, which is the quantity directly analogous to $\kappa_{\rm Cep}$. The key observational signature of the pulsar channel is the suppressed density scaling ($\Gamma = 0.39$ vs Newtonian 0.75), reflecting the screened TEP shear response saturating where the unscreened Newtonian drift would grow with the potential gradient.