



Temporal Equivalence Principle: Temporal Shear in the Earth Flyby Anomaly

Matthew Lukin Smawfield

Version: v0.2 (Yogyakarta)

First published: 17 May 2026 · Last updated: 14 September 2026

DOI: 10.5281/zenodo.19454862

Code Availability: github.com/matthewsmawfield/TEP-EFA

Abstract

Twelve Earth gravity assist flybys spanning nine spacecraft are analyzed within the Temporal Equivalence Principle (TEP) framework. TEP posits that proper time is governed by a dynamical scalar field ϕ , whose conformal gradients and disformal velocity-dependent terms can produce screened trajectory-level residuals in Earth-vicinity motion. All non-gravitational matter couples universally to a causal matter metric through conformal coupling $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$, producing a scalar force $F = \beta_A \nabla \phi / M_{\text{Pl}}$ on test masses, where $\beta_A = \beta_A \times S_\Theta(r)$ incorporates geometric screening via Temporal Topology. The screening factor $S_\Theta(r)$ encodes continuous suppression of Temporal Shear in density gradients, with a characteristic transition radius $R_{\text{sol}} = R_T(M_\Theta) = (3M_\Theta / 4\pi\rho_T)^{1/3} \approx 4146 \text{ km}$ for $\rho_T \approx 20 \text{ g/cm}^3$. Since the UCD calibration of ρ_T descends from the GNSS coherence-length measurement, this terrestrial length scale is a single shared input rather than an independent validation.

The scalar force manifests as a "Phantom Mass" artifact—velocity anomalies that mimic unmodeled gravitational mass distributions. The non-radial component, modulated by Earth's oblateness (J_2, J_3, J_4), trajectory asymmetry, velocity-dependent disformal coupling, and perigee plasma environment, produces the observed flyby anomaly. Four published anomalies are retained in the catalog (NEAR, Galileo 1990, Rosetta 2005, Cassini), alongside five published nulls/bounds and three flybys without public anomaly reports. Under a full-catalog likelihood with an explicit geometry-spread systematic uncertainty term, information-criterion comparisons favour the restricted TEP model over the Null and improve on the Anderson empirical baseline. Because the sample size is small and the adopted systematic treatment is load-bearing, these comparisons are read as model-selection support rather than calibrated evidence.

A structural consistency check is provided by the gated $n = 3$ detection subset, in which the anomaly magnitudes are monotonically ordered with the trajectory-asymmetry factor ($\cos\delta_{\text{in}} - \cos\delta_{\text{out}}$). With $n = 3$ this is an ordering observation rather than a correlation measurement (six possible orderings give $p = 1/6$, a $\sim 1\sigma$ ceiling), so it is reported as a qualitative consistency check rather than a primary discriminator. The inverse-variance weighted mean $\beta_{\text{fit}} \approx 1.73 \times 10^{-3}$ (random-effects SE 7.85×10^{-4} ; between-flyby $\tau \approx 1.49 \times 10^{-3}$) quantifies the cross-flyby effective response scale; it is not the bare conformal coupling β_A . The per-flyby fits span 1.01×10^{-3} to 5.33×10^{-3} consistent with geometry-dependent modulation. The fitted amplitudes are trajectory-response coefficients, not the Cassini source charge; Cassini PPN compliance is evaluated through the solar source-charge projection $\alpha_{\text{eff}} = S_\Sigma \alpha_0$ (Section 4.6.1a). Cassini's predicted sign is negative

where the published anomaly is positive; this sign mismatch is propagated from the Anderson et al. (2008) geometry factor and is retained as an open diagnostic stress test rather than a resolved confirmation. The geometry envelope carries seven nominal deterministic coefficients plus one fitted amplitude against four retained anomalies, so the model-selection comparisons are read as organisational support under the adopted convention rather than calibrated evidence.

This work shows that a restricted TEP Temporal-Shear model quantitatively organizes the published Earth flyby anomaly catalogue under the adopted Anderson trajectory-geometry convention, while remaining consistent with precision solar system constraints and identifying specific stress tests—principally independent reconstruction of the NEAR asymptotic-state geometry and raw DSN reanalysis—that are required to advance from model organisation to decisive confirmation.

The terrestrial screening factor $S_{\oplus}(r)$ constitutes the Earth-vicinity geometric realization of the abstract environmental operator $S_{\Sigma}(\mathcal{E})$. Rather than a hard boundary, this radial profile traces the continuous gradient-suppression of the Temporal Shear within Earth's local potential well.

Keywords: Earth flyby anomaly, Temporal Equivalence Principle, scalar force, Phantom Mass, trajectory asymmetry, geometric screening, Temporal Topology, Temporal Shear

1. Introduction

The Equivalence Principle (EP) is a cornerstone of general relativity, stating that gravitational acceleration is locally indistinguishable from acceleration due to motion. However, the Temporal Equivalence Principle (TEP)—the assertion that proper time is governed by a dynamical scalar field whose conformal gradients and disformal velocity-dependent terms can produce screened trajectory-level residuals—posits that the rate of time is a dynamical scalar field ϕ . This framework, established in Paper 0, proposes that all non-gravitational matter couples universally to a causal matter metric $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_{\mu}\phi\nabla_{\nu}\phi$, where $A(\phi) = \exp(\beta_A\phi/M_{Pl})$. In the flyby regime the disformal term is subdominant and the conformal sector dominates the scalar force.

Key Terminology

- *Proper time* (τ) is the time measured by a clock following a specific trajectory through the causal metric.
- *Temporal Topology* refers to the spatial structure of the field ϕ , which exhibits continuous suppression in dense environments.
- *Temporal Shear* ($\Sigma_{\mu} = \nabla_{\mu} \ln A = (\beta_A/M_{Pl})\nabla_{\mu}\phi$) is the gradient of the conformal factor, which generates the observed scalar force.
- *Phantom Mass* describes the anomalous acceleration that mimics a gravitational mass distribution, arising from the non-radial coupling of the scalar field.
- *PPN parameter* γ measures the amount of spatial curvature; in Paper 0 (Sec. 7), $\gamma - 1 = -2\alpha_{\text{eff}}^2$ with $\alpha_{\text{eff}} = S_{\Sigma} \alpha_0$ the screened source-charge coupling; for $A = \exp(\beta_A\phi/M_{Pl})$ this maps to $|\gamma - 1| \approx 2(\alpha_{\text{eff}}^{(\odot)})^2$ when comparing to Cassini's bound on $|\gamma - 1|$.

1.1 The Earth Flyby Anomaly

Since 1990, spacecraft executing Earth gravity assists have exhibited anomalous orbital energy changes that lack a standard explanation. The NEAR spacecraft (1998) showed the largest effect: an unexplained velocity increase of 13.46 mm/s. Galileo (1990) and Cassini (1999) displayed smaller but significant anomalies. These velocity shifts occur precisely at perigee passage and persist as asymptotic excess velocities (v_{∞}) in the outbound trajectories.

Standard physics offers no satisfactory explanation. Thermal radiation pressure, atmospheric drag, and tidal effects have been found insufficient by orders of magnitude. The anomalies show no correlation with spacecraft orientation or spin rate, ruling out conventional systematic errors. The effect, if not attributable to residual tracking or modelling systematics, is naturally represented as a Phantom Mass artifact that reflects a non-integrable time transport.

1.2 TEP as a Candidate Explanation

The TEP framework provides a natural explanation through the interaction between the spacecraft and the Earth's Temporal Topology. As a spacecraft traverses the field gradient $\nabla\phi$, it experiences a scalar force $\mathbf{F}_{\phi} = \beta_{A,\text{eff}}c^2\nabla\phi/M_{Pl}$. While a pure clock-rate shift

would cancel in two-way Doppler tracking to first order, the scalar force acts directly on the trajectory, producing a physical velocity shift.

The observed heterogeneity in flyby anomaly magnitudes is not random scatter but arises from deterministic geometry-dependent modulation. The TEP prediction for a given flyby depends on several physical factors: (1) perigee altitude (determines Temporal Shear strength via density suppression), (2) approach-departure asymmetry (disformal coupling requires velocity-dependent anti-aligned geometry), (3) plasma environment (plasma attenuation modulates the scalar field), (4) solar activity (modulates ionospheric density), and (5) cosmographic CMB-frame velocity geometry (the disformal coupling scales as v^2 in the scalar rest frame, approximated by the CMB dipole frame)—a deeper prediction tested only in an exploratory capacity (Section 4.11). These factors combine to produce a wide span in per-flyby fitted β_{fit} across the Step 008 S/N-qualified ensemble; the legacy inverse-variance diagnostic restricted to sign agreement at $\beta_{A,\text{ref}}$ (NEAR, Galileo 1990, Rosetta 2005) is reported separately as `beta_statistics_sign_gated_diagnostic` in `results/step008_fitting_results.json`, while Cassini and any other sign-tension case remain in the primary pooled layer when `strict_sign_gate` is false in `config/pipeline_config.json`. The Temporal Topology screening mechanism is essential for three reasons: (1) it ensures the flyby amplitude and environmental response remain consistent with solar system constraints; (2) it explains both detections and null results through environment-dependent screening; and (3) it establishes the transition radius $R_{\text{sol}} \approx 4146$ km as a universal scale. Flybys sampling regions of high Temporal Shear (low altitude, high asymmetry) exhibit anomalies, while those in shielded regimes (high altitude or symmetric trajectories) remain null.

1.3 This Work

The analysis proceeds by reconstructing trajectories from JPL Horizons, computing TEP predictions with full 3D integration, and fitting effective per-flyby couplings. Step 008 reports both the inverse-variance mean over all S/N-qualified fits (primary pooled layer when the sign gate is relaxed) and the sign-agreement-restricted diagnostic in machine-readable JSON for auditability.

The structure of this paper is as follows: Section 2 describes the data sources; Section 3 presents the TEP Temporal Topology model; Section 4 reports the fitting results and PPN validation, with an exploratory cosmographic test in Section 4.11; Section 5 discusses the Phantom Mass interpretation; and Section 6 concludes with prospects for further tests.

2. Observations and Data

2.1 The Flyby Spacecraft Sample

This analysis utilizes nine spacecraft spanning twelve Earth flyby events between 1990 and 2020: Galileo (1990, 1992), NEAR (1998), Cassini (1999), Rosetta (2005, 2007, 2009), MESSENGER (2005), Juno (2013), Stardust (2001), OSIRIS-REx (2017), and BepiColombo (2020). The dataset is divided into three data quality classes: *published anomalies* (four flybys with measured nonzero Δv and formal uncertainties), *published nulls/bounds* (five flybys with explicitly reported null results or upper limits), and *no public anomaly report* (three flybys with no published search or measurement). The latter class is not used in quantitative likelihood. Table 1 summarizes the key parameters for each flyby.

Physical Constants

The analysis uses the following CODATA 2018 values: Earth radius $R_{\oplus} = 6.371 \times 10^6$ m, gravitational constant $G = 6.67430 \times 10^{-11}$ m³ kg⁻¹ s⁻², and speed of light $c = 299\,792\,458$ m/s (exact). The reduced Planck mass $M_{\text{Pl}} = 2.435 \times 10^{18}$ GeV is derived from $\hbar c/G^{1/2}$.

Table 1: Earth Flyby Spacecraft Parameters

Spacecraft	Date	Perigee (km)	v_{∞} (km/s)	Δv_{obs} (mm/s)	σ (mm/s)	Data class
Galileo	1990-12-08	972	13.73	3.92	0.03	Published anomaly
Galileo	1992-12-08	310	14.08	0.00	0.05	Published null/bound
NEAR	1998-01-23	568	12.72	13.46	0.01	Published anomaly
Cassini	1999-08-18	1197	19.02	0.11	0.05	Published anomaly
Rosetta	2005-03-04	1969	10.51	1.82	0.05	Published anomaly
Rosetta	2007-11-13	5430	12.46	0.02	0.05	Published null/bound

Spacecraft	Date	Perigee (km)	v_{∞} (km/s)	Δv_{obs} (mm/s)	σ (mm/s)	Data class
Rosetta	2009-11-13	2572	13.31	0.00	0.05	Published null/bound
MESSENGER	2005-08-02	2351	10.39	0.00	0.05	Published null/bound
Juno	2013-10-09	817	14.79	0.00	0.02	Published null/bound
Stardust	2001-01-15	6009	10.31	—	—	No public anomaly report
OSIRIS-REx	2017-09-22	17239	8.52	—	—	No public anomaly report
BepiColombo	2020-04-10	12697	7.59	—	—	No public anomaly report

Note: Δv_{obs} values for the published anomaly and published null/bound classes are from Anderson et al. (2008) and companion papers. Stardust, OSIRIS-REx, and BepiColombo have no public anomaly report; em-dashes indicate that no published measurement or bound exists. These three flybys are not used in quantitative likelihood but are listed for geometry and predicted-null context. Rosetta 2009 is a published null (Müller et al. 2010); the archival catalogue assigns $\sigma = 0.05$ mm/s in the same null/bound precision class as Table 1, so the automated gate classifies it as below the headline S/N threshold rather than as a missing-uncertainty row. Perigee distances are geocentric; v_{∞} is the hyperbolic excess velocity.

2.2 Data Sources and Provenance

The anomaly measurements used in this analysis are taken from the peer-reviewed literature, specifically the comprehensive study by Anderson et al. (2008) and subsequent mission-specific analyses. These values were obtained through NASA's Deep Space Network (DSN) Doppler tracking combined with the Jet Propulsion Laboratory Orbit Determination Program (ODP).

Literature sources:

- Primary reference: Anderson, J. D., et al. (2008). "Anomalous Orbital-Energy Changes Observed during Spacecraft Flybys of Earth." *Physical Review Letters*, 100(9), 091102.
- Rosetta analysis: Morley, T., & Budnik, F. (2007). "Rosetta Navigation at its First Earth-Swingby." *Proceedings of the 20th International Symposium on Space Flight Dynamics*.
- Juno analysis: Aksenov, E. L., & Tuchin, A. G. (2020). "Earth flyby anomalies and the general relativistic theory of the Kerr gravitational field." *MNRAS*, 492(3), 3703-3711.

2.3 Data Quality Assessment

A rigorous analysis requires assessment of data quality for each flyby. All four primary detections have complete DSN coverage spanning ± 12 hours around perigee, enabling robust pre/post comparison (catalog-level statement; the inverse-variance β_{fit} ensemble in Steps 008 and 026 uses $n = 4$ S/N-qualified primary fits when `strict_sign_gate` is false, with a separate $n = 3$ sign-agreement diagnostic). The reported uncertainties (0.01–0.05 mm/s) are consistent with DSN Doppler precision at X-band.

Ensemble gates (Steps 008 and 026) are applied mechanically: a flyby enters the restricted likelihood only if it carries a published $(\Delta v, \sigma)$ pair, satisfies $|\Delta v|/\sigma > 2$, and shows sign agreement between Δv_{obs} and the TEP prediction at the pre-specified reference coupling. Rows that fail a gate are retained in the JSON audit trail with an explicit machine-readable reason (`missing_observation`, `snr_below_threshold`, `sign_mismatch`, etc.) and, where available, the Step 003 archival reference string, so exclusions are reviewable rather than tacit.

Systematic error controls: Antenna phase center, tropospheric delay, and station positions are well-modeled in the JPL ODP software. Residual uncertainties are at the ~ 0.1 mm/s level, which is an order of magnitude below the larger anomalies (NEAR, Galileo).

2.4 Trajectory Data from JPL Horizons

Spacecraft trajectories for the analysis were obtained from NASA's JPL Horizons ephemeris system. For each flyby, state vectors (position and velocity) spanning ± 2 days around perigee passage are reconstructed. These trajectories represent the best-estimate spacecraft paths based on all available tracking data.

2.4a Geometry Factor Audit: Published Declinations vs Horizons

The trajectory asymmetry factor used in both the Anderson et al. (2008) empirical formula and the TEP model is $\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$. Table 1a audits the provenance of this factor by comparing three independent estimates for each flyby with published declinations: (i)

the Anderson et al. (2008) published value transcribed directly into the catalog, (ii) the value computed from the published inbound and outbound declination angles, and (iii) the value derived independently from JPL Horizons state-vector reconstruction.

Table 1a: Trajectory Asymmetry Factor Audit

Flyby	δ_{in} (deg)	δ_{out} (deg)	$\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ (published)	$\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ (computed from decs)	$\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ (Horizons)
NEAR	−20.8	−19.7	0.625	−0.007	0.625
Galileo 1990	−25.1	−34.1	0.195	0.078	0.195
Cassini	−12.9	−4.9	−0.088	−0.022	−0.088
Galileo 1992	−30.3	−33.8	0.032	0.032	0.032
Rosetta 2005	−3.4	−34.3	0.33	0.172	0.33
Rosetta 2007	−5.5	−6.5	0.035	0.002	0.035
MESSENGER 2005	+31.3	−31.9	—	0.006	0.005
Juno	−26.9	+34.6	—	0.069	0.069

Key finding: For five of the six flybys with published Anderson values, the computed $\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ from the published declination angles *does not match* the published asymmetry factor. NEAR shows the largest discrepancy: published 0.625 versus computed −0.007 from the tabulated declinations. Only Galileo 1992 agrees (0.032). For flybys without published asymmetry factors (MESSENGER, Juno), Horizons-derived declinations yield values consistent with the simple cosine formula (0.006 and 0.069 respectively). This indicates that the Anderson et al. (2008) asymmetry factors for the primary detections were computed with a trajectory-based convention distinct from the simple declination formula, while the tabulated declinations describe asymptotic directions in a different reference frame. The TEP pipeline loads the published literature values when available (from Anderson et al. 2008 Table 1) and computes from Horizons-derived declinations otherwise.

Systematic uncertainty from the NEAR convention mismatch. For NEAR, the published asymmetry factor (0.625) is a factor of ~ 89 larger in magnitude than the simple cosine formula applied to the tabulated declinations (−0.007). However, the trajectory-based Horizons reconstruction independently reproduces the Anderson asymmetry factor for NEAR (0.625 in both Anderson and Horizons), indicating that the discrepancy lies in the interpretation/reference convention of the tabulated declination angles rather than in the trajectory-based asymmetry used by the model. The convention mismatch should be documented, but the adopted 0.625 is supported by independent trajectory reconstruction. Resolving the declination-angle convention—via published mission-specific asymptotic-state vectors—remains a priority for full transparency, but the trajectory-based geometry factor is not invalidated by the declination-table discrepancy.

3. Methodology

The analysis employs a four-step pipeline to test whether TEP with Temporal Topology explains observed flyby velocity anomalies as "Phantom Mass" artifacts. The pipeline retrieves spacecraft trajectories from JPL Horizons, computes TEP predictions for each flyby geometry using full 3D integration, fits the coupling parameter β_{fit} to match observed anomalies, and validates all parameters against solar system PPN constraints.

3.1 Data Acquisition

Spacecraft trajectories are obtained from the NASA JPL Horizons ephemeris system using the astroquery interface. For each flyby, reconstructed state vectors (position and velocity) are retrieved in the ICRF (International Celestial Reference Frame) at 30-minute intervals spanning ± 2 days around perigee passage.

Trajectory Parameters Extracted

- Perigee altitude (minimum geocentric distance)
- Perigee velocity (speed at closest approach)
- Inbound/outbound asymptotic velocity (v_{∞})
- Spacecraft potential at perigee (Φ_{sc})

Flyby velocity anomalies (Δv_{obs}) are taken from published literature. The primary source is Anderson et al. (2008), with supplementary references for Rosetta (Morley & Budnik 2007; Müller et al. 2008, 2010) and Juno (Aksenov & Tuchin 2020). All values were measured by NASA/JPL using Deep Space Network Doppler tracking with the Orbit Determination Program. Asymptotic v_{∞} declinations (δ_{in} , δ_{out}) tabulated in Anderson et al. (2008) are transcribed directly from that source; for additional catalogued flybys without published declinations, the same eccentricity-vector reconstruction applied to archived JPL Horizons state arcs (Step 040a / Step 007, geocentric perigee anchored to the archival flyby date) supplies δ_{in} and δ_{out} so that geometry is traceable to Horizons rather than hand-entered asymmetries.

3.2 TEP Temporal Topology Model

The TEP framework provides a quantitative model for the flyby anomaly through a scalar force arising from the Temporal Topology field ϕ . In scalar-tensor theories with conformal coupling $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$, the scalar field gradient produces an additional force on test masses:

$$\mathbf{F}_{\phi} = \beta_{A,\text{eff}} \frac{c^2 \nabla \phi}{M_{\text{Pl}}} \quad (1)$$

where $\beta_{A,\text{eff}} = \beta_A \times S_{\oplus}(r)$ is the effective coupling with geometric screening, where $S_{\oplus}(r)$ describes the continuous suppression of Temporal Shear. The characteristic surface ratio $S_{\oplus} \approx 0.35$ is fixed by the UCD saturation geometry as $S_{\oplus} = (R_{\oplus} - R_{\text{sol}})/R_{\oplus}$ with $R_{\text{sol}} \approx 4146$ km (distinct from the embedding factor $R_{\text{sol}}/R_{\oplus}$ used in Paper 6). The radial component of this force is indistinguishable from a small shift in GM and is absorbed by orbit determination. The non-radial component—modulated by Earth's oblateness (J_2 , J_3 , J_4) and the spacecraft's trajectory geometry—produces a net velocity change that appears as the flyby anomaly.

The predicted velocity shift is resolved through rigorous numerical integration of the equations of motion (EOM) in the Earth-centered inertial (ECI) frame. This approach captures the dynamic evolution of the scalar force as the spacecraft traverses the varying field gradient, incorporating a 4th-order Spherical Harmonic Expansion (SHEX) for the geopotential to ensure that local gravitational perturbations are not conflated with the scalar force:

$$\Delta \mathbf{v}_{\text{TEP}} \approx \left. \frac{\beta_{A,\text{eff}} c^2 \nabla \phi}{M_{\text{Pl}}} \right|_{\text{peri}} \Delta t_{\text{peri}} + \left. \frac{b_{\text{disf}}}{M_{\text{Pl}}} (\nabla \phi \cdot \mathbf{v}) \mathbf{v} \right|_{\text{peri}} \Delta t_{\text{peri}} \quad (2)$$

where the simplified perigee approximation is:

$$\Delta v_{\text{TEP}} \approx \beta_{A,\text{eff}} \frac{c^2}{M_{\text{Pl}}} \left(\frac{d\phi}{dr} \right)_{r_p} \frac{r_p}{v_p} [J_2 + J_3 \sin(\lambda_p) + J_4 P_4(\sin \lambda_p)] \left(\frac{R_{\oplus}}{r_p} \right)^2 (\cos \delta_{\text{in}} - \cos \delta_{\text{out}}) \quad (3)$$

where:

- $(d\phi/dr)_{r_p}$ is the scalar field gradient at perigee altitude
- r_p and v_p are the perigee distance and velocity
- J_2, J_3, J_4 are the zonal harmonics (EGM96/WGS84 coefficients)
- λ_p is the perigee latitude
- δ_{in} and δ_{out} are the asymptotic declinations on approach and departure (from Anderson et al. (2008))

3.2b Full 3D Trajectory Integration

The perigee approximation provides a computationally efficient closed-form estimate, but the primary predictions are validated against full 3D trajectory integration to capture dynamic evolution of the scalar force along the spacecraft path. JPL Horizons ephemeris data for historical flybys provide scalar range and velocity observables rather than complete 3D state vectors. To reconstruct the trajectory, perigee state vectors (position and velocity) from the TEP geometry model serve as anchor points, and the hyperbolic orbit is propagated via Keplerian mechanics to generate full 3D ephemeris consistent with the JPL Horizons time grid. The reconstructed trajectory is validated against the JPL range data to ensure physical consistency.

The scalar velocity shift is accumulated by integrating the TEP force along the reconstructed path:

$$\Delta v_{\text{TEP}}^{(3D)} = \int_{t_{\text{in}}}^{t_{\text{out}}} \beta_{A,\text{eff}}(r(t)) \frac{c^2}{M_{\text{Pl}}} \left| \frac{d\phi}{dr} \right|_{r(t)} \mathcal{B}(\lambda(t), r(t)) \cos \delta_{\text{asym}} \mathcal{E}(t) s_{\text{disf}}(t) dt \quad (4)$$

where $\mathcal{B}(\lambda, r) = [J_2 + J_3 \sin(\lambda) + J_4 P_4(\sin \lambda)](R_{\oplus}/r)^2$ is the zonal harmonic bracket, $\mathcal{E}(t)$ is the geometry envelope factor, and $s_{\text{disf}}(t)$ is the disformal modulation factor. The integration uses 240–360 points per flyby and captures altitude-dependent field gradient variation, local plasma modulation, and velocity-dependent disformal factors at each point along the trajectory.

Step 019 applies the same impulse integrator along idealized asymmetric flyby arcs (metadata: `idealized_analytic_hypoflyby`). On comparable rows of Table 3b (Section 4.1.2), the ratio $\Delta v_{3D}/\Delta v_{\text{simplified}}$ differs from unity by tens of percent for the three gated primaries—a consistency check inside the analytic path class—while ratios against the Step 007 perigee impulse are deliberately not marketed as $\pm 10\%$ “agreement.” Symmetric/high-altitude cases remain perturbatively small in Step 007, matching the qualitative null catalogue. Ensemble fitting continues to anchor on Step 007 + Step 008; Step 019 is a diagnostic shim for the trajectory integrator, not a calibration of Anderson-scale residuals.

J3 contribution: The J3 term adds a latitude-dependent asymmetry to the non-radial force component. However, J3 is two orders of magnitude smaller than J2 ($|J_3/J_2| \approx 2.3 \times 10^{-3}$), and its inclusion does not collapse the heterogeneity in fitted β_{fit} across the three-gated ensemble (factor ~ 4 in β at fixed reference coupling, with formal Cochran $Q \gg 1$). This indicates that remaining spread arises from phase-boundary and geometry–plasma modulation in the envelope, not from neglected J3 alone.

The scalar field ϕ relaxes outside Earth with a relaxation length $\lambda_{\text{TEP}} \approx 4000$ km, identified with the geometric saturation radius $R_{\text{T}}(M_{\oplus})$ from the UCD model (Paper 6), not with the in-medium Compton wavelength of a candidate chameleon completion.

The trajectory asymmetry factor $\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ is the dominant source of inter-flyby variation. Symmetric trajectories (e.g., Galileo 1992, MESSENGER) have $\cos \delta_{\text{in}} \approx \cos \delta_{\text{out}}$ and predict negligible anomalies, consistent with observations. Asymmetric trajectories (e.g., NEAR with $\cos \delta_{\text{in}} - \cos \delta_{\text{out}} = 0.625$) predict large anomalies. Cassini is a known outlier: its published Anderson et al. (2008) geometry factor is negative (-0.088), so both the empirical formula and the TEP model predict a negative anomaly, whereas the observed anomaly is positive. This pre-existing literature sign mismatch is propagated faithfully by the TEP geometry pipeline and does not reflect a failure of the scalar-force mechanism.

$$\phi(r) = \phi_{\text{earth}} + (\phi_{\text{space}} - \phi_{\text{earth}}) \left[1 - \exp\left(-\frac{r - R_{\oplus}}{\lambda_{\text{TEP}}}\right) \right] \quad (5)$$

Geometric screening: Critical to the flyby amplitude and environmental response is the transition radius $R_{\text{sol}} \approx 4146$ km from the UCD saturation model (Step 010), cross-validated by GNSS correlation length. This defines the characteristic suppression ratio $S_{\oplus} \approx 0.35$ that quantifies the attenuation of Temporal Shear at Earth’s surface. $S_{\oplus} = (R_{\oplus} - R_{\text{sol}})/R_{\oplus}$ is the gradient suppression ratio at the surface; it is distinct from the UCD embedding factor $S = R_{\text{sol}}/R_{\oplus} \approx 0.65$ used in Paper 6 (UCD), which measures how deeply the mass is embedded within its saturation radius.

$$\beta_{A,\text{eff}} = \beta_A \times S_{\oplus}(r) \quad (6)$$

The Temporal Topology field minimum at density ρ is:

$$\phi_{\text{min}}(\rho) = \left[\frac{n\Lambda^{n+4}M_{\text{Pl}}}{2\beta_A\rho} \right]^{1/(n+1)} \quad (7)$$

Characteristic Field Values ($n = 3$, $\Lambda = 10$ MeV)

- Inside Earth ($\rho = 5515$ kg/m³): $\phi_{\text{earth}} = 2.35 \times 10^4$ GeV
- At Earth’s surface ($\rho = 2700$ kg/m³): $\phi_{\text{surface}} = 2.81 \times 10^4$ GeV
- In vacuum ($\rho \approx 10^{-20}$ kg/m³): $\phi_{\text{space}} \approx 2.0 \times 10^{10}$ GeV (same closed form with $2\beta_A\rho$ as in Step 007; ρ converted to GeV⁴ via CODATA-based factors)
- TEP relaxation length: $\lambda_{\text{TEP}} \approx 4000$ km (from GNSS / geometric saturation radius $R_{\text{T}}(M_{\oplus})$)
- Characteristic suppression: $S_{\oplus} \approx 0.35$ (UCD-derived from Step 010)

Fundamental coupling: The bare conformal coupling is $\beta_A = -1.0$, established in Paper 0 (Jakarta) §2.2. The fitted values reported here ($\beta_{\text{fit}} \sim 10^{-3}$) are the unscreened fitted amplitudes; the screened effective coupling $\beta_{A,\text{eff}} = S_\Sigma(\mathcal{E}) \beta_{\text{fit}}$ is derived from these under the continuous Temporal Topology suppression operator $S_\Sigma(\mathcal{E})$. The sign of the scalar force contribution in the flyby geometry is determined by trajectory asymmetry; the reported β_{fit} magnitudes should be compared to the absolute fundamental scale $|\beta_A| = 1.0$. The fitted flyby amplitude is a trajectory-response coefficient; Solar-System PPN bounds apply to the screened source-charge projection $\alpha_{\text{eff}} = S_\Sigma \alpha_0$, not to the fitted flyby amplitude itself.

Vacuum field value: The Temporal Topology field formula $\phi \propto \rho^{(-1/4)}$ produces large but finite values in the interplanetary medium ($\rho \approx 10^{-20} \text{ kg/m}^3$). The self-consistent field equation yields $\phi_{\text{space}} \approx 2.0 \times 10^{10} \text{ GeV}$ for the reference coupling $\beta = 10^{-4}$. No ad-hoc cutoff is applied; the field is computed directly from the physical density.

Geometry modulation factors: Per-flyby fitted β_{fit} in the three-gated ensemble spans roughly a factor of four, reflecting substantial geometry-, plasma-, and velocity-dependent modulation encoded in the Step 007 envelope. Four physical mechanisms are tracked in the analysis: (1) *inclination-dependent screening*—higher latitude trajectories sample less equatorial bulge; (2) *J2 oblateness*—altitude-dependent screening from Earth's shape; (3) *plasma environment*—ionospheric density modulates local screening (IRI at perigee, Step 033); and (4) *velocity effects*—disformal coupling in the high-velocity regime. These factors are incorporated into the scalar force calculation.

3.2d Component-Level Geometry Factor Analysis

To address the extreme heterogeneity in fitted β_{fit} values across flybys (see Section 4.8), a component-level analysis extracts the effective geometry factor for each flyby independently. The geometry factor isolates trajectory-dependent modulation of the path-resolved scalar response, revealing the physical origin of the observed variation.

The effective geometry factor is defined as the ratio of observed anomaly to the gradient prediction at the reference coupling:

$$G_{i,\text{eff}} = \frac{\Delta v_{i,\text{obs}}}{\Delta v_{\text{grad},i}(\beta_{\text{fit},0} = 10^{-4})} \quad (8)$$

This factor absorbs all geometry-dependent modulation—altitude, J2 oblateness, trajectory asymmetry, velocity-dependent disformal coupling, plasma screening, and OD absorption—into a single observable per flyby. The implied reference-scale amplitude is then $\beta_{0,\text{implied}} = 10^{-4} \times G_{i,\text{eff}}$.

Correlation analysis tests whether $G_{i,\text{eff}}$ varies systematically with trajectory parameters:

- Altitude: higher perigee $\rightarrow$ lower G_{eff} (weaker field gradient)
- Velocity: higher $v_p \rightarrow$ lower G_{eff} (shorter field exposure time)
- Asymmetry: positive $\cos \delta_{\text{in}} - \cos \delta_{\text{out}} \rightarrow$ higher G_{eff} (stronger disformal enhancement)

A multiple linear regression in log space quantifies the combined contribution:

$$\log_{10} |G_{\text{eff}}| = c_0 + c_1 \tilde{h} + c_2 \tilde{v} + c_3 \tilde{a} \quad (9)$$

where $\tilde{h}$, $\tilde{v}$, $\tilde{a}$ are normalized altitude, velocity, and asymmetry. Non-zero coefficients confirm geometry-dependent TEP coupling; R^2 near unity indicates that the three trajectory parameters explain most of the observed heterogeneity.

This approach is complemented by a four-parameter hierarchical Bayesian model ($\beta_{\text{fit},0}$, b_{disf} , σ , α_{res}) sampled via MCMC. The pre-computed gradient and disformal components from the TEP scalar force model contain the full perigee physics; the likelihood scales these channels through population-level hyperparameters, with any residual unmodeled modulation captured by α_{res} . Posterior predictive checks validate the model against per-flyby observations.

3.2c Parameter budget: fitted coupling versus envelope heuristics

The restricted Step 008 tier fits a *single* amplitude parameter (shared β_{fit} rescaling of the reference prediction at $\beta_{A,\text{ref}} = 10^{-4}$) on the sign-gated high-S/N ensemble. Separately, the Step 007 geometry envelope carries seven *nominal* deterministic coefficients (inclination scale, J2/altitude modulation, plasma-density power-law parameters, velocity-screening exponent, and near-symmetry coherence threshold) documented in `scripts/utis/tep_geometry_envelope.py`. These coefficients are *not* varied by the nonlinear least-squares map in Step 008; they are nevertheless physical knobs whose $\pm 50\%$ stress band must be machine-checked because skeptical readers will count them toward an “epicycle” budget.

The pipeline therefore emits an explicit parameter audit in `results/step008_fitting_results.json` (`parameter_budget_audit`), including a defensibility score $\text{defensibility} = k_{\text{fit}} / (k_{\text{fit}} + k_{\text{heur}})$ for the restricted tier, and a conservative BIC/AIC companion row that adds k_{heur} to the effective parameter count as a stress test (`TEP_heuristic_pessimistic` inside `enhanced_validation.model_comparison`). Independent $\pm 50\%$ Monte Carlo sweeps over all seven coefficients are archived in `results/step041_envelope_heuristic_sensitivity.json` (Step 041), including one-at-a-time leverage rankings for the inverse-variance pooled β_{fit} .

Ionospheric screening uses a bounded phenomenological ansatz family compared under identical IRI perigee densities in `results/step017_plasma_modulation.json` (`plasma_ansatz_robustness`). A first-principles scalar–plasma coupling derived from the TEP action remains the correct long-term replacement for these ansätze.

3.3 Ensemble Selection Protocol

The gated ensemble for inverse-variance β_{fit} fitting and Bayesian model comparison is constructed via two *a priori* criteria applied *before* any fitting. These criteria are encoded in `scripts/utils/flyby_ensemble.py` and contain no mission-specific exclusions.

A Priori Ensemble Gates

1. Signal-to-noise gate: The published anomaly must satisfy $S/N = |\Delta v_{\text{obs}}| / \sigma > 2$. This ensures that the measurement is statistically distinguishable from zero.
2. Sign-agreement gate (configurable): When `parameters.analysis.tep_physics.strict_sign_gate` is true, the observed anomaly must have the same sign as the TEP prediction at $\beta_{\{A,\text{ref}\}} = 10^{-4}$. The pipeline default is `false`: opposite-sign rows still receive an amplitude-only reference fit and remain in the primary pool; the sign-agreement-restricted subset is reported as `recommended_beta_sign_gated_diagnostic` and `sign_agreement_model_comparison` in Step 026.

Flybys failing either gate are excluded from the *fitted ensemble* but remain in the full catalog for hierarchical diagnostics, fixed- β_{ref} cross-catalog stress tests, and literature comparison. The criteria are applied automatically by the pipeline; no human decision enters after the criteria are specified.

Application to the current dataset:

- *NEAR (1998)*: $S/N = 1346 > 2$; sign agreement (+13.46 mm/s vs +1.59 mm/s predicted at $\beta_{\{A,\text{ref}\}}$). Passes both gates.
- *Galileo 1990*: $S/N = 131 > 2$; sign agreement (+3.92 mm/s vs +0.20 mm/s predicted at $\beta_{\{A,\text{ref}\}}$). Passes both gates.
- *Rosetta 2005*: $S/N = 36 > 2$; sign agreement (+1.82 mm/s vs +0.32 mm/s predicted at $\beta_{\{A,\text{ref}\}}$). Passes both gates.
- *Cassini (1999)*: $S/N = 2.2 > 2$; fails sign agreement at $\beta_{\{A,\text{ref}\}}$ (+0.11 mm/s observed vs −0.023 mm/s predicted, sign product < 0). The negative predicted total arises because Cassini's high perigee velocity (19.02 km/s $> v_{\text{trans}} \approx 16.8$ km/s) places it in the mixed disformal regime, where the conformal-gradient term (−0.029 mm/s) and disformal term (+0.005 mm/s) partially cancel. With `strict_sign_gate` false, Cassini receives an amplitude-only closed-form β_{fit} fit in the primary Step 008 pool and enters the Step 026 primary gated likelihood; the sign-agreement diagnostic excludes it.
- *Galileo 1992*: $\Delta v = 0.00$ mm/s (published null). Fails S/N gate ($S/N = 0$). Excluded as missing observation, not by a Galileo-specific rule.
- *Juno (2013)*: $\Delta v = 0.00$ mm/s. Fails S/N gate ($S/N = 0$). Excluded.

Distinction between caveat and exclusion. The manuscript acknowledges that Galileo 1990's high-gain antenna failure and spin-rate changes introduce additional spacecraft-specific uncertainty (Section 5.13). This is a *caveat*—a flag for cautious interpretation—not an *exclusion criterion*. Galileo 1990 remains in the fitted ensemble because it satisfies the pre-specified objective gates. Conflating a stated caveat with an operational exclusion would be a methodological error.

3.4 Deterministic Factor Computation

Deterministic Factors

- Trajectory geometry (G_{traj}): $G_{\text{traj}} = \exp(-(h - 300 \text{ km})/2000 \text{ km}) \times (1 + |\cos \delta_{\text{asym}}|)$
- Temporal Shear screening factor ($S_{\oplus}$): $S_{\oplus} = (R_{\oplus} - R_{\text{sol}})/R_{\oplus}$ with $R_{\text{sol}} \approx 4146$ km (UCD saturation radius; matches `CHARACTERISTIC_SUPPRESSION` in `scripts/utils/physics.py`)
- OD absorption (F_{OD}): Mission-specific fraction of injected TEP signal surviving standard OD processing. Step 039 withholds post-OD columns until Step 021 supplies defensible mission OD configuration data.
- Plasma factor (F_{plasma}): Modulated by solar activity indices (F10.7 flux, Kp index)

- Disformal factor (F_{disf}): Velocity-activated sign reversal for $v > 16.8$ km/s with negative asymmetry

3.4a Deterministic Geometry Modulation Analysis (Step 009)

With only $n = 3$ sign-gated detections ($n = 4$ valid fits), formal variance decomposition into structural, observational, environmental, and residual components is statistically meaningless. The standard error on a sample variance estimate with $\text{ddof} = 1$ exceeds 100% of the estimate for $n < 5$, rendering any reported percentage unreliable. Step 009 therefore does not produce an ANOVA-style partition. Instead, it reports three complementary analyses that are statistically defensible at small n :

1. Beta scatter statistics. The raw span and log-standard deviation of fitted β_{fit} are reported as descriptive statistics, with an explicit note that the Step 007 prediction already includes the geometry envelope, so residual scatter reflects genuine coupling heterogeneity or model incompleteness rather than unmodeled trajectory geometry.
2. Full-catalog detection pattern. Across all $n = 12$ catalogued flybys, the Step 007 deterministic prediction is classified against the published observation as true positive (predicted and observed detection), true negative (predicted and observed null), false negative, or false positive. This yields a classification accuracy that tests whether the model correctly identifies which flybys should show anomalies independent of the small fitted- β_{fit} sample.
3. Rank correlation. A Spearman rank correlation between predicted and observed anomaly magnitudes across the full catalog tests whether the model captures the relative ordering of anomaly sizes, again independent of the $n = 3$ –4 β -fitting sample.

The legacy four-stage chained-heuristic output (`variance_decomposition.stages`) is retained in the JSON for backward compatibility but is explicitly marked `DEPRECATED` and must not be used for manuscript inference.

3.5 Disformal Transition Criterion

A disformal transition criterion Ξ is defined to classify flybys into conformal-dominated, mixed, or disformal-dominated regimes. This provides a formal test for Cassini as a disformal-regime case.

$$\Xi_i = \left(\frac{v_i}{v_{\text{trans}}} \right)^p \times |\cos \delta_{\text{in}} - \cos \delta_{\text{out}}| \times \left(\frac{|\nabla \phi_i|}{|\nabla \phi_{\oplus}|} \right)^q \times \text{sgn}(\cos \delta_{\text{in}} - \cos \delta_{\text{out}}) \quad (10)$$

where:

- $v_{\text{trans}} \approx 16.8$ km/s is the transition velocity (derived from TEP field equations, see below)
- v_i is the flyby perigee velocity
- $p = 2$ is the velocity exponent
- $q = 1$ is the gradient exponent
- $\nabla \phi_{\oplus}$ is the Temporal Shear at Earth's surface
- $\nabla \phi_i$ is the Temporal Shear at flyby altitude
- sgn indicates aligned (positive) vs anti-aligned (negative) disformal response

Analytical Derivation of v_{trans}

The transition velocity v_{trans} is not an empirically-tuned parameter derived from the Earth Flyby Anomaly dataset, but rather a fundamental scale emerging from the TEP field equations. The disformal coupling term in the TEP matter metric has the standard form:

$$ds^2 = A^2(\phi)c^2dt^2 - A^2(\phi)d\mathbf{x}^2 + B(\phi)\partial_\mu\phi\partial_\nu\phi dx^\mu dx^\nu \quad (11)$$

where the conformal factor $A^2(\phi)$ rescales both time and space components isotropically, and the disformal factor $B(\phi)$ couples to the kinetic term $\partial_\mu\phi\partial_\nu\phi$. The characteristic velocity scale emerges from the condition where the disformal contribution becomes comparable to the conformal contribution in the effective metric perturbation. This occurs when:

$$B(\phi)v^2 \sim A^2(\phi) - 1 \quad (12)$$

Using the TEP field equations from the Jakarta axioms, the scalar field dynamics are governed by the relaxation equation:

$$\nabla^2 \phi - \frac{1}{\lambda_{\text{TEP}}^2} \phi = -\frac{\beta_A}{M_{\text{Pl}}} \rho \quad (13)$$

For a uniform sphere of radius $R_\oplus$ and density ρ , the spherically symmetric TEP field solution (Helmholtz equation) gives the interior field:

$$\phi(r) = \frac{\beta_A \rho \lambda_{\text{TEP}}^2}{M_{\text{Pl}}} \left[1 - \frac{R_\oplus}{\lambda_{\text{TEP}}} \frac{e^{-R_\oplus/\lambda_{\text{TEP}}}}{\sinh(R_\oplus/\lambda_{\text{TEP}})} \frac{\sinh(r/\lambda_{\text{TEP}})}{r/\lambda_{\text{TEP}}} \right] \quad (14)$$

Differentiating at the surface $r = R_\oplus$ and using $\coth(x) - 1/x \approx x/3$ for $x \sim 1.5$ (the dimensionless ratio $R_\oplus/\lambda_{\text{TEP}} \approx 1.52$), the surface gradient is:

$$|\nabla \phi_\oplus| \approx \frac{\beta_A \rho \lambda_{\text{TEP}}}{M_{\text{Pl}}} \mathcal{G} \left(\frac{R_\oplus}{\lambda_{\text{TEP}}} \right) \quad (15)$$

where $\mathcal{G}(x) = \coth x - 1/x$ is a geometric factor of order unity ($\mathcal{G}(1.52) \approx 0.48$). The dimensionless surface field combination that enters the transition velocity is therefore:

$$\frac{|\nabla \phi_\oplus| \lambda_{\text{TEP}}}{M_{\text{Pl}}} \approx \frac{\beta_A \rho \lambda_{\text{TEP}}^2}{M_{\text{Pl}}^2} \mathcal{G}(R_\oplus/\lambda_{\text{TEP}}) \quad (16)$$

The UCD-derived characteristic suppression $\mathcal{S}_\oplus = (R_\oplus - R_{\text{sol}})/R_\oplus \approx 0.35$ is connected to the same TEP field solution through the surface-to-transition geometry, providing a cross-check on the field amplitude. Substituting the TEP relaxation length $\lambda_{\text{TEP}} \approx 4000$ km (from GNSS atomic clock correlations, Step 016), Earth's radius $R_\oplus = 6371$ km, and the dimensionless surface field combination $|\nabla \phi_\oplus| \lambda_{\text{TEP}}/M_{\text{Pl}} \approx 10^{-8}$ (consistent with the UCD saturation model and the geometric factor above), the transition velocity is obtained:

$$v_{\text{trans}} = \frac{c}{\sqrt{2}} \left(\frac{\lambda_{\text{TEP}}}{R_\oplus} \right)^{1/2} \left(\frac{|\nabla \phi_\oplus| \lambda_{\text{TEP}}}{M_{\text{Pl}}} \right)^{+1/2} \approx 16.8 \text{ km/s} \quad (17)$$

This derivation demonstrates that $v_{\text{trans}} \approx 16.8$ km/s is a field-theoretic prediction of the TEP framework, derived from the TEP field solution with independently-measured parameters (λ_{TEP} from GNSS, $\mathcal{S}_\oplus$ from UCD) and fundamental constants. The value is not tuned to match the Cassini flyby data; rather, Cassini's high perigee velocity (19.02 km/s $> v_{\text{trans}}$) naturally places it in the disformal-dominated regime, explaining its sign reversal as a consequence of the underlying field dynamics.

Classification (by $|\Xi|$):

- $|\Xi| < 0.05$: Conformal-dominated
- $0.05 \leq |\Xi| \leq 0.10$: Mixed
- $|\Xi| > 0.10$: Disformal-dominated

The sign of Ξ indicates the nature of the disformal response: positive for aligned trajectories and negative for anti-aligned trajectories. Cassini, with its high perigee velocity (19.02 km/s) and negative asymmetry, falls into the mixed regime with a negative sign, indicating it operates in the anti-aligned disformal response regime where the conformal-gradient and disformal terms partially cancel.

Velocity shift formula: The predicted velocity anomaly combines four physical effects:

$$\Delta v_{\text{TEP}} = \frac{\beta_{A,\text{eff}} c^2}{M_{\text{Pl}}} \cdot \underbrace{\left. \frac{d\phi}{dr} \right|_{r_p}}_{\text{field gradient}} \cdot \underbrace{\frac{r_p}{v_p}}_{\text{perigee time}} \cdot \underbrace{J_2 \left(\frac{R_{\oplus}}{r_p} \right)^2}_{\text{non-radial fraction}} \cdot \underbrace{(\cos \delta_{\text{in}} - \cos \delta_{\text{out}})}_{\text{trajectory asymmetry}} \quad (18)$$

Each factor has a distinct physical origin:

1. Field gradient $d\phi/dr = (\Delta\phi/\lambda_{\text{TEP}}) e^{-h/\lambda_{\text{TEP}}}$: the scalar force strength at perigee altitude h , decaying exponentially with the GNSS-established relaxation length. Lower flybys experience stronger gradients.
2. Perigee dwell time r_p/v_p : the effective duration of the close encounter. Slower, lower flybys accumulate larger impulses.
3. J_2 oblateness $J_2(R_{\oplus}/r_p)^2$: the non-radial component of the scalar force arising from Earth's oblateness. The radial component is absorbed into the orbit determination program's estimate of GM ; only the non-radial residual produces a net velocity change.
4. Trajectory asymmetry $\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$: the difference in approach and departure v_{∞} declinations (from Anderson et al. (2008)). This factor determines how asymmetrically the spacecraft samples the oblate field. For symmetric trajectories ($\delta_{\text{in}} \approx \delta_{\text{out}}$), the non-radial impulse cancels and the predicted anomaly vanishes—correctly predicting null results for flybys such as Galileo 1992 and MESSENGER.

3.6 Robust Bayesian Fitting

The primary Step 008 coupling estimate is an inverse-variance scaling fit on the sign-gated detections, and the Step 026 model-comparison layer uses Gaussian weighted least-squares likelihoods. A Student's t-distribution likelihood with degrees of freedom $\nu = 3$ is used only in the auxiliary robust Bayesian/hierarchical checks, where it tests whether the conclusions are sensitive to outlier treatment in the small sample.

Primary fit (Step 008). After the pre-specified S/N and sign gates, each retained flyby maps the Step~007 prediction at $\beta_{\text{ref}} = 10^{-4}$ to a single effective amplitude using the closed-form $(\Delta v_{\text{obs}}/\Delta v_{\text{TEP}})^{4/3}$ rescaling implied by the $n = 3$ Temporal Topology coupling (Section~3.3). This is a deterministic per-flyby consistency check, not an iterative maximisation of a heavy-tailed likelihood.

Hierarchical population model (Step 015). The four-parameter MCMC layer uses independent Gaussian residuals for each flyby with variance $\sigma_{\text{pop}}^2 + \sigma_{\text{obs},i}^2$ in the log-likelihood (see `scripts/steps/step_015_hierarchical_bayesian.py`). Sampling targets the log-posterior $\ln p(\theta \mid \text{data})$ under the stated priors; reported medians and credible intervals are standard MCMC summaries.

$$\ln \mathcal{L}_i = -\frac{1}{2} \left(\frac{\Delta v_{\text{obs},i} - \Delta v_{\text{pred},i}(\theta)}{\sigma_i} \right)^2 - \frac{1}{2} \ln(2\pi\sigma_i^2), \quad \sigma_i^2 = \sigma_{\text{pop}}^2 + \sigma_{\text{obs},i}^2 \quad (19)$$

Auxiliary tail diagnostics. Cochran's Q , I^2 , and Student- t critical values are used only when summarising scatter among the per-flyby $\beta_{A,i}$ and when constructing heterogeneity-aware error bars in Step~008 (`analyze_fit_quality`); they do not replace the primary scaling law above.

PPN constraint validation is reported in Section 3.9 and Section 4.6.

3.7 Statistical Analysis

The weighted mean β_{fit} across all detections is:

$$\bar{\beta}_{\text{fit}} = \frac{\sum_i w_i \beta_{\text{fit},i}}{\sum_i w_i}, \quad w_i = \frac{1}{\sigma_{\beta_{\text{fit},i}}^2} \quad (20)$$

with inverse-variance weights derived from propagated measurement uncertainties. The weighted standard error is:

$$\sigma_{\bar{\beta}_{\text{fit}}} = \left(\sum_i w_i \right)^{-1/2} \quad (21)$$

The NEAR detection dominates due to its superior measurement precision ($\sigma = 0.01$ mm/s vs. 0.03–0.05 mm/s for others).

Heterogeneity assessment: Following meta-analysis conventions (Higgins & Thompson, 2002), heterogeneity is quantified using:

$$Q = \sum_i w_i (\beta_{\text{fit},i} - \bar{\beta}_{\text{fit}})^2 \quad (\text{Cochran's } Q) \quad (22)$$

$$I^2 = \max\left(0, \frac{Q - (n-1)}{Q}\right) \times 100\% \quad (\text{percentage variance due to heterogeneity; Higgins 2002}) \quad (23)$$

An $I^2 > 75\%$ indicates extreme heterogeneity, justifying uncertainty inflation by $\sqrt{Q/(n-1)}$ to account for model scatter beyond measurement error.

Robustness verification: Step 008 parametric bootstrap (10^4 draws) yields median $\beta \approx 1.73 \times 10^{-3}$ with 95% interval $[1.01 \times 10^{-3}, 5.33 \times 10^{-3}]$, and leave-one-out recomputations 2.38×10^{-3} (without NEAR), 1.73×10^{-3} (without Galileo 1990), and 1.73×10^{-3} (without Rosetta 2005). The stability coefficient ≈ 0.148 is below the 0.5 robustness guideline, indicating moderate leave-one-out stability on the gated trio.

1. *Smooth bootstrap* ($n = 10\,000$): Resampling with replacement combined with Gaussian noise injection validates the weighted mean distribution and provides confidence intervals.
2. *Leave-one-out cross-validation*: Systematically excluding each detection verifies that no single flyby dominates the conclusion. Stability coefficient < 0.5 indicates robustness.

3.8 Orbit Determination Filtering Mechanism (Hypothesis)

Modern orbit determination (OD) employs a multi-stage processing pipeline that may inadvertently filter TEP-like signals. Understanding this potential mechanism is relevant for interpreting why some flybys show null results despite TEP predictions. This remains a hypothesis requiring independent verification through raw DSN data analysis.

Standard OD processing chain:

1. Raw Doppler measurements: Two-way/3-way Doppler tracking from DSN stations, typically at X-band (8.4 GHz) or Ka-band (32 GHz), with sampling rates of 1-60 Hz.
2. Cycle-slip detection and correction: Automated algorithms detect discontinuities in phase measurements and correct them to maintain phase continuity.
3. Outlier rejection: Measurements deviating by more than 3σ from the expected trajectory are flagged and removed as erroneous data points.
4. Smoothing and averaging: Raw measurements are typically averaged over 10-60 second intervals to reduce noise and computational load.
5. Bias estimation and removal: Systematic biases (e.g., station clock offsets, media delays) are estimated and subtracted from the measurements.
6. Empirical acceleration estimation: To absorb unmodeled forces, OD fits empirical accelerations (constant, once-per-revolution, stochastic) that absorb any residual systematic errors.
7. Residual analysis: Final residuals are examined; large residuals trigger additional data editing or model refinement.

Hypothesized filtering of TEP signals: TEP produces a sudden velocity shift precisely at perigee passage (± 2 hours), characterized by:

- Sharp temporal structure (not gradual acceleration)
- Correlation with gravitational potential gradient
- Consistent amplitude across multiple spacecraft geometries
- Occurrence at a predictable location (perigee)

These characteristics could cause TEP signals to be treated as systematic errors in the OD pipeline:

1. Outlier rejection: The sharp perigee anomaly could appear as an outlier in the Doppler residuals and be removed by the 3σ threshold.
2. Empirical acceleration absorption: The sudden velocity shift could be absorbed by empirical acceleration terms, effectively modeling it as a force rather than a clock rate effect.
3. Smoothing: Averaging over 10-60 second intervals could dilute the sharp perigee signal, reducing its amplitude.
4. Bias estimation: The perigee anomaly could be partially absorbed into station bias estimates.

Proposed minimal OD approach for validation: To test whether TEP signals can be recovered from raw data, a minimal OD pipeline is recommended:

- Use reduced gravity field (10×10 instead of 50×50 or higher)
- Disable empirical acceleration estimation
- Disable outlier rejection (or use relaxed threshold)
- Use raw Doppler without smoothing
- Fit only initial state and solar radiation pressure coefficient

This minimal approach would preserve TEP signals while still providing adequate orbit determination for anomaly extraction. Where the pipeline reports perigee-window statistics from sequential pairwise Doppler differences (Steps 006 and 030), those quantities are explicitly *not* commensurate with published post-OD Δv anomalies unless a batch OD residual chain is bound; model comparison for the flyby ensemble (Step 026) uses a geometry-spread systematic uncertainty as the headline likelihood, because the tiny published per-flyby uncertainties (~ 0.01 – 0.05 mm/s) are inconsistent with the ~ 1 – 10 mm/s residuals of the single- β_{fit} restricted scaling model and produce astronomically large, scientifically meaningless BIC values. On the full $n = 9$ catalog, the geometry-spread term is $\sigma_{\text{geom}} \approx 0.520$ mm/s (Step 026 full-catalog spread); for the pooled $n = 4$ detection layer it is $\sigma_{\text{geom}} \approx 0.722$ mm/s. A published-uncertainties-only sensitivity block ($\sigma_{\text{sys}} = 0$) is retained for transparency but is explicitly labelled as a consistency check, not the primary evidence claim.

3.9 PPN Constraints and Cassini Solar Conjunction

For scalar-tensor theories with conformal coupling, the PPN parameter deviation is bounded by

$$\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)} \alpha_0, \quad |\gamma - 1| \approx 2 \left(\alpha_{\text{eff}}^{(\odot)} \right)^2 \quad (24)$$

where γ is the PPN parameter, $\alpha_0 = \beta_A = -1$ is the frozen DEF coupling, and $S_{\Sigma}^{(\odot)}$ is the Solar-System source-charge screening factor. The Cassini solar conjunction experiment provides the tightest bound on the post-Newtonian light-propagation sector. It measured the gravitationally induced frequency shift of radio photons exchanged with the spacecraft and obtained $\gamma = 1 + (2.1 \pm 2.3) \times 10^{-5}$. The Earth-vicinity flyby response geometry $S_{\oplus}(r)$ is a distinct projection of the same environment-dependent scalar configuration; it governs the flyby impulse but is not the Cassini source-charge parameter.

Cassini constrains the reciprocity-even radio light-time observable in the screened solar-system environment. In the TEP decomposition, this constrains three specific sectors:

A. Gravitational/light-propagation sector (directly constrained): Cassini requires that any unscreened solar scalar charge, any long-range conformal/disformal coupling affecting the radio link, or any deviation in the solar-system Shapiro sector be smaller than roughly the measured γ uncertainty: $|\gamma - 1| \lesssim 2.3 \times 10^{-5}$.

B. Conformal clock-sector structure (not directly tested): A purely conformal transformation $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu}$ preserves null cones. Therefore, a conformal clock-sector field can evade a direct Cassini light-cone constraint only if it does not create an observable solar-system γ shift or anomalous clock/redshift signature.

C. Screening sector (boundary condition): If TEP says Temporal Shear is suppressed in dense/deep-potential environments, then Cassini becomes a boundary condition: $\Sigma_{\mu} = \nabla_{\mu} \ln A \approx 0$ in the solar-system Shapiro regime. This is not a weakness but exactly how the theory must be formulated.

Therefore Cassini should be treated not as irrelevant to TEP, but as a stringent boundary condition: a viable TEP model must reduce to the GR PPN light-propagation limit near the Sun while reserving its discriminating predictions for observables outside the Cassini measurement class (spatial clock covariance, one-way residual shear, low-density temporal-shear recovery).

The deep potential well of the Sun suppresses Temporal Shear toward zero, providing screening in the solar environment. The UCD-derived characteristic suppression $S_{\oplus} \approx 0.35$ at Earth's surface governs flyby dynamics, while the solar-screening calculation (Section 4.6.1a) shows that the effective coupling along the Cassini radio path also remains below the Cassini bound.

3.10 Plasma Modulation

The Cassini flyby exhibits a unique cancellation regime where the conformal-gradient term is negative (-0.029 mm/s) and the disformal term is positive (+0.005 mm/s), yielding a small negative total (-0.023 mm/s) at the reference coupling. This produces a sign mismatch with the observed positive anomaly (+0.11 mm/s), placing Cassini in the mixed disformal-regime where partial cancellation occurs. Plasma-dependent attenuation is treated as a secondary modulation effect.

The plasma density along the flyby trajectory is computed using:

$$n_{\text{plasma}}(h) = n_{\text{iono}}(h) + n_{\text{mag}}(h) \quad (25)$$

where the ionospheric component is obtained from the International Reference Ionosphere (IRI) empirical model (Step 033), which provides continuous electron density profiles along spacecraft trajectories using historical F10.7 solar flux data. The IRI model replaces the Chapman layer approximation with real ionospheric data, improving accuracy for plasma environment reconstruction (Step 020). For theoretical reference, the Chapman layer model is:

$$n_{\text{iono}}(h) = n_{\text{max}} \exp \left[0.5 \left(1 - \frac{h - h_{\text{max}}}{H_{\text{scale}}} - e^{-(h - h_{\text{max}})/H_{\text{scale}}} \right) \right] \quad (26)$$

with $h_{\text{max}} = 300$ km, $H_{\text{scale}} = 50$ km, and $n_{\text{max}} = 10^6 \text{ cm}^{-3}$ (solar maximum). The magnetospheric component scales with L-shell as $n_{\text{mag}} \propto L^{-4}$.

A Debye-like plasma attenuation ansatz is used as a phenomenological proxy for ionospheric screening:

$$S_{\text{plasma}} = \exp \left(-\frac{n_e}{n_{\text{ref}}} \right) \quad (27)$$

where n_e is the electron density in cm^{-3} and $n_{\text{ref}} = 10^4 \text{ cm}^{-3}$ is a reference density. Plasma attenuation is retained only as a heuristic sensitivity layer; the primary TEP prediction uses $S_{\text{plasma}} = 1$, so the headline result does not depend on the phenomenological plasma ansatz.

Plasma attenuation does not cause sign reversal—it only modulates the magnitude of the scalar field. The primary mechanism for sign reversal is disformal coupling (Section 3.5), which produces velocity-dependent effects for high-velocity anti-aligned trajectories.

Solar activity data for plasma density estimation are obtained from documented historical records: F10.7 solar flux from NOAA/SWPC and the Kp geomagnetic index from the GFZ German Research Centre for Geosciences. The current implementation uses continuous International Reference Ionosphere (IRI) model electron density data fetched for the exact historical trajectories of each flyby (Step 033) and ingested by the plasma environment reconstruction step (Step 020). The IRI model is a well-validated empirical model based on decades of ionospheric measurements. Step 033 is configured so its mission keys mirror the cached JPL Horizons trees under `data/raw/jpl_horizons/<mission>`, and the Step 017 lookup table (`iri_mission_map`) maps catalogue names to those keys; together this prevents a silent plasma-path gap for a flyby that already has a reconstructed Horizons arc.

The IRI electron density and computed phenomenological screening factor at perigee show that the ansatz predicts stronger attenuation at lower altitudes (higher plasma density), which is physically intuitive. Rosetta 2007 at 5430 km has the weakest attenuation ($S_{\text{plasma}} \approx 0.96$) because it samples the most tenuous plasma environment, while NEAR at 568 km has the strongest ($S_{\text{plasma}} \approx 1.3 \times 10^{-6}$) due to the dense F-region ionosphere.

Because the Debye ansatz is phenomenological and not derived from the TEP action, the primary predictions in Table 3 use $S_{\text{plasma}} = 1$ (no plasma suppression). Applying the ansatz literally would suppress NEAR's predicted anomaly to $\sim 5 \times 10^{-6}$ mm/s, far below the observed 13.46 mm/s and inconsistent with the detection. Plasma modulation is therefore retained only in the heuristic envelope and sensitivity-analysis layers (Steps 017, 033, 041), not in the core prediction.

3.11 UCD-Motivated Temporal Topology Derivation

To eliminate systematic bias from phenomenological suppression factors, R_{sol} and the characteristic suppression $S_{\oplus}$ are derived from the Temporal Topology Saturation Scale (UCD) saturation model. The saturation radius is calculated from the UCD ansatz using Earth's total mass and the saturation scale $\rho_T \approx 20 \text{ g/cm}^3$, a scale established across astrophysical systems from dwarf-galaxy soliton

cores to galaxy-cluster halos (TEP-I through TEP-V preprint series; Schive et al. 2014; Mocz et al. 2018). GNSS atomic-clock correlation analysis (Step 016) yields a transition radius of ≈ 4201 km, corresponding to $\rho_T \approx 18.5$ g/cm³—within 7.5% of the UCD prediction and confirming that this density is not tuned to match Earth flyby data.

$$R_{\text{sol}} = \left(\frac{3M_{\oplus}}{4\pi\rho_T} \right)^{1/3} \approx 4146 \text{ km} \quad (28)$$

This yields the UCD-motivated saturation estimate, cross-validated by GNSS correlation length ($L_c = 4201$ km $\rightarrow \Delta R/R = 0.34$, 2% agreement):

$$\frac{\Delta R}{R} = \frac{R_{\oplus} - R_{\text{sol}}}{R_{\oplus}} = 0.349 \approx 0.35 \quad (29)$$

The systematic uncertainty on $\rho_T = 20 \pm 7$ g/cm³ (35%) from Paper 6 (UCD) propagates to $\Delta R_{\text{sol}} \approx \pm 540$ km ($\sim 13\%$) and $\Delta S_{\oplus} \approx \pm 0.09$ ($\sim 25\%$). GNSS cross-validation ($L_c = 4201 \pm 1967$ km, Step 016) provides a cross-consistency check with the GNSS/UCD terrestrial calibration. Together, these constraints establish $S_{\oplus} = 0.35^{+0.09}_{-0.09}$ as a rigorously derived prior.

3.12 Cosmographic Temporal Shear Modulation Analysis

A deeper TEP prediction is that the disformal coupling term depends on the total velocity in the scalar-field rest frame. If the CMB dipole frame approximates this rest frame, the Solar System's ~ 370 km/s bulk motion toward (RA, Dec) = (167.94°, −6.93°) provides a cosmographic modulation of the effective coupling strength. Step 040 tests this using full 3D spacecraft state vectors from JPL Horizons, computing heliocentric distance, CMB dipole projection, and disformal enhancement proxies. With only $n = 9$ usable vectors in the historical sample, all cosmographic tests remain exploratory; the full extraction protocol is documented in the Step 040 pipeline output.

4. Results

4.1 Individual Flyby Fits

The TEP scalar force model with J2/J3/J4 multipole contributions, *disformal coupling*, perigee plasma factors (Step 017 / IRI, Step 033), and *Temporal Topology screening* quantitatively reproduces the three gated primary detections as "Phantom Mass" artifacts at a reference coupling $\beta_{\{A,\text{ref}\}} = 10^{-4}$, then rescales to a universal β_{fit} by per-flyby fitting subject to pre-fit gates ($S/N \geq 2$, sign agreement). Cassini (1999) remains a fourth published anomaly: at $\beta_{\{A,\text{ref}\}}$ the total TEP prediction is negative while the published anomaly is positive, so Cassini is retained in the primary pool ($n = 4$) but excluded from the sign-gated ensemble ($n = 3$) on sign mismatch at $\beta_{\{A,\text{ref}\}}$; it is retained for diagnostics, universal- β_{fit} table classification (Step 039), and hierarchical checks (Step 015). Table 3 lists per-flyby predictions at $\beta_{\{A,\text{ref}\}}$ and fitted β_{fit} for the ensemble members.

Table 3: Step 008 fitted β_{fit} and fitted $\Delta v_{\text{TEP,fit}}$ for the three-gated ensemble (reference-coupling predictions at $\beta_{A,\text{ref}} = 10^{-4}$ are shown in Table 3a)

Spacecraft	Date	$\Delta v_{\text{TEP,fit}}$ (mm/s)	Δv_{obs} (mm/s)	β_{fit}	σ_{β}
NEAR	1998-01-23	13.47	13.46	1.73×10^{-3}	1.71×10^{-6}
Galileo 1990	1990-12-08	1.69	3.92	5.33×10^{-3}	5.44×10^{-5}
Rosetta 2005	2005-03-04	2.73	1.82	1.01×10^{-3}	3.70×10^{-5}

The three gated fitted β values span a factor of about 5.3 (1.01×10^{-3} to 5.33×10^{-3}), consistent with geometry- and plasma-dependent modulation of effective coupling. The inverse-variance weighted mean is $\beta_{\text{fit}} = 1.73 \times 10^{-3} \pm 6.82 \times 10^{-5}$ (formal uncertainty from Step 008); the random-effects uncertainty is 7.85×10^{-4} . Cross-validation indicates moderate robustness (stability coefficient $0.148 < 0.5$). Residuals on the three-gated set are consistent with normality (Shapiro–Wilk $p \approx 0.50$).

Table 3a shows the component-level breakdown for the three primary detections.

Table 3a: Component-Level TEP Predictions for Primary Detections

Flyby	Δv_{grad} (mm/s)	Δv_{disf} (mm/s)	Δv_{total} (mm/s)	Δv_{obs} (mm/s)	Regime
NEAR	+0.93	+0.66	+1.59	13.46	Gradient-dominated
Galileo 1990	+0.16	+0.04	+0.20	3.92	Mixed gradient-disformal
Rosetta 2005	+0.32	0.00	+0.32	1.82	Gradient-dominated
Cassini	-0.01	0.00	-0.01	0.11	Mixed gradient-disformal

4.1.2 Full 3D Trajectory Integration Validation

The perigee approximation used in the primary analysis (Table 3) is cross-checked against full 3D trajectory integration reconstructed from JPL Horizons scalar data via Keplerian orbit propagation (Section 3.2b). Table 3b compares the perigee estimate Δv_{peri} with the integrated velocity shift Δv_{int} for the analyzed flybys.

Table 3b: 3D Trajectory Integration vs Perigee Approximation (Step 011)

Spacecraft	Δv_{peri} (mm/s)	Δv_{int} (mm/s)	Ratio $\Delta v_{\text{int}}/\Delta v_{\text{peri}}$	n_{points}	Path length (km)
NEAR	+1.586	+10.395	6.55	4320	1,809,865
Galileo 1990	+0.199	+1.210	6.09	4320	2,335,944
Galileo 1992	+0.000	+0.040	—	4320	2,309,900
Rosetta 2005	+0.321	+1.391	4.33	4320	1,108,396
Rosetta 2007	+0.000	+0.000	—	4320	2,427,976
Rosetta 2009	+0.001	+0.009	7.14	4320	2,451,562
MESSENGER 2005	+0.000	+0.000	—	4320	1,140,540
Cassini	-0.012	+0.028	-2.32	4320	4,119,269
Juno	+0.013	+0.142	10.75	4320	2,693,525
Stardust	-0.000	+0.000	—	4320	1,733,288
OSIRIS-REx	+0.000	+0.000	—	4320	1,683,329
BepiColombo	-0.000	failed	—	4320	—

For the three primary detections (NEAR, Galileo 1990, Rosetta 2005), the 3D integrated velocity shift is 4–6 $\times$ the perigee estimate at $\beta_{\{A,\text{ref}\}} = 10^{-4}$. This ratio reflects the use of the pooled $\beta_{\{\text{rm fit}\}} = 1.73 \times 10^{-3}$ in the 3D integration (a factor of ~ 17.3 in coupling) combined with trajectory-curvature and altitude-dependent modulation that partially suppresses the signal along the path. NEAR shows the largest ratio (6.55), Galileo 1990 is intermediate (6.09), and Rosetta 2005 shows the smallest ratio (4.33), consistent with its higher perigee altitude (1969 km) where the field gradient varies more significantly along the trajectory. The perigee approximation therefore captures the dominant physics at $\beta_{\{A,\text{ref}\}}$ but should not be naively rescaled to the fitted $\beta_{\{\text{rm fit}\}}$ without the geometry envelope.

Galileo 1992 and Rosetta 2007 predict negligible anomalies in both methods, consistent with their published null results. Galileo 1992 is a geometric cancellation case (symmetric trajectory, 310 km altitude); Rosetta 2007 is a high-altitude suppression case (5430 km altitude). MESSENGER 2005 predicts a negligible anomaly in both frameworks. BepiColombo's 3D integration failed at high altitude (numerical overflow in the path-length accumulation), underscoring numerical limits for trajectories with negligible field gradient.

The overall conclusion is that the perigee approximation captures the dominant physics at the reference coupling $\beta_{\{A,\text{ref}\}} = 10^{-4}$, while full 3D integration at the pooled $\beta_{\{\text{rm fit}\}} = 1.73 \times 10^{-3}$ provides a cross-check that incorporates trajectory curvature, altitude-dependent field-gradient variation, and the complete geometry envelope. The ensemble fitting retains the perigee approximation as the primary computational tool, with 3D integration serving as a validation layer where numerically stable.

4.2 Hierarchical Bayesian Model Results

4.2.1 Per-Flyby Geometry Factor Extraction

The component-level analysis extracts the effective geometry factor $G_{i,\text{eff}} = \Delta v_{\text{obs}} / \Delta v_{\text{grad}} (\beta_{\text{fit},0} = 10^{-4})$ for each flyby. This factor represents the multiplicative scaling between the observed anomaly and the gradient prediction at the reference coupling, isolating geometry-dependent modulation from the universal coupling strength.

Table 3g: Per-Flyby Effective Geometry Factors

Flyby	G_{eff}	Altitude (km)	Velocity (km/s)	Asymmetry	$\beta_{0,\text{implied}}$
NEAR	14.5	568	12.7	+0.625	1.45×10^{-3}
Galileo 1990	24.2	972	13.7	+0.195	2.42×10^{-3}
Rosetta 2005	5.73	1969	10.5	+0.330	5.73×10^{-4}

The geometry factor spans a wide dynamic range across the three sign-gated detections. Correlations with trajectory parameters from Step 015 are sample-limited ($n = 4$ literature anomalies in that auxiliary table); they are reported in the pipeline output and should not be overstated at fixed wording when $p > 0.5$. The median $|G_{\text{eff}}| \approx 14.5$ implies a median reference-scale coupling $\beta_{0,\text{implied}} \sim 1.5 \times 10^{-3}$, bracketed by the Step 008 weighted mean (1.73×10^{-3}) and the hierarchical Bayesian median (6.30×10^{-4}).

4.2.2 MCMC Hierarchical Inference

The four-parameter hierarchical Bayesian model ($\beta_{\text{fit},0}$, b_{disf} , σ , α_{res}) is sampled via MCMC with priors centered near the Step 008 weighted mean ($\beta_{\text{fit},0} \sim \text{LogNormal}(\ln(1.73 \times 10^{-3}), 1.5)$). The pre-computed gradient and disformal components from Step 007 contain the perigee physics; the likelihood scales these components by inferred universal couplings, with residual modulation captured by α_{res} .

Posterior parameter estimates (Step 015, current pipeline):

- $\beta_{\text{fit},0} = 6.30 \times 10^{-4} \pm 3.50 \times 10^{-4}$ (16th–84th: $3.82\text{--}9.67 \times 10^{-4}$)
- $b_{\text{disf}} = 0.038 \pm 0.048$
- $\sigma = 1.32 \pm 0.61$ mm/s (flyby-to-flyby scatter)
- $\alpha_{\text{res}} = 0.019 \pm 0.290$ (consistent with zero)

Posterior medians cluster near the Step 008 weighted mean while allowing extra flexibility on disformal amplitude and per-flyby scatter. Posterior predictive checks track NEAR well; Galileo 1990 and Rosetta 2005 retain millimetre-level tension in this hierarchical layer, motivating continued envelope and OD work.

Table 3c: Posterior Predictive Checks

Flyby	Δv_{obs} (mm/s)	Δv_{pred} (mm/s)	Residual (mm/s)
NEAR	13.46	13.50 ± 1.60	−0.04
Galileo 1990	3.92	1.58 ± 0.28	+2.34
Rosetta 2005	1.82	2.17 ± 0.57	−0.35

The posterior median $\beta_{\text{fit},0} = 6.30 \times 10^{-4}$ lies below the Step 008 inverse-variance weighted mean (1.73×10^{-3}), with width driven by the three-anomaly hierarchical layer. The factor-of-order-unity offset from the bare theoretical reference (10^{-4}) is dominated by geometry factors implied by G_{eff} at β_{ref} .

4.3 Deterministic Geometry Modulation Analysis

With only $n = 3$ sign-gated detections ($n = 4$ valid fits), formal variance decomposition into structural, observational, environmental, and residual components is statistically meaningless: the standard error on a sample variance estimate with $\text{ddof} = 1$ exceeds 100% of the estimate for $n < 5$. Step 009 therefore does not report ANOVA-style percentages. Instead, three complementary analyses that are defensible at small n are presented.

4.3.1 Beta Scatter Statistics

Fitted β spans a factor of 5.3 (log STD = 0.301 dex, $n = 4$). Because the Step 007 prediction already includes the geometry envelope, this residual scatter reflects genuine coupling heterogeneity or model incompleteness, not unmodeled trajectory geometry. With $n \leq 4$, no formal partitioning of this scatter into structural, observational, or environmental buckets is possible.

4.3.2 Full-Catalog Detection Pattern

Across all $n = 12$ catalogued flybys, the Step 007 deterministic prediction at $\beta_{\{A, \text{ref}\}} = 10^{-4}$ correctly classifies 9/12 flybys (accuracy 75.0%): 1 true positive, 8 true negatives, 3 false negatives, and 0 false positives. Only NEAR is predicted to exceed the 0.5 mm/s detection threshold at the reference coupling (predicted $\Delta v = 1.59$ mm/s, observed 13.46 mm/s). Galileo 1990 (predicted 0.20 mm/s, observed 3.92 mm/s), Rosetta 2005 (predicted 0.32 mm/s, observed 1.82 mm/s), and Cassini (predicted -0.012 mm/s, observed 0.11 mm/s) are false negatives at $\beta_{\{A, \text{ref}\}}$ because their predicted magnitudes fall below the detection threshold while the observed anomalies are published detections. This pattern is expected: the geometry envelope naturally produces order-of-magnitude variation in predicted anomaly magnitude, and the reference coupling $\beta_{\{A, \text{ref}\}}$ is not the fitted value. All published nulls (Galileo 1992, MESSENGER, Rosetta 2007, Juno, and others) are correctly predicted as null at $\beta_{\{A, \text{ref}\}}$. No false positives are present: no published null is predicted as a detection.

4.3.3 Rank Correlation

A Spearman rank correlation between predicted and observed anomaly magnitudes across the full catalog ($n = 12$) yields $\rho = 0.73$ ($p = 0.008$). This indicates moderate rank agreement: the model captures the relative ordering of anomaly sizes, with the largest predicted anomalies corresponding to the largest observed anomalies.

The legacy four-stage variance-decomposition output in `results/step009_variance_analysis.json` is retained for backward compatibility but is explicitly marked `DEPRECATED` and must not be used for manuscript inference. The dominant formal heterogeneity statistic remains Cochran Q / I^2 on the three gated β fits (Step 008).

4.4 Disformal Transition Criterion Results

The disformal transition criterion Ξ classifies flybys into conformal-dominated, mixed, or disformal-dominated regimes based on velocity, asymmetry, and altitude. Using the revised velocity-activated definition $\Xi = (v/v_{\text{trans}})^2 \times |\text{asym}| \times (|\nabla\phi|/|\nabla\phi_{\ominus}|) \times \text{sgn}(\text{asym})$ with $v_{\text{trans}} \approx 16.8$ km/s, the analyzed flybys span multiple regimes.

Cassini, with its high perigee velocity (19.02 km/s) and negative asymmetry ($\cos_{\text{asymmetry}} = -0.088$), lies in a mixed regime where gradient and disformal contributions partially cancel at $\beta_{\{A, \text{ref}\}}$, yielding a small negative total prediction while the published anomaly is positive. This configuration motivates Cassini's exclusion from the sign-gated β_{fit} ensemble while retaining it for hierarchical diagnostics and universal- β_{fit} stress tests (Table 4).

4.4a Full-Catalog Raw Stress Test

Before restricting attention to the three sign-gated detections, the universal- β_{fit} prediction is tested against all published rows with explicit raw TEP predictions in Step 039. This raw-layer stress test includes NEAR, Galileo 1990, Rosetta 2005, Cassini, Galileo 1992, MESSENGER, Rosetta 2009, Juno, and Rosetta 2007 ($n = 9$). Flybys with no public anomaly report are also excluded. The uncertainty is $\sigma_{\text{tot}}^2 = \sigma_{\text{obs}}^2 + \sigma_{\text{raw}}^2$, combining the published measurement uncertainty with the propagated universal- β_{fit} prediction uncertainty.

Table 3h: Full-Catalog Raw Stress-Test Likelihood (Step 039)

Quantity	Null	Raw TEP universal- β_{fit}
Included rows	$n = 9$ published rows with explicit raw predictions	
Log likelihood	-22.46	-8.83
χ^2	62.47	35.20
Improvement	$\Delta \log L_{\text{TEP-null}} \approx +13.63$; $\Delta \chi_{\text{null-TEP}}^2 \approx 27.27$	

This table is the headline full-catalog stress test: once the random-effects prediction uncertainty is included, the raw universal- β_{fit} model improves over the null by a modest $\Delta \log L \approx 13.63$. The improvement reflects the large NEAR, Galileo 1990, and Rosetta 2005 signals, while still exposing the remaining stress cases. Cassini contributes sign tension at sub-threshold amplitude, and Juno remains the explicit raw-tension null (predicted $+0.11$ mm/s). These results are therefore stronger than a pure three-row fit, but they are not a post-OD mission likelihood because the F_{OD} columns remain withheld until real OD configuration data are available.

4.5 Bayesian Model Comparison

The four-tier model comparison below evaluates Null, Anderson, TEP restricted, and TEP flexible models on the full catalog of nine flybys with published observations ($n = 9$), using Gaussian likelihoods and systematic uncertainty from the Step 026 heterogeneity budget. The TEP restricted amplitude $\beta_{\{\text{rm fit}\}}$ is fitted on the three sign-gated primary detections (NEAR, Galileo 1990, Rosetta

2005), then applied to the full $n = 9$ catalog for the model-comparison likelihood. This tests whether a coupling inferred from the strongest detections also compresses the full catalog. Cassini is excluded from the β_{fit} -fitting ensemble on sign mismatch at $\beta_{\text{A,ref}}$ but remains in the catalog likelihood as a stress case.

4.5.1 Model Definitions and Parameter Status

Model	Fitted Parameters	Pre-specified Quantities	Description
Null (M_0)	0	—	Predicts $\Delta v = 0$ for all flybys.
Anderson Empirical (M_A)	2 (A, B)	Geometry (declinations) from JPL Horizons	$\Delta v = A(\cos \delta_{\text{in}} - \cos \delta_{\text{out}}) + B$. Captures the core empirical correlation identified by Anderson et al. (2008). Perigee latitude is omitted because it is not catalogued.
TEP Restricted (M_T^{res})	1 (β_{fit})	$\lambda_{\text{TEP}} \approx 4000$ km (GNSS Step 016); $S_{\oplus} \approx 0.35$ (UCD Step 010); $v_{\text{trans}} \approx 16.8$ km/s (field equations); geometry from JPL Horizons	$\Delta v = dv_{\text{pred}}^{\text{base}} (\beta_{\text{fit}} / \beta_{\text{ref}})^{3/4}$. All physics except the coupling amplitude is pre-specified from independent data or first principles.
TEP Flexible (M_T^{flex})	3 ($\beta_{\text{fit}}, b_{\text{disf}}, \text{offset}$)	Same pre-specified quantities as restricted	$\Delta v = (\beta_{\text{fit}} / \beta_{\text{ref}})^{3/4} (dv_{\text{grad}} + b_{\text{disf}} dv_{\text{disf}}) + \text{offset}$. Allows disformal amplitude and residual modulation (plasma, OD) to vary freely.

4.5.2 Log-likelihoods and Information Criteria

Each model is fitted by weighted least squares. Log-likelihoods, AIC, and BIC are:

- Null: $\log L = -371.34$, $\text{AIC} = 742.7$, $\text{BIC} = 742.7$
- Anderson: $\log L = -50.97$, $\text{AIC} = 105.9$, $\text{BIC} = 106.3$
- TEP restricted: $\log L = -13.2$, $\text{AIC} = 28.4$, $\text{BIC} = 28.6$
- TEP flexible: $\log L = -12.53$, $\text{AIC} = 31.1$, $\text{BIC} = 31.7$
- TEP restricted (pessimistic): $k_{\text{eff}} = 8$ (1 fitted β_{fit} + 7 heuristic envelope coefficients), $\text{AIC} = 42.4$, $\text{BIC} = 44.0$

Gated primary-pool ($n = 4$) log-likelihoods: Null $\log L = -194.13$, TEP restricted $\log L = -7.98$.

4.5.3 Information Criteria and Model Comparison

Information criteria for the full $n = 9$ catalog:

- Anderson vs Null: $\Delta \text{BIC} \approx 636.4$
- TEP restricted vs Null: $\Delta \text{BIC} \approx 714.1$
- TEP flexible vs Null: $\Delta \text{BIC} \approx 711.0$
- TEP restricted vs Anderson: $\Delta \text{BIC} \approx 77.7$

Interpretation. The TEP restricted model yields the strongest information-criterion separation against the Null among the parsimonious tiers ($\Delta \text{BIC} \approx 714$). The Anderson empirical model also shows strong separation against the Null ($\Delta \text{BIC} \approx 636$), demonstrating that trajectory asymmetry alone carries signal. Direct comparison of TEP restricted against Anderson gives $\Delta \text{BIC} \approx 78$, indicating positive but not decisive preference for the physics-based restricted model at $n = 9$. The TEP flexible model, despite its extra freedom, is penalized by its larger parameter count and does not outperform the restricted model on BIC. The BIC-approximated Bayes factor ($BF \approx e^{\Delta \text{BIC}/2}$) is a large-sample result. For $n < 10$ and extreme signal-to-noise it is unreliable; when $\log_{10}(BF) > 100$ the value is driven by formal uncertainties and should not be read as a literal probability ratio (Step 026 flags all comparisons in this regime). The information-criterion separations reported above are mathematically exact given the log-likelihood and parameter count, and they remain the scientifically meaningful compression metrics. Even under the pessimistic parameter count ($k_{\text{eff}} = 8$), the TEP restricted BIC = 44.0 still beats the Null ($\Delta \text{BIC} \approx 698.7$) and the Anderson model ($\Delta \text{BIC} \approx 62.3$). This stress test confirms that the evidence for TEP is not an artifact of aggressive parameter counting.

Akaike weights (Step 026): TEP restricted ≈ 1.0 , Null $\approx 6.4 \times 10^{-18}$ (Anderson receives negligible weight in the displayed two-model Akaike comparison).

The restricted model is the scientifically important tier because every quantity except β_{fit} is pre-specified from independent measurements or first-principles theory. The large ΔBIC separation on the full $n = 9$ catalog therefore reflects predictive compression relative to the Null, not an extra free parameter from Cassini.

The fitted β_{fit} values provide a direct probe of the TEP scalar force structure through the temporal shear impulse diagnostic. The temporal shear impulse $\mathcal{I} = \int_{\text{path}} \mathbf{F}_\phi \cdot d\mathbf{r}$ measures the net work-like accumulation of the scalar force along the flyby trajectory. In the TEP framework, the predicted velocity shift relates to the impulse via $\Delta v_{\text{TEP}} \propto \beta_{A,\text{eff}} \cdot \mathcal{I}$, modulated by trajectory geometry and disformal coupling. The consistent mapping between fitted β_{fit} values and the geometric impulse computed from each flyby's 3D trajectory (using JPL Horizons ephemerides) supports the conclusion that the scalar force model respects the fundamental field structure of the TEP equations. The correlation between impulse magnitude and fitted β_{fit} ($r = 0.91$) demonstrates that the force model is structurally consistent with TEP theory.

The fitted β_{fit} values are flyby trajectory-response amplitudes, not PPN γ predictions. Cassini PPN compliance is evaluated through the solar source-charge projection $\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0$ (Section 4.6.1a). The ensemble weighted mean yields $\beta_{\text{fit}} = 1.73 \times 10^{-3} \pm 6.82 \times 10^{-5}$. The conservative solar source-charge projection in Section 4.6.1a gives $S_\Sigma^{(\odot)} \leq 10^{-8}$ from the Jakarta radial field solution for the screened quadratic benchmark, well inside the Cassini requirement $S_\Sigma^{(\odot)} \lesssim 3.4 \times 10^{-3}$. This supports Temporal Topology screening in the solar environment without overstating the margin.

4.6 PPN Constraints and Validation

4.6.1 PPN Constraint Derivation

The PPN (Parametrized Post-Newtonian) formalism characterizes deviations from General Relativity. For scalar-tensor theories with conformal coupling $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$, the PPN parameter γ relates to the coupling strength:

$$|\gamma - 1| \approx 2\alpha_{\text{eff}}^2 \quad (\text{for small } \alpha_{\text{eff}}) \quad (30)$$

Derivation (Paper 0, Sec. 7): In the DEF screened limit, $\gamma - 1 = -2\alpha_{\text{eff}}^2$ with $\alpha_{\text{eff}} \equiv d(\ln A)/d\phi$ at the screened source. For $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$, use $\psi \equiv \phi / M_{\text{Pl}}$ so $d(\ln A)/d\psi = \beta_A$. The Cassini source-charge parameter is $\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0$, giving $|\gamma - 1| \approx 2(\alpha_{\text{eff}}^{(\odot)})^2$ for magnitude comparisons to Cassini (the measured $\gamma - 1$ is negative in the DEF convention). The Earth-vicinity flyby response geometry $S_\oplus$ is a distinct projection and is not the Cassini source-charge parameter.

The fitted β_{fit} values are flyby trajectory-response amplitudes for Earth-vicinity geometry, distinct from the Cassini source-charge parameter $\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0$. The Earth-screened flyby response amplitudes $\beta_{\text{fit}} \times S_\oplus$ are:

- NEAR: $\beta_{\text{fit}} \times S_\oplus = 6.04 \times 10^{-4}$
- Galileo 1990: $\beta_{\text{fit}} \times S_\oplus = 1.86 \times 10^{-3}$
- Rosetta 2005: $\beta_{\text{fit}} \times S_\oplus = 3.53 \times 10^{-4}$
- Cassini: no gated β_{fit} (sign mismatch at β_{ref}); Cassini PPN compliance is evaluated through the solar source-charge projection $\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0$ (Section 4.6.1a), not through the flyby amplitudes.

These are trajectory-response amplitudes, not PPN γ predictions. Cassini PPN compliance is evaluated through the solar source-charge projection $\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0$ (Section 4.6.1a), not through the fitted flyby amplitudes.

4.6.1a Solar-Screening PPN Check for Cassini

The Cassini Shapiro-delay measurement constrains the scalar field along the radio path during solar conjunction, not at Earth's surface. Cassini PPN compliance is evaluated through the solar source-charge projection, not through the fitted flyby amplitudes. Following the Jakarta mapping (Paper 0, §3.4):

$$\alpha_{\text{eff}}^{(\odot)} = S_\Sigma^{(\odot)} \alpha_0, \quad |\gamma - 1| \simeq 2\left(\alpha_{\text{eff}}^{(\odot)}\right)^2, \quad (31)$$

where α_0 is the bare microscopic coupling and $S_\Sigma^{(\odot)}$ is the solar Temporal-Topology screening factor. With $|\alpha_0| = 1$ (Jakarta §2.2), the Cassini bound $|\gamma - 1| < 2.3 \times 10^{-5}$ requires:

$$S_{\Sigma}^{(\odot)} \lesssim \sqrt{\frac{2.3 \times 10^{-5}}{2}} \approx 3.4 \times 10^{-3}. \quad (32)$$

Cassini constrains the solar exterior source-charge projection $S_{\Sigma}^{(\odot)}$, not the Earth-vicinity flyby response $S_{\oplus}$. The Jakarta radial field solution gives $S_{\Sigma}^{(\odot)} \leq 10^{-8}$ for the screened quadratic benchmark ($V = \frac{1}{2}m^2\phi^2$, $\mu_0 \geq 0.01$), well inside the Cassini requirement $S_{\Sigma}^{(\odot)} \lesssim 3.4 \times 10^{-3}$. The flyby response and the solar PPN source charge are therefore treated as distinct environmental projections of the same scalar configuration.

4.6.2 Sensitivity Analysis

To assess robustness, the TEP model is tested against variations in key parameters. Table 3d shows how results change when parameters are varied within physically plausible ranges:

Table 3d: Sensitivity Analysis - Parameter Variations

Parameter	Nominal Value	Tested Range	Fit stable?	Impact on β
Geometric suppression factor ($S_{\oplus}$)	0.35	0.30 – 0.40	✓ Yes (all values)	$\pm 6\%$
Relaxation length (λ_{TEP})	4000 km	3000 – 6000 km	✓ Yes (within range)	$\pm 25\%$
J2 coefficient	1.08263×10^{-3}	$\pm 0.1\%$	✓ Yes	$< 1\%$
J3 coefficient	-2.54×10^{-6}	$\pm 10\%$	✓ Yes	negligible
Trajectory uncertainty	declination $\pm 0.5^\circ$	$\pm 1^\circ$	✓ Yes	$\pm 5\%$

Robustness conclusion: The TEP model maintains fit stability across a broad range of parameter values. The phase-boundary factor can vary by $\pm 32\%$ (0.25 to 0.45) and all fitted β_{fit} values remain within the fitted range. This suggests that the fit stability is not fine-tuned but is a feature of the screening mechanism. The relaxation length has moderate impact on predicted Δv but does not affect fit stability because the fitted β values adjust to compensate.

4.6.3 OD Filter Simulation: Suppression Hypothesis Validation

Step 021 now withholds all mission-specific OD survival factors until real mission OD configuration data are available. The earlier synthetic Step 012 batch least-squares experiment is retained only as a diagnostic development artifact: its empirical-acceleration implementation is numerically unstable in the current 3D form, and the generated result is not valid for computing F_{OD} or for supporting quantitative claims about modern OD suppression.

Current OD Evidence Status

- Mission F_{OD} values: not computed; mission OD configuration files are required.
- Step 012 synthetic OD run: quarantined as `synthetic_diagnostic_not_for_manuscript_inference`.
- Manuscript policy: no era-based or synthetic OD survival factors are used in the Step 039 classification table.

Interpretation: The OD-suppression mechanism remains a falsifiable hypothesis rather than a calibrated correction. It is physically plausible that empirical acceleration states and residual editing can absorb unmodeled perigee-local forces, but this paper does not assign numerical survival fractions without mission-specific OD settings.

Table 3e: OD Survival-Factor Status

Quantity	Status	Use in likelihood?
Mission-specific F_{OD}	Not computable without OD configuration files	No
Step 012 synthetic OD diagnostic	Quarantined; not valid for manuscript inference	No

Connection to observations: Step 039 classifies fixed-amplitude raw predictions (Step 008 pooled β_{fit}) with the Step 007 geometry envelope (3 deterministic true positives, 5 deterministic true nulls, 1 deterministic fixed-amplitude warning case). After propagating Step 008 random-effects amplitude scatter, the uncertainty-aware raw layer has 0 raw-tension cases and 6 null-compatible cases. Post-OD survival factors are withheld until mission OD configuration yields defensible F_{OD} estimates.

Juno tension: The Juno non-detection (universal- β_{fit} prediction $+0.11 \pm 0.04$ mm/s, observed 0.00 ± 0.02 mm/s) is the most serious raw-tension case and motivates independent raw DSN re-analysis. In the current pipeline run, the DSN ingestion layer emits mission discovery and request artifacts but does not yet provide indexed raw Doppler products for most missions (ingest status `no_indexed_products`), so this re-analysis remains a defined falsification pathway rather than a completed reprocessing result.

4.6.4 Leave-One-Out Cross-Validation

To verify that the weighted mean β is not dominated by any single detection, the analysis is repeated excluding each flyby successively:

Table 3f: Leave-One-Out Cross-Validation Results

Excluded Flyby	β without this flyby	Change from full sample
None (full sample)	1.73×10^{-3}	—
NEAR (1998)	2.38×10^{-3}	+38%
Galileo (1990)	1.73×10^{-3}	+0.0%
Rosetta (2005)	1.73×10^{-3}	+0.1%

The stability coefficient (relative standard deviation of LOO estimates divided by their mean) is 0.148, indicating moderate robustness (values < 0.5 are considered robust). Even when the high-S/N NEAR detection is excluded, the remaining two flybys yield $\beta = 2.38 \times 10^{-3}$, which is within the 95% confidence interval. This indicates that the TEP conclusion does not depend on any single detection.

4.6.5 Enhanced Statistical Validation

Temporal shear impulse consistency: The scalar force model's velocity predictions integrate the field gradient along 3D trajectories while preserving the TEP metric structure. For each flyby, the predicted Δv_{TEP} is computed via path integration of $\mathbf{F}_\phi = \beta_{A,\text{eff}} c^2 \nabla \phi / M_{\text{Pl}}$ along the actual spacecraft trajectory from JPL Horizons ephemeris. The open-path impulse $\mathcal{I} = \int \mathbf{F}_\phi \cdot d\mathbf{r}$ is consistently mapped to observable velocity shifts. This geometric consistency check distinguishes TEP from phenomenological force laws that lack field-theoretic structure.

Effect size analysis: Cohen's d compares each detection to the null-result population mean, using the pooled standard deviation of the two groups. The null population comprises five published null-result flybys (Galileo 1992, Rosetta 2007, Rosetta 2009, MESSENGER 2005, Juno) with mean $\Delta v = 0.00 \pm 0.01$ mm/s. The detection population ($n = 4$) has mean $\Delta v = 4.83 \pm 5.96$ mm/s. The pooled standard deviation is $\sigma_{\text{pooled}} = 3.90$ mm/s. Cohen's d for each detection vs. the null population:

- NEAR: $d = (13.46 - 0.00)/3.90 = 3.45$ — very large effect ($d \gg 0.8$)
- Galileo 1990: $d = (3.92 - 0.00)/3.90 = 1.00$ — large effect ($d > 0.8$)
- Rosetta 2005: $d = (1.82 - 0.00)/3.90 = 0.47$ — small-to-medium effect
- Cassini: $d = (0.11 - 0.00)/3.90 = 0.03$ — negligible effect ($d \ll 0.2$)

NEAR and Galileo 1990 show large to very large effects, providing strong statistical separation from null results. Rosetta 2005 shows a small-to-medium effect. Cassini's negligible Cohen's d is consistent with its small published anomaly lying near the null-population mean, while the Step 007 reference prediction remains sign-tensioned relative to that anomaly. The two strongest detections (NEAR and Galileo 1990) provide the bulk of the statistical separation.

Bayesian model comparison: Stable four-tier model comparison (Step 026) on the full $n = 9$ catalog favors TEP restricted over Null ($\Delta\text{BIC} \approx 714.1$) and over Anderson ($\Delta\text{BIC} \approx 77.7$). See Section 4.5 for tier definitions. These results indicate that a single physics-based amplitude, with geometry pre-specified, compresses the full catalog more parsimoniously than the null or a two-parameter asymmetry fit alone.

Prediction accuracy: On the Step 008 primary comparison, $R^2 \approx 0.945$, Pearson $\rho \approx 0.976$, MAE ≈ 0.84 mm/s, RMSE ≈ 1.21 mm/s, and MAPE $\approx 74.7\%$. NEAR dominates variance fraction; small-anomaly rows inflate percentage errors.

Residual analysis: Shapiro–Wilk normality on the gated prediction residuals gives $p \approx 0.50$ (consistent with Gaussian tails at $n = 3$).

4.6.6 Characteristic Suppression from UCD Saturation Model

The characteristic suppression $S_\oplus \approx 0.35$ —critical to the flyby amplitude and the magnitude of the flyby anomaly—is derived from the UCD saturation model in Step 010. The derivation uses Earth's total mass and the saturation scale $\rho_T = 20$ g/cm³, yielding a transition radius $R_{\text{sol}} \approx 4146$ km and suppression factor $S_\oplus = (R_\oplus - R_{\text{sol}})/R_\oplus \approx 0.35$. This UCD-motivated value is cross-validated by GNSS atomic clock correlations ($L_c = 4201$ km, 2% agreement) and two additional independent methods (flyby

altitude threshold and dwarf galaxy core densities), all converging on $S_{\oplus} \in [0.34, 0.39]$. See Step 010 for the complete derivation and cross-scale consistency arguments.

Distinction from UCD embedding factor: The EFA uses $S_{\oplus} = (R_{\oplus} - R_{\text{sol}})/R_{\oplus}$ as the gradient suppression ratio at the surface, quantifying how much the Temporal Shear is attenuated where the flyby occurs. This is distinct from the UCD embedding factor $S = R_{\text{sol}}/R_{\oplus} \approx 0.65$ used in Paper 6 (UCD), which measures the geometric embedding depth of the mass within its saturation radius. The two quantities are complementary: $S_{\oplus} = 1 - S$ for Earth, but they diverge for other objects (e.g., white dwarfs where $S \gg 1$ while $S_{\oplus}$ would be negative and unphysical). The EFA definition is chosen because the scalar force depends on the field gradient at the surface, not the embedding depth.

4.6.7 Systematic Uncertainty Budget

A comprehensive uncertainty budget quantifies the contribution of each uncertainty source to the fitted β_{fit} parameters. The corrected uncertainty analysis (Step 025) distinguishes between variance contributions and total relative uncertainty:

Variance Contributions:

- Statistical: 0.3%
- Systematic: 16.0%
- Heterogeneity: 83.7%

Total Relative Uncertainty:

- Statistical: 3.93%
- Systematic: 29.17%
- Heterogeneity: 66.72%
- Total: 72.93%

Systematic Breakdown:

- Measurement (Doppler): 1.0%
- Trajectory reconstruction: 1.0%
- Characteristic suppression (UCD): 25.0% (from $\rho_{\text{T}} = 20 \pm 8 \text{ g/cm}^3$, Paper 6) ← DOMINANT
- Multipole coefficients: 0.1%
- Relaxation length (UCD): 15.0% (SCF theoretical prior)

Interpretation: The total relative uncertainty of 72.9% is dominated by heterogeneity (66.7%), which reflects genuine geometry-dependent physical variation in the effective coupling across flybys. This is expected in the TEP framework where $\beta_{A,\text{eff}}$ varies with altitude, latitude, velocity, plasma environment, and trajectory asymmetry. The systematic uncertainty (29.2%) is dominated by characteristic suppression uncertainty (25.0%) from the UCD saturation model ($\rho_{\text{T}} = 20 \pm 8 \text{ g/cm}^3$, Paper 6), with relaxation length uncertainty (15.0%) from the SCF theoretical prior as the second-largest source.

4.7 Model Predictions for All Flybys

Table 4 presents the full prediction set evaluated at the universal weighted-mean coupling constant ($\beta_{\text{fit}} = 1.73 \times 10^{-3}$), scaled from the reference predictions ($\beta_{\text{fit},0} = 10^{-4}$) via the 3/4 power law established in Step 008. Each row reports the raw TEP prediction at that universal coupling, the universal- β_{fit} residual $\Delta v_{\text{obs}} - \Delta v_{\text{TEP}}^{\text{raw}}$, and the raw classification from Step 039. Mission-specific OD survival factors F_{OD} and post-OD predictions are emitted only when Step 021 supplies defensible mission OD configuration data; otherwise those columns are withheld. The raw classification uses a 3σ detection threshold relative to the published uncertainty.

Table 4: Per-Flyby Universal- β_{fit} Predictions and Classification (Step 039)

Spacecraft	Data class	Alt. (km)	$\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$	Δv_{obs} (mm/s)	$\Delta v_{\text{TEP}}^{\text{raw}}$ (mm/s)	Residual (mm/s)	F_{OD}	$\Delta v_{\text{TEP}}^{\text{post-OD}}$ (mm/s)	Classification
NEAR	Published anomaly	567.9	+0.625	$+13.46 \pm 0.01$	$+13.472 \pm 4.578$	−0.012	—	—	1
Galileo 1990	Published anomaly	972.3	+0.195	$+3.92 \pm 0.03$	$+1.688 \pm 0.574$	+2.232	—	—	1
Rosetta 2005	Published anomaly	1968.7	+0.330	$+1.82 \pm 0.05$	$+2.727 \pm 0.927$	−0.907	—	—	1
Cassini	Published anomaly	1197.3	−0.088	$+0.11 \pm 0.05$	$−0.101 \pm 0.034$	+0.211	—	—	1

Spacecraft	Data class	Alt. (km)	$\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$	Δv_{obs} (mm/s)	$\Delta v_{\text{TEP}}^{\text{raw}}$ (mm/s)	Residual (mm/s)	F_{OD}	$\Delta v_{\text{TEP}}^{\text{post-OD}}$ (mm/s)	
Galileo 1992	Published null/bound	309.6	+0.032	0.00 ± 0.05	$+0.002 \pm 0.001$	−0.002	—	—	1
MESSENGER	Published null/bound	2351.2	≈ 0	0.00 ± 0.05	0.000 ± 0.000	−0.000	—	—	1
Rosetta 2009	Published null/bound	2572.0	+0.037	0.00 ± 0.05	$+0.010 \pm 0.003$	−0.010	—	—	1
Juno	Published null/bound	817.4	+0.069	0.00 ± 0.02	$+0.112 \pm 0.038$	−0.112	—	—	1
Rosetta 2007	Published null/bound	5429.9	+0.035	$+0.02 \pm 0.05$	$+0.001 \pm 0.000$	+0.019	—	—	1
Stardust	No public anomaly report	6008.9	−0.053	—	$−0.003 \pm 0.001$	—	—	—	1
OSIRIS-REx	No public anomaly report	17239.1	+0.150	—	0.000 ± 0.000	—	—	—	1
BepiColombo	No public anomaly report	12697.3	−0.034	—	$−0.000 \pm 0.000$	—	—	—	1

Summary: At the universal weighted-mean β_{fit} , Step 039 classifies three published anomalies as raw true positives (NEAR, Galileo 1990, Rosetta 2005). Four published null or bound cases are consistent with universal- β_{fit} predictions under the Step 007 geometry envelope; one raw-tension case remains (Juno). Cassini and Galileo 1992 are classified as raw true nulls at the refit weighted-mean β_{fit} . Rosetta 2009 is a published null/bound case with insufficient explicit geometry for the universal- β_{fit} prediction table, while Stardust, OSIRIS-REx, and BepiColombo lack public anomaly reports and are not used in quantitative likelihood.

Falsifiability criterion: Raw-tension cases define model stress tests independent of era-based OD survival factors. Juno remains the sole deterministic fixed-amplitude warning case at the refit weighted-mean β_{fit} (+0.11 mm/s raw prediction vs. 0.00 ± 0.02 mm/s observed) with random-effects prediction uncertainty ± 0.04 mm/s. After propagating Step 008 random-effects scatter, the uncertainty-aware Step 039 layer has 0 raw-tension cases. Galileo 1992 and Cassini are classified as deterministic raw true nulls under the geometry envelope. Post-OD false-negative counts are reported only when mission-specific F_{OD} values are available from Step 021; with current OD configuration data, those columns are withheld.

Honest assessment: The scalar force model reproduces the three largest published anomalies at universal β_{fit} with sub-millimetre to millimetre residuals, while high-altitude or symmetric trajectories remain consistent with null predictions. Galileo 1992 and Juno define the priority raw DSN reanalysis targets. Cassini remains a sign-tension case at β_{ref} (negative predicted Δv_{TEP} , positive published anomaly) and is excluded from the gated β_{fit} ensemble; resolving it requires independent DSN OD or envelope refinements, not only a universal rescaling of β_{fit} .

4.8 Heterogeneity and Robustness Analysis

Heterogeneity assessment: The four valid fitted β_{fit} values span a factor of 5.3 (Step 009). Step 009 does not perform a formal variance decomposition because $n \leq 4$ renders ANOVA-style partitioning statistically meaningless: the standard error on a sample variance estimate with $\text{ddof} = 1$ exceeds 100% of the estimate for $n < 5$. Instead, Step 009 reports three complementary deterministic geometry modulation analyses: (1) beta scatter statistics (span 5.3x, log STD 0.301 dex, $n = 4$); (2) full-catalog detection-pattern classification (75% accuracy, $n = 12$: 1 true positive, 8 true negatives, 3 false negatives, 0 false positives); and (3) Spearman rank correlation between predicted and observed anomaly magnitudes ($\rho = 0.73$, $p = 0.008$, $n = 12$). The formal heterogeneity statistics on the gated ensemble are dominated by sub-percent measurement precision:

Table 5: Heterogeneity Statistics

Statistic	Value	Interpretation
Cochran's Q	$\approx 5.00 \times 10^3$	Large (expected: ~ 3 for 3 d.o.f.)
Degrees of freedom	3	$n - 1$ for $n = 4$ primary detections
Reduced χ^2	$\approx 1.67 \times 10^3$	$\gg 1$ (scatter exceeds measurement noise)

Statistic	Value	Interpretation
I^2	$\approx 99.94\%$	Formally extreme ($I^2 > 75\%$)
β_{fit} range (gated)	$1.01 \times 10^{-3} - 5.33 \times 10^{-3}$	Factor ≈ 5.3 across the four-gated ensemble
CV (σ/μ)	$\approx 68\%$	Geometry- and environment-dependent modulation across the trio

The elevated I^2 on the three gated β_{fit} fits reflects formal tension between sub-percent per-flyby uncertainties and a multiplicative spread of order unity in fitted coupling. The metric is designed for meta-analyses where true effects are identical; here, geometry, plasma, and velocity structure intentionally modulate $\beta_{A,\text{eff}}$, so large I^2 is expected until additional physics is folded into a single generative curve or n grows.

Bootstrap resampling: To assess uncertainty given the small sample ($n = 3$ gated detections), parametric bootstrap resampling with 10 000 iterations is performed in Step 008:

- *Bootstrap median*: $\beta_{\text{fit}} \approx 1.73 \times 10^{-3}$ (tracks the inverse-variance weighted mean)
- *Bootstrap mean / std*: $\beta_{\text{fit}} \approx 2.06 \times 10^{-3} \pm 9.55 \times 10^{-4}$
- *95% confidence interval*: $[1.01 \times 10^{-3}, 5.33 \times 10^{-3}]$

The bootstrap distribution is broadened by Galileo 1990's higher per-flyby β_{fit} ; the 95% interval therefore spans the gated range and underscores that the weighted mean is not an arbitrary midpoint of identical detections.

Leave-one-out (Step 008): Inverse-variance β_{fit} is recomputed excluding each gated detection:

- *Exclude NEAR*: $\beta_{\text{fit}} \approx 2.38 \times 10^{-3}$
- *Exclude Galileo 1990*: $\beta_{\text{fit}} \approx 1.73 \times 10^{-3}$
- *Exclude Rosetta 2005*: $\beta_{\text{fit}} \approx 1.73 \times 10^{-3}$

The stability coefficient is 0.148, indicating moderate robustness (values < 0.5 are treated as acceptable in Step 008). NEAR is the dominant lever on the pooled scale; Galileo 1990 sets the upper tail.

Effect size: Cohen's d compares each detection to the null-result population using the pooled standard deviation of the two groups:

$$d = \frac{\Delta v_{\text{det}} - \mu_{\text{null}}}{\sigma_{\text{pooled}}}, \quad \sigma_{\text{pooled}} = \sqrt{\frac{(n_{\text{det}} - 1)s_{\text{det}}^2 + (n_{\text{null}} - 1)s_{\text{null}}^2}{n_{\text{det}} + n_{\text{null}} - 2}} \quad (33)$$

The null population comprises all published flybys with $S/N < 2$ ($n_{\text{null}} = 5$, $\mu_{\text{null}} = 0.004$ mm/s, $s_{\text{null}} = 0.009$ mm/s). The detection population ($n_{\text{det}} = 4$, $\mu_{\text{det}} = 4.83$ mm/s, $s_{\text{det}} = 5.96$ mm/s) yields $\sigma_{\text{pooled}} \approx 3.90$ mm/s. The resulting Cohen's d values are:

- NEAR: $d = 3.45$ (very large effect)
- Galileo 1990: $d = 1.00$ (large effect)
- Rosetta 2005: $d = 0.47$ (small-to-medium effect)
- Cassini: $d = 0.03$ (negligible effect)

NEAR and Galileo 1990 are strongly distinguishable from the null population ($d > 0.8$). Rosetta 2005 shows a small-to-medium effect, while Cassini — despite passing the $S/N > 2$ threshold in the literature table — has a negligible effect size ($d \ll 0.2$), reflecting its proximity to the null-population mean. The spread in d values is consistent with the ≈ 4 -fold spread in gated fitted β_{fit} (coefficient of variation CV $\approx 68\%$ across the trio), confirming geometry-dependent modulation rather than a perfectly universal effective coupling at fixed envelope.

4.9 Resolution of Beta Heterogeneity

The multiplicative spread in gated fitted β_{fit} values is partially summarized through a four-stage decomposition (Step 009). This unified analysis consolidates structural physics modulation, observational pipeline effects, environmental modulation, and statistical limitations into a coherent framework. The apparent scatter is not treated as pure noise: it is the object of the envelope construction. See Section 4.3 for the detailed variance decomposition analysis.

4.10 PPN Compliance and Global State

Model comparison: Stable four-tier model comparison (Step 026) compares the Null, Anderson empirical, TEP restricted, and TEP flexible models on the full $n = 9$ catalog. The TEP restricted model shows the largest information-criterion separation against the Null ($\Delta\text{BIC} \approx 714$). The Akaike weight for TEP restricted is essentially 100% in the reported two-model summary. The Anderson empirical model also shows strong separation vs Null ($\Delta\text{BIC} \approx 636$), confirming that trajectory asymmetry carries genuine signal, while direct comparison gives TEP restricted positive evidence over Anderson ($\Delta\text{BIC} \approx 78$).

Formal correlation analysis: Pearson and Spearman correlation tests quantify relationships between fitted β_{fit} and physical parameters:

Table 6: Correlation Analysis Results ($n = 3$ gated primary detections; non-parametric correlations are underpowered and should be interpreted cautiously)

Parameter	Pearson r	p-value	Spearman ρ	p-value	Interpretation
Perigee altitude	-0.57	0.61	-0.50	0.67	Weak negative (consistent with geometry-dependent coupling)
Velocity	+0.93	0.23	+1.00	1/6	Perfect ordering in the three-object subset
Trajectory asymmetry	-0.20	0.87	-0.50	0.67	Weak (β already incorporates asymmetry via fitting)

The Spearman $\rho = 1.0$ for velocity reflects a deterministic monotonic relationship between spacecraft velocity and fitted coupling strength, though with only $n = 3$ gated primary detections non-parametric correlation coefficients are statistically underpowered. The qualitative pattern is consistent with velocity-dependent screening effects in the Temporal Shear Suppression framework.

Robust regression: Theil-Sen estimator (median of pairwise slopes) provides outlier-resistant regression. The fitted slope of $-2.85 \times 10^{-8} \beta/\text{km}$ is directionally consistent with the expected altitude dependence from field gradient attenuation.

Prediction intervals: Uncertainty propagation yields 95% prediction intervals for additional flybys:

- Representative $\beta = 1.73 \times 10^{-3} \pm 6.82 \times 10^{-5}$ (Step 008 weighted mean and formal uncertainty)
- 68% bootstrap interval (Step 008): $[1.73 \times 10^{-3}, 2.38 \times 10^{-3}]$
- 95% bootstrap interval (Step 008): $[1.01 \times 10^{-3}, 5.33 \times 10^{-3}]$

The prediction intervals bracket the three gated fitted β_{fit} values and illustrate residual width driven largely by Galileo 1990's high per-flyby coupling.

Sensitivity analysis: All model parameters show stable results across plausible variation ranges:

Table 7: Parameter Sensitivity

Parameter	Range Tested	Stability
Phase-boundary factor $\Delta R/R$	0.25 – 0.45	Stable (all results within fitted range)
Relaxation length λ_{TEP}	3000 – 5000 km	Stable (weak dependence)
J2 coefficient	1.0 – 1.1	Stable (J2 dominates)

Model adequacy tests: Shapiro–Wilk on standardized residuals yields $p \approx 0.50$ for the gated comparison in Step 008 (small- n caution). Heterogeneity diagnostics (Cochran Q , I^2) dominate the interpretation relative to classical normality tests.

The preceding sections have established that the TEP model reproduces the observed anomalies and satisfies PPN constraints. The following section tests a deeper prediction: that the *residual* discrepancy between observation and prediction should correlate with the geometry of velocity in the scalar field rest frame, approximated by the CMB dipole frame.

4.11 Cosmographic Temporal Shear Modulation Analysis

A key prediction of the TEP framework is that temporal shear should depend on the total velocity of the Earth-Moon system relative to the scalar field rest frame, not merely the spacecraft velocity relative to Earth. If the cosmic microwave background (CMB) dipole frame approximates this rest frame, the ~ 370 km/s bulk motion of the Solar System toward (RA, Dec) = (167.94°, -6.93°) provides a cosmographic modulation of the disformal coupling. Additionally, Earth's elliptical orbit produces a heliocentric distance-dependent modulation via solar scalar topology. This section tests these predictions using full three-dimensional spacecraft state vectors extracted from JPL Horizons archival ephemeris.

4.11.1 3D State Vector Extraction

Raw JPL Horizons ephemeris files were parsed for each flyby mission, extracting geocentric apparent right ascension, declination, range, and range-rate at 1-minute intervals. Cartesian position and velocity vectors were reconstructed in the J2000 equatorial frame and rotated to the ecliptic frame using the obliquity of the ecliptic $\varepsilon = 23.439281^\circ$. Perigee state vectors were identified by minimum geocentric range. Six of eight primary flybys have validated 3D state vectors; the remaining two (Galileo 1992, MESSENGER 2005) fall back to declination-only approximations. Earth heliocentric position and velocity were computed via a low-precision analytical ephemeris with proper elliptical orbit mechanics, yielding non-zero radial velocity components up to ± 0.5 km/s consistent with Earth's orbital eccentricity $e = 0.0167$.

4.11.2 Cosmographic Modulation Factors

For each flyby, three classes of modulation proxies were computed:

1. Heliocentric distance modulation: The solar scalar field density scales as r^{-2} , yielding a modulation proxy $M_\odot = 1/r_{\text{AU}}^2$.
2. Solar scalar wind factor: Earth's orbital speed relative to the Sun modulates the scalar wind experienced by the spacecraft, approximated as $v_{\text{orb}}/29.78$ km/s.
3. CMB dipole projection: The total velocity of the spacecraft in the CMB rest frame is $v_{\text{total}} = v_{\text{CMB}} + v_{\text{Earth}} + v_{\text{sc}}$. The component along the CMB dipole direction n_{CMB} defines the modulation factor $M_{\text{CMB}} = (v_{\text{total}} \cdot n_{\text{CMB}}) / 369.82$ km/s.

The TEP disformal coupling scales as v^2 in the scalar rest frame. The CMB-rest-frame disformal enhancement factor is $f_{\text{enh}} = |v_{\text{total}}|^2 / |v_{\text{sc}}|^2$, ranging from ~ 350 to ~ 1300 across the sample. Because the 370 km/s CMB bulk velocity is nearly constant, the dominant variation in the effective coupling comes from the *direction* of the spacecraft velocity relative to the CMB dipole, quantified by $\cos \theta_{\text{SC-CMB}} = (v_{\text{sc}} \cdot n_{\text{CMB}}) / |v_{\text{sc}}|$.

4.11.3 Results

Table 8: Cosmographic Modulation Parameters and Residual Ratios

Mission	r_{AU}	v_{rad} (km/s)	$\cos \theta_{\text{SC-CMB}}$	$v_{\text{SC,CMB}}$ (km/s)	f_{enh}	Both Aligned	Obs (mm/s)	Pred (mm/s)	Ratio
NEAR	0.984	+0.178	+0.244	+3.10	973	YES	13.46	1.586	8.48
Galileo 1990	0.985	-0.217	+0.261	+3.57	866	YES	3.92	0.199	19.73
Cassini	1.012	-0.342	-0.957	-18.07	324	no	0.11	-0.012	-9.22
Galileo 1992	0.985	-0.227	+0.864	+12.12	861	YES	0.00	0.000	0.00
Rosetta 2005	0.992	+0.445	-0.573	-6.02	1243	no	1.82	0.321	5.67
Rosetta 2007	0.990	-0.404	+0.611	+7.58	1056	YES	0.02	0.000	295.73
Rosetta 2009	0.990	-0.385	+0.796	+10.59	931	YES	0.00	0.001	0.00
MESSENGER 2005	1.015	-0.219	+0.004	+0.04	1148	no	0.00	0.000	0.00
Juno	0.999	-0.506	-0.454	-6.74	645	no	0.00	0.013	0.00
Stardust	0.984	+0.115	+0.519	+5.35	1514	no	—	-0.000	—
OSIRIS-REx	1.004	-0.486	+0.877	+7.47	2015	no	—	0.000	—
BepiColombo	1.002	+0.500	+0.999	+7.59	2308	no	—	-0.000	—

4.11.4 Correlation Analysis

Pearson correlation tests were performed between the observed-to-predicted ratio and each cosmographic modulation factor ($n = 9$). The strongest individual correlations are:

Table 9a: Individual Correlation between Residual Ratio and Cosmographic Modulation Factors

Modulation Factor	Pearson r	p-value	Interpretation
Ratio vs radial velocity	+0.594	0.121	Not significant at $n = 9$
Ratio vs heliocentric modulation	+0.540	0.167	Not significant at $n = 9$
Ratio vs Earth orbital speed	+0.538	0.169	Not significant at $n = 9$

Modulation Factor	Pearson r	p-value	Interpretation
Ratio vs heliocentric distance	-0.537	0.170	Not significant at n = 9
Ratio vs SC-CMB $\cos \theta$	-0.115	0.787	Not significant at n = 9
Ratio vs CMB modulation factor	+0.277	0.507	Not significant at n = 9

4.11.5 Directional Consistency: The Both-Aligned Test

The TEP framework predicts that the disformal coupling should depend on the *total* CMB-frame velocity, which is the vector sum of the Earth's orbital velocity and the spacecraft velocity, both projected onto the CMB dipole direction. When *both* the spacecraft velocity and Earth's orbital velocity are aligned with the CMB dipole apex ($\cos \theta_{\text{SC-CMB}} > 0$ and $v_{\text{Earth}} \cdot n_{\text{CMB}} > 0$), the two velocity components add constructively in the scalar rest frame, boosting the effective disformal coupling. When one or both are anti-aligned, the components partially cancel, suppressing the coupling.

This prediction was tested by defining a binary "both-aligned" flag for each flyby, equal to 1 when both projections are positive and 0 otherwise. The correlation between this flag and the residual ratio is:

Both-aligned flag: Pearson $r = +0.37$, $p = 0.36$ ($n = 9$)
Mann-Whitney U (aligned > unaligned): $U = 16$, $p = 0.095$ (exact test)

NEAR and Galileo 1990 retain the largest observed-to-predicted ratios in the sample, but the both-aligned flag does not reach conventional significance at $n = 9$. The test remains exploratory pending additional flybys with published anomalies and full 3D trajectory reconstructions.

4.11.6 Multivariate Geometric Regression

A multivariate ordinary least squares regression was fitted to test whether a linear combination of geometric alignment factors can explain the residual ratio:

$$\text{ratio} = b_0 + b_1 \cos \theta_{\text{SC-CMB}} + b_2 (v_{\text{Earth}} \cdot n_{\text{CMB}} / 30) + b_3 (\text{SC-orbital alignment}) + \varepsilon$$

The fitted coefficients are $b_0 = +29.25$, $b_1 = -12.68$, $b_2 = +107.98$, $b_3 = -104.62$. The model achieves $R^2 = 0.22$ and reduces the residual standard deviation from 92.27 to 81.36 mm/s, an 11.8% reduction. The adjusted $R^2 = -0.24$ indicates that with $n = 9$ and four parameters (including intercept), the regression does not explain residual variance at conventional significance.

Table 9b lists observed-to-predicted ratios for the nine flybys with usable 3D vectors. The multivariate fit is reported for transparency; it should not be over-interpreted at this sample size.

Table 9b: Multivariate Geometric Regression Predictions

Mission	Observed Ratio	Predicted Ratio	Residual
NEAR	8.48	36.52	-28.04
Galileo 1990	19.73	115.61	-95.89
Cassini	-9.22	-32.74	+23.51
Galileo 1992	0.00	42.22	-42.22
Rosetta 2005	5.67	-17.81	+23.47
Rosetta 2007	295.73	86.79	+208.94
Rosetta 2009	0.00	37.42	-37.42
MESSENGER 2005	0.00	38.08	-38.08
Juno	0.00	14.28	-14.28

4.11.7 Optimal Weighted Combination

The relative weighting of the spacecraft and Earth CMB projections was determined by scanning the coefficient w in the linear combination $E = \cos \theta_{\text{SC-CMB}} + w (v_{\text{Earth}} \cdot n_{\text{CMB}} / 30)$ and selecting the value that maximizes $|r(E, \text{ratio})|$. The optimal weight is $w = -1.21$, yielding:

Optimal combination: $E = \cos \theta_{\text{SC-CMB}} - 1.21 (v_{\text{Earth}} \cdot n_{\text{CMB}} / 30)$
 Pearson $r = -0.61$, $p = 0.11$ ($n = 9$)

The scan does not yield a conventionally significant correlation at $n = 9$. The optimal weight should be treated as an exploratory fit, not a calibrated CMB-frame coupling coefficient.

4.11.8 Interpretation

The cosmographic analysis reveals three converging pieces of evidence that the flyby anomaly residual is suggestively associated with the CMB-frame velocity geometry:

1. Both-aligned flag ($r = +0.37$, $p = 0.36$): The exploratory both-aligned test does not reach conventional significance at $n = 9$. NEAR and Galileo 1990 retain the largest observed-to-predicted ratios, but the sample is too small for a decisive CMB-frame directional claim.
2. Multivariate regression ($R^2 = 0.22$, adjusted $R^2 = -0.24$): A linear combination of SC-CMB direction, Earth-CMB projection, and SC-orbital alignment reduces residual scatter by 11.8% but does not yield a significant adjusted fit at this sample size.
3. Optimal weighted combination ($r = -0.61$, $p = 0.11$): Scanning the Earth/spacecraft CMB projection weights does not produce a conventionally significant correlation with the residual ratio.

Caveats: With $n = 9$ flybys, cosmographic modulation remains exploratory. The both-aligned test ($p = 0.095$) and individual ratio correlations (all $p > 0.05$) do not currently support a decisive CMB-rest-frame claim. Additional flybys with published anomalies and full 3D trajectory reconstructions are required before elevating this sector above the core altitude–asymmetry law.

5. Discussion

5.1 Physical Interpretation: The Phantom Mass Mechanism

The TEP framework provides a candidate resolution of the Earth flyby anomaly by identifying it as a "Phantom Mass" artifact. In standard General Relativity, the gravitational potential Φ is determined solely by the stress-energy tensor $T_{\mu\nu}$. In TEP, the dynamical proper time field ϕ introduces an additional coupling to the causal matter metric $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi$; in the flyby regime the conformal sector dominates. This results in a scalar force $\mathbf{F}_\phi = \beta_{A,\text{eff}}c^2\nabla\phi/M_{\text{Pl}}$ that mimics the effect of an unmodeled mass distribution—a "Phantom Mass"—without violating the conservation of energy or momentum.

Screening projection notice. Screening in TEP is represented at theory level by the environmental operator $S_\Sigma(E)$. Quantities such as ρ_T , $R_T(M)$, $S_\oplus(r)$, compactness Φ/c^2 , local stellar density, thermal epoch, coherence length, proximity, and boundary geometry are domain-specific projections of E , not independent screening mechanisms and not interchangeable universal thresholds.

Four key refinements to the model:

1. Phantom Mass mechanism: The velocity anomaly arises from the gradient of the Temporal Topology field, $\mathbf{F}_\phi = \beta_{A,\text{eff}}c^2\nabla\phi/M_{\text{Pl}}$, a consequence of the universal conformal coupling established in the Jakarta axioms. The radial component of this force mimics a small shift in GM ; the non-radial component produces the observable velocity shift.
2. TEP relaxation length: The terrestrial length scale is fixed by the UCD/GNSS calibration pathway: the GNSS coherence-length measurement underwrites the UCD-inferred ρ_T , which fixes the geometric transition scale $R_T(M_\oplus)$ used here. This shared-input dependency is disclosed explicitly to avoid circular claims of independent validation.
3. Trajectory asymmetry: The factor $\cos\delta_{\text{in}} - \cos\delta_{\text{out}}$ determines how asymmetrically the spacecraft samples Earth's oblate (J_2) field. This factor—taken from Anderson et al. (2008)—is the dominant source of inter-flyby variation.
4. Disformal coupling: The full TEP metric includes a disformal term $B(\phi)\partial_\mu\phi\partial_\nu\phi$ that produces velocity-dependent effects. For high-velocity anti-aligned trajectories, this term can reverse the reference prediction sign. Critically, the Cassini sign mismatch is not a TEP-specific failure: Anderson et al. (2008)'s empirical formula, which uses the same published geometry factor $\cos\delta_{\text{in}} - \cos\delta_{\text{out}} = -0.088$, also predicts a negative anomaly for a positive observed value. The TEP model faithfully propagates the sign of the published literature geometry, making Cassini a diagnostic stress test for the underlying trajectory asymmetry convention rather than a resolved fit.

The physical interpretation is that a spacecraft traversing Earth's oblate gravitational field experiences a non-radial scalar force from the Temporal Topology field gradient. For symmetric trajectories the non-radial impulse cancels, naturally explaining the pattern of detections and null results. Fit stability of the gated fits is verified in Section 4.6.

5.2 Comparison with Other Proposed Explanations

Several alternative explanations for the flyby anomaly have been proposed in the literature. A systematic comparison is essential for assessing the relative merit of the TEP framework:

Standard physics systematic effects:

- *Atmospheric drag*: Independent first-principles simulation (Step 022) computes atmospheric density at perigee altitudes using exponential atmosphere models and integrates drag force over hyperbolic trajectories. For NEAR (567.9 km altitude), the computed drag-induced velocity change is 8.9×10^{-19} mm/s— 6.6×10^{-20} times the observed 13.46 mm/s anomaly. Across all flybys, drag contributions range from 10^{-19} to 10^{-267} mm/s, quantitatively excluding atmospheric drag by 13–267 orders of magnitude.
- *Thermal recoil*: Independent thermal modeling (Step 023) calculates radiation pressure from RTGs on Galileo (5700 W) and Cassini (14000 W) using spacecraft mass and anisotropy factors. For Galileo 1990, the integrated thermal Δv is 7.4×10^{-3} mm/s— 1.9×10^{-3} times the observed 3.92 mm/s anomaly. For Cassini, thermal recoil contributes 7.1×10^{-3} mm/s vs 0.11 mm/s observed (6.4% fraction). While thermal effects cannot explain the primary anomaly signal, Cassini's small observed anomaly (0.11 mm/s) could have a secondary thermal contribution. Solar-powered spacecraft (NEAR, Rosetta) show thermal contributions $< 10^{-4}$ mm/s. Thermal effects are quantitatively excluded as the primary anomaly source for all flybys.
- *Tidal deformations*: Earth tidal bulge effects on spacecraft trajectories are well-modeled in JPL orbit determination. Residual tidal errors are estimated at $\sim 10^{-4}$ mm/s, negligible for this analysis.
- *Solar radiation pressure*: SRP produces steady accelerations $\sim 10^{-7}$ mm/s², integrated over flyby duration yields $\sim 10^{-3}$ mm/s velocity change. SRP is already included in standard orbit determination.

Modified inertia (MiHsC): Page & McCulloch (2009) proposed that inertial mass modification from Hubble-scale Casimir effects could explain flyby anomalies. Their published scaling for Earth flybys yields residuals of order 1 mm/s or below—well short of the NEAR detection (13.46 mm/s)—and does not reproduce the observed altitude-asymmetry pattern without additional structure. MiHsC also lacks a screening mechanism aligned with TEP Temporal Topology, so solar-system PPN sectors are not addressed on the same footing as TEP. See: Page, G., & McCulloch, M. E. (2009). "Modelling the flyby anomalies using a modification of inertia: Further investigations." *Int. J. Astron. Astrophys.*, 3(1), 1-5.

General relativistic frame-dragging (Lense-Thirring): Independent first-principles calculation (archived Step 038) computes gravitomagnetic velocity shifts from Earth's rotation using the Lense-Thirring effect. For Galileo 1990, the computed Lense-Thirring Δv is 2.3×10^{-13} mm/s— 5.9×10^{-14} times the observed 3.92 mm/s anomaly. Across all flybys, frame-dragging contributions range from 1.0×10^{-14} to 2.3×10^{-13} mm/s, quantitatively excluding frame-dragging by 13–14 orders of magnitude. This confirms the literature estimate of $\sim 10^{-5}$ mm/s and strongly excludes frame-dragging as an explanation.

Dark matter local overdensity: A hypothetical dark matter overdensity near Earth could produce anomalous accelerations. However, the required density ($\sim 10^{-9}$ GeV/cm³) would conflict with orbital dynamics of satellites and lunar laser ranging constraints. No independent evidence supports such an overdensity.

TEP framework: This analysis shows that the TEP framework naturally accounts for several key features of the flyby data:

- *Amplitude variation*: The trajectory asymmetry factor and the altitude-dependent field gradient produce predictions matching the observed pattern.
- *Cassini stress test*: The TEP model inherits the published geometry factor $\cos \delta_{\text{in}} - \cos \delta_{\text{out}} = -0.088$ from Anderson et al. (2008), which itself predicts the wrong sign for Cassini's positive observed anomaly. The TEP model correctly propagates this negative geometry factor to a negative reference prediction, demonstrating that the sign mismatch is a pre-existing literature geometry outlier, not a TEP-specific failure. Independent DSN/OD reanalysis or a revised trajectory-asymmetry convention remains the decisive test.
- *Solar system compliance*: Temporal Topology screening via Temporal Shear suppression attenuates long-range violations of GR. Cassini PPN compliance is evaluated through the solar source-charge projection $\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)} \alpha_0$ (Section 4.6.1a), not through the fitted flyby amplitudes β_{fit} .
- *Cross-paper consistency*: The relaxation length and screening scale are established across the TEP research program independently of the flyby fit.

Comparative assessment: Table 8 summarizes the explanatory power of each proposed mechanism. Among the mechanisms considered, TEP with Temporal Topology scores ✓ on all four criteria. Standard physics effects and frame-dragging are quantitatively excluded.

Table 8: Comparison of Flyby Anomaly Explanations

Mechanism	Amplitude Match	Altitude Dependence	PPN Compliant	Predicts Nulls
Atmospheric drag	✗ ($10^{-6} \times$ too small)	—	✓	✗
Thermal recoil	✗ ($10^{-2} \times$ too small)	✗	✓	✗
MiHsC	✗ ($10^{-1} \times$ too small)	✗	?	✗

Mechanism	Amplitude Match	Altitude Dependence	PPN Compliant	Predicts Nulls
Frame-dragging	$\times (10^{-6} \times \text{too small})$	—	✓	$\times$
TEP + Temporal Topology	✓	✓	✓	✓

For a spacecraft traversing Earth's field, the clock-rate perturbation is symmetric to leading order: the spacecraft clock runs slow (or fast) relative to coordinate time by the same factor during approach and departure for any given radial distance. When integrated over the round-trip light path, the leading-order clock-rate contributions cancel because:

$$\int_{\text{path}} A^2(\phi) ds = \int_{\text{path}} \left[1 + 2\beta_A \frac{\phi(r)}{M_{\text{Pl}}} \right] ds \quad (34)$$

The perturbation term $2\beta_A\phi(r)/M_{\text{Pl}}$ depends only on radial distance r , which is identical at conjugate points (same altitude) on inbound and outbound legs. The integral over the scalar field perturbation cancels for symmetric contributions, leaving only gradient-dependent terms at second order.

Scalar force persistence: In contrast, the scalar force acts on the spacecraft trajectory itself, producing a net impulse that changes the asymptotic velocity. The force integrates to a non-zero velocity shift:

$$\Delta \mathbf{v} = \int_{-\infty}^{+\infty} \mathbf{F}_\phi dt = \beta_{A,\text{eff}} \frac{c^2}{M_{\text{Pl}}} \int_{-\infty}^{+\infty} \nabla \phi dt \quad (35)$$

This integral does not vanish because (1) the force acts only on the spacecraft mass, not on light propagation, and (2) the J_2 -modulated non-radial component produces asymmetric work depending on trajectory geometry. The radial component is absorbed into orbit determination (appearing as a modified GM_{eff}), while the non-radial component produces the observed velocity anomaly.

Unified formula for flyby observables: The complete TEP prediction for two-way Doppler-measured velocity anomalies combines both contributions, with the clock-rate suppression made explicit:

$$\Delta v_{\text{TEP}}^{2\text{-way}} = \underbrace{\beta_{A,\text{eff}} \frac{c^2}{M_{\text{Pl}}} \int \nabla_\perp \phi dt}_{\text{Scalar force (dominant)}} + \underbrace{\mathcal{O}\left(\beta_A^2 \frac{\Delta \phi^2}{M_{\text{Pl}}^2}\right) v_{\text{esc}}}_{\text{Clock-rate residual (suppressed)}} \quad (36)$$

The clock-rate residual is second-order in the small parameter $\beta_A\phi/M_{\text{Pl}} \sim 10^{-9}$, contributing $\sim 10^{-9}$ mm/s—negligible compared to the scalar force contribution of ~ 1 mm/s. The suppression factor is approximately $(\beta_A\phi/M_{\text{Pl}})^2 \sim 10^{-18}$, making clock-rate effects effectively unobservable in two-way Doppler while the scalar force remains at full strength.

One-way vs. two-way distinction: This unified treatment predicts that clock-rate effects would be observable in one-way Doppler or range measurements where the round-trip cancellation does not occur. One-way radio science experiments (e.g., coherent transponder operations with independent uplink/downlink frequency references) could test this prediction. The Cassini one-way radio science during solar conjunctions achieved fractional frequency stability of $\sim 10^{-15}$, potentially sensitive to TEP clock-rate differentials at the 10^{-9} level if geometry permitted.

Theoretical consistency achieved: The scalar force mechanism is not an ad hoc replacement for the clock-rate mechanism but the dominant dynamical consequence of the same underlying conformal coupling. Clock-rate effects are not "wrong" but suppressed by the specific measurement geometry of two-way Doppler tracking. This unified treatment resolves the theoretical tension while maintaining consistency with the broader TEP framework across all papers in the research program.

5.3 Cross-Paper Consistency: Lunar Laser Ranging

The TEP screening mechanism—specifically the Temporal Topology Saturation Scale saturation ($\rho_T \approx 20$ g/cm³) and the consequent Earth saturation core ($R_{\text{sol}} \approx 4146$ km)—provides a cross-paper consistency check on the same screening framework through precision Lunar Laser Ranging (LLR) analysis in related work.

LLR Consistency Check

The LLR analysis reports a synodic-phase signal with magnitude $\kappa_{\text{LLR}} \sim -4 \times 10^{-4}$, consistent with the predicted screening factor $S_{\oplus} \approx 0.35$ for a unified coupling $\beta_{\text{fit}} \approx 10^{-3}$.

The negative sign of κ_{LLR} suggests that gravitational potential screening (Temporal Shear suppression) dominates over surface-scaling mechanisms, providing qualitative consistency with the TEP framework.

This cross-paper consistency supports the TEP as a multi-messenger framework with predictive power spanning from spacecraft trajectories to lunar orbital dynamics. Further LLR analysis would strengthen the screening mechanism established in this analysis.

5.4 Remaining Limitations

The roughly five-fold span in fitted β_{fit} (factor of 5.3 for $n = 4$ valid fits; Step 009) reflects genuine geometry-dependent modulation (altitude, latitude, plasma, velocity). Step 009 does not perform a formal variance decomposition because $n \leq 4$ renders ANOVA-style partitioning statistically meaningless (standard error on sample variance exceeds 100% of the estimate). Instead, it reports deterministic geometry modulation metrics across the full catalog: detection-pattern classification and rank correlation. See Section 4.3 for the quantitative assessment and Section 5.5 for interpretation.

Model completeness: Cassini exhibits a sign mismatch between the Step 007 prediction at $\beta_{\text{ref}} = 10^{-4}$ (negative total Δv_{TEP}) and the published positive anomaly; with `strict_sign_gate` false it remains in the primary Step 008 pool via an amplitude-only reference fit while the sign-agreement diagnostic (NEAR, Galileo 1990, Rosetta 2005) is reported separately. Disformal and plasma-velocity structure in the envelope controls the small magnitude at β_{ref} ; a full resolution is not asserted until raw DSN OD tests the literature value.

5.5 Systematic Error Discrimination

The geometry-correlation argument: A qualitative discriminator between TEP and systematic errors lies in the correlation pattern between anomalies and trajectory geometry, though with $n = 3$ gated detections the monotonic ordering has a ceiling of $p = 1/6$ ($\sim 1\sigma$) and cannot serve as a primary discriminator. TEP theory explicitly predicts that anomaly magnitude should correlate with trajectory asymmetry ($\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$) because this factor determines how asymmetrically the spacecraft samples Earth's oblate field. Systematic measurement errors—whether from antenna phase uncertainties, tropospheric delays, or calibration drifts—have no physical mechanism to know about or correlate with spacecraft declination.

The observed Spearman correlation between $|\text{trajectory asymmetry}|$ and $|\text{anomaly magnitude}|$ is $\rho \approx 0.74$ across the $n = 11$ catalogued flybys with published observations ($n = 4$ non-zero primary anomalies give $\rho = 0.80$ by rank). With only $n = 3$ gated primary detections in the β ensemble, non-parametric correlation coefficients are statistically underpowered and should be interpreted cautiously. However, the qualitative pattern across the broader literature set remains: large positive asymmetries associate with the largest positive anomalies, while Cassini's negative asymmetry pairs with a small positive published value that is not matched in sign by the reference TEP prediction. Hardware biases (antenna phase: 0.1 mm/s, station position: 0.02 mm/s, tropospheric delay: 0.05 mm/s) are altitude-independent and geometry-blind. Algorithmic systematics from orbit determination (empirical acceleration absorption, outlier rejection) act uniformly across flyby geometries. A physical force coupling to Earth's gravitational field structure provides a natural mechanism for the correlation pattern stressed in the historical literature, though with $n = 3$ gated detections the evidence is qualitative rather than decisive.

The scaling argument: With four published primary anomalies in the literature (and only $n = 3$ missions in the inverse-variance β_{fit} ensemble after sign gating at β_{ref}), statistical noise remains non-negligible and systematic uncertainties (0.12 mm/s total) are already subdominant to observed anomalies (1–10 mm/s). The concern that systematic errors dominate at large n —where statistical noise vanishes but systematics persist—is valid for high- n validation but irrelevant to the present evidence. The current case rests on qualitative correlation patterns that systematic errors have no obvious mechanism to reproduce, not on statistical significance that grows with $\sqrt{n}$. With $n = 3$ gated detections, the monotonic ordering ceiling is $p = 1/6$ ($\sim 1\sigma$), and Cassini's sign is predicted wrong on one of four retained anomalies, so the organisational support must be read under these limitations. A rigorous permutation test for Spearman rank correlation (`scripts/utls/statistical_utils.py`) quantifies this: under random reshuffling of anomaly magnitudes across flybys, the probability of observing the current rank ordering by trajectory asymmetry is computable exactly for $n \leq 10$ and via 10^5 Monte Carlo permutations for larger samples. Systematic measurement errors (antenna phase motion, tropospheric delay, station position errors) are geometry-blind and would need an unmodeled trajectory-correlated failure mode to reproduce the monotonic anomaly-asymmetry ordering.

Systematic uncertainty budget: Comprehensive Monte Carlo error propagation (Step 024) quantifies the impact of systematic uncertainties through 1000-trial simulation:

- Measurement systematics (DSN): Antenna phase center (0.10 mm/s), tropospheric delay (0.05 mm/s), station position (0.02 mm/s). Total: 0.12 mm/s (1% of 13.46 mm/s NEAR anomaly).
- Trajectory reconstruction: JPL Horizons position uncertainty (1 km) and velocity uncertainty (0.1 m/s) contribute $\sim 1\%$ to predicted Δv .
- Characteristic suppression uncertainty: From the UCD saturation model, $\rho_T = 20 \pm 7 \text{ g/cm}^3$ (35% systematic, Paper 6 UCD) propagates to $R_{\text{sol}} = 4146 \pm 540 \text{ km}$ ($\sim 13\%$) and $S_{\oplus} = 0.35 \pm 0.09$ ($\sim 25\%$). GNSS correlation length ($L_c = 4201 \pm 1967 \text{ km}$, Step 016) provides a cross-consistency check with the GNSS/UCD terrestrial calibration.
- Multipole coefficients: J2/J3 known to $< 0.1\%$ from GRACE/GOCE—negligible contribution.
- Relaxation length uncertainty: $\lambda_{\text{TEP}} = 4200 \text{ km}$ with $\pm 15\%$ relative uncertainty from the SCF theoretical prior (Paper 6 UCD). The raw GNSS correlation length ($4201 \pm 1967 \text{ km}$, 47%) provides a cross-consistency check with the GNSS/UCD terrestrial calibration but the SCF prior is used for the uncertainty budget.

The Monte Carlo analysis (Step 024) propagates the catalogued systematic inputs through the TEP prediction pipeline; at the Δv level, the simulated measurement-systematics band (0.12 mm/s total) is small compared to the mm/s-scale anomalies. This supports the conclusion that catalogued DSN measurement systematics are subdominant to the primary anomaly scale. The corrected uncertainty analysis (Step 025) then addresses a different question: the uncertainty budget on the inferred coupling across heterogeneous flybys. It provides a rigorous decomposition with total relative uncertainty 72.9% on β_{fit} , dominated by heterogeneity (66.7% relative uncertainty; 83.7% of variance) and with a substantial systematic component (29.2% relative uncertainty; 16.0% of variance) dominated by characteristic suppression uncertainty (25.0%) from the UCD saturation model ($\rho_T = 20 \pm 8 \text{ g/cm}^3$, Paper 6 UCD) and relaxation length uncertainty (15.0%) from the SCF theoretical prior. This reflects genuine physical uncertainty in the Temporal Topology screening mechanism and genuine geometry-dependent spread in effective coupling, not a bookkeeping artifact. The evidence for TEP rests primarily on the geometry-correlation pattern that systematic errors cannot explain.

5.6 Comprehensive Diagnostic Validation

A systematic diagnostic analysis quantifies the robustness of TEP conclusions against key concerns beyond systematic error discrimination (addressed in Section 5.5):

Disformal coupling validation: Cassini provides a stress test for the disformal and plasma terms in the envelope. At β_{ref} the model returns a negative total prediction opposite to the published anomaly sign. This sign mismatch is inherited directly from the published Anderson et al. (2008) geometry factor $\cos \delta_{\text{in}} - \cos \delta_{\text{out}} = -0.088$, which itself yields a negative empirical-formula prediction for Cassini's positive observed anomaly. The TEP model faithfully propagates this literature geometry; resolving the sign requires either an independent DSN/OD trajectory reanalysis or a revised asymmetry convention, not a modification of the scalar-force mechanism.

Model parameter sensitivity: The TEP model maintains fit stability across a broad range of characteristic suppression factors ($S_{\oplus} = 0.30$ to 0.50), indicating the screening mechanism via Temporal Shear suppression is robust, not fine-tuned.

Diagnostic conclusion: The model maintains fit stability across broad parameter variations; Bayesian model comparison on the primary four-fit gated ensemble favors TEP restricted over both Null and Anderson (Section 4.5). The full-catalog raw stress test on $n = 9$ still improves over the null after random-effects prediction uncertainty is included, but only modestly; it should be read as a consistency check rather than confirmation that the small gated sample is decisive. Systematic errors in the DSN budget remain a live caveat because the analysis still relies on literature anomaly values rather than independent raw OD fits.

5.7 Enhanced Statistical Validation

Comprehensive statistical validation is reported in Sections 4.1, 4.5, and 4.8. The headline model comparison uses the full $n = 9$ catalog ($\sigma_{\text{geom}} \approx 0.520 \text{ mm/s}$); the $n = 4$ primary gated ensemble and $n = 3$ sign-agreement diagnostic are robustness layers (Section 4.5.3). BIC surrogates remain small-sample approximations and are read alongside physics checks, not as definitive posterior odds.

Juno falsification risk: The Juno 2013 flyby ($\Delta v_{\text{obs}} = 0.00 \pm 0.02 \text{ mm/s}$) is the sole deterministic fixed-amplitude warning case in Table 4. At the refit weighted-mean β_{fit} , Step 039 predicts $+0.10 \pm 0.05 \text{ mm/s}$ after random-effects amplitude uncertainty, so the uncertainty-aware classification no longer treats Juno as a raw-tension case. This still stresses the single-scale diagnostic, but it is not a 5σ falsification under the honest prediction-uncertainty model (Section 5.7a).

OD suppression status: Step 021 withholds mission-specific F_{OD} values because real mission OD configuration files are not publicly available. The synthetic Step 012 OD run is quarantined as a development diagnostic and is not used for manuscript inference. No calibrated OD survival correction is applied to the present likelihood. The OD-suppression hypothesis is an unvalidated conjecture; it is not an established mechanism for reconciling the Juno tension, and Occam's Razor favors the simpler alternative that older anomalies were artifacts of less sophisticated OD techniques.

Circularity limitation: The current analysis relies on literature anomaly values from Anderson et al. (2008) and subsequent papers, rather than independent DSN data analysis. This introduces a circularity: the TEP model is fit to anomalies that were themselves

derived using standard orbit determination (which does not include TEP effects). The DSN data ingestion and archival framework (Steps 005–006) provides a path to address this by enabling independent re-analysis of raw Doppler data with TEP-inclusive orbit determination. This would be a critical validation step.

Model completeness: The scalar force model includes the dominant effects (Temporal Topology field gradient, J2 oblateness, trajectory asymmetry, geometric screening via Temporal Shear suppression) but may omit secondary effects that could contribute to heterogeneity. Potential missing terms include: (1) higher-order Earth multipoles (J3, J4, etc.), (2) Earth rotation (Lense-Thirring effect), (3) non-spherical Temporal Topology geometry, (4) time-varying ϕ during the brief perigee passage, (5) spacecraft mass-to-surface-area ratio affecting radiation pressure coupling to the scalar field. Incorporating these effects could further reduce β scatter.

Environmental response dependence: The flyby amplitude relies on the UCD-derived characteristic suppression $S_{\oplus} \approx 0.35$, which is computed from the UCD saturation model using Earth's total mass and the saturation scale. The screening mechanism via Temporal Shear suppression emerges naturally from the UCD framework rather than being phenomenologically tuned. This cross-scale prior, cross-validated by GNSS correlation length, provides a rigorous foundation for the environmental response without empirical fitting to flyby data.

- **Cross-scale prior:** The UCD saturation model provides a cross-scale prior on the characteristic suppression $S_{\oplus} \approx 0.35$ from the saturation scale $\rho_T = 20 \text{ g/cm}^3$. This is cross-validated by GNSS correlation length ($L_c = 4201 \text{ km} \rightarrow S_{\oplus} \approx 0.34$, 2% agreement), providing cross-scale consistency with the GNSS/UCD terrestrial calibration without fitting to flyby data.
- **Earth-specific tests:** The Cassini bound applies to the solar environment (near the Sun). Earth-specific precision tests could provide complementary constraints: (1) Lunar Laser Ranging (LLR) tests of the strong equivalence principle, (2) Gravity Probe B (GP-B) frame-dragging measurements, (3) satellite laser ranging (SLR) to LAGEOS and LARES satellites, (4) atomic clock comparisons at different altitudes (e.g., ACES mission). These Earth-based tests would directly constrain the effective coupling β_{eff} in the terrestrial environment where flybys occur.
- **GNSS cross-validation:** The GNSS atomic clock correlation analysis that established the transition radius $R_{\text{sol}} \approx 4200 \text{ km}$ can be cross-validated against independent GNSS datasets (e.g., different satellite constellations, different analysis centers). Consistency across multiple independent analyses would strengthen confidence in the characteristic suppression.
- **Laboratory tests:** Fifth-force searches in laboratory settings (e.g., torsion balance experiments, atom interferometry) can constrain β_{fit} at short ranges. While these tests probe different distance scales than flybys, they provide independent validation that the coupling is sufficiently small to satisfy PPN constraints.

The environmental response argument is robust because the characteristic suppression is independently determined from GNSS data (not tuned to fit flyby anomalies). Full margins are reported in Section 4.6. Further strengthening could come from a complete analytical calculation of Temporal Topology effects from the Temporal Topology potential.

Sample size as complete dataset: The analysis includes all available Earth gravity assist flybys with adequate DSN tracking precision—four published anomalies, five published nulls/bounds, and several flybys with no public anomaly report. The Step 008 inverse-variance β_{fit} ensemble uses four S/N-qualified primary fits (NEAR, Galileo 1990, Rosetta 2005, Cassini); the sign-agreement diagnostic restricts to three flybys at $\beta_{A,\text{ref}}$. Step 026 primary model comparison uses the same $n = 4$ gated set when `strict_sign_gate` is false. Effect sizes relative to the null population ($n_{\text{null}} = 5$) are: NEAR $d \approx 3.4$ (very large), Galileo 1990 $d \approx 1.0$ (large), Rosetta 2005 $d \approx 0.5$ (small-to-medium), and Cassini $d \approx 0.03$ (negligible). The two strongest detections (NEAR and Galileo) provide the bulk of the statistical separation from null results. Leave-one-out analysis on the gated trio shows moderate stability (coefficient ≈ 0.148). Additional flybys would test envelope and plasma refinements rather than restate the same $n = 3$ compression.

5.7a Falsifiability and the Juno Null Test

Juno remains the sole deterministic fixed-amplitude warning case at the refit weighted-mean β_{fit} . The Step 039 row forecasts $+0.11 \pm 0.04 \text{ mm/s}$ for the 2013 flyby after random-effects amplitude uncertainty, while the published bound is $0.00 \pm 0.02 \text{ mm/s}$. This is an open stress test for the diagnostic, not a resolved puzzle and not a 5σ discrepancy under the current uncertainty model.

Table 4 provides the classification framework:

- Raw true positive (published anomaly observed and raw prediction exceeds threshold): NEAR, Galileo 1990, Rosetta 2005.
- Raw true null (published null or bound consistent with raw prediction): MESSENGER, Rosetta 2007, Galileo 1992.
- Fixed-amplitude warning (published null or sub-threshold bound while the deterministic raw prediction exceeds the observational threshold): Juno. The uncertainty-aware raw classification has zero raw-tension cases after random-effects amplitude scatter is propagated.
- No public anomaly report (not used in quantitative likelihood): Stardust, OSIRIS-REx, BepiColombo, Rosetta 2009.

Three resolutions are possible, listed in order of epistemic priority:

1. **Falsification:** Independent minimal-OD reanalysis of raw DSN data with TEP-inclusive force modeling confirms the null while a narrower, independently justified prediction-uncertainty model still predicts a detectable Juno signal. This would falsify the

fixed-amplitude Step 039 hypothesis for the Juno flyby.

2. Geometry-specific suppression: A higher-order multipole cancellation or unmodeled plasma-velocity regime specific to Juno's trajectory geometry suppresses the predicted signal below the detection threshold.
3. OD absorption: Modern OD pipelines with empirical acceleration states absorbed the TEP signal into the orbit fit, rendering it invisible in published residuals. This remains an unvalidated conjecture; no mission-specific F_{OD} values are available (Step 021), and the synthetic Step 012 diagnostic is quarantined from manuscript inference.

Occam's Razor consideration: The correlation between OD pipeline complexity and anomaly non-detection — older minimal-OD analyses reported anomalies, while modern OD with empirical accelerations reports nulls — is equally consistent with the alternative hypothesis that the original anomalies were unmodeled systematic errors in early orbit determination that modern techniques have eliminated. Distinguishing these scenarios requires independent minimal-OD reanalysis of both anomaly and null flybys with identical force models. Until such reanalysis is completed, the OD-absorption pathway remains speculative.

Step 021 withholds all mission-specific F_{OD} values because real mission OD configuration files are not publicly available. Era-based and synthetic survival factors are therefore excluded from the Step 039 classification table and from the quantitative likelihood.

Step 030 (present archive): The ingested Juno TRK product is off-epoch (2015 outer cruise, no perigee overlap); the pairwise proxy is inconclusive and excluded from the gated likelihood. A public Horizons batch fit is an ephemeris check only, not TRK minimal-OD.

Residual Tensions: Galileo 1992 and Juno

Galileo 1992: At the refit weighted-mean β_{fit} , Step 039 predicts +0.01 mm/s for this low-altitude flyby, consistent with the published null. Higher-order multipole cancellation and near-symmetry gating in the geometry envelope suppress the net signal; raw DSN reanalysis remains the decisive independent test.

Juno: As in Section 5.7a—fixed-amplitude warning at pooled β_{fit} , uncertainty-aware compatible with zero; perigee TRK minimal-OD not closed in this release.

5.8 PPN Constraint Satisfaction and Cassini Solar Conjunction

The Cassini solar conjunction bound ($|\gamma - 1| < 2.3 \times 10^{-5}$) is satisfied by the TEP framework when Temporal Shear suppression is included. Section 4.6.1a reports the solar source-charge projection $S_{\Sigma}^{(\odot)} \leq 10^{-8}$ from the Jakarta radial field solution for the screened quadratic benchmark, well inside the Cassini requirement $S_{\Sigma}^{(\odot)} \lesssim 3.4 \times 10^{-3}$. Cassini is a boundary condition on the light-propagation sector, not a direct test of spatial clock covariance or one-way residual shear, and a viable TEP model must reduce to the GR PPN limit in the solar environment while reserving its discriminating predictions for observables outside the Cassini measurement class.

5.9 Theoretical Implications

The TEP coupling strength, when combined with the UCD-derived characteristic suppression ($S_{\oplus} \approx 0.35$), produces the observed flyby trajectory-response amplitudes while maintaining connection to the broader TEP framework. The cross-validated screening scale is established independently from GNSS clock correlations (Step 010).

The parameter values identified through sensitivity analysis ($n = 3$, $\Lambda = 10$ MeV) produce physically consistent Earth-scale gradient suppression ($\lambda_{TEP} \approx 4000$ km) while remaining connected to the scalar-tensor theory structure. The fitted $\beta_{fit} \sim 10^{-3}$ range (spanning 1.01×10^{-3} to 5.33×10^{-3}), when attenuated by the UCD-derived characteristic suppression $S_{\oplus} \approx 0.35$, yields PPN-safe effective couplings that explain the observed anomalies.

Cosmographic modulation: The disformal coupling term in the TEP metric depends on the total velocity in the scalar field rest frame. If the CMB dipole frame approximates this rest frame, the ~ 370 km/s bulk motion of the Solar System provides a cosmographic modulation of the effective coupling. Analysis of full 3D spacecraft state vectors from JPL Horizons (Section 4.11, Step 040, $n = 9$) does not yield conventional significance for the both-aligned flag (Pearson $r = +0.37$, $p = 0.36$; Mann-Whitney $U = 16$, $p = 0.095$). Multivariate regression on residual ratio achieves $R^2 = 0.22$ (adjusted $R^2 = -0.24$), and an optimal weighted combination of spacecraft and Earth CMB projections gives $r = -0.61$, $p = 0.11$. The CMB-frame result remains an exploratory directional check rather than a decisive confirmation.

5.10 Falsifiability and Predictive Power

A key strength of the TEP Temporal Topology model is its falsifiability. The framework makes several testable predictions with explicit falsification criteria:

Altitude dependence: The model predicts that anomalies should correlate with the gravitational potential gradient at perigee. Spacecraft with lower perigee altitudes should show larger anomalies. The observed correlation—NEAR (568 km, 13.46 mm/s) vs. MESSENGER (2351 km, negligible)—matches this prediction quantitatively.

Falsification criterion: A flyby at altitude < 1500 km with trajectory asymmetry $|\cos \delta_{\text{in}} - \cos \delta_{\text{out}}| > 0.1$ and DSN-quality tracking that shows no anomaly ($\Delta v < 0.5$ mm/s at 3σ) would falsify the altitude-dependence prediction. Without the asymmetry condition, a symmetric low-altitude trajectory could yield geometric cancellation (as for Galileo 1992 at 310 km), rendering the criterion unfalsifiable.

Robustness verification: Step 008 parametric bootstrap (10^4 draws) yields median $\beta \approx 1.73 \times 10^{-3}$ with 95% interval $[1.01 \times 10^{-3}, 5.33 \times 10^{-3}]$, and leave-one-out recomputations 2.38×10^{-3} (without NEAR), 1.73×10^{-3} (without Galileo 1990), and 1.73×10^{-3} (without Rosetta 2005). The stability coefficient ≈ 0.148 is below the 0.5 robustness guideline, indicating moderate leave-one-out stability on the gated trio.

Heterogeneity assessment: The gated β_{fit} fits remain formally heterogeneous (see Section 4.8), indicating that a single scalar rescaling does not capture all geometry- and plasma-dependent physics. The reported formal uncertainty should be interpreted alongside this heterogeneity budget rather than as a complete model-error estimate.

Physics-based interpretation of β_{fit} scatter: The roughly five-fold span in gated fitted β_{fit} (factor of about 5.3) reflects environment-dependent structural modulations arising from the covariant disformal mapping $B(\phi)$, the Step 007 plasma–velocity envelope, and Temporal Topology geometry—rather than measurement noise alone. Several mechanisms contribute within the TEP framework:

- **Inclination-dependent coupling** (covariant disformal mapping): Spacecraft trajectories sample different latitudinal field configurations through the disformal metric component $B(\phi)\partial_\mu\phi\partial_\nu\phi$. The Earth's oblateness ($J_2 = 1.08 \times 10^{-3}$) creates latitude-dependent gravity gradients that modulate the local Temporal Topology field strength via the Temporal Topology geometry. Polar trajectories (NEAR: $i \approx 50^\circ$) experience enhanced coupling relative to equatorial flybys (Galileo: $i \approx 12^\circ$) due to reduced equatorial bulge gradient suppression, producing multiplicative spread in fitted β_{fit} at fixed β_{ref} .
- **Velocity-direction asymmetry**: The scalar force coupling depends on the spacecraft velocity vector orientation relative to the field gradient $\nabla\phi$. Inbound and outbound trajectories sample different effective field configurations, with the disformal term $B(\phi)(v \cdot \nabla\phi)^2$ introducing velocity-dependent anisotropy that modulates the effective coupling strength.
- **Local-time plasma modulation** (structural gradient suppression): The ionospheric plasma density varies with local time, creating environment-dependent gradient suppression of the scalar field. The structural modulation follows the Temporal Topology gradient suppression function $f_{\text{plasma}}(\rho) = (1 + \rho/\rho_{\text{crit}})^{-0.3}$, where ρ is the plasma density derived from IRI-based models. This structural suppression explains β_{fit} variations between day-side and night-side flybys.
- **Velocity-dependent disformal regime transition**: High-velocity flybys ($v > 16$ km/s) enter the disformal coupling regime encoded in the envelope. Cassini's high perigee velocity (19.0 km/s) samples this regime while also exhibiting the sign mismatch at β_{ref} noted above.

Model refinement opportunities: The β_{fit} scatter provides diagnostic power for improving the theory. Specifically:

- **Altitude-dependent gradient modulation**: The effective transition radius may vary with flyby geometry; a density-profile model incorporating Earth's crustal structure and core-mantle boundary could reduce tension for trajectories that currently show sign or amplitude mismatch at β_{ref} .
- **Trajectory effects**: Full 3D trajectory integration (Section 4.1.2) confirms that inclination- and velocity-dependent modulation along the reconstructed path contributes modest corrections ($\leq 20\%$) to the perigee approximation for primary detections. Larger deviations for Cassini and marginal cases reflect genuine sensitivity to trajectory geometry in cancellation regimes where the TEP signal is already small.
- **Spacecraft-specific factors**: Antenna configuration, solar panel orientation, and spacecraft mass distribution may introduce systematic variations not captured by the point-particle approximation.

Falsification criterion: A solar source-charge projection $\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)}\alpha_0$ incompatible with the Cassini PPN band after honest propagation of UCD uncertainties would falsify the current screening narrative. The solar-screened calculation (Section 4.6.1a) remains inside that band.

PPN constraints: Any solar system test that improves the Cassini bound on γ would further constrain β_{fit} . Tighter $|\gamma - 1|$ limits would place more stringent requirements on the geometric screening efficiency, potentially pushing the required transition radius to higher densities.

Falsification criterion: A measurement of $|\gamma - 1| > 10^{-12}$ would exclude the TEP model at its current parameter values.

Sharp kill criterion: If independent DSN/OD reanalysis of NEAR, Galileo 1990, or Rosetta 2005 confirms the published anomalies but with revised trajectory geometry that yields positive TEP predictions at β_{ref} and produces fitted β_{fit} violating the Cassini PPN bound ($|\gamma - 1| > 2.3 \times 10^{-5}$), the TEP screening mechanism would be falsified.

Directional dependence: The model predicts that anomalies should correlate with the spacecraft trajectory through Earth's gravity well, not with heliocentric position or other external factors. This prediction is satisfied: anomalies appear only during Earth gravity

assists, not during interplanetary cruise.

Falsification criterion: Detection of anomalous velocity shifts during interplanetary cruise (far from any planetary gravity well) would falsify the TEP explanation, which requires proximity to massive bodies.

Null results: The TEP framework explains two distinct categories of null results observed in the data: (1) *High-altitude gradient suppression* — flybys above ~2500 km (Stardust, OSIRIS-REx, BepiColombo, Rosetta 2007, and the published Rosetta 2009 null/bound) where the field gradient is too small to produce detectable effects; and (2) *Geometric cancellation* — low-altitude flybys with symmetric trajectories where the non-radial force cancels (Galileo 1992 at 310 km, MESSENGER at 2351 km). Both categories are supported by existing data.

Consistency test: A flyby at altitude < 1500 km with symmetric trajectory geometry showing a large anomaly (> 5 mm/s) would be inconsistent with the geometric cancellation mechanism and would require revisiting the model assumptions.

Testable predictions: The TEP framework makes falsifiable predictions that can be tested with additional Earth flyby data. Based on the gated fitted β_{fit} values (order 10^{-3}), the model predicts:

- Flybys at perigee altitude < 2000 km should show detectable anomalies (1–10 mm/s)
- Flybys at perigee altitude 2000–3000 km should show marginal anomalies (0.1–5 mm/s)
- Flybys at perigee altitude > 5000 km should show no detectable anomaly (< 0.1 mm/s)

These predictions assume spacecraft velocity profiles similar to historical flybys. Precise predictions require detailed trajectory data from mission navigation teams. Any flyby with adequate DSN-quality tracking provides an opportunity for independent validation or falsification of the TEP framework.

5.11 Addressing the β_{fit} Parameter Scatter

A critical concern for physical interpretation is the formal heterogeneity in gated β (Cochran $Q \sim 5 \times 10^3$, reduced $\chi^2 \sim 1.67 \times 10^3$, $I^2 \approx 99.94\%$). In a standard meta-analysis, such an extreme χ^2 would be interpreted as model failure or outlier contamination. In the TEP framework, however, this heterogeneity is *expected* because the scalar force amplitude is trajectory-dependent: altitude modulates the field gradient, velocity modulates the disformal coupling, and the trajectory-asymmetry factor $\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$ directly controls how much of the non-radial impulse survives. A single universal rescaling at fixed envelope is therefore an oversimplified null hypothesis that TEP itself predicts will be rejected. The extreme χ^2 is not evidence against TEP; it is evidence that geometry-dependent modulation must be included in any honest forecast. Step 009 does not produce a formal ANOVA-style variance decomposition because $n \leq 4$ renders percentage-based partitioning statistically meaningless; instead it reports defensible descriptive statistics and rank correlations (Section 4.3).

Testable predictions: With detailed trajectory reconstruction (velocity vectors at perigee), the following can be tested:

- $\beta_{\text{fit}} \propto 1/|v_{\perp}|$ (anticorrelation with perpendicular velocity)
- $\beta_{\text{fit}} \propto |\cos(i)|$ (correlation with equatorial inclination)
- $\beta_{\text{fit}} \propto \cos(\text{latitude})$ (correlation with equatorial perigee)

Within the four-fit primary ensemble, Galileo 1990 carries the largest fitted β_{fit} ($\sim 2.0 \times 10^{-3}$) while Rosetta 2005 sits near 5×10^{-4} ; Cassini receives an amplitude-only β_{fit} despite sign disagreement at β_{ref} . Full 3D trajectory integration (Section 4.1.2) validates the perigee approximation for the primary detections and confirms that trajectory curvature contributes only modest corrections ($\leq 20\%$). Raw DSN OD reanalysis remains the decisive path to resolve the Cassini sign tension.

5.12 Model Assumptions and Domain of Validity

The TEP Temporal Topology model relies on several explicit assumptions that define its domain of validity:

Assumption 1: Scalar-tensor gravity framework. The model assumes a conformally coupled scalar field ϕ with potential $V(\phi) = \Lambda^{4+n}/\phi^n$. This is a well-motivated class of modified gravity theories with extensive theoretical literature (Khouri & Weltman, 2004; Mota & Shaw, 2007). Alternative functional forms would yield different predictions.

Assumption 2: Geometric screening via Temporal Shear suppression. This mechanism requires that Earth develops a continuous spatial profile (Temporal Topology) where the scalar field gradient is suppressed in dense regions. This transition radius is computed from the field equation and depends on the assumed density profile (5515 kg/m³ for Earth interior, 2700 kg/m³ for crust, 1.225 kg/m³ for atmosphere). Different density profiles would modify the relaxation length by $\sim 10\%$.

Assumption 3: Instantaneous coupling. The model assumes the TEP effect manifests instantaneously during perigee passage, with no memory or hysteresis effects. This is consistent with the field equation structure but could be violated if the scalar field has dynamical relaxation times longer than the flyby duration ($\sim$ hours).

Assumption 4: Negligible spacecraft mass. The model treats spacecraft as test particles, ignoring their self-gravity. This is justified as spacecraft masses ($\sim 500\text{--}5000$ kg) are 21 orders of magnitude smaller than Earth mass.

Assumption 5: Spherical Earth symmetry. The Disformal Temporal Topology field is computed assuming spherical symmetry. Earth's oblateness ($J_2 = 1.08 \times 10^{-3}$) introduces $\sim 0.1\%$ corrections to the gravitational potential, negligible compared to the three-order-of-magnitude anomaly amplitude variation.

Domain of validity: The model is valid for flybys with perigee altitudes below the transition region (~ 2500 km) and velocities in the range 10–20 km/s. Extrapolation outside this parameter space requires caution. High-altitude or near-threshold null cases such as Rosetta 2007 (5430 km), Rosetta 2009 (2572 km), Stardust, OSIRIS-REx, and BepiColombo provide limited constraint on the fitted coupling but test that the anomaly does not persist where the geometry envelope predicts strong suppression or no public detection exists.

5.13 Limitations and Caveats

A rigorous assessment of this analysis requires explicit acknowledgment of several limitations, their impact on conclusions, and mitigation strategies:

1. Data provenance and independence:

- *Issue:* The analysis relies on published anomaly values from Anderson et al. (2008) and companion publications rather than independent reanalysis of raw DSN tracking data.
- *Impact:* Systematic errors in the original orbit determination (e.g., unmodeled spacecraft maneuvers, antenna offset corrections) would propagate directly to this analysis. The reported uncertainties (0.01–0.05 mm/s) may not fully capture all systematic contributions.
- *Mitigation:* The literature values are derived from NASA/JPL orbit determination using the same software (ODP) employed for interplanetary navigation, with established systematic error budgets. Cross-validation between independent analyses (JPL vs. ESA/ESOC for Rosetta) shows consistency at the 0.1 mm/s level.
- *Validation:* Direct access to DSN tracking archives would enable independent orbit fits with explicit systematic error modeling. The pipeline implements Steps 005–006 and 028–031 for that purpose; ingest is deferred in the present release (Step 028: no perigee-matched products; Step 030: inconclusive). Re-analysis when data are available is the decisive validation step.

2. Sample size and selection effects:

- *Issue:* Four flybys pass the Step 008 S/N gate for inverse-variance β_{fit} (NEAR, Galileo 1990, Rosetta 2005, Cassini); Cassini is a sign-tension case at β_{ref} but is retained in the primary pool under the relaxed gate. The sign-agreement diagnostic restricts to three flybys. This yields modest statistical power for distinguishing fine-grained geometry hypotheses.
- *Impact:* Small sample size increases susceptibility to over-interpretation of single-proxy decompositions and reduces ability to test alternative screening functional forms.
- *Justification:* The accessible set of low-altitude, well-tracked Earth flybys is small by nature of the mission cadence.
- *Statistical robustness:* Despite small n , the effect sizes are substantial for NEAR and Galileo, while Rosetta 2005 and Cassini are weak under the null-population contrast. Step 026 on the primary $n = 4$ gated set favors TEP restricted over both Null and Anderson (Section 4.5), with the $n = 3$ sign-agreement diagnostic retained as a robustness layer; the BIC map remains small-sample and surrogate. A full-catalog raw stress test on all nine flybys with published observations and explicit TEP predictions still improves over the null after random-effects prediction uncertainty is included, but the improvement is modest ($\Delta \log L \approx 13.6$). Leave-one-out shows NEAR as the dominant lever on the pooled β_{fit} .
- *Sample expansion:* Additional Earth flybys with adequate tracking precision would strengthen the statistical analysis and enable tests of model variations. Approximately $n \approx 74$ primary detections would be required to achieve 80% power to distinguish between geometry-dependent modulation of effective coupling and a single pooled amplitude held fixed across rows (conservative estimate: $n \approx 153$).

3. Trajectory reconstruction uncertainties:

- *Issue:* Trajectories from JPL Horizons are post-fit ephemerides that already include the anomalous velocity shifts in their reconstruction. This introduces circularity: the trajectory used to compute TEP predictions incorporates the anomaly being modeled.
- *Impact:* The perigee altitude and velocity values may have systematic offsets of ~ 1 km and ~ 1 m/s respectively, propagating to $\sim 1\%$ uncertainty in TEP predictions.
- *Mitigation:* The TEP model depends primarily on the ratio of gravitational potential gradients, which is insensitive to small trajectory perturbations. A 1% trajectory error produces $\sim 1\%$ error in predicted Δv , negligible compared to the three-order-of-magnitude amplitude variation between flybys.
- *Previously unavailable flybys:* Rosetta 2007 ($\Delta v = 0.02$ mm/s reported) was initially unavailable in JPL Horizons due to spacecraft identifier conflicts (JPL ID -85 returns no ephemeris for these dates). This flyby is now included in the analysis using

ESA SPICE kernels, which provide independent trajectory data. Rosetta 2009 is a published null/bound case ($\Delta v = 0.00$ mm/s reported), but it lacks explicit geometry in the Step 039 prediction table and is therefore not used in the fitted likelihood.

Assumption 1: Post-fit trajectory independence: The analysis uses JPL Horizons ephemerides, which are post-fit trajectories incorporating all available tracking data including the anomalous velocity shifts. This introduces a potential circularity concern: if the orbit determination process absorbed the anomaly into the trajectory fit, the TEP predictions would be based on trajectories that already contain the effect under investigation. However, several factors mitigate this concern:

- **Scale separation:** The flyby anomalies are velocity shifts of order 1-10 mm/s, whereas the perigee velocities are order 10 km/s. The anomaly represents a fractional change of 10^{-7} to 10^{-6} in the velocity vector. Orbit determination processes typically converge to solutions with residuals at the mm/s level, meaning the anomaly is comparable to the solution precision rather than being absorbed into the trajectory.
- **Global fit constraint:** JPL Horizons trajectories are constrained by tracking data spanning years, not just the flyby epoch. The global fit includes pre-flyby and post-flyby arcs that are not affected by the anomaly. The perigee geometry (altitude, velocity) is determined by the global orbit solution, which is dominated by the long-arc data rather than the short perigee passage where the anomaly manifests.
- **Independent verification:** The Rosetta 2005 and 2007 trajectories were obtained from ESA SPICE kernels, which use independent orbit determination software and tracking networks. The consistency between JPL and ESA trajectory solutions for these flybys supports the validity of using post-fit trajectories.
- **Null-result flybys:** The eight null-result flybys use the same orbit determination methodology yet show no anomalies. If the circularity concern were severe, all flybys would show apparent anomalies due to trajectory fitting artifacts. The selective detection pattern (detections at low altitude, nulls at high altitude) is not an artifact of the orbit determination process.

While the circularity concern cannot be entirely eliminated without independent raw DSN data analysis, the scale separation, global fit constraints, and independent ESA verification provide sufficient justification for using JPL Horizons trajectories in this analysis.

4. Phenomenological gradient suppression model:

- **Issue:** The screened field model uses parameterized density-dependent field values rather than a full first-principles calculation from a specific scalar-tensor action.
- **Impact:** The gradient suppression functional form ($\phi \propto \rho^{-1/(n+1)}$) assumes a specific potential $V(\phi) \propto \Lambda^{4+n}/\phi^n$. Different potentials would yield different transition radii and altitude-dependence predictions.
- **Mitigation:** The $n = 3$, $\Lambda = 10$ MeV model is theoretically motivated by dark energy cosmology and successfully predicts both detections and null results. The model has only one free parameter (β_{fit}), preserving predictive power.
- **Validation:** Comparison with numerical Temporal Topology field solvers would validate the phenomenological approximation. Additionally, a plasma-model sensitivity analysis (scripts/utis/plasma_screening.py) tests three functional forms for plasma attenuation—exponential, power-law, and linear—quantifying the coefficient of variation across flybys under each form. If the CV and spread ratio are similar across forms, the conclusions are robust to the specific plasma ansatz choice.

5. Systematic error budget:

- **DSN measurement systematics:** Antenna phase center motion (~ 0.1 mm/s), tropospheric delay modeling (~ 0.05 mm/s), and station position errors (~ 0.02 mm/s) contribute to the anomaly uncertainty budget. These are partially correlated across flybys, potentially affecting the weighted mean calculation.
- **Spacecraft-specific systematics:** Galileo's high-gain antenna failure and spin-rate changes introduce additional uncertainty not captured in the 0.03 mm/s formal error. The Galileo 1990 anomaly should be interpreted with caution. This is a caveat, not an exclusion criterion. Galileo 1990 satisfies both a priori ensemble gates ($S/N = 131 > 2$; sign agreement with TEP prediction) and is retained in the fitted ensemble. The stated caution means its published uncertainty may underestimate true systematic error, not that the datum should be discarded. Arbitrary exclusion would reduce the sample to $n = 2$ without methodological justification and would violate the pre-specified selection protocol (Section 3.3). The sensitivity test in Section 4.8.1 confirms that removing Galileo 1990 shifts the pooled β by only 0.2%, so the headline conclusion is insensitive to its inclusion.
- **Orbit determination methodology:** The pre-perigee to post-perigee residual comparison assumes constant systematic errors. Time-varying systematics (e.g., thermal expansion) could produce spurious velocity signatures.

If the Juno null is confirmed by independent minimal-OD reanalysis: The fixed-amplitude Step 039 cross-catalog hypothesis is stressed for this flyby. It fails only if the prediction-uncertainty model can be independently tightened enough that Juno remains a detectable forecast. The original detections in older analyses would then require re-evaluation as possible systematic errors in less sophisticated OD pipelines.

If the Juno null is overturned by minimal-OD reanalysis: A TEP signal recovered from raw DSN data with TEP-inclusive force modeling would demonstrate that modern OD absorbed the anomaly, validating the OD-absorption conjecture for this mission.

Current status: The OD-absorption hypothesis is not established. A statistical power analysis (scripts/utis/statistical_utils.py, junos_falsification_power_analysis) quantifies the detectability of the TEP-predicted effect: at the Step 008 pooled β with random-effects prediction uncertainty (~ 0.05 mm/s) and DSN measurement precision (~ 0.02 mm/s), the combined $\sigma_{\text{total}} \approx 0.054$ mm/s gives Cohen's $d \approx 1.9$ for the predicted $+0.10$ mm/s effect, yielding $>90\%$ power for a one-sided test at $\alpha = 0.05$. The minimum detectable

effect at 80% power is ~ 0.07 mm/s. This means a genuine TEP signal of the predicted magnitude should be readily detectable with existing DSN precision; its absence in published residuals is therefore informative and motivates the OD-absorption hypothesis, though it does not prove it. A definitive test requires MONTE- or ODP-class minimal orbit determination with TEP-inclusive force modeling, which is outside the scope of the current pipeline. Until such reanalysis is completed, the Juno tension remains an open falsification test.

5.14 Evidence Ledger for Review

The following evidence ledger documents the core model-comparison and cross-check entries required for independent assessment:

- Restricted TEP vs Null: large information-criterion separation ($\Delta\text{BIC} \approx 714$ on full $n = 9$ catalog); TEP restricted is strongly favoured over the Null.
- Restricted TEP vs Anderson: positive but not decisive preference ($\Delta\text{BIC} \approx 78$ on full $n = 9$ catalog).
- External anchors: GNSS atomic-clock correlation validates $\lambda_{\text{TEP}} \approx 4000$ km (Step 016); UCD saturation model gives $S_{\oplus} \approx 0.35$ (Step 010); Cassini solar conjunction constrains $|\gamma - 1|$ (Section 4.6).
- Null-catalog stress test: full $n = 9$ raw-catalog likelihood improves by $\Delta \log L \approx 13.63$ over the null after random-effects prediction uncertainty is included (Step 039).
- Failure-mode accounting: disformal regime confirmation via $v_{\text{trans}} \approx 16.8$ km/s and sign-rule consistency checks (Step 007/017).
- Machine-checked manuscript: this document is generated from `site/components/*.html` and audited against pipeline JSON outputs by Step 027.

5.15 Summary

These limitations are explicitly acknowledged to ensure intellectual honesty. They do not invalidate the central conclusion—that TEP with Temporal Shear suppression within continuous Temporal Topology provides a quantitative explanation for the flyby anomaly—but indicate areas requiring additional scrutiny. The framework makes falsifiable predictions that can be tested with additional flyby data.

6. Conclusions

This study investigated whether the Temporal Equivalence Principle (TEP), incorporating Temporal Shear Suppression, can explain the Earth flyby anomaly—unexplained velocity shifts observed during spacecraft gravity assists. The analysis of twelve Earth flyby events spanning nine spacecraft (Galileo 1990/1992, NEAR, Cassini, Rosetta 2005/2007/2009, MESSENGER, Juno, Stardust, OSIRIS-REx, BepiColombo) yields the following key findings:

1. Three-gated TEP ensemble: Inverse-variance fitting (Step 008) enters NEAR (13.46 ± 0.01 mm/s), Galileo 1990 (3.92 ± 0.03 mm/s), and Rosetta 2005 (1.82 ± 0.05 mm/s) after pre-fit gates ($S/N \geq 2$, sign agreement at $\beta_{\text{ref}} = 10^{-4}$). At β_{ref} , Rosetta 2005 predicts $\Delta v_{\text{TEP}} \approx 0.32$ mm/s vs 1.82 mm/s observed; the full-model fit yields $\beta_{\text{fitted}} \approx 1.01 \times 10^{-3}$. Cassini (0.11 ± 0.05 mm/s) remains a fourth published anomaly but is excluded from the β_{fit} ensemble because $\Delta v_{\text{TEP}}(\beta_{\text{ref}}) < 0$ while the published anomaly is > 0 . The three gated β_{fit} values span roughly a factor of 5.3 (1.01×10^{-3} to 5.33×10^{-3}), consistent with geometry- and plasma-dependent modulation. The flyby trajectory-response amplitudes β_{fit} are distinct from the bare conformal coupling $\beta_A = -1.0$ and from the Cassini source charge $\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)} \alpha_0$. Cassini PPN compliance is evaluated through the solar source-charge projection (Section 4.6.1a), not through the fitted flyby amplitudes.
2. TEP parameter estimate: The inverse-variance weighted mean $\beta_{\text{fit}} \approx 1.73 \times 10^{-3} \pm 6.82 \times 10^{-5}$ summarizes the gated trio. This is the flyby trajectory-response amplitude, distinct from the bare conformal coupling $\beta_A = -1.0$ and from the Cassini source charge α_{eff} . Formal heterogeneity (reduced $\chi^2 \gg 1$, $I^2 \approx 100\%$) signals that a single rescaling does not exhaust geometry-plasma physics. Bootstrap resampling (10^4 draws) and leave-one-out recomputations in Step 008 show moderate stability (coefficient ≈ 0.148), with NEAR as the dominant lever.
3. PPN compliance via Temporal Topology screening: The Cassini solar conjunction experiment provides the tightest bound on the post-Newtonian light-propagation sector, measuring $\gamma = 1 + (2.1 \pm 2.3) \times 10^{-5}$. This constrains the solar-system Shapiro/light-propagation sector but does not directly test spatial clock-sector covariance, one-way residual shear, or low-density temporal-shear recovery. The fitted β_{fit} values are flyby trajectory-response amplitudes, not the Cassini source charge $\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)} \alpha_0$. Cassini PPN compliance is evaluated through the solar source-charge projection (Section 4.6.1a): the Jakarta radial field solution gives $S_{\Sigma}^{(\odot)} \leq 10^{-8}$ for the screened quadratic benchmark, well inside the Cassini requirement $S_{\Sigma}^{(\odot)} \lesssim 3.4 \times 10^{-3}$. This supports the claim that TEP can reduce to the GR PPN light-propagation limit in the screened solar environment while reserving its discriminating predictions for observables outside the Cassini measurement class.
4. TEP suppression by modern orbit determination: Analysis of the expanded dataset reveals that published null results (MESSENGER, Rosetta 2007) are consistent with universal- β_{fit} null predictions, while Juno is the sole raw-tension case at

- universal β_{fit} . Post-OD survival factors are withheld until mission OD configuration yields defensible F_{OD} estimates. Rosetta 2009 is treated as a published null/bound case with insufficient explicit geometry for the universal- β_{fit} prediction table; Stardust, OSIRIS-REx, and BepiColombo have no public anomaly report and are not used in quantitative likelihood.
5. Multiple independent lines of evidence: Altitude-dependent anomaly pattern (see point 6), historical timeline and the OD filtering mechanism motivate the hypothesis that modern orbit determination can treat TEP-like signals as systematic errors. However, mission-specific survival factors are not computed in this paper: Step 021 withholds F_{OD} until real OD configuration data are available, and the synthetic Step 012 OD run is quarantined from manuscript inference.
 6. Temporal Topology screening support: The model predicts null results for high-altitude flybys where gradient suppression attenuates TEP effects, while explaining large anomalies for low-altitude encounters. Altitude alone is not the relevant TEP discriminator; the signal is governed by the full trajectory-geometry response, which combines altitude, asymmetry, velocity and field structure.
 7. Systematic uncertainty compression: Transitioning from empirical characteristic suppression factors to a UCD-derived estimate via the Self-Consistent Field (SCF) solver and the corrected uncertainty analysis (Step 025) provides a rigorous uncertainty budget. The total relative uncertainty is 72.9%, with heterogeneity contributing 66.7% and systematic uncertainty contributing 29.2%. The systematic component is dominated by characteristic suppression uncertainty (25.0%, Paper 6 UCD) and relaxation length uncertainty (15.0%, SCF theoretical prior). This shift from "parameter fitting" to "systematic prediction" with proper variance decomposition strengthens the theoretical foundation of the TEP analysis.
 8. Robust statistical checks: The primary gated fit and model-comparison layer use inverse-variance/Gaussian weighted likelihoods, while Student's t likelihoods are retained in auxiliary Bayesian checks to test sensitivity to outliers. Residual diagnostics on the gated trio are consistent with normality (Shapiro–Wilk $p \approx 0.50$), so heterogeneity diagnostics, not normality alone, dominate the statistical interpretation. (NEAR, Galileo 1990, Rosetta 2005; Cassini analysed separately).
 9. Cosmographic CMB-frame directional consistency: Full 3D spacecraft state vectors from JPL Horizons were tested for CMB-frame velocity geometry (Step 040, $n = 9$). The exploratory both-aligned flag yields Pearson $r = +0.37$, $p = 0.36$, and Mann-Whitney $U = 16$, $p = 0.095$. Multivariate regression on residual ratio achieves $R^2 = 0.22$ (adjusted $R^2 = -0.24$), and an optimal weighted combination of spacecraft and Earth CMB projections gives $r = -0.61$, $p = 0.11$. These results are sample-limited and remain exploratory.

Significance

The TEP interpretation of the Earth flyby anomaly provides a coherent theoretical framework connecting spacecraft dynamics to fundamental physics. The flyby trajectory-response amplitude $\beta_{\text{fit}} \sim 1.7 \times 10^{-3}$ (weighted mean) is a trajectory-response coefficient, not the bare coupling β_A or the Cassini source charge α_{eff} . With geometric screening via Temporal Shear suppression, the flyby response is consistent with solar system constraints while explaining the anomalous velocity shifts.

Unlike ad hoc modifications to gravity, the TEP framework preserves all successes of general relativity in solar system tests while explaining anomalous behavior in the specific regime of planetary gravity assists. The geometric screening via Temporal Shear suppression, calibrated by independent UCD saturation analysis, is essential for the flyby amplitude and environmental response: without it, the required β_{fit} would not produce the observed trajectory-dependent residuals.

Statistical evidence strength: The validation analysis provides substantial statistical support for TEP:

- Effect sizes: Cohen's d relative to the published null population ($n_{\text{null}} = 5$) yields very large effects for NEAR ($d \approx 3.4$) and Galileo 1990 ($d \approx 1.0$), a small-to-medium effect for Rosetta 2005 ($d \approx 0.5$), and a negligible effect for Cassini ($d \approx 0.03$). The coefficient of variation $CV \approx 68\%$ across the three gated β_{fit} fits reflects genuine geometry-dependent modulation at fixed envelope.
- Model comparison: On the full $n = 9$ catalog, Step 026 gives a large information-criterion separation between TEP restricted and Null ($\Delta\text{BIC} \approx 714$) and positive but not decisive preference over Anderson empirical ($\Delta\text{BIC} \approx 77.7$)
- Akaike model weight on TEP restricted ≈ 1 in the reported two-model summary; Anderson and flexible tiers are documented in Step 026 outputs.
- Robustness: Bootstrap resampling, leave-one-out recomputations, and auxiliary robust checks are reported in Step 008/013.
- Prediction accuracy: Step 008 reports $R^2 \approx 0.945$ between predicted and observed anomalies for the primary comparison table; prediction intervals are bootstrap-derived.
- Residual analysis: Shapiro–Wilk $p \approx 0.50$ on the gated trio; heterogeneity diagnostics dominate formal tests.
- Sensitivity analysis: All parameters stable across plausible ranges; fit stability maintained

The catalog of Earth flyby events (four published anomalies, five published nulls/bounds, and several without public anomaly reports) provides the empirical substrate. Step 026's headline likelihood uses the full $n = 9$ catalog with a geometry-spread systematic uncertainty ($\sigma_{\text{geom}} \approx 0.520$ mm/s) because the tiny published per-flyby uncertainties would otherwise produce astronomically large, scientifically meaningless values. Information-criterion comparison under this adopted likelihood favours TEP restricted over Null and yields positive but assumption-sensitive separation over Anderson; the magnitude of the separation depends on the treatment of systematic uncertainty and is interpreted as model-selection support rather than calibrated evidence.

Several potential concerns have been investigated and addressed through rigorous statistical analysis (Step 024, Step 025, Step 026):

Systematic error discrimination: The evidence against systematic error origins is primarily organisational rather than statistical. TEP theory explicitly predicts that anomaly magnitude should correlate with trajectory asymmetry ($\cos \delta_{\text{in}} - \cos \delta_{\text{out}}$); systematic measurement errors have no mechanism to produce such correlations. However, with only $n = 3$ gated detections, the monotonic ordering of anomaly magnitudes with trajectory asymmetry has a ceiling of $p = 1/6$ ($\sim 1\sigma$): six orderings are possible and the observed one is favourable, but this is a qualitative consistency check, not a primary discriminator. Furthermore, Cassini's predicted sign is negative where the published anomaly is positive (one of four retained anomalies), and the geometry envelope carries seven nominal deterministic coefficients plus one fitted amplitude against four anomalies, so the model-selection support must be read as organisational under the adopted convention rather than calibrated evidence. The broader catalog rank correlation (Spearman $\rho = 0.73$, $p = 0.008$, $n = 12$) provides complementary but independent support. See Section 5.5 for comprehensive systematic uncertainty budget.

Data provenance: The analysis relies on published anomaly values from Anderson et al. (2008) rather than independent DSN re-analysis. This is addressed by: (a) cross-referencing multiple literature sources for consistency, (b) demonstrating that TEP predictions match the observed anomaly pattern (altitude dependence, trajectory geometry), (c) providing a framework for raw DSN data re-analysis to independently test the suppression hypothesis. The current repository state provides mission discovery, structured requests, and an ingestion scaffold; in typical runs most missions do not yet have indexed local raw Doppler products available for reprocessing (pipeline ingest status `no_indexed_products`). The DSN pathway is therefore an explicit falsification route, not a completed independent reanalysis across the catalog.

β_{fit} scatter as physical modulation: Fitted $\beta_{A,\text{eff}}$ spans roughly a factor of 5.3 for $n = 4$ valid fits (Step 009), reflecting geometry-dependent modulation: altitude (J_2 gradient suppression), perigee latitude (inclination-dependent coupling), plasma environment (ionospheric gradient modulation), and velocity (disformal regime). Step 009 does not report a formal variance decomposition because $n \leq 4$ renders ANOVA-style partitioning statistically meaningless (standard error on sample variance exceeds 100% of the estimate). Instead, it reports deterministic geometry modulation metrics: beta scatter statistics, full-catalog detection-pattern classification (75% accuracy, $n = 12$), and Spearman rank correlation ($\rho = 0.73$, $p = 0.008$). Cross-validation on the gated trio reports stability coefficient ≈ 0.148 . The UCD-derived characteristic suppression $S_{\oplus} \approx 0.35$ provides a cross-scale prior. See Section 4.3 for the quantitative geometry modulation assessment and Section 5.5 for detailed interpretation.

Cassini status: At $\beta_{\text{ref}} = 10^{-4}$ the Step 007 envelope yields $\Delta v_{\text{TEP}} \approx -0.023$ mm/s while the published anomaly is $+0.11$ mm/s, so Cassini has sign mismatch at β_{ref} and is retained in the primary pool ($n = 4$) but excluded from the sign-gated ensemble ($n = 3$). Addressing Cassini therefore requires independent DSN OD or envelope refinements, not only a universal rescaling of β_{fit} .

Juno falsification pressure: At the refit weighted-mean β_{fit} , Step 039 predicts $\Delta v_{\text{TEP}}^{\text{raw}} \approx +0.11$ mm/s with uncertainty ~ 0.038 mm/s while the published bound is 0.00 ± 0.02 mm/s. This raw-tension case is the headline stress test on universal β_{fit} before mission-specific OD survival factors are supplied (Step 021). Independent raw DSN re-analysis with TEP-inclusive orbit determination is required to adjudicate the tension.

Sample size as complete dataset: The analysis includes all accessible Earth gravity assist flybys with adequate DSN tracking between 1990–2020. The rarity of suitable flyby events (low altitude, Doppler tracking, no major maneuvers) means only a handful of literature anomalies exist; three enter the gated β_{fit} ensemble. Step 008 bootstrap 95% CI $[1.01 \times 10^{-3}, 5.33 \times 10^{-3}]$ brackets the gated fits. Additional flybys would test envelope refinements rather than restate the same $n = 3$ compression.

PPN compliance: The UCD-derived characteristic suppression $S_{\oplus} \approx 0.35$ is determined from the UCD saturation model. The flyby trajectory-response amplitudes β_{fit} are distinct from the Cassini source charge $\alpha_{\text{eff}}^{(\odot)} = S_{\Sigma}^{(\odot)} \alpha_0$; Cassini PPN compliance is evaluated through the solar source-charge projection (Section 4.6.1a), where the Jakarta radial field solution gives $S_{\Sigma}^{(\odot)} \leq 10^{-8}$ for the screened quadratic benchmark, well inside the Cassini requirement $S_{\Sigma}^{(\odot)} \lesssim 3.4 \times 10^{-3}$.

Model comparison: Step 026 on the full $n = 9$ catalog gives a large information-criterion separation between TEP restricted and Null ($\Delta \text{BIC} \approx 714$) and a positive but not decisive preference over Anderson ($\Delta \text{BIC} \approx 77.7$). The Anderson empirical model also shows strong separation vs Null ($\Delta \text{BIC} \approx 636$), confirming that trajectory asymmetry carries genuine signal. The TEP flexible model (3 parameters) is penalized by its parameter count and does not outperform the restricted model on BIC. Residual diagnostics on the gated trio are consistent with normality (Shapiro–Wilk $p \approx 0.50$).

Independent validation pathways: Several approaches can independently test the TEP hypothesis without relying on the published anomaly values:

- Raw DSN data re-analysis: Analysis of raw DSN tracking archives from NASA's Planetary Data System using minimal orbit determination (reduced gravity field expansion, unfiltered Doppler, no continuity penalties) would test whether TEP signals are filtered by modern orbit determination methods. This would provide an important test of the suppression hypothesis.
- Additional flyby analysis: Earth gravity assist missions provide opportunities for independent detection. Analysis with both standard and minimal orbit determination methods would test the suppression prediction.

- GNSS clock correlation: GNSS atomic clock correlation analysis provides a cross-consistency check with the GNSS/UCD terrestrial calibration on the transition radius ($R_{\text{sol}} \approx 4200$ km). This shared-input dependency is disclosed explicitly to avoid circular claims of independent validation.
- Lunar Laser Ranging: Precision LLR analysis in related work reports a synodic-phase signal consistent with the screening mechanism and Temporal Topology Saturation Scale (UCD) framework. Independent LLR validation would strengthen the screening mechanism established in this analysis.

Data Availability

Spacecraft trajectories are available through the JPL Horizons ephemeris service. Literature anomaly values are from Anderson et al. (2008) and companion publications. Analysis code and processed data products are available at <https://github.com/matthewsmawfield/TEP-EFA> with archived DOI at 10.5281/zenodo.19454862.

Acknowledgments

The NASA Deep Space Network and Jet Propulsion Laboratory provided the precision Doppler tracking that enabled flyby anomaly detection. The JPL Horizons system provided trajectory reconstruction. This work utilizes published literature values from the Orbit Determination Program analyses by Anderson et al. and collaborators. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The author declares no conflicts of interest.

Additional Considerations

Several avenues for extending this analysis are identified:

Raw DSN data re-analysis: Analysis of raw DSN tracking archives from NASA's Planetary Data System using minimal orbit determination (reduced gravity field expansion, unfiltered Doppler, no continuity penalties) would test whether TEP signals are filtered by modern orbit determination methods. This provides an important test of the suppression hypothesis.

Extended spacecraft sample: Additional flyby events would increase the sample size beyond the current four published anomaly cases (three sign-gated for the inverse-variance β_{fit} ensemble). A sample of $n \approx 74$ primary detections would provide sufficient statistical power to distinguish between geometry-dependent modulation of β_{fit} and a single universal coupling constant at 80% power (conservative estimate: $n \approx 153$).

- Full numerical Temporal Shear Suppression solver: Implementation of a numerical Temporal Topology field solver (e.g., using the shooting method or relaxation techniques) validates the phenomenological gradient suppression model used in this analysis. This enables prediction of the Temporal Topology profile without the phase-boundary approximation and could explain the observed β_{fit} scatter through detailed density-dependent effects.
- Inclination-dependent modeling: Incorporation of Earth's oblateness (J_2) and latitude-dependent density variations into the TEP model could explain part of the observed β_{fit} scatter. Spacecraft with different orbital inclinations sample different gravitational field geometries, which modulate the Temporal Topology field strength.
- Disformal coupling exploration: Extension to scalar-tensor theories with disformal coupling terms introduces velocity-dependent gradient suppression that could explain the correlation between fitted β_{fit} and flyby velocity. This provides a more general framework for understanding the geometry-dependence of the TEP effect.
- Local-time plasma effects: Investigation of ionospheric plasma density variations with local time could explain time-dependent modulation of the TEP signal. Day-side vs. night-side flybys experience different plasma environments that may modify the effective gradient suppression.

References

1. Anderson, J. D., Campbell, J. K., Ekelund, J. E., Ellis, J., & Jordan, J. F. 2008, "Anomalous Orbital-Energy Changes Observed during Spacecraft Flybys of Earth," *Phys. Rev. Lett.*, 100, 091102
2. Anderson, J. D., & Nieto, M. M. 2009, "Astrometric solar-system anomalies," in *Relativity in Fundamental Astronomy*, IAU Symp. 261, 189
3. Antreasian, P. G., & Guinn, J. R. 1998, "Investigations into the Unexpected Delta-V during the Earth Gravity Assist of NEAR," Paper AAS 98-428
4. Bertotti, B., Iess, L., & Tortora, P. 2003, "A test of general relativity using radio links with the Cassini spacecraft," *Nature*, 425, 374
5. Brax, P., van de Bruck, C., Davis, A.-C., Khoury, J., & Weltman, A. 2004, "Detecting dark energy in orbit: The cosmological chameleon," *Phys. Rev. Lett.*, 93, 200405

6. Einstein, A. 1915, "Die Feldgleichungen der Gravitation," *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 844
7. Halsey, D., et al. 2012, "Anomalous Earth flybys: Status and developments," *Adv. Space Res.*, 50, 362
8. Khoury, J., & Weltman, A. 2004, "Chameleon cosmology," *Phys. Rev. D*, 69, 044026
9. Lämmerzahl, C., Preuss, O., & Dittus, H. 2006, "Is the physics within the Solar system understood?" in *Lasers, Clocks and Drag-Free Control*, 75, 75
10. McCulloch, M. E. 2008, "Modelling the Pioneer anomaly as modified inertia," *MNRAS*, 389, L57
11. Meeus, J. 1998, *Astronomical Algorithms*, 2nd edn. (Richmond: Willmann-Bell)
12. Mota, D. F., & Shaw, D. J. 2007, "Strongly coupled chameleon fields," *Phys. Rev. Lett.*, 97, 151102
13. Nieto, M. M., & Anderson, J. D. 2007, "Search for a solution of the Pioneer anomaly," *Contemp. Phys.*, 48, 41
14. Page, G., & McCulloch, M. E. 2009, "Modelling the flyby anomalies using a modification of inertia: Further investigations," *Int. J. Astron. Astrophys.*, 3, 1
15. Schive, H.-Y., Chiueh, T., & Broadhurst, T. 2014, "Understanding the Core-Halo Relation of Quantum Wave Dark Matter from 3D Simulations," *Phys. Rev. Lett.*, 113, 261302
16. Turyshv, S. G., & Toth, V. T. 2010, "The Pioneer anomaly," *Living Rev. Relativ.*, 13, 4
17. Will, C. M. 2014, "The confrontation between general relativity and experiment," *Living Rev. Relativ.*, 17, 4
18. Folkner, W. M., et al. 2022, "Planetary ephemeris DE440," *IPN Progress Report*, 42-284, 1
19. JPL Horizons, "NASA/JPL Horizons System" <https://ssd.jpl.nasa.gov/horizons/> (accessed 2024)
20. Morley, T., & Budnik, F. 2007, "Rosetta Navigation at its First Earth-Swingby," *Proceedings of the 20th International Symposium on Space Flight Dynamics*
21. Müller, J., Soffel, M., & Klioner, S. A. 2008, "Geodesy and relativity," *Journal of Geodesy*, 82, 133
22. Müller, J., et al. 2010, "Relativistic models for spacecraft tracking," *Acta Astronautica*, 67, 975
23. Aksenov, E. L., & Tuchin, A. G. 2020, "Earth flyby anomalies and the general relativistic theory of the Kerr gravitational field," *MNRAS*, 492, 3703
24. Ciufolini, I., & Pavlis, E. C. 2004, "A confirmation of the general relativistic prediction of the Lense-Thirring effect," *Nature*, 431, 958
25. IERS Conventions 2010, IERS Technical Note No. 36, eds. Petit, G. & Luzum, B.
26. Brax, P., & Burrage, C. 2014, "Constraining screened modified gravity with the CASPER experiment," *Phys. Rev. D*, 90, 104009
27. Lemoine, F. G., et al. 1998, "The Development of the NASA GSFC and NIMA Joint Geopotential Model," in *Proceedings of the International Symposium on Gravity, Geoid, and Marine Geodesy*, Tokyo, Japan
28. Pavlis, N. K., et al. 2012, "The development and evaluation of the Earth Gravitational Model 2008 (EGM2008)," *J. Geophys. Res.*, 117, B04406
29. Mocz, P., Vogelsberger, M., Robles, V., et al. 2018, "Galaxy Halos from Fuzzy Dark Matter," *Phys. Rev. Lett.*, 121, 141102
30. Moyer, T. D. 2000, *Formulation for Observed and Computed Values of Deep Space Network Data Types*, JPL Publication 00-7
31. Burrage, C., & Sakstein, J. 2016, "Tests of Ambient Symmetry Restoration," *Living Rev. Relativ.*, 21, 1
32. Upadhye, A., Hu, W., & Khoury, J. 2012, "Quantum stability of chameleon field theories," *Phys. Rev. Lett.*, 109, 041301
33. Joyce, A., Jain, B., Khoury, J., & Trodden, M. 2015, "Beyond the cosmological standard model," *Phys. Rept.*, 568, 1
34. Kass, R. E., & Raftery, A. E. 1995, "Bayes Factors," *J. Am. Stat. Assoc.*, 90, 773
35. Clifton, T., Ferreira, P. G., Padilla, A., & Skordis, C. 2012, "Modified gravity and cosmology," *Phys. Rept.*, 513, 1
36. Higgins, J. P., & Thompson, S. G. 2002, "Quantifying heterogeneity in a meta-analysis," *Stat. Med.*, 21, 1539

TEP Research Series

- Smawfield, M. L. *Temporal Equivalence Principle: Dynamic Time & Emergent Light Speed*. Preprint v0.12 (Jakarta). Zenodo. DOI: 10.5281/zenodo.16921911
- Smawfield, M. L. *Global Time Echoes: Distance-Structured Correlations in GNSS Clocks*. Preprint v0.27 (Jaipur). Zenodo. DOI: 10.5281/zenodo.17127229
- Smawfield, M. L. *Global Time Echoes: 25-Year Analysis of CODE Precise Clock Products*. Preprint v0.20 (Cairo). Zenodo. DOI: 10.5281/zenodo.17517141
- Smawfield, M. L. *Global Time Echoes: Raw RINEX Consistency Test*. Preprint v0.6 (Kathmandu). Zenodo. DOI: 10.5281/zenodo.17860166
- Smawfield, M. L. *Temporal-Spatial Coupling in Gravitational Lensing: A Reinterpretation of Dark Matter Observations*. Preprint v0.8 (Tortola). Zenodo. DOI: 10.5281/zenodo.17982540
- Smawfield, M. L. *Global Time Echoes: Empirical Synthesis*. Preprint v0.6 (Singapore). Zenodo. DOI: 10.5281/zenodo.18004832
- Smawfield, M. L. *Temporal Topology Saturation Scale: Cross-Scale Consistency of ρ_T* . Preprint v0.8 (New Delhi). Zenodo. DOI: 10.5281/zenodo.18064365

Smawfield, M. L. *The Soliton Wake: Exploring RBH-1 as a Temporal Topology Candidate*. Preprint v0.4 (Doha). Zenodo. DOI: 10.5281/zenodo.18059250

Smawfield, M. L. *Global Time Echoes: Optical-Domain Consistency Test via Satellite Laser Ranging*. Preprint v0.4 (Mombasa). Zenodo. DOI: 10.5281/zenodo.18064581

Smawfield, M. L. *What Do Precision Tests of General Relativity Actually Measure?*. Preprint v0.7 (Istanbul). Zenodo. DOI: 10.5281/zenodo.18109760

Smawfield, M. L. *Temporal Equivalence Principle: Suppressed Density Scaling in Globular Cluster Pulsars*. Preprint v0.9 (Caracas). Zenodo. DOI: 10.5281/zenodo.18165798

Smawfield, M. L. *The Cepheid Bias: Resolving the Hubble Tension*. Preprint v0.10 (Kingston upon Hull). Zenodo. DOI: 10.5281/zenodo.18209702

Smawfield, M. L. *Temporal Equivalence Principle: A Unified Resolution to the JWST High-Redshift Anomalies*. Preprint v0.7 (Kos). Zenodo. DOI: 10.5281/zenodo.19000827

Smawfield, M. L. *Temporal Equivalence Principle: Temporal Shear Recovery in Gaia DR3 Wide Binaries*. Preprint v0.6 (Santiago). Zenodo. DOI: 10.5281/zenodo.19102061

Data Availability & Reproducibility

This work follows open-science practices. All results are fully reproducible from the repository’s raw inputs using the documented pipeline. All numerical results, figures, and statistics are generated by deterministic Python scripts processing real mission ephemerides and peer-reviewed, published Doppler-derived anomaly measurements.

Cross-corpus theory registry (TEP-EFA ↔ manuscript series)

The flyby pipeline fixes a single set of conventions so numerical results agree with the manuscript HTML in `site/components/`. The following table is the authoritative map between this repository and the numbered theory papers under `manuscripts/` (Paper 0 = Jakarta, Paper 6 = UCD, Paper 10 = Caracas COS, Paper 12 = JWST Kos, etc.).

Canonical EFA quantities aligned to the manuscript corpus		
Quantity	EFA implementation	Manuscript anchor(s)
Matter metric / conformal factor	<code>A(φ) = exp(β_A φ / M_{Pl})</code> (reduced Planck mass in GeV)	Paper 0 (Jakarta) axioms; Papers 4–5, 8–9, 12 (notation)
Scalar force (flyby sector)	$F_φ = β_{A,eff} c^2 ∇φ / M_{Pl}$; $β_{A,eff} = β_A × S_⊙$	Paper 0; Paper 4 (Phantom Mass); methodology §3.2
Density minimum $φ_{min}(ρ)$	$φ = Λ [n Λ^{n+4} M_{Pl} / (2 β_A ρ_{GeV4})]^{1/(n+1)}$ in <code>step_007_tep_model.py</code> , <code>step_011_trajectory_integration.py</code> , <code>step_019_3d_field_integration.py</code>	Paper 10 Appendix C (aligned to this form, May 2026); Paper 6 (scaling $φ ∝ ρ^{-1/(n+1)}$ only). Paper 0 states screening/PPN mapping; it does not fix the closed $φ(ρ)$ line in the main text.
PPN $γ$ (magnitude checks)	<code>ppn_gamma_deviation</code> : report $ γ - 1 ≈ 2 β_{A,eff}^2$ vs Cassini	Paper 0 Sec. 7: $γ - 1 = -2 α_{eff}^2$ (DEF); Papers 5, 11, 12 (screened limit narrative)
Screening / UCD radius	$R_{sol} ≈ 4146$ km, $S_⊙ = (R_⊙ - R_{sol})/R_⊙ ≈ 0.35$; $ρ_T ≈ 20$ g cm ⁻³	Paper 6 (UCD); Step 010 / <code>physics.py</code>
Scalar field equation (sign reference)	Pipeline uses explicit <code>field_gradient</code> / TEP screening relaxation outside Earth; trace source uses $β_A, M_{Pl}$ as in Step 007 comments	Paper 12 Appendix A.1.2: $K(φ)□φ - V' = - (β/M_{Pl})T^{(matter)}$ (Einstein-frame convention; overall sign of T follows chosen action)

Residual ambiguities: Individual papers sometimes use illustrative potentials or linearized V_{eff} without the Einstein-frame factor 2; any updated analytic appendix should match the table above before reusing EFA numerical $φ_{Earth}$, $φ_{space}$, or $Δφ$ in secondary

calculations.

Repository & Code

The repository contains a deterministic, version-controlled analysis pipeline with analysis steps for Earth flyby trajectory data. All steps are orchestrated by `scripts/run_all.py` with comprehensive logging.

Repository Structure

```
TEP-EFA/
├── data/                                # Raw and processed data
│   ├── raw/                            # Raw DSN tracking, trajectories
│   │   ├── dsn_tracking/               # Deep Space Network archives
│   │   ├── flyby_trajectories/         # JPL Horizons ephemeris data
│   │   └── spice_kernels/              # Navigation SPICE kernels
│   └── processed/                      # Pipeline outputs (JSON/CSV)
├── scripts/
│   ├── steps/                          # Analysis pipeline steps
│   │   ├── step_001_download_spice.py
│   │   ├── step_002_spice_to_json.py
│   │   ├── step_003_archival_data_mining.py
│   │   ├── step_004_jpl_horizons_fetch.py
│   │   ├── step_005_dsn_data_ingestion.py
│   │   ├── step_006_dsn_framework.py
│   │   ├── step_007_tep_model.py
│   │   ├── step_008_fitting.py
│   │   ├── step_009_variance_analysis.py
│   │   ├── step_010_tep_first_principles.py
│   │   ├── step_011_trajectory_integration.py
│   │   ├── step_012_od_filter_simulation.py
│   │   ├── step_013_cross_validation.py
│   │   ├── step_014_sensitivity_analysis.py
│   │   ├── step_015_hierarchical_bayesian.py
│   │   ├── step_016_gnss_validation.py
│   │   ├── step_017_plasma_modulation.py
│   │   ├── step_018_space_weather.py
│   │   ├── step_019_3d_field_integration.py
│   │   ├── step_020_plasma_environment_reconstruction.py
│   │   ├── step_021_mission_specific_od_absorption.py
│   │   ├── step_022_atmospheric_drag_simulation.py
│   │   ├── step_023_thermal_recoil_modeling.py
│   │   ├── step_024_systematic_error_monte_carlo.py
│   │   ├── step_025_corrected_uncertainty.py
│   │   ├── step_026_stable_model_comparison.py
│   │   ├── step_027_claim_consistency_audit.py
│   │   ├── step_028_dsn_processing.py
│   │   ├── step_029_read_trk234.py
│   │   ├── step_030_juno_reanalysis.py
│   │   ├── step_031_pds_search.py
│   │   ├── step_032_tep_suppression.py
│   │   ├── step_033_iri_trajectory_profile.py
│   │   ├── step_034_covariant_holonomy.py
│   │   ├── step_035_cross_corpus_export.py
│   │   ├── step_036_final_report.py
│   │   ├── step_037_visualizations.py
│   │   ├── step_038_extract_3d_vectors.py
│   │   ├── step_039_flyby_prediction_table.py
│   │   ├── step_040_cosmographic_shear.py
│   │   ├── step_041_envelope_heuristic_sensitivity.py
│   │   └── step_042_time_resolved_cosmography.py
│   ├── utils/                          # Utility functions
│   └── build_markdown.js                # Manuscript builder
└── site/
    └── components/                     # Manuscript HTML sections
```

└─ config/	# Pipeline configuration
└─┬─ pipeline_config.json	
└─ logs/	# Per-step execution logs
└─ requirements.txt	# Python dependencies
└─ README.md	# Documentation
└─ LICENSE	# CC-BY-4.0

Data Provenance

Data Source	Provider	Access Method	Size	Location
JPL Horizons Ephemeris	NASA/JPL	Astroquery API	~2 MB	data/raw/flyby_trajectories/
DSN Doppler Archives	NASA DSN	Literature values	~500 KB	Anderson et al. (2008)
Flyby Anomaly Catalog	Peer-reviewed literature	Manual compilation	~50 KB	results/step003_archival_flyby_catalog.json
SPICE Kernels	NASA NAIF	Auto-downloaded	~100 MB	data/raw/spice_kernels/

Pipeline Architecture

The analysis pipeline comprises 42 deterministic steps organized into logical groups. Each step is a standalone Python script in `scripts/steps/` that produces JSON outputs and detailed logs in `logs/step_*.log`.

Complete Step Inventory & Runtime

Group	Step	Script	Description	Runtime
Phase 1: Data Acquisition & Preparation (001-006)				
Data	001	step_001_download_spice.py	SPICE kernel download (NAIF archive)	~30s
Data	002	step_002_spice_to_json.py	SPICE to JSON conversion	~1s
Data	003	step_003_archival_data_mining.py	Archival flyby catalog compilation	~2s
Data	004	step_004_jpl_horizons_fetch.py	JPL Horizons ephemeris data fetch	~5s
Data	005	step_005_dsn_data_ingestion.py	DSN tracking data ingestion	~1s
Data	006	step_006_dsn_framework.py	DSN raw data acquisition framework	~1s
Phase 2: Core Physics & Geometry Modulation Analysis (007-010)				
Core	007	step_007_tep_model.py	TEP Temporal Topology model with screening	~1s
Core	008	step_008_fitting.py	β parameter fitting with PPN validation	~1s
Core	009	step_009_variance_analysis.py	Deterministic geometry modulation analysis (formal variance partitioning deferred to n ≥ 10)	~2s
Core	010	step_010_tep_first_principles.py	UCD saturation derivation	~10s
Phase 3: Trajectory & Observational Pipeline (011-012)				
Traj	011	step_011_trajectory_integration.py	Numerical trajectory integration	~5s
OD	012	step_012_od_filter_simulation.py	Synthetic OD diagnostic: noise-only control, perigee state bias vs TEP truth, station/two-way sensitivity (not F_OD)	~10s
Phase 4: Validation & Robustness (013-016)				
Valid	013	step_013_cross_validation.py	Cross-validation analysis	~5s

Group	Step	Script	Description	Runtime
Valid	014	step_014_sensitivity_analysis.py	Parameter sensitivity analysis	~2s
Valid	015	step_015_hierarchical_bayesian.py	Hierarchical Bayesian model	~30s
Valid	016	step_016_gnss_validation.py	GNSS atomic clock validation	~1s
Phase 5: Extended Physics (017-019)				
Phys	017	step_017_plasma_modulation.py	Plasma-dependent gradient modulation	~2s
Phys	018	step_018_space_weather.py	Space weather correlation analysis	~1s
Phys	019	step_019_3d_field_integration.py	3D integrator diagnostic (idealized analytic hypoflyby; see JSON metadata)	~1s
Phase 6: Plasma & Environmental (020-023)				
Plasma	020	step_020_plasma_environment_reconstruction.py	Plasma environment reconstruction	~3s
OD	021	step_021_mission_specific_od_absorption.py	Mission-specific OD absorption	~1s
Env	022	step_022_atmospheric_drag_simulation.py	Atmospheric drag simulation	~1s
Env	023	step_023_thermal_recoil_modeling.py	Thermal recoil modeling	~1s
Phase 7: Statistical Analysis (024-026)				
Stat	024	step_024_systematic_error_monte_carlo.py	Systematic error Monte Carlo analysis	~5s
Stat	025	step_025_corrected_uncertainty.py	Corrected uncertainty analysis	~1s
Stat	026	step_026_stable_model_comparison.py	Stable model comparison	~2s
Phase 8: Advanced Topics (027-028)				
Audit	027	step_027_claim_consistency_audit.py	Claim consistency audit	~1s
DSN	028	step_028_dsn_processing.py	DSN processing framework	~1s
Phase 9: DSN Reanalysis (029-032)				
DSN	029	step_029_read_trk234.py	Read TRK-2-34 data format	~1s
DSN	030	step_030_juno_reanalysis.py	Juno 2013: archive ingest, pairwise-Doppler residual proxy (& optional Horizons range/velocity batch; not MONTE/ODP OD)	~5s
DSN	031	step_031_pds_search.py	NASA PDS archive search	~2s
DSN	032	step_032_tep_suppression.py	TEP suppression analysis	~1s
Phase 10: Advanced Analysis (033-035)				
IRI	033	step_033_iri_trajectory_profile.py	Continuous IRI trajectory profiles	~5s
Holo	034	step_034_covariant_holonomy.py	Covariant temporal shear impulse	~1s
Export	035	step_035_cross_corpus_export.py	Cross-corpus parameter export	~1s
Phase 11: Reporting & Extended Validation (036-042)				
Report	036	step_036_final_report.py	Final report generation	~1s
Fig	037	step_037_visualizations.py	Publication-quality figure generation	~3s
Vec	038	step_038_extract_3d_vectors.py	Extract 3D state vectors from JPL Horizons	~2s
Pred	039	step_039_flyby_prediction_table.py	Flyby prediction table with uncertainty-aware classification	~1s
Cosmo	040	step_040_cosmographic_shear.py	Cosmographic temporal shear modulation test	~1s

Group	Step	Script	Description	Runtime
Env	041	<code>step_041_envelope_heuristic_sensitivity.py</code>	Geometry envelope heuristic sensitivity	~1s
Cosmo	042	<code>step_042_time_resolved_cosmography.py</code>	Time-resolved cosmography (after Horizons + optional Juno DSN sidcar)	~2s

Total Runtime Summary

Component	Steps	Runtime
Data Acquisition (001-006)	6	~40s
Core Physics & Variance (007-010)	4	~14s
Trajectory & Observational (011-012)	2	~8s
Validation & Robustness (013-016)	4	~38s
Extended Physics (017-019)	3	~4s
Plasma & Environmental (020-023)	4	~6s
Statistical Analysis (024-026)	3	~8s
Advanced Topics (027-028)	2	~2s
DSN Reanalysis (029-032)	4	~9s
Advanced Analysis (033-035)	3	~7s
Reporting & Extended Validation (036-042)	7	~11s
Total	42	~2 min

Reproduction Instructions

Quick Start (Full Reproduction)

```
# 1. Clone repository
git clone https://github.com/matthewsmawfield/TEP-EFA.git
cd TEP-EFA

# 2. Install dependencies
pip install -r requirements.txt

# 3. Run full pipeline (generates all results & figures)
python scripts/run_all.py

# 4. Results are located in:
#   - results/           (JSON data products and figures)
#   - logs/             (Detailed execution logs)
#   - site/dist/         (Built static site)
```

System Requirements

Component	Minimum	Recommended	Tested On
CPU	2 cores	4+ cores	Apple M4 Pro (14-core)
RAM	4 GB	8 GB	24 GB (M4 Pro)
Storage	500 MB	1 GB	NVMe SSD
Runtime	~2 min	~1 min	~40s (M4 Pro)

Key Analysis Outputs

- `results/step003_archival_flyby_catalog.json` — Literature flyby catalog with provenance

- `results/step007_tep_predictions.json` — TEP model predictions for all modeled flybys
- `results/step008_fitting_results.json` — β fitting results with PPN validation
- `results/step027_claim_consistency_audit.json` — Machine-readable manuscript/pipeline claim audit, including evidence-frame pass/fail status
- `results/step036_final_report.json` — Comprehensive results with Temporal Topology screening
- `results/step039_flyby_prediction_table.json` — Per-flyby raw pooled- β classification (post-OD columns withheld until Step 021 supplies F_{OD})
- `results/step032_tep_suppression_analysis.json` — Legacy empirical suppression diagnostic; superseded by Step 039 for manuscript inference
- `results/step037_figure1_altitude_anomaly.png` — Altitude vs anomaly correlation
- `results/step037_figure2_beta_comparison.png` — Fitted β comparison by spacecraft
- `results/step037_figure3_ppn_constraints.png` — PPN constraint analysis
- `results/step037_figure4_screening_profile.png` — Temporal Topology profile

Log Files

Each step produces detailed logs:

- `logs/pipeline.log` — Master pipeline execution log
- `logs/step_*.log` — Individual step logs

Software Dependencies

Package	Version	Purpose
Python	3.10+	Language runtime
NumPy	1.24+	Numerical computing
SciPy	1.10+	Statistical functions
Matplotlib	3.7+	Visualization
Astroquery	0.4.6+	JPL Horizons interface
spiceypy	5.1+	SPICE kernel handling
PyIRI	(package current)	Ionospheric electron density for Step 033 trajectory profiles
pytest	7+	Smoke tests (<code>pytest</code> from repository root)

All dependencies are specified in `requirements.txt`.

Validation & Testing

The pipeline includes comprehensive validation:

- Bootstrap Resampling: $n=10,000$ iterations for uncertainty quantification
- Leave-One-Out Cross-Validation: Tests robustness against single-flyby exclusion
- Heterogeneity Assessment: Cochran's Q and I^2 statistics for model scatter
- GNSS clock correlation: The GNSS atomic clock correlation analysis provides a cross-consistency check with the GNSS/UCD terrestrial calibration on the transition radius ($R_{sol} \approx 4146$ km). This shared-input dependency is disclosed explicitly to avoid circular claims of independent validation.
- Claim-consistency audit: Step 027 machine-checks manuscript claims against pipeline outputs, including model-comparison values, Table 3 fitted parameters, variance decomposition, Cassini exclusion status, Step 039 Juno classification, and evidence framing. The audit fails if the manuscript reverses the TEP-vs-Null comparison, omits the random-effects uncertainty, or promotes Juno from a deterministic warning case to an uncertainty-aware falsification.
- Uncertainty discipline: Formal inverse-variance summaries, random-effects scatter, geometry-spread model comparison, and published-uncertainty stress tests are reported as separate layers rather than blended into a single headline number.
- Automated smoke tests: `pytest` (see `tests/` and `pytest.ini`) checks repository layout and step logger conventions.

Reproducibility Checklist

To verify successful reproduction:

1. All configured pipeline steps complete with "SUCCESS" status
2. Primary JSON products are present in `results/`
3. Figure files are present in `results/` (PNG)
4. Key result: gated inverse-variance β_{fit} span 1.01×10^{-3} to 5.33×10^{-3} (NEAR, Galileo 1990, Rosetta 2005; weighted mean $\approx 1.73 \times 10^{-3}$ from `results/step008_fitting_results.json`)
5. Key result: $\beta_{\{A, \text{eff}\}} \sim 6 \times 10^{-4}$ with Temporal Topology screening
6. Key result: $|\gamma-1| \approx 7 \times 10^{-7}$ (safely below Cassini bound 2.3×10^{-5})
7. Key result: $I^2 \approx 100\%$ extreme heterogeneity (supports β scatter hypotheses)
8. Key result: Altitude-anomaly correlation $\rho = -0.18$ (not significant, $n = 9$); the TEP envelope predicts near-zero for high-altitude null trajectories and large for low-altitude high-asymmetry encounters, which is the relevant test
9. Key result: 5 published null or bound cases are consistent with fixed-amplitude Step 039 predictions (Step 008 pooled β_{fit}) under the Step 007 geometry envelope; 1 deterministic fixed-amplitude warning case remains (Juno, predicted $+0.11$ mm/s), but 0 raw-tension cases remain after random-effects prediction uncertainty; Rosetta 2009 is a published null/bound case with insufficient explicit geometry for the Step 039 table; 3 flybys (Stardust, OSIRIS-REx, BepiColombo) have no public anomaly report

Data Availability Statement

Spacecraft trajectories are available through the NASA JPL Horizons ephemeris service. Literature anomaly values are from Anderson et al. (2008) and companion publications. Analysis code and processed data products are available at <https://github.com/matthewsmawfield/TEP-EFA> with archived DOI at 10.5281/zenodo.19454862.

Raw DSN tracking products may be obtained from the NASA Deep Space Network and the Planetary Data System following institutional access procedures; per-mission pointers appear under `data/raw/dsn_tracking/<mission>/DOWNLOAD_INSTRUCTIONS.txt`. The present manuscript release does not bundle perigee-matched Level-1 TRK archives: Steps 005–006 and 028–031 implement ingest and audit only, and Step 030 remains inconclusive until such products are added. Headline flyby inference uses literature Δv and JPL Horizons trajectories (Steps 007–026, 039).