



Temporal Equivalence Principle: Native `hi_class` Conformal Implementation, Linear Perturbation Closure, and CMB Acoustic Peak Preservation

Matthew Lukin Smawfield

Version: v0.6 (Cambridge)

First published: 8 June 2026 · Last updated: 7 August 2026

DOI: 10.5281/zenodo.20682752

Code Availability: github.com/matthewsmawfield/TEP-HC

Abstract

Standard cosmology explains the Cosmic Microwave Background (CMB) acoustic peaks, the pre-recombination sound horizon, and the thermal scaling relevant to Big Bang Nucleosynthesis (BBN) within an FLRW expansion history conventionally extrapolated toward a physical singularity. This paper demonstrates that the CMB acoustic-sector and conformal thermal/sound-horizon scalings are preserved with high fidelity under a static conformal temporal-transport geometry governed by the Temporal Equivalence Principle (TEP).

In the TEP framework, matter clocks and photon phases evolve in a causal matter metric defined by a conformal scalar field $\tilde{g}_{\mu\nu} = A(\phi)^2 g_{\mu\nu}$. Because this conformal transport geometry is mathematically isomorphic to the FLRW scale factor $a(t)$, standard Boltzmann solvers like `hi_class` and `CLASS` can be used as conformal-frame calculators for the background/acoustic-sector mapping tested here. The parameter traditionally identified as Dark Energy (Ω_Λ) is operationally reinterpreted within this implementation as the homogeneous temporal-shear background contribution filling the same background budget slot, Ω_ϕ .

This paper implements the native TEP interpretation directly in `hi_class`. Within the broader TEP interpretation, by recognizing that the spatial metric does not stretch, the "Big Bang" is reinterpreted not as a physical density singularity, but as a TEP temporal horizon—an asymptotic boundary where the observational clock map $A_{\text{clock}} \rightarrow 0$. Direct Boltzmann integration verifies this background/acoustic mathematical isomorphism, verifying the internal conformal-frame preservation of the acoustic sector — the pre-recombination sound-horizon ratio is $r_s^{\text{mTEP}}/r_s^{\text{m}\Lambda\text{CDM}} = 0.999994$ (corresponding to a < 6 ppm deviation) and the acoustic-peak morphology remains intact without invoking early-universe spatial expansion.

Beyond the background mapping, the paper closes the linear pure-conformal scalar perturbation sector by deriving the runtime Bellini–Sawicki functions $\alpha_M = -2\alpha_A$, $\alpha_B = 2\alpha_A$, $\alpha_K = -5\alpha_A^2$, and $\alpha_T = 0$, for which the physical no-ghost discriminant satisfies $D = \alpha_K + \frac{3}{2}\alpha_B^2 = \alpha_A^2$. An active-perturbation `hi_class` run evolving $\delta\phi$ through the full Einstein–Boltzmann hierarchy produces posteriors statistically indistinguishable from the background-only chain, demonstrating that

the implemented linear pure-conformal scalar perturbation sector is stable and observationally negligible at the current homogeneous-amplitude bound.

A joint `hi_class` Cobaya MCMC (Planck 2018 low- ℓ TT/EE + lensing + BAO + Pantheon+) tests the screened TEP conformal background within the native `hi_class` implementation, while companion TEP-C0 (Paper 26) reports Pantheon+ nested-sampling evidence for the physical no- Λ temporal-shear branch: Bayes factor approximately 4.6 for the conservative $z_{\text{los}} = 5$ branch, approximately 61.8 for the fixed $z_{\text{los}} = 100$ benchmark, and approximately 40.3 for the broad free- z_{los} branch. These late-time model-comparison results are not re-derived in HC; HC supplies the native `hi_class` acoustic and perturbation closure. Within the broader TEP corpus, Paper 11 interprets the Hubble tension as a late-time, environment-dependent clock-transport effect caused by the mass-screening of the scalar field, rather than through a crisis in early-universe physics. HC does not independently re-analyse the distance-ladder data; it imports the interpretation.

Keywords: cosmology theory, cosmic microwave background, static conformal geometry, scalar-tensor theories, conformal gravity, `hi_class`, Horndeski, temporal equivalence principle, proper time, Cobaya, Planck 2018

1. Introduction

1.1 Contextualizing the TEP Corpus

The Temporal Equivalence Principle (TEP) has been constrained across many orders of magnitude in mass density, from terrestrial laboratory scales ($\rho \sim 20 \text{ g/cm}^3$) to the cosmological mean ($\rho \sim 10^{-29} \text{ g/cm}^3$). Previous papers in this series have established:

- *Terrestrial scales (Paper 1)*: Terrestrial atomic clock networks show 4,200 km phase correlations consistent with the candidate Temporal Topology saturation scale $\rho_T \approx 20 \text{ g/cm}^3$.
- *Galactic scales (Paper 6, UCD)*: SPARC rotation curves validate the potential-dependent proper-time mapping.
- *Stellar scales (Paper 13, WB)*: Gaia DR3 wide binaries exhibit the predicted environment-dependent kinematic transition.
- *Cosmological scales (Paper 12, JWST)*: High-redshift anomalies align with environment-dependent time dilation.

1.2 The Cosmological Horizon

The Hubble tension and the JWST high-redshift galaxy anomalies represent the two most persistent challenges in modern cosmology. Standard Λ CDM relies on a stretching spatial metric, which extrapolates to a physical singularity at $a \rightarrow 0$ and tightly restricts the available proper time for early galaxy assembly.

Previous TEP work (Paper 11, H_0 ; Paper 12, JWST) argued that these anomalies are resolved within the TEP framework. The TEP temporal-horizon picture replaces the FLRW singularity with an asymptotic boundary, removing the finite-age assembly bottleneck, and the H_0 tension is addressed as a local environmental mass-screening effect on kinematic distance probes.

1.3 Purpose of This Paper

To rigorously test the CMB acoustic-sector component of the Static Conformal thesis, this paper demonstrates that the acoustic-sector integrals can be reproduced by an exactly conformal temporal mapping without explicit spatial expansion. A full light-element abundance calculation is not performed in HC. TEP-TH supplies the nonsingular temporal-horizon geometry and recombination-visibility treatment, while TEP-BBN supplies the native chemical-evolution framework and local CMB-thermalization proof of concept. The present HC claim is limited to conformal thermal/sound-horizon scaling, acoustic preservation, and the native `hi_class` perturbation implementation.

Because the TEP conformal scalar field $\tilde{g}_{\mu\nu} = A(\phi)^2 g_{\mu\nu}$ is mathematically isomorphic to the FLRW scale factor $a(t)$, the TEP Static Conformal geometry can be natively evaluated by deploying the `hi_class` Boltzmann solver as a conformal-frame calculator. This requires:

1. Mapping the TEP conformal geometry onto the Boltzmann framework, establishing the implementation-level correspondence between the parameter conventionally written as Ω_Λ and the homogeneous temporal-shear background contribution, Ω_ϕ , occupying the same background-budget role as Ω_Λ .
2. Native implementation in `hi_class` to evaluate the conformal temporal shear field directly.
3. A joint MCMC parameter estimation ($H_0, \Omega_b h^2, \Omega_{\text{cdm}} h^2, n_s, A_s, \tau, A_{\text{planck}}, \epsilon_T$) against Planck 2018, BAO, and Pantheon+ data to quantitatively demonstrate that the acoustic peaks are preserved with sound-horizon ratio $r_s^{\text{TEP}} / r_s^{\text{ACDM}} = 0.999994$ (corresponding to a < 6 ppm deviation) in a static conformal geometry.

This work does not independently analyse Pantheon+ supernovae or derive the full nonsingular temporal-horizon closure; those are addressed in companion papers TEP-C0 (Paper 26) and TEP-TH (Paper 27).

The critical question: Can a static conformal geometry mathematically reproduce the CMB acoustic peaks?

The claim-discipline framework for the TEP corpus, including the scope limitations of canonical precision tests, is established in TEP-EXP (Paper 9).

2. Theoretical Architecture: The EFT Mapping

2.1 The Bi-Metric Action

The TEP framework posits that matter couples to a screened metric $\tilde{g}_{\mu\nu}$ related to the Einstein-frame metric $g_{\mu\nu}$ via a disformal transformation:

$$\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi \quad (1)$$

where:

- $A(\phi) = \exp(\beta_A\phi/M_{\text{Pl}})$ is the conformal factor, with $\beta_A = -1.0$ (the locked lab-scale convention used across the TEP corpus)
- $B(\phi)$ controls disformal deformation of the causal structure
- ϕ is the dynamical proper-time field

Following TEP-TH (Paper 27), two distinct projections of the temporal field are distinguished in cosmology: $A_{\text{clock}}(z)$ is the exact observational clock/redshift mapping that generates the apparent distance-redshift relation, while $A_{\text{dyn}}(z)$ is the physical dynamical response that modifies expansion and perturbations. In the homogeneous background limit evaluated here, the hi_class conformal-frame mapping corresponds to A_{clock} ; the dynamical response A_{dyn} is parameterized by ϵ_T and modifies the Hubble parameter and acoustic observables at late times without requiring early-universe thermal screening (TEP-TH, Paper 27; TEP-BBN, Paper 29).

Metric signature convention: $(+, -, -, -)$ throughout.

2.2 Formal Bellini-Sawicki Alpha Correspondence

hi_class requires the EFT property functions α_i that encode metric modifications at linear perturbation level.

2.2.1 Planck Mass Running (α_M)

The conformal coupling directly determines the running of the effective Planck mass:

$$\alpha_M \equiv \frac{d \ln M_{\text{eff}}^2}{d \ln a} = -\frac{d \ln A^2(\phi)}{d \ln a} = -\frac{2\beta_A}{M_{\text{Pl}}} \frac{\phi'}{\mathcal{H}} \quad (2)$$

where $\mathcal{H} = aH$ is the conformal Hubble parameter and primes denote derivatives with respect to conformal time.

2.2.2 Tensor Speed Excess (α_T)

The disformal term $B(\phi)$ alters the gravitational wave propagation speed. Multi-messenger constraints from GW170817/GRB 170817A require:

$$|c_g - c_\gamma|/c \lesssim 10^{-15} \Rightarrow \alpha_T \approx 0 \text{ (today)} \quad (3)$$

However, $B(\phi)$ may be non-zero at recombination ($z \approx 1100$) provided it relaxes to zero by $z \sim 0$. This statement concerns the general disformal EFT only; the production HC acoustic/perturbation branch analysed below is the pure-conformal ($\alpha_T = 0$) branch.

2.2.3 Braiding (α_B) and Kineticity (α_K)

These functions govern scalar field clustering and metric mixing:

$$\alpha_B = -\frac{\mathcal{H}'\phi'}{\mathcal{H}^2} \cdot f_B(\phi, X) \quad (4)$$

$$\alpha_K = \frac{\phi'^2}{\mathcal{H}^2 M_{\text{Pl}}^2} \cdot f_K(\phi, X) \quad (5)$$

where $X = -\nabla_\mu \phi \nabla^\mu \phi / 2$ and f_B, f_K are functions derived from the TEP action:

- $f_B(\phi, X)$ encodes the disformal coupling to the energy-momentum tensor trace.
- $f_K(\phi, X)$ encodes the kinetic term non-canonicity from the TEP proper-time field.

The explicit functional forms follow from the bi-metric action (Equation 1) and are determined by the conformal factor $A(\phi)$ and the disformal function $B(\phi)$. Their derivation is detailed in the TEP theoretical framework (Papers 1 and 11 of the TEP corpus). The schematic f_B, f_K forms describe the general disformal EFT bookkeeping; the production perturbation run restricts to the pure-conformal branch, where these functions reduce to the closed runtime identities given in Appendix A.5.

The background-only configuration uses α_B and α_K as formal EFT bookkeeping; during native `tep_mode` background-only integration f_B and f_K evaluate identically to zero, verifying that this is a strict geometric mapping at the background level. The active-perturbation closure configuration (Section 4.5) passes `gravity_model = tep` and `M2_evolution = yes` to `hi_class`, which evolves the implemented linear pure-conformal scalar fluctuation sector $\delta\phi$ through the exact Bellini–Sawicki EFT functions derived from the conformal geometry, with sound speed $c_s^2 = 1.0$ and no-ghost condition $D = \alpha_A^2$ enforced by construction. The theoretical basis for this closure is developed in the foundational TEP formalism (Papers 1 and 11).

2.3 The Static Conformal Isomorphism

The defining feature of the TEP framework is that it recasts the role normally played by a physically expanding spatial metric in the background/acoustic sector. In standard Λ CDM cosmology, the proper distance between co-moving galaxies physically increases over time, parameterized by the scale factor $a(t)$.

In the static conformal interpretation tested here, intergalactic separations are not treated as primitively expanding; the apparent expansion is reconstructed through temporal transport. The causal matter metric $\tilde{g}_{\mu\nu}$ is modulated by the conformal clock-rate field $A(\phi)$. The Temporal Equivalence Principle relies on this distinct, dynamical proper-time field and is fundamentally separate from the standard Einstein Equivalence Principle (EEP), which concerns the universality of free fall and local Lorentz invariance in metric theories of gravity. Photons propagating through this gradient experience a shift in phase and frequency, leading to the exact geometric relation:

$$1 + z = \frac{A_0}{A_{\text{em}}} = A_{\text{clock}}^{-1}(z) \quad (6)$$

The exact observational clock map $A_{\text{clock}}(z) = (1 + z)^{-1}$ defines the redshift–distance relation in the static conformal geometry. Because the mathematical transport geometry of A_{clock} across a static background is formally isomorphic to the transport geometry of a photon in an expanding FLRW metric with reconstructed scale factor $a_{\text{eff}}(z)$, standard cosmological integrators (like `hi_class` and `CLASS`) can be deployed natively as conformal-frame calculators for the background/acoustic-sector mapping tested here. The physical dynamical response $A_{\text{dyn}}(z)$ modifies the Hubble parameter and acoustic observables at late times (TEP-TH, Paper 27; TEP-BBN, Paper 29).

- The primary acoustic peaks ($100 \lesssim \ell \lesssim 2000$) generated at $z \sim 1089$ are preserved with extreme precision, because the mathematics of the acoustic horizon r_s depend only on the conformal integration path, not on physical spatial stretching.
- The parameter conventionally identified as Dark Energy (Ω_Λ) is reinterpreted within this implementation as the homogeneous temporal-shear background contribution, Ω_ϕ , occupying the same background-budget role as Ω_Λ in the reference FLRW calculation.
- The Big Bang singularity is reinterpreted, within TEP, as a temporal-horizon boundary of the conformal clock-rate field where $A_{\text{clock}} \rightarrow 0$ relative to the present epoch, driving the reconstructed scale factor $a_{\text{eff}} \rightarrow 0$ and creating the mathematical appearance of infinite density in standard FLRW reconstructions. The full nonsingular closure—finite curvature invariants, geodesic completeness, and the origin of the CMB blackbody spectrum—is developed in TEP-TH (Paper 27) and TEP-BBN (Paper 29), where the non-exact covariance/topology correction C_T provides the transport closure beyond exact conformal shear.
- *Thermodynamic Cooling*: The adiabatic cooling of the CMB photon gas ($T \propto 1/a$ in standard cosmology) is preserved as $T \propto 1/A$. The energy density shifts natively via the conformal temporal shear without requiring physical spatial volume dilution.

Operational definition of Ω_ϕ . In this implementation, Ω_ϕ is operationally defined as the homogeneous conformal-sector contribution that fills the same background budget slot occupied by Ω_Λ in the reference FLRW integration. A first-principles stress-energy

derivation of ρ_ϕ , p_ϕ , and the effective equation of state belongs to the broader TEP-C0 action-level treatment.

Archived EFT reference. The Bellini–Sawicki α_i functions mapped from the TEP bi-metric action (step-3 fiducial) are archived in `results/03_alpha_functions.json`. Production CMB constraints use the native conformal implementation (Section 4), evaluating the strict isomorphism directly without relying on linear-perturbation mapping approximations.

3. Software Implementation: `hi_class` and the Unscreened Regime

3.1 The `hi_class` Architecture

`hi_class` extends the CLASS Boltzmann solver to handle general scalar-tensor theories via the EFT formalism. This work uses `hi_class` v3.2.3 with the modified gravity (SMG) module enabled.

3.2 Native TEP Conformal Background Implementation

The native TEP background-only Hubble modification is implemented directly in `hi_class` via the `tep_mode` flag. When enabled, the background expansion history is modified as:

$$H_{\text{TEP}}(z) = H_{\text{ACDM}}(z) \times M(z), \quad M(z) = \frac{A(z)}{1 - \alpha_A(z)} \quad (7)$$

where $S(z) = \exp[-(z/z_T)^{n_T}]$ is the redshift suppression factor, $A(z) = \exp[\epsilon_T \ln(1+z) S(z)]$ is the covariant conformal factor, and $\alpha_A = -d \ln A / d \ln(1+z)$. The production integration directly evaluates the exact geometric relation $M = A/(1 - \alpha_A)$, ensuring computational fidelity to the underlying formal derivation (see Appendix A.3a). The transition function $f_T(z) = \ln(1+z) S(z)$ appearing in the exponent is the shared TEP-C0 implementation (Paper 26; `core/cosmology.py`: `f_T`, `conformal_factor_native`, `jordan_frame_M`):

$$f_T(z) = \ln(1+z) S(z). \quad (8)$$

The suppression factor $S(z)$ defines the physical profile of the conformal field: it drives $f_T \rightarrow 0$ for $z \gg z_T$, ensuring the field profile flattens asymptotically as it approaches the TEP temporal horizon; the $\ln(1+z)$ factor enforces $f_T(0) = 0$, fixing the local reference frame so H_0 is anchored to the local observer. The function peaks at intermediate redshift ($z \sim z_T$), where the effective homogeneous temporal-shear contribution mimics apparent acceleration, and flattens out in the deep past.

Implementation note: an earlier development build used the *complement* $f_T = 1 - \exp[-(z/z_T)^{n_T}]$, which instead saturates to unity for $z \gg z_T$. This inverted the scalar field profile and corrupted the acoustic peak evaluation (see Appendix A.3). The default TEP conformal parameters are:

```
tep_mode = yes
epsilon_T = 0.0066
z_T = 5.0
n_T = 2.0
```

Parameter-Scale and Amplitude Convention

Turnover scales. z_T^{HC} denotes the homogeneous/acoustic `hi_class` profile scale used here. z_T^{los} denotes the C0 line-of-sight supernova transport turnover. These scales are related projections of the temporal sector but are not numerically interchangeable.

Amplitudes. ϵ_T^{HC} denotes the native `hi_class` homogeneous conformal amplitude reported here (0.00602 ± 0.00493). ϵ_T^{los} denotes the late-time line-of-sight transport amplitude fitted in TEP-C0. ϵ_T^{CMB} denotes the C0 background/acoustic diagnostic amplitude. ϵ_{dyn} denotes the dynamical temporal-horizon response in TEP-TH, while $\epsilon_{\text{field}} = 0.0175$ denotes the primordial spectral-flow parameter constrained by n_s in TEP-TH. These are related projections of the same temporal sector, but they are not numerically interchangeable parameters.

The background conformal mapping is implemented through $M(z) = A/(1 - \alpha_A)$. The associated linear scalar perturbation sector is closed separately through the Bellini–Sawicki runtime functions described in Section 3.3 and Appendix A.5, and validated in the

active $\delta\phi$ run of Section 4.5. This implementation is the `hi_class` analogue of the CLASS native TEP module used in TEP-C0 (Paper 26).

3.3 Perturbation Stability and Closure

The present analysis natively closes the scalar perturbation sector by evaluating the exact runtime Bellini-Sawicki Effective Field Theory (EFT) parameters in the pure conformal limit. While early formulations of the TEP acoustic mapping relied on the working assumption that scalar spatial fluctuations ($\delta\phi$) decouple or are heavily suppressed at recombination, the implementation developed here leverages the `hi_class` SMG framework to evolve the full Einstein-Boltzmann hierarchy actively.

Because the TEP framework maps its causal matter metric via the exact conformal relation $\tilde{g}_{\mu\nu} = A(\phi)^2 g_{\mu\nu}$, the corresponding EFT functions (α_i) in the Jordan frame are analytically fixed by the background derivative $\alpha_A \equiv -d \ln A / d \ln(1+z)$. In the pure conformal limit ($\beta_A = -1.0$), the kinematicity parameter evaluates to $\alpha_K = -5\alpha_A^2$. Although a negative kinematicity frequently triggers ghost instabilities in canonical scalar-tensor theories, the full physical no-ghost discriminant D in the Horndeski framework encompasses the braiding parameter, which in this limit is identically $\alpha_B = 2\alpha_A$.

Evaluating the full discriminant yields $D = \alpha_K + \frac{3}{2}\alpha_B^2 = \alpha_A^2$. The no-ghost discriminant remains positive-definite ($D = \alpha_A^2 > 0$) for all redshifts $0 < z < 1100$ and for ϵ_T values within the 95% posterior from the MCMC run ($0 < \epsilon_T \lesssim 0.016$). Across the full sampled posterior, $D_{\min} = 1.7 \times 10^{-19}$ and $|\alpha_M|_{\max} = 0.041$, confirming that the scalar sector is ghost-free and gradient-stable throughout the observationally viable parameter space.

ϵ_T (95% range)	$D_{\min}$	$ \alpha_M _{\max}$
3.4×10^{-6}	1.7×10^{-19}	9.0×10^{-6}
0.007	1.0×10^{-12}	0.018
0.016	4.0×10^{-12}	0.041

By forcing the integration of the implemented linear pure-conformal $\delta\phi$ scalar field through this exact geometric relation, the numerical solver verifies that the continuous screening transition smoothly manages the emergence into the late universe without triggering gradient instabilities or pathological phantom energies. At the fiducial amplitude ($\epsilon_T = 0.0066$), integrating the active perturbations yields an Integrated Sachs-Wolfe (ISW) residual of less than 0.001% across the entire acoustic spectrum ($100 \leq l \leq 2000$). The background acoustic isomorphism is therefore preserved to the reported numerical precision under active linear perturbation evolution, validating that the active linear scalar fluctuations in the temporal shear field remain stable and do not distort the CMB damping tail.

3.3b Covariant Frame Alignment: Matter-Frame Hubble Friction

The TEP action defines the causal metric for matter particles as $\tilde{g}_{\mu\nu} = A(\phi)^2 g_{\mu\nu}$. Under this conformal rescaling, the matter-frame conformal Hubble rate acquires the exact relation

$$\tilde{\mathcal{H}} = \mathcal{H} - \frac{\dot{A}}{A} = \mathcal{H}(1 - \alpha_A), \quad (9)$$

where $\alpha_A \equiv -d \ln A / d \ln(1+z)$. In the Jordan-frame EFT, matter particles follow geodesics of $\tilde{g}_{\mu\nu}$, so their Euler equations must be evaluated with $\tilde{\mathcal{H}}$ rather than the Einstein-frame rate $\mathcal{H}$. The standard `hi_class` implementation uses $\mathcal{H}$ for all species, which is appropriate for Einstein-frame observers (photons and neutrinos) but does not include the conformal-frame distinction for collisionless matter. We implement this distinction as follows.

A new background index `index_bg_H_conformal` is defined in `include/background.h` and populated in `source/background.c` after the TEP Hubble modification:

```
/* TEP matter-frame conformal Hubble rate */
if (pba->tep_mode == TRUE && pba->epsilon_T != 0.0) {
    double z = 1.0/a - 1.0;
    double alpha_A = tep_alpha_A(pba, z);
    pvecback[pba->index_bg_H_conformal] = pvecback[pba->index_bg_H] * (1.0 - alpha_A);
}
else {
    pvecback[pba->index_bg_H_conformal] = pvecback[pba->index_bg_H];
}
```

In `source/perturbations.c`, we compute a new local variable `a_prime_over_a_matter = pvecback[index_bg_H_conformal] * a` and substitute it for `a_prime_over_a` in the Euler friction terms of CDM, baryons, interacting dark matter, and decaying cold dark matter. Because massless propagation is conformally invariant, photons natively retain the Einstein-frame rate; this is a direct consequence of the universal matter coupling rather than an exception to it. For neutrinos, retaining the Einstein-frame rate is an approximation strictly valid only while they remain highly relativistic. The metric equations naturally retain the Einstein-frame rate, preserving the CMB acoustic physics. The modification is minimal and upstream: a single background quantity and a single local variable replacement in the fluid velocity derivatives.

3.3c Scale-Dependent Bellini-Sawicki Mapping

The gradient-screening envelope $f(g) = [1 + (g/g_t)^n]^{-1}$ governs the suppression of the temporal-shear coupling in regions of steep gravitational potential. In the cosmological context, the gradient operator translates to the Fourier wavenumber k through the Poisson equation $\nabla^2 \Phi \sim k^2 \Phi_k$, so the envelope becomes

$$\mathcal{S}_\Sigma(k) = \frac{(k/k_t)^{n_t}}{1 + (k/k_t)^{n_t}}, \quad (10)$$

where k_t is the screening threshold scale and n_t the steepness. The Bellini-Sawicki parameter α_M , which controls the running of the effective Planck mass and therefore the strength of the scalar fifth force, must acquire this k -dependence:

$$\alpha_M(k, a) = \alpha_M^{\text{bg}}(a) \times [1 - \mathcal{S}_\Sigma(k)]. \quad (11)$$

This is implemented by modifying the function `get_gravity_coefficients_smg` in `gravity_smg/perturbations_smg.c` to accept the mode wavenumber k as an argument. The screening is applied to the local `run` variable (carrying α_M) immediately after it is extracted from the background workspace:

```
if (pba->tep_mode == TRUE && pba->tep_k_t > 0.0 && k > 0.0) {
    double S_Sigma = tep_screening_factor(k, pba->tep_k_t, pba->tep_n_t);
    double alpha_M_bg = *run;
    if (alpha_M_bg != 0.0) {
        *run = alpha_M_bg * (1.0 - S_Sigma);
    }
}
```

The screening function `tep_screening_factor` is defined in `source/background.c` and exposed in `include/background.h`. Two new parameters, `tep_k_t` and `tep_n_t`, are added to the `background` structure and parsed in `source/input.c`. The function signature of `get_gravity_coefficients_smg` is updated in `gravity_smg/include/perturbations_smg.h`, and all three call sites (in `perturbations_einstein_scalar_smg`, `perturbations_qs_functions_at_tau_and_k_qs_smg`, and `get_x_x_prime_qs_smg`) pass the mode wavenumber k .

At cosmic scales ($k \rightarrow 0$), $\mathcal{S}_\Sigma \rightarrow 0$ and $\alpha_M(k) \rightarrow \alpha_M^{\text{bg}}$, preserving the background expansion. At structure scales ($k \gg k_t$), $\mathcal{S}_\Sigma \rightarrow 1$ and $\alpha_M(k) \rightarrow 0$, stripping the scalar fifth force and yielding $G_{\text{eff}} \rightarrow G_N$ inside collapsing halos. Both the conformal friction and the scale-dependent screening use only the background parameters already constrained by the distance data (ϵ_T, z_T, n_T); k_t and n_t are physical parameters of the gradient envelope with no additional tuning required for σ_8 .

3.4 Pipeline Architecture

The full analysis pipeline, executed via `scripts/run_all.py`, consists of:

1. *Step 0 (Setup)*: Environment configuration and dependency check.
2. *Step 1 (Install)*: Install Cobaya, Planck 2018 likelihoods, and `hi_class` with the native TEP patch (`external/patches/hiclass_tep_native.patch`).
3. *Step 2 (Background)*: Compute the TEP-modified background expansion history $H(z)$ and density evolution.
4. *Step 3 (Alpha Functions)*: Compute Bellini-Sawicki coefficients from the TEP theoretical mapping (archived for reference).
5. *Step 4 (CMB Spectra)*: Run `hi_class` with native `tep_mode` at the Planck 2018 best-fit point. Compare TT, TE, and EE spectra against standard Λ CDM.
6. *Step 5 (Matter-Frame Scan)*: Dual-scan reconstruction of the acoustic scale in screened and unscreened limits.
7. *Step 6 (Cobaya Config)*: Generate the Cobaya YAML configuration for the MCMC pipeline with native TEP parameters.
8. *Step 7 (MCMC)*: Execute the Cobaya MCMC with `hi_class`, using real Planck + BAO + Pantheon+ likelihoods, for both the background-only configuration and the active-perturbation closure configuration (`gravity_model = tep`).

9. *Step 8 (Posteriors)*: Analyze MCMC chains with burn-in removal and weighted statistics.

10. *Step 9 (Synthesis)*: Combine all results into summary JSON and markdown.

Publication figures are generated separately via `python scripts/generate_figures.py` (not part of `run_all.py`). Both figures are written to `results/figures/` with filenames matching their publication numbering. Figure 3 requires step 04b. Include them in the static site with `cd site && npm run build`.

The perturbation-level screening implemented via the effective α -functions in the Boltzmann hierarchy represents the linear cosmological realization of the abstract environmental operator $\mathcal{S}_{\Sigma}(\mathcal{E})$. The continuous running of these parameters dynamically enforces the spatial flattening of the Temporal Topology across structure formation epochs without committing to a specific microphysical scalar potential.

4. MCMC Parameter Estimation Pipeline

4.1 The Cobaya Framework

Cobaya provides a Python interface to CLASS/hi_class with extensive MCMC sampling capabilities. The transition from SciPy/Pandas pipelines to Cobaya enables:

- Native hi_class integration without file-based I/O bottlenecks
- Parallel tempering and adaptive Metropolis-Hastings sampling
- Direct Planck likelihood wrapper integration
- Seamless GetDist posterior visualization

4.2 Likelihood Configuration

The pipeline uses the following Planck 2018 likelihoods:

Likelihood	Description	ℓ Range
<code>planck_2018_lowl.TT</code>	Low- ℓ temperature	2–29
<code>planck_2018_lowl.EE</code>	Low- ℓ polarization	2–29
<code>planck_2018_lensing.native</code>	CMB lensing reconstruction	8–400
<code>bao.sdss_dr12_consensus_final</code>	BAO SDSS DR12 consensus	—
<code>sn.pantheonplus</code>	Type Ia supernovae (Pantheon+)	—

4.3 Free Parameters and Priors

The MCMC pipeline samples standard Λ CDM parameters alongside the TEP amplitude parameter ϵ_T :

Parameter	Prior	Description
$\Omega_b h^2$	$\mathcal{U}(0.005, 0.1)$	Baryon density
$\Omega_{\text{cdm}} h^2$	$\mathcal{U}(0.01, 0.99)$	Cold dark matter density
H_0	$\mathcal{U}(40, 100)$	Hubble constant
τ	$\mathcal{U}(0.01, 0.8)$	Optical depth
A_s	$\mathcal{U}(10^{-10}, 5 \times 10^{-9})$	Scalar amplitude
n_s	$\mathcal{U}(0.94, 1.0)$	Scalar spectral index
A_{planck}	$\mathcal{U}(0.9, 1.1)$	Planck calibration nuisance
ϵ_T	$\mathcal{U}(-1, 1)$	TEP amplitude parameter (background Hubble modification)

4.4 Pipeline Execution

```
# Cobaya YAML configuration
theory:
  class:
    path: /path/to/hi_class
```



```

extra_args:
  output: tCl,pCl,lCl,mPk
  lensing: yes
  modes: s,t
  non_linear: halofit
  # Native TEP background-only Hubble modification
  tep_mode: 'yes'
  z_T: 5.0
  n_T: 2.0
  # epsilon_T is sampled in params below — do not duplicate here

likelihood:
  planck_2018_lowl.TT: null
  planck_2018_lowl.EE: null
  planck_2018_lensing.native: null
  bao.sdss_dr12_consensus_final: null
  sn.pantheonplus: null

params:
  logA:
    prior: {min: 2.5, max: 3.5}
    ref: {dist: norm, loc: 3.044, scale: 0.014}
    proposal: 0.01
    drop: true
  A_s:
    value: 'lambda logA: 1e-10*np.exp(logA)'
  n_s:
    prior: {min: 0.94, max: 1.0}
    ref: {dist: norm, loc: 0.966, scale: 0.004}
    proposal: 0.004
  H0:
    prior: {min: 40, max: 100}
    ref: {dist: norm, loc: 67.4, scale: 0.5}
    proposal: 1.5
  omega_b:
    prior: {min: 0.005, max: 0.1}
    ref: {dist: norm, loc: 0.0224, scale: 0.0002}
    proposal: 0.0003
  omega_cdm:
    prior: {min: 0.01, max: 0.99}
    ref: {dist: norm, loc: 0.12, scale: 0.001}
    proposal: 0.0015
  tau_reio:
    prior: {min: 0.01, max: 0.8}
    ref: {dist: norm, loc: 0.054, scale: 0.007}
    proposal: 0.01
  A_planck:
    prior: {min: 0.9, max: 1.1}
    ref: {dist: norm, loc: 1.0, scale: 0.0025}
    proposal: 0.005
  epsilon_T:
    prior: {min: -1.0, max: 1.0}
    ref: {dist: norm, loc: 0.006, scale: 0.005}
    proposal: 0.0005
    latex: '\epsilon_T'
  sigma8:
    latex: '\sigma_8'

sampler:
  mcmc:
    burn_in: 0
    max_tries: 10000
    max_samples: 500000
    Rminus1_stop: 0.05
    Rminus1_cl_stop: 0.2
    output_every: 10
    drag: true
    seed: 42

```

4.5 Perturbation-Mode Validation

In addition to the background-only MCMC configuration above, the pipeline includes an active perturbation closure configuration (`data/cobaya/tep_hiclass_perturbations.yaml`) that passes `gravity_model = tep` and `M2_evolution = yes` to the `hi_class` SMG module. This forces the Boltzmann solver to evolve the implemented linear pure-conformal scalar fluctuation sector $\delta\phi$ through the exact Bellini-Sawicki EFT functions derived from the conformal geometry (Appendix A.5).

A 4-chain MPI MCMC with this configuration (58,180 accepted steps; Planck 2018 low- ℓ TT/EE + lensing + BAO + Pantheon+; cross-chain Gelman–Rubin $R - 1 = 0.007$ at termination) ran successfully with finite posterior at every step. The resulting parameter constraints are:

$$\epsilon_T = 0.00547 \pm 0.00429, \quad (12)$$

with $H_0 = 66.77 \pm 1.73$ km/s/Mpc, $n_s = 0.9956 \pm 0.0043$, $\Omega_b h^2 = 0.02144 \pm 0.00257$, $\Omega_{\text{cdm}} h^2 = 0.1155 \pm 0.0042$, $\tau = 0.0497 \pm 0.0075$, $A_{\text{planck}} = 1.088 \pm 0.013$, and $S_8 = 0.868 \pm 0.025$. Because this implementation run uses low- ℓ TT/EE plus lensing rather than the full high- ℓ TTTEEE acoustic likelihood, the n_s posterior should be interpreted as a robustness diagnostic rather than as a final spectral-index constraint. Direct comparison with the background-only chain ($\epsilon_T = 0.00602 \pm 0.00493$) yields $\Delta\epsilon_T = -0.00055$ (-0.08σ), $\Delta H_0 = +0.09$ km/s/Mpc ($+0.04\sigma$), $\Delta n_s = +0.00009$ ($+0.02\sigma$), and $\Delta S_8 = +0.0009$ ($+0.03\sigma$). The maximum parameter disagreement across all eight cosmological parameters is 0.07σ , and $\Delta\chi^2 = -0.46$. This confirms that the late-time ISW contribution from the dynamical scalar field is observationally negligible at the current bound, and that the ϵ_T posterior is driven by background acoustic-peak shifts rather than by perturbation-sector physics. Figure 1 shows the background versus active-perturbation posterior comparison.




Figure 1. Marginalized posterior triangle plot from the 4-chain MPI run. Native `hi_class tep_mode` with $z_{\text{HC}} = 5.0$, $n_T = 2.0$. **Blue:** background-only TEP chain with $\delta\phi$ frozen. **Red:** active-perturbation TEP chain with $\delta\phi$ evolved through the implemented Bellini–Sawicki EFT sector ($\alpha_M = -2\alpha_A$, $\alpha_B = 2\alpha_A$, $\alpha_K = -5\alpha_A^2$, $\alpha_T = 0$). Parameters shown: ϵ_T^{HC} (native `hi_class` homogeneous conformal amplitude), H_0 , n_s , σ_8 . Contours show 68% and 95% credible regions. The near-complete overlap demonstrates that active scalar perturbations are observationally negligible at the current homogeneous-amplitude bound. Maximum posterior shift: $\lesssim 0.07\sigma$ across all eight cosmological parameters. Cross-chain Gelman–Rubin $R - 1 = 0.007$ at 58,180 accepted steps.

The `hi_class` configuration uses native `tep_mode` with the transition function $f_T(z) = \ln(1+z) \exp[-(z/z_T)^{n_T}]$ and fixed `z_T = 5.0`, `n_T = 2.0`, with `epsilon_T` sampled freely in `params`. This configuration natively explores the parameter space of the static conformal field, leveraging the strict isomorphism to evaluate the acoustic physics exactly. The production configuration is `data/cobaya/tep_hiclass_suite.yaml` (reference alternate: `data/cobaya/tep_native_mcmc.yaml`).

Pipeline status. The native-`tep_mode` joint MCMC against Planck 2018 low- ℓ TT/EE + lensing + BAO (SDSS DR12) + Pantheon+ was run using the structurally corrected `hi_class` engine, allowing Ω_Λ to natively fill the background cosmological budget. The primary production chain (`tep_hiclass_suite`; 18,720 accepted steps, 13,104 post-burn-in with 30% burn-in discard; single chain, so Gelman–Rubin $R-1$ is undefined; the sampler-internal $R-1$ reported by Cobaya reached 0.045 at termination) gives a Λ CDM-compatible background while measuring the TEP amplitude parameter:

$$\epsilon_T = 0.00602 \pm 0.00493, \quad (13)$$

with $H_0 = 66.68 \pm 1.82$ km/s/Mpc, $\Omega_b h^2 = 0.0214 \pm 0.0027$, $\Omega_{\text{cdm}} h^2 = 0.1150 \pm 0.0042$, $\tau = 0.050 \pm 0.008$, $A_{\text{planck}} = 1.088 \pm 0.012$, and $S_8 = 0.867 \pm 0.026$. The result is consistent with the TEP dual-domain expectation: the homogeneous amplitude ϵ_T remains small ($\sim 10^{-3}$) on the largest scales, where the CMB bound from TEP-C0 (Paper 26) is much tighter. This recombination-era value is expected to be smaller than the primordial temporal-horizon amplitude ($\epsilon_t = 0.0175$) derived in TEP-TH (Paper 27), because the temporal field evolves and is progressively screened as the universe cools.

Multi-chain validation. A parallel 4-chain run (`tep_native`; configuration `data/cobaya/tep_native_mcmc.yaml`) using a Gaussian A_{planck} prior (loc = 1.0, scale = 0.0025) was executed. After an initial short run (2,993 post-burn-in samples; $R-1$ for $\epsilon_T = 0.098$), the chain was extended to 8,960 accepted samples (max_samples increased from 500k to 1M), reaching convergence at $R-1 = 0.032$ (parameter-level) and $R-1 = 0.098$ (class-level). The converged extended chain yields $\epsilon_T = 0.0066 \pm 0.0049$ and $H_0 = 67.15 \pm 1.36$ km/s/Mpc, consistent with the primary chain at 0.10σ and 0.24σ respectively. The widened-prior chain (below) provides a fully converged multi-chain determination ($\epsilon_T R-1 = 0.013$, all parameters $R-1 < 0.05$), and the two agree to 0.31σ . Together they confirm the single-chain result is not an artefact of the sampling configuration.

High- ℓ TTTEEE attempt. A dedicated production run adding `planck_2018_highl_plik.TTTEEE` to the baseline likelihood was attempted. The `tep_mode`-modified `hi_class` produces CMB power spectra that trigger numerical overflow (divide by zero, invalid value) in the bundled `clipy` Python interface during the high- ℓ binning operation (`cmbonly.py: matmul`). Both the full `plik` and the `plik_lite` variants exhibit the same failure. This is a pre-existing incompatibility between the `tep_mode`-patched `hi_class` build and the Planck `clipy` high- ℓ interface, not a statistical or convergence issue.

To circumvent this blockage, a `CAMB`-based TEP approximation was implemented. Because `CAMB` lacks a native `tep_mode`, the TEP Jordan-frame Hubble rate $H_{\text{TEP}}(z) = H_{\Lambda\text{CDM}}(z) \cdot M(z)$ (with $M(z) = A(z)/(1 - \alpha_A)$) was mapped onto an effective dark-energy equation of state $w_{\text{eff}}(a)$ that reproduces the same background expansion. The effective $w(a)$ is fed into `CAMB` via `DarkEnergyPPF.set_w_a_table`, which supports phantom crossing. Exploratory testing confirms that `CAMB` generates clean, numerically stable high- ℓ spectra with this prescription: no NaN or Inf are produced up to $\ell = 2508$, and the `plik_lite` likelihood evaluates successfully. A grid scan over $\epsilon_T \in [0, 0.02]$ (fixed cosmology $H_0 = 66.63$, $\Omega_b h^2 = 0.0212$, $\Omega_{\text{cdm}} h^2 = 0.1154$, $\tau = 0.049$, $A_s = 2.1 \times 10^{-9}$, $n_s = 0.965$) gives a parabolic χ^2 surface with minimum at $\epsilon_T \approx 0.010$ and $\Delta\chi^2 \approx -260$ relative to ΛCDM ($\epsilon_T = 0$). This best-fit value lies within $\sim 1\sigma$ of the low- ℓ `hi_class` posterior ($\epsilon_T = 0.00602 \pm 0.00493$), confirming consistency across acoustic scales. Because the $w(a)$ mapping is an approximation to the true conformal transformation (perturbation-sector differences are not captured exactly), this high- ℓ diagnostic is reported as a consistency check. The native `hi_class tep_mode` high- ℓ TTTEEE production likelihood remains pending. The low- ℓ + lensing + BAO + Pantheon+ constraint therefore remains the primary acoustic result of this paper. A future definitive high- ℓ production run will require either a patched `clipy` wrapper, a native `CAMB tep_mode` implementation, or an alternate high- ℓ likelihood (e.g. ACT DR6, SPT).

Planck calibration prior sensitivity. The nuisance parameter A_{planck} (absolute CMB calibration) is implemented as a hard uniform prior on $[0.9, 1.1]$ in the primary chain. The posterior mean is $A_{\text{planck}} = 1.088 \pm 0.012$ with maximum sampled value 1.1000, indicating saturation against the upper prior bound. To test whether this truncation biases the cosmological inference, a dedicated sensitivity test was executed with the prior widened to $[0.9, 1.25]$ (Step 20: `scripts/steps/step_20_aplanck_sensitivity.py`, configuration `data/cobaya/tep_hiclass_aplanck_sens.yaml`). The converged run (16,321 total samples; all parameters Gelman–Rubin $R-1 < 0.05$; maximum $R-1 = 0.036$) yields $A_{\text{planck}} = 1.229 \pm 0.026$, confirming the old posterior was truncated by approximately 3.0σ . The TEP amplitude from the widened run is $\epsilon_T = 0.0063 \pm 0.0048$ ($R-1 = 0.036$), consistent with the primary chain at 0.33σ and with the multi-chain validation at 0.40σ . The correlation between A_{planck} and ϵ_T is $r = -0.19$, and splitting at $A_{\text{planck}} = 1.15$ gives a difference in ϵ_T of only -0.21σ . Even a 0.1 upward shift in A_{planck} would move H_0 by only ~ 0.2 km s $^{-1}$ Mpc $^{-1}$, well below its posterior width. The χ^2 does decrease monotonically toward the old boundary, but there is no evidence of a degeneracy cascade with ϵ_T . The TEP constraint on the homogeneous amplitude is robust against A_{planck} prior systematics. The widened- A_{planck} run is used only as a numerical robustness test of the ϵ_T posterior; the resulting calibration value should not be interpreted as a physically preferred Planck calibration model.

The companion paper TEP-C0 (Paper 26) provides the primary late-time constraints: Pantheon+ nested sampling favors the physical no- Λ TEP M1 branch over baseline Λ CDM with BF approximately 4.6 (conservative $z_{\text{los}} = 5$), approximately 61.8 (fixed $z_{\text{los}} = 100$ benchmark), and approximately 40.3 (broad free- z_{los}). Those model-comparison results are not re-derived here; they are used as the late-time empirical context for the `hi_class` acoustic-preservation implementation.

5. Results and Cosmological Constraints

5.1 The Acoustic Spectra

The physically meaningful test of the native TEP integration is to evaluate whether the conformal field exactly replicates the acoustic physics of the early universe without invoking spatial expansion. Because the conformal field mathematically mimics the FLRW scale factor, the recombination-era physics is evaluated natively within the static frame. Throughout this paper, "exact isomorphism" refers to the conformal background/acoustic mapping: the equality of the relevant sound-horizon and photon-transport integrals under the identification of the TEP conformal factor with the FLRW scale factor. The associated linear pure-conformal scalar perturbation sector is closed separately through the runtime Bellini–Sawicki EFT functions and validated by the active $(\delta\phi)$ `hi_class` run reported in Section 4.5 and Appendix A.5. This perturbative closure applies to the pure-conformal sector implemented here; the fully disformal, nonlinear, and environmentally inhomogeneous screening sectors remain extensions of the present calculation.

5.1.1 Sound-horizon and acoustic-peak preservation

Running `hi_class` native `tep_mode` against standard CLASS Λ CDM at the Planck 2018 best-fit point, with $\epsilon_T = 0.0066$, $z_T = 5$, $n_T = 2$, yields:

- *Sound horizon preserved to ~ 6 ppm:* $r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}} = 0.999994$. The comoving sound horizon integrates identically in the static conformal frame. The remaining ~ 6 ppm offset is a numerical/implementation-level residual associated with the finite precision of the conformal-frame background mapping and output reconstruction. Analytically, the exact mapping used in the production implementation is $M = A/(1 - \alpha_A)$, whose first-order expansion is $A(1 + \alpha_A)$. Direct verification confirms that the discrepancy is not a failure of the conformal isomorphism.
- *Acoustic-peak morphology unchanged:* with r_s , the baryon loading, and the photon-baryon driving at $z \approx 1089$ all operating identically under the conformal clock-rate, the relative peak heights and the damping tail closely match Λ CDM.

The central result is therefore not that all cosmological observables are already closed, but that the CMB acoustic scale itself is not uniquely diagnostic of physical spatial expansion.

5.1.1b Acoustic-scale preservation metrics

Table 2 quantifies the acoustic-scale preservation in three multipole ranges. The sound-horizon ratio $r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}} = 0.999994$ is independent of multipole because r_s is a single integrated quantity; the ℓ -centroid shift tracks the expected -0.185% angular-diameter-distance projection, and the active-perturbation residual remains below 1.9% across the acoustic range.

Multipole range	ℓ_{peak} shift $\Delta\ell$	$r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}}$	Max residual (%)
$100 \leq \ell \leq 500$	-0.06	0.999994	0.67
$500 \leq \ell \leq 1000$	-0.36	0.999994	1.34
$1000 \leq \ell \leq 2000$	-0.64	0.999994	1.82

5.1.2 The residual is a late-time projection, largely degenerate with H_0

The field profile is active over intermediate redshift ($z \sim 1\text{--}15$, peaking near z_T), changing the apparent angular distance to last scattering. At the fiducial $\epsilon_T = 0.0066$ this shifts the angular acoustic scale by $\Delta\theta_s/\theta_s = +0.185\%$. This rigid rescaling produces a coherent, oscillatory $\Delta C_\ell/C_\ell$ pattern whose envelope reaches $\sim 1.8\%$ across $100 < \ell < 2000$. This is *not* a change in early-universe physics: it is a pure angular-diameter-distance projection, largely degenerate with H_0 .

5.1.3 Polarization Spectra (C_ℓ^{TE} , C_ℓ^{EE})

The TE and EE spectra inherit the same exact behavior: the recombination-era polarization source is natively preserved by the conformal integration, and the only effect is the common θ_s projection shared with TT. Running `hi_class` with active scalar perturbations ($\delta\phi$ evolved through the exact Bellini–Sawicki EFT) and comparing against the background-only TEP run and Planck 2018 Λ CDM, the maximum active-perturbation residuals in the polarization channels are:

Multipole range	Max TE residual (%)	Max EE residual (%)
$100 \leq \ell \leq 500$	1.16	1.62
$500 \leq \ell \leq 1000$	2.14	2.46
$1000 \leq \ell \leq 2000$	3.19	3.62

These residuals are consistent with the pure angular-diameter-distance projection already documented for TT; there is no additional polarization-specific distortion from the dynamical scalar sector at the current precision.

5.2 Cosmological Constraints: Late-Time Evidence and the CMB Bound

The cosmological constraints on TEP come from two complementary regimes, established in the companion paper TEP-C0 (Paper 26).

Late-time evidence (supernovae). A nested-sampling model comparison over the full 1701×1701 Pantheon+ statistical-plus-systematic covariance finds substantial Bayesian preference for the TEP geometry over Λ CDM:

Model	Bayes factor vs Λ CDM	Interpretation
TEP M1 fixed ($z_T = 100$)	~ 32.1	Strong
TEP M1 free ($z_T \in [0.1, 150]$)	~ 36.2	Strong
w CDM	~ 21.1	Strong
CPL ($w_0 w_a$)	~ 35.2	Strong
Einstein-de Sitter	4.3×10^{-126}	Rejected (sanity check)

The TEP M1 branch improves the Pantheon+ likelihood relative to baseline Λ CDM with BF approximately 4.6 (conservative $z_{\text{los}} = 5$), approximately 61.8 (fixed $z_{\text{los}} = 100$ benchmark), and approximately 40.3 (broad free- z_{los}) (TEP-C0, Paper 26). Those model-comparison results are not re-derived here; they are used as the late-time empirical context for the `hi_class` acoustic-preservation implementation. The model-comparison result is consistent with the TEP claim that the Etherington distance-duality relation is a mathematically native feature of the static conformal field. TEP shows that the supernova distance-redshift relation can be fit without treating late-time acceleration as primitive spatial acceleration.

z_T distinction. The $z_T = 5$ profile used in this paper is the homogeneous acoustic-sector benchmark for the `hi_class` conformal implementation. It should not be confused with the C0 supernova-sector transport benchmark ($z_T^{\text{los}} = 100$), where z_T^{los} parameterizes the effective line-of-sight temporal-shear transition in the Pantheon+ distance law. The C0 free- z_T robustness test uses $z_T \in [0.1, 150]$.

Amplitude dictionary. ϵ_T^{los} denotes the late-time line-of-sight transport amplitude fitted in TEP-C0. ϵ_T^{CMB} denotes the C0 background/acoustic diagnostic amplitude. ϵ_T^{HC} denotes the native `hi_class` homogeneous conformal amplitude reported here (0.00602 ± 0.00493). ϵ_{dyn} denotes the dynamical response in TEP-TH. $\epsilon_{\text{field}} = 0.0175$ denotes the primordial spectral-flow parameter constrained by n_s in TEP-TH. These are related projections of the same temporal sector, but they are not numerically interchangeable.

Homogeneous (CMB) bound. The low- ℓ Planck likelihoods used in this paper's `hi_class` MCMC (TT/EE + lensing) yield $\epsilon_T = 0.00602 \pm 0.00493$, consistent with zero at $\sim 1.2\sigma$. The homogeneous CMB amplitude and the primordial spectral-flow parameter are different observational projections of the same temporal sector and are not numerically interchangeable. Their differing numerical values therefore do not require a phenomenological epoch-screening mechanism. In the canonical TEP-TH/TEP-BBN architecture, no early-time screening function is imposed to restore a standard hot-Big-Bang thermal history. The low- ℓ +lensing+BAO+Pantheon+ chains reported here serve as implementation and robustness tests of the native `hi_class` module. The native `hi_class tep_mode` high- ℓ TTTEE production likelihood remains pending because of a `clipy / hi_class` interface incompatibility (Section 5.1). However, the high- ℓ sector is not untested: a `CAMB`-based effective-background mapping was run successfully, producing stable high- ℓ spectra and a `plik_lite` likelihood evaluation consistent with the low- ℓ posterior. This `CAMB` diagnostic is reported as a consistency check. The low- ℓ constraint therefore remains the primary acoustic result of the present paper.

Native-TEP joint MCMC. This paper's primary contribution is the verified `hi_class` implementation, demonstrating ppm-level sound-horizon preservation and acoustic-sector equivalence in the native static conformal geometry.

5.3 Structure Growth and the Matter Power Spectrum

The full `hi_class` Boltzmann closure with active SMG perturbations yields a linear growth amplitude in agreement with Planck and weak-lensing measurements:

- *Linear growth amplitude:* $\sigma_8 = 0.825 \pm 0.016$ at the fiducial TEP parameters ($\epsilon_T = 0.0066$, $z_T = 5$, $n_T = 2$), compared to $\sigma_8 = 0.823$ for standard Λ CDM at the same cosmology. The TEP value is a native output of the full SMG EFT solver with runtime Bellini-Sawicki mappings; no phenomenological suppression factor is applied.
- *Growth-function diagnostic:* The linear growth rate $f\sigma_8$ from the `hi_class` matter power spectrum (computed via $P(k)$ at $z = 0$ and $z = 0.5$) shows excellent agreement between the screened TEP model and Planck+BOSS best-fit Λ CDM:

Model	$f\sigma_8(z = 0)$	$f\sigma_8(z = 0.5)$
Λ CDM	0.823	0.410
TEP (background-only)	0.824	0.413
TEP (active $\delta\phi$)	0.824	0.413

The TEP deviation from Λ CDM is 0.001 (0.1%) at $z = 0$ and 0.003 (0.7%) at $z = 0.5$, well within current BOSS and eBOSS uncertainties ($\sim 3\%$). This confirms that the TEP covariant action, when mapped rigorously into the Bellini-Sawicki EFT and solved with the full Boltzmann hierarchy, naturally restores structure growth to the observationally consistent range.

- *Scale-dependent screening signature:* The gradient-screening envelope $\mathcal{S}_\Sigma(k) = [1 + (k/k_t)^n]^{-1}$ injects a characteristic scale dependence into the matter power spectrum. At $k < k_t$ (cosmic scales), $\mathcal{S}_\Sigma \rightarrow 0$ and the background expansion is preserved. At $k > k_t$ (structure scales), $\alpha_M(k) \rightarrow 0$ and the scalar fifth force is suppressed, yielding $G_{\text{eff}} \rightarrow G_N$ inside collapsing halos. The fractional deviation $\Delta P(k)/P_{\Lambda\text{CDM}}(k)$ exhibits a smooth transition around k_t , producing a testable spectral feature for DESI and Euclid clustering data.

The σ_8 result demonstrates that the TEP covariant action, when mapped rigorously into the Bellini-Sawicki EFT and solved with the full Boltzmann hierarchy, naturally restores structure growth to the observationally consistent range. Simplified EdS-only growth ODEs, which lack the SMG EFT perturbation closure, are insufficient for this sector; the full Boltzmann closure is required.

5.4 The Hubble Tension in TEP

The TEP framework offers a proposed reconciliation of the Hubble tension without invoking an early-universe crisis. The homogeneous background is exactly mathematically isomorphic to Λ CDM ($H_0 \approx 67$ km/s/Mpc from the CMB), while the apparent local $H_0 \approx 73$ km/s/Mpc arises from an environment-dependent clock-transport bias along the local distance ladder (Cepheid/SN Ia calibration in unscreened stellar atmospheres). The tension is a measurement-environment effect, bypassing the need for early-universe expansion.




Figure 2. Corpus-level H_0 comparison: HC acoustic posterior and Paper 11 local-ladder correction. **HC joint MCMC:** native `hi_class` `tep_mode` homogeneous posterior with $\epsilon_T^{\text{HC}} = 0.00602 \pm 0.00493$, $z_{\text{HC}} = 5.0$, $n_T = 2.0$. **Planck 2018:** independent CMB constraint. **SH0ES uncorrected / TEP-corrected:** the local distance-ladder values are imported from Paper 11; HC does not independently re-analyse the Cepheid/SN Ia calibration data. This figure illustrates consistency of the HC early-universe sector with the broader TEP Hubble-tension interpretation; it is not a standalone HC distance-ladder test.

5.5 The Mathematical Limit of the Conformal Field

To explicitly map the action of the conformal field on the acoustic horizon, the acoustic scale is evaluated in a mathematically idealized geometry ($\Omega_m = 1.0, \Omega_\Lambda = 0.0$) using the `hi_class` native `tep_mode` implementation.

Regime I: Screened HC branch ($z_{\text{HC}} = 5$)

In the standard TEP model, the profile $\exp[-(z/z_{\text{HC}})^{n_T}]$ ensures the conformal field correctly matches the apparent late-time acceleration inferred from Pantheon+. The integration verifies this background/acoustic mathematical isomorphism:

ϵ_T	$100\theta_s$	r_s [Mpc]	$\Delta D_C/D_C$	Interpretation
0.00	1.0403	144.526	0.00%	Pure EdS reference (no TEP)
0.05	1.0548	144.519	-1.38%	r_s preserved; θ_s shifts from D_C projection

The sound horizon r_s remains exact because the conformal field geometry accurately tracks the mathematics of the acoustic horizon without requiring physical stretching of space.

Regime II: Unscreened diagnostic branch ($z_{\text{HC}} \rightarrow \infty$)

Removing the empirical profile and forcing the conformal factor to grow as a pure unsuppressed power law $A(z) = (1+z)^{\epsilon_T}$ exposes the mathematical divergence of the bare field:

ϵ_T	$100\theta_s$	r_s [Mpc]	$\Delta D_C/D_C$	Interpretation
0.00	1.0403	144.526	0.00%	Pure EdS reference
0.05	0.7565	100.584	-4.30%	r_s mathematically squeezed by divergence

This mathematical limit demonstrates that the $z_{\text{HC}} \sim 5$ empirical fitting function accurately defines the physical profile of the conformal field, allowing it to mimic Dark Energy while preserving the CMB acoustic horizon without expanding space.




Figure 3. Jordan-frame acoustic-scale screening diagnostic. Background-only native `hi_class` `tep_mode` diagnostic in an Einstein-de Sitter geometry ($\Omega_m = 1.0, \Omega_\Lambda = 0.0$). **Left panel:** Screened HC branch ($z_{\text{HC}} = 5$, $\epsilon_T^{\text{HC}} = 0.05$, $n_T = 2.0$); the acoustic scale is protected and stays near the Planck 2018 reference. **Right panel:** Unscreened diagnostic branch ($z_{\text{HC}} \rightarrow \infty$, $\epsilon_T^{\text{HC}} = 0.05$, $n_T = 2.0$); removing the empirical profile exposes unphysical acoustic-horizon squeezing. The dashed line is the Planck 2018 acoustic reference value; the comparison is a diagnostic acoustic-scale recovery, not a full TT/TE/EE likelihood plot.

6.1 Falsifiable Predictions

The following near-term observational tests would strengthen or falsify the TEP-HC acoustic-sector and perturbation claims.

1. **Small-scale CMB polarization from active scalar perturbations.** The TEP-HC linear scalar perturbation sector predicts a characteristic phase shift in the TE cross-spectrum at $\ell \gtrsim 1000$ ($\sim 0.1\%$ – 0.3%) due to the $\alpha_B \neq 0$ running of the conformal scalar. CMB-S4 and Simons Observatory, targeting $\ell \sim 3000$ with μK -arcmin sensitivity, can test this deviation directly. A null result at $> 3\sigma$ would push the TEP scalar amplitude into the screened limit.
2. **B-mode polarization from native tensor modes.** The TEP-TH tensor analysis yields $r(k_{\text{pivot}}) = 9 \times 10^{-6}$ and $r_{\text{max}} = 6.26 \times 10^{-4}$, well below current BICEP/Keck bounds ($r < 0.036$) but above the cosmic-variance floor for CMB-S4 ($r \sim 10^{-4}$ – 10^{-3}). A B-mode detection with $r \sim 10^{-3}$ would be inconsistent with the TEP native tensor prediction; a non-detection down to $r \sim 10^{-4}$ would remain compatible and would sharpen the bound on p in the temporal-horizon profile.
3. **Acoustic-peak consistency under `tep_mode` background.** The TEP background factor $M(z) = A/(1 - \alpha_A)$ preserves the acoustic-peak locations to within the numerical integration tolerance of the Boltzmann solver (currently $\lesssim 10$ ppm). The current `hi_class` implementation reports $r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}} = 0.999994$, where the residual ~ 6 ppm is understood as a numerical/implementation-level artifact of finite-precision conformal-frame background reconstruction. Any claimed physical deviation from the exact conformal isomorphism must be shown to exceed the standard solver interpolation noise (~ 10 ppm) before it can be interpreted as a falsification of the background mapping, rather than as a C++ integration artifact.
4. **Growth-factor amplitude from native EFT.** The TEP-HC active-perturbation run yields $\sigma_8 = 0.825 \pm 0.016$, giving a structurally higher linear growth amplitude than the standard Planck ΛCDM baseline (0.811 ± 0.006). As Stage-IV weak-lensing surveys (Euclid, Rubin/LSST) resolve their internal calibration systematics, the S_8 consensus is expected to trend upward toward the TEP prediction if the conformal-frame Hubble-friction modification is physically realized. This is framed as a diagnostic trend rather than a binary kill-switch: TEP does not predict an exact S_8 threshold but rather a directionally higher growth amplitude arising from the same Bellini–Sawicki EFT sector that preserves the acoustic scale.

Each test targets a distinct sector of the TEP-HC claim: active scalar perturbations (test 1), tensor modes (test 2), background isomorphism (test 3), and growth amplitude (test 4). A combined null result across all four would require a significant revision of the conformal-frame mapping.

6. Conclusion: The Static Conformal Universe

This paper implements and validates the native Temporal Equivalence Principle (TEP) conformal modification directly within the `hi_class` Boltzmann solver framework. By leveraging the mathematical isomorphism between the FLRW expanding scale factor $a(t)$ and the TEP conformal scalar field $A(\phi)$, this analysis demonstrates that the early-universe acoustic-sector observables can be reproduced at high fidelity under a static conformal temporal-transport mapping.

6.1 Summary of Results

1. *The Mathematical Isomorphism:* The TEP conformal factor $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$ dictates the clock-rates and photon phases in the causal matter metric $\tilde{g}_{\mu\nu} = A(\phi)^2 g_{\mu\nu}$. Because this scalar field evolves identically to the standard spatial scale factor $a(t)$, standard Boltzmann solvers can be used as conformal-frame calculators for the background/acoustic-sector mapping tested here. The parameter traditionally defined as Dark Energy (Ω_Λ) is operationally reinterpreted within this implementation as the homogeneous temporal-shear background contribution filling the same background budget slot, Ω_ϕ .
2. *CMB Acoustic Preservation (Pure-Conformal Consistency):* Because of the static conformal isomorphism, the `hi_class` native integration demonstrates that the background/acoustic-sector observables are preserved to parts-per-million ($r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}} = 0.999994$). This is a pure-conformal consistency check: the conformal transport geometry recovers the standard FLRW acoustic results in the regime where the conformal field dominates and disformal corrections are suppressed. It is not an independent confirmation of TEP, but rather a necessary consistency requirement that any viable conformal-frame alternative must satisfy. The background acoustic observables alone do not uniquely force the spatial-expansion interpretation; they can be naturally accommodated by the evolving background scalar field $A(\phi)$. The active-perturbation closure (Section 4.5) confirms that evolving the implemented linear pure-conformal scalar fluctuation sector $\delta\phi$ through the exact Bellini–Sawicki EFT produces parameter posteriors indistinguishable from the background-only chain to 0.07σ , with cross-chain Gelman–Rubin $R - 1 = 0.007$ at 58,180 accepted steps.
3. *The Temporal Horizon:* The result of this paper does not require the CMB acoustic peaks to originate from a physically expanding spatial metric beginning at a density singularity. In the TEP interpretation tested here, the same conformal transport integrals normally written in terms of the FLRW scale factor $a(t)$ are reproduced by the temporal conformal field $A(\phi)$. The limit conventionally described as $a \rightarrow 0$ is therefore reinterpreted, at the level of clock transport and photon phase evolution, as a temporal-horizon limit $A_{\text{clock}} \rightarrow 0$ relative to the present epoch. This is the precise sense in which the present calculation removes the Big Bang singularity from the acoustic-sector interpretation: the sound horizon, photon-baryon driving, and acoustic-peak morphology are preserved without requiring physical spatial stretching back to a zero-scale-factor origin. In this precise but physically important sense, the CMB acoustic sector no longer requires a physical zero-scale-factor Big Bang; it can be equivalently represented as a conformal temporal-horizon limit of the clock-rate field. Because this temporal horizon is asymptotic, cosmological epochs typically defined by a finite "time since the Big Bang" are instead fundamentally mapped by their thermodynamic temperature and the exact conformal clock-rate, shifting the measurement of cosmic history from a linear stopwatch to a thermodynamic state. Crucially, as established in TEP-BBN, the observed radiation temperature (T_{obs}) and the local matter temperature (T_{loc}) are distinct physical quantities governed by temporal transport. The full nonsingular closure—geodesic completeness, curvature invariants, entropy evolution, steady-state abundances, and the origin of the CMB blackbody

spectrum—is delivered in TEP-TH (Paper 27) and TEP-BBN (Paper 29), where Proposition 1 establishes finite matter-frame curvature at the temporal horizon and the pipeline validates the eternal equilibrium observables.

4. *Cosmological Constraints:* A joint hi_class Cobaya MCMC (Planck 2018 low- ℓ TT/EE + lensing + BAO + Pantheon+) yields a close match to the conformal field parameters. The companion paper TEP-C0 (Paper 26) provides robust late-time evidence: BF approximately 4.6 (conservative $z_{\text{los}} = 5$), approximately 61.8 (fixed $z_{\text{los}} = 100$ benchmark), and approximately 40.3 (broad free- z_{los}), reducing the phenomenological need to treat late-time acceleration as primitive spatial expansion.

6.2 Resolving the Cosmological Crises

Standard Λ CDM cosmology currently faces two severe crises: the Hubble Tension (H_0) and the unexpectedly massive high-redshift galaxy candidates observed by JWST. The Static Conformal Universe offers a unified TEP interpretation of both without invoking early-universe modifications.

- *The Hubble Tension:* In the broader TEP corpus, Paper 11 argues that the temporal shear field is environmentally screened by mass. Supernovae exist in empty voids (where the field is unscreened, yielding a high inferred H_0), while Cepheids exist in dense galaxies (where the field is partially screened, yielding a lower inferred H_0). Paper 11 argues that the tension is an artifact of environmental mass-screening on local kinematic distance probes, not an early-universe physics crisis. HC does not independently re-analyse the distance-ladder data; it imports the interpretation.
- *JWST High-Redshift Galaxies:* Within the broader TEP interpretation, the temporal-horizon picture removes the finite-age assembly bottleneck by replacing the FLRW singularity with an asymptotic conformal-clock boundary. The massive galaxies observed by JWST therefore form strictly within standard astrophysical accretion models over vast timescales (Paper 12).

6.3 Synthesis of the Paradigm Shift

This analysis implements and explicitly validates the native TEP implementation within a rigorous Boltzmann solver framework. The acoustic indistinguishability of the static conformal background from Λ CDM at recombination demonstrates that the early-universe background physics cannot easily distinguish between a stretching spatial metric and an evolving conformal clock-rate field.

Late-universe Pantheon+ data in TEP-C0 favor the physical no- Λ TEP branch over baseline Λ CDM with BF $\simeq 4.6$ (conservative), BF $\simeq 61.8$ (benchmark), and BF $\simeq 40.3$ (free- z_T), providing a concrete conformal-frame alternative to the background expansion interpretation. The active-perturbation closure reported in Section 4.5 confirms that the implemented linear pure-conformal scalar fluctuation sector is ghost-free and observationally negligible, with the implemented $\delta\phi$ -enabled Einstein–Boltzmann chain agreeing with the background-only chain to 0.07σ across all cosmological parameters. By treating time itself as a dynamical, mass-screened scalar field, TEP seeks to unify early-universe acoustic physics, late-time "acceleration", the H_0 tension, and JWST anomalies into a single, cohesive static geometric framework. The present paper provides both the hi_class background/acoustic benchmark and the perturbation-closure validation.

References

- Smawfield, M. (Paper 1). *Global Time Echoes: Distance-Structured Correlations in GNSS Clocks*. TEP Corpus. DOI: 10.5281/zenodo.17127229.
- Smawfield, M. (Paper 6). *Universal Critical Density: Unifying Atomic, Galactic, and Compact Object Scales*. TEP Corpus. DOI: 10.5281/zenodo.18064365.
- Smawfield, M. (Paper 11). *The Cepheid Bias: Resolving the Hubble Tension*. TEP Corpus. DOI: 10.5281/zenodo.18209702.
- Smawfield, M. (Paper 12). *The Temporal Equivalence Principle: A Unified Resolution to the JWST High-Redshift Anomalies*. TEP Corpus. DOI: 10.5281/zenodo.19000827.
- Smawfield, M. (Paper 13). *The Temporal Equivalence Principle: Temporal Shear Recovery in Gaia DR3 Wide Binaries*. TEP Corpus. DOI: 10.5281/zenodo.19102061.
- Smawfield, M. (Paper 26). *Temporal Equivalence Principle: A Covariant Alternative to Cosmic Expansion*. TEP Corpus. DOI: 10.5281/zenodo.20370143.
- Smawfield, M. (Paper 27). *Temporal Equivalence Principle: Temporal Horizon Cosmology and the Absence of a Physical Big Bang Singularity*. v0.3 (Thika). DOI: 10.5281/zenodo.20723059.
- Smawfield, M. (Paper 29). *Temporal Equivalence Principle: Dynamical Proper Time and the Illusion of Primordial Deuterium*. Paper 29 (Dubai). DOI: 10.5281/zenodo.21841148.
- Bellini, E., & Sawicki, I. 2014, JCAP, 07, 050. *Maximal freedom at minimum cost: linear large-scale structure in scalar-tensor theories*.
- Brax, P., Burrage, C., Davis, A.-C., & Gubitosi, G. 2019, Phys. Rev. D, 100, 083515. *Screening mechanisms in scalar-tensor theories*.

Cobaya Team. 2023, *Cobaya: Code for Bayesian Analysis of physical theories*. arXiv:2305.02971.

Hu, B., Raveri, M., Frusciante, N., & Silvestri, A. 2014, Phys. Rev. D, 89, 103530. *EFTCAMB/EFTCosmoMC: Numerical Notes*.

Knox, L., & Millea, M. 2020, Phys. Rev. D, 101, 043533. *Hubble constant hunter's guide*.

Lagos, M., Bellini, E., Jimenez, J. B., et al. 2018, JCAP, 03, 021. *hi_class: Horndeski in the Cosmic Linear Anisotropy Solving System*.

Lewis, A., Challinor, A., & Lasenby, A. 2000, Astrophys. J., 538, 473. *Efficient computation of cosmic microwave background anisotropies*.

Planck Collaboration. 2020, A&A, 641, A1. *Planck 2018 results. I. Overview and cosmological parameters*.

Planck Collaboration. 2020, A&A, 641, A6. *Planck 2018 results. VI. Cosmological parameters*.

Riess, A. G., Casertano, S., Yuan, W., et al. 2022, ApJ, 934, L7. *A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km/s/Mpc Uncertainty from the Hubble Space Telescope and the SH0ES Team*.

Sawicki, I., & Bellini, E. 2015, Phys. Rev. D, 92, 084061. *Stability of dark energy and the generalized no-slip condition*.

Zumalacárregui, M., & García-Bellido, J. 2014, Phys. Rev. D, 89, 064046. *Transforming gravity: from derivative couplings to matter to second-order scalar-tensor theories beyond the Horndeski Lagrangian*.

Appendix A: Technical Implementation Details

A.1 hi_class Installation and Configuration

A.1.1 Building with TEP Support

hi_class is installed automatically by pipeline Step 1 (`step_00b_install.py`), which clones hi_class and applies the native TEP patch from `external/patches/hiclass_tep_native.patch` to `source/background.c`, `source/input.c`, and `include/background.h`. Manual rebuild:

```
cd external/hi_class/hi_class
make clean && make
```

Screening projection notice. Screening in TEP is represented at theory level by the environmental operator $S_{\Sigma}(E)$. Quantities such as ρ_T , $R_T(M)$, $S_{\oplus}(r)$, compactness Φ/c^2 , local stellar density, coherence length, proximity, and boundary geometry are domain-specific projections of E , not independent screening mechanisms and not interchangeable universal thresholds.

A.2 Cobaya Installation

```
pip install cobaya
cobaya-install planck_2018_lowl.TT planck_2018_lowl.EE \
  planck_2018_lensing.native bao.sdss_dr12_consensus_final \
  sn.pantheonplus --path /path/to/likelihoods
```

A.3 TEP Module C Code Structure and Implementation Note

The native conformal modification is implemented directly in hi_class `source/background.c`, controlled by the `.ini` flags `tep_mode`, `epsilon_T`, `z_T`, `n_T`. The relevant functions are:

1. `tep_f_transition(pba, z)`: returns the suppression factor $S(z) = \exp[-(z/z_T)^{n_T}]$; the full transition is $f_T(z) = \ln(1+z) S(z)$ (see `core/cosmology.py:f_T`).
2. `tep_gamma_factor(pba, z)`: returns the exact covariant conformal factor $A(z) = \exp[\epsilon_T \ln(1+z) S(z)]$ (not linearised).
3. The Hubble rate and its conformal-time derivative are mathematically mapped using the exact geometric relation $M(z) = A/(1 - \alpha_A)$ in `background_functions` and in the initial-Hubble setter, explicitly evaluating the static conformal geometry.

Implementation note (corrected bug). An earlier build used $f_T = 1 - \exp[-(z/z_T)^{n_T}]$ (the complement of the suppression function). This incorrectly inverted the scalar field profile, erroneously mapping the peak kinetic energy to the early universe rather than intermediate redshifts, which logically corrupted the acoustic integration. In addition, the post-processing step that read the spectra used a hard-coded output index and could silently load a stale file from an earlier run. Both issues are fixed: the transition function now uses the shared TEP-C0 implementation (`core/cosmology.py`), correctly matching the field profile to the Pantheon+

apparent acceleration, and the analysis resolves the most recent `hi_class` output deterministically. *Sign convention (TEP disformal metric)*: the distance integrand is multiplied by $A(z)$ for null-geodesic propagation in the conformal frame. The legacy SMG alpha-function stub (`smg_tep_*`) has been retired; production physics lives in the patched `background.c` (`external/patches/hiclass_tep_native.patch`).

A.3a Derivation of the Conformal-Frame Factor $M(z)$

This appendix derives the background conformal mapping $M(z)$ from the bi-metric action (Equation 1) using a single frame convention held fixed throughout, demonstrating the exact geometric relation implemented natively in the codebase.

Setup and convention. Matter, photons, and rods couple to the conformal metric $\tilde{g}_{\mu\nu} = A^2(\phi) g_{\mu\nu} + B(\phi) \nabla_\mu \phi \nabla_\nu \phi$. For the homogeneous background the disformal term contributes only through the time-time component and is absorbed into the lapse; the evolution is governed by the conformal part, so $B \rightarrow 0$ is imposed here (the disformal sector re-enters at the perturbative/GW level via α_T , Section 2.2.2). The conformal part gives the standard map between the Einstein-frame scale factor a_E and cosmic time t_E and their conformal counterparts:

$$\tilde{a} = A(\phi) a_E, \quad d\tilde{t} = A(\phi) dt_E. \quad (14)$$

These two relations *define* the convention; every subsequent equation is derived from them. The transition factor $S(z) = \exp[-(z/z_T)^{n_T}]$ correctly forces $A(z) \rightarrow 1$ at the local endpoint, so the code's redshift grid can be identified with the physical conformal-frame redshift. The explicit $(z_E, \tilde{z})$ distinction is retained only to derive the frame relation.

Physical Hubble rate. The expansion rate measured by conformal-frame clocks and rulers is $\tilde{H} \equiv \tilde{a}^{-1} d\tilde{a}/d\tilde{t} = d \ln \tilde{a} / d\tilde{t}$. Using $d/d\tilde{t} = A^{-1} d/dt_E$ and $\ln \tilde{a} = \ln A + \ln a_E$,

$$\tilde{H} = \frac{1}{A} \frac{d}{dt_E} (\ln A + \ln a_E) = \frac{1}{A} \left(\frac{d \ln A}{dt_E} + H_E \right), \quad (15)$$

where $H_E = d \ln a_E / dt_E$ is the Einstein-frame rate. Because the TEP conformal factor $A(z)$ is evaluated as a function of the observable physical redshift (the matter-frame redshift $1 + z = \tilde{a}_0 / \tilde{a}$), the coupling α_A computed in the codebase is fundamentally the derivative with respect to the *matter-frame* scale factor:

$$\frac{d \ln A}{dt_E} = \frac{d \ln A}{d \ln \tilde{a}} \frac{d \ln \tilde{a}}{dt_E} = \alpha_A (A \tilde{H}), \quad \alpha_A \equiv \frac{d \ln A}{d \ln \tilde{a}} = - \frac{d \ln A}{d \ln(1+z)}, \quad (16)$$

which matches the definition in Section 3.2. Substituting $d \ln A / dt_E = \alpha_A A \tilde{H}$ into (15) yields $\tilde{H} = \alpha_A \tilde{H} + H_E / A$, or equivalently $\tilde{H}(1 - \alpha_A) = H_E / A$. Identifying the reference FLRW background rate with the conformally mapped Einstein-frame rate used by the implementation, $H_E = A^2 H_{\Lambda\text{CDM}}$, gives the exact geometric relation:

$$\boxed{\tilde{H}(z) = \frac{A(z)}{1 - \alpha_A(z)} H_{\Lambda\text{CDM}}(z)} \quad \implies \quad M_{\text{exact}}(z) = \frac{A}{1 - \alpha_A}. \quad (17)$$

Implementation Status. The production codebase (`hiclass_tep_native.patch`) evaluates this exact geometric relation $M = A/(1 - \alpha_A)$ directly, guaranteeing mathematical fidelity to the conformal evaluation without requiring first-order approximations.

A.4 Saturation Scale in Cosmological Units

The candidate Temporal Topology saturation scale $\rho_T \approx 20 \text{ g/cm}^3$ converts to cosmological units as:

$$\rho_T = 20 \text{ g/cm}^3 = 2 \times 10^4 \text{ kg/m}^3 \approx 1.1 \times 10^{34} \text{ eV/cm}^3 \quad (18)$$

In Planck units ($\hbar = c = G = 1$):

$$\rho_T \approx 4 \times 10^{-93} M_{\text{Pl}}^4 \quad (19)$$

Compare to cosmic mean density today ($\rho_{\text{crit},0} \approx 10^{-123} M_{\text{Pl}}^4$). The hierarchy ensures the vast cosmological voids evaluate the pure unsuppressed conformal field, accurately simulating the expansion of space.

Intermediate environments and operational parameter bounds. Between the terrestrial laboratory and the cosmic mean, the screening transition is continuous. At stellar atmospheric densities ($\rho \sim 10^{-6} \text{ g/cm}^3$), the field is partially screened; at interplanetary densities ($\rho \sim 10^{-23} \text{ g/cm}^3$), it is essentially unscreened. Certain orbital datasets—notably the Galileo GNSS clock ensemble—fall outside the operational parameters established for valid TEP-GNSS screening analysis (Paper 1), because their orbital altitude and local gravitational environment do not satisfy the strict kinematic isolation required to isolate the conformal phase drift from standard relativistic corrections. These exclusions are documented in the TEP-GNSS pipeline and do not affect the cosmological bound, which operates in the deep unscreened regime where $\rho \ll \rho_T$.

A.5 Stability Sector Closure

To formally verify the stability of the active scalar perturbations, the native `hi_class` SMG module was extended to evaluate the exact analytical limits of the TEP conformal geometry at runtime.

In a generalized Horndeski treatment, the solver enforces the following physical stability conditions:

1. $c_s^2 \geq 0$ (no gradient instabilities)
2. $D = \alpha_K + \frac{3}{2}\alpha_B^2 \geq 0$ (no ghosts)
3. $|\alpha_M|$ bounded (sub-luminal Planck-mass running)
4. $\alpha_T \approx 0$ (gravitational wave speed constraints)

For the conformal modification implemented here, the EFT parameters map strictly to the dynamical background derivative α_A :

$$\alpha_M = -2\alpha_A \quad (20)$$

$$\alpha_B = 2\alpha_A \quad (21)$$

$$\alpha_K = -5\alpha_A^2 \quad (22)$$

$$\alpha_T = 0 \quad (23)$$

The substitution of the kineticity and braiding terms into the physical no-ghost discriminant yields an exact identity:

$$D = (-5\alpha_A^2) + \frac{3}{2}(2\alpha_A)^2 = \alpha_A^2 \quad (24)$$

This identity proves that the continuous conformal field transition is analytically protected against ghost instabilities, as the discriminant remains strictly positive-definite. In the pure-conformal implementation used here, the runtime SMG closure fixes the scalar sound-speed sector to the luminal conformal-frame limit ($c_s^2 = 1$), and the integration verifies that this choice produces no gradient instability across the sampled redshift range. The no-ghost condition is independently protected by the discriminant identity ($D = \alpha_A^2$). A fully derived sound-speed expression for the disformal and nonlinear screening sectors is outside the scope of the present pure-conformal closure.

The production codebase forces these limits natively during the calculation of the SMG perturbation coefficients, guaranteeing mathematical fidelity to the conformal evaluation without requiring pre-tabulated interpolation or analytical approximations.

Unified TEP Parameter Dictionary

The TEP corpus uses related but distinct symbols across its papers. This dictionary maps every parameter, its definition, the paper where it is primary, and its fiducial or fitted value.

Symbol	Definition	Primary Paper	Fiducial / Fitted Value
$A_{\text{clock}}(z)$	Exact observational clock/redshift map: $A_{\text{clock}} = (1+z)^{-1}$	TEP-TH	$(1+z)^{-1}$ (exact)
$A_{\text{dyn}}(z)$	Dynamical shear response: $(1+z/z_t)^{-\epsilon_t}$	TEP-TH	Modifies late-time evolution
α_A	Temporal-shear conformal amplitude in Jordan-frame notation	TEP-HC	-0.0028 (Planck best-fit)
$\alpha_M, \alpha_B, \alpha_K, \alpha_T$	Runtime Bellini–Sawicki EFT functions: $\alpha_M = -2\alpha_A$, $\alpha_B = 2\alpha_A$, $\alpha_K = -5\alpha_A^2$, $\alpha_T = 0$	TEP-HC	Derived from α_A
ϵ_T^{los}	Late-time line-of-sight transport amplitude (C0 supernova fit)	TEP-C0	$\mathcal{U}[0, 1.0]$ (prior); posterior peaked near ~ 0.89
ϵ_T^{CMB}	C0 background/acoustic diagnostic amplitude	TEP-C0	-0.0015 ± 0.0037
ϵ_T^{HC}	Native hi_class homogeneous conformal amplitude	TEP-HC	0.00602 ± 0.00493
ϵ_{dyn}	Dynamical temporal-horizon response	TEP-TH	Determined by late-time shear
ϵ_{field}	Primordial spectral-flow parameter constrained by n_s	TEP-TH	0.0175 (from $n_s = 0.965$)
z_T^{los}	C0 line-of-sight supernova transport turnover	TEP-C0	5 (conservative), 100 (benchmark), free (broad)
z_T^{HC}	Homogeneous/acoustic hi_class profile scale	TEP-HC	Fitted jointly with ϵ_T
p	Temporal-horizon conformal exponent: $A_{\text{clock}} \sim \eta^{-p}$	TEP-TH	$0 < p \leq \frac{1}{2}$ (regular branch)
$r_s^{\text{TEP}}/r_s^{\Lambda\text{CDM}}$	Pre-recombination sound-horizon ratio	TEP-HC	0.999994 (< 6 ppm deviation)
$D = \alpha_K + \frac{3}{2}\alpha_B^2$	No-ghost discriminant (physical branch: $D = \alpha_A^2$)	TEP-HC	$\alpha_A^2 > 0$ (positive definite)
$r(k_{\text{pivot}})$	Native tensor-to-scalar ratio at Planck pivot	TEP-TH	9×10^{-6}
r_{max}	Maximum tensor-to-scalar ratio across transition profile	TEP-TH	6.26×10^{-4}
H_0	Hubble parameter (TEP-C0 joint MCMC)	TEP-C0	$66.70 \pm 0.58 \text{ km s}^{-1} \text{ Mpc}^{-1}$
S_8	$\sigma_8 \sqrt{\Omega_m/0.3}$ (TEP-HC joint MCMC)	TEP-HC	0.867 ± 0.026
σ_8^{HC}	Native hi_class matter-fluctuation amplitude	TEP-HC	0.825 ± 0.016

Note: Parameters with superscript labels ($^{\text{los}}$, $^{\text{HC}}$) are related projections of the same temporal sector but are not numerically interchangeable. The turnover scales z_T^{los} and z_T^{HC} describe different physical regimes; the amplitudes ϵ_T^{los} , ϵ_T^{CMB} , ϵ_T^{HC} , and ϵ_{field} are constrained by different observables.

Appendix B: Data Availability & Reproducibility

This work follows open-science practices. All results are fully reproducible from raw data using the documented pipeline. All numerical results, figures, and statistics are generated by deterministic Python scripts processing public observational data.

Repository and Code

GitHub Repository: github.com/matthewsmawfield/TEP-HC

The repository contains a deterministic, version-controlled cosmological analysis pipeline for CMB acoustic peak preservation tests and MCMC parameter estimation with TEP screening.

All MCMC chains, hi_class patch files, posterior samples, and the exact `cobaya` YAML configuration files are released in the Zenodo repository (DOI: 10.5281/zenodo.20682752) under CC-BY 4.0. The `run_all.py` orchestration script and all step scripts are provided in the GitHub repository.

Repository Structure

```
TEP-HC/
├── data/
│   ├── cobaya/           # Cobaya MCMC chains
│   ├── external/        # hi_class submodule
│   └── hi_class/         # TEP-CLASS implementation
├── scripts/
│   └── steps/            # Analysis pipeline steps
├── core/                 # TEP shared constants and parameters
├── site/
│   └── components/       # Manuscript HTML sections
├── requirements.txt
├── CITATION.bib
└── README.md
```

Software Environment

Key packages: NumPy, SciPy, Matplotlib, Cobaya, hi_class. The pipeline has been tested on Python 3.10+.

License

All code and manuscripts are released under CC-BY-4.0.