



Temporal Topology Saturation Scale: Cross-Scale Consistency of ρ_T

Matthew Lukin Smawfield

Version: v0.8 (New Delhi)

First published: 28 December 2025 · Last updated: 30 September 2026

DOI: 10.5281/zenodo.18064365

Abstract

Dark matter observations across cosmological scales exhibit a regularity: the characteristic radius at which Newtonian dynamics fails scales as $R \propto M^{1/3}$, implying a universal saturation/proximity scale (historically labelled a critical density), observationally proxied by ρ_T . This scaling appears in galaxy rotation curves (SPARC database), ultra-diffuse galaxies (DF2/DF4), and the Milky Way's dark-matter onset. Within TEP, this pattern is interpreted as evidence of a candidate saturation scale in the conformal time-field sector, where field saturation occurs at a characteristic proximity scale.

Terrestrial calibration—derived from a newly identified distance-structured correlation in GNSS atomic clocks—provides an empirical anchor. A 25-year CODE analysis yields $L_c \approx 4,200$ km for Earth's mass ($M_\oplus \approx 6 \times 10^{27}$ g), consistent with multi-center results (CODE, IGS, ESA). The characteristic length L_c is operationally identified with the projected Temporal Topology covariance scale associated with $R_T(M_\oplus)$, the geometric saturation scale for Earth's mass; a soliton interpretation is one candidate microscopic realization, not assumed in the calibration. This implies $\rho_T \approx 20$ g/cm³. This calibration exhibits 25-year temporal stability and survives raw RINEX validation, constraining processing-artifact explanations.

Galactic-scale consistency is tested using the SPARC rotation curve database (167 galaxies). The empirical dark matter onset scaling is $\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ (stat) ± 0.07 (definition), consistent with the $M^{1/3}$ expectation within $\sim 0.3\sigma$. For the Milky Way, the SPARC-calibrated $M^{1/3}$ relation predicts a dark-matter onset radius $R_{\text{DM}} \approx 3$ kpc, consistent with the observed transition from baryonic to dark-matter-dominated rotation at $R \sim 3\text{--}5$ kpc. For ultra-diffuse galaxies DF2 and DF4, the model predicts Temporal Topology saturation radii exceeding tidal radii, consistent with observed dark matter deficiency via an illustrative field-envelope disruption mechanism (a dynamical timescale for re-equilibration remains to be computed).

Temporal Topology screening provides a candidate explanation for why precision GR tests remain satisfied. A hierarchy of 26 astrophysical objects spanning 15 orders of magnitude in density is assembled; regression on the 11 dense objects ($\rho > \rho_T$) yields $S \propto \rho^{0.334}$ ($R^2 = 0.99995$), algebraically expected from the $R_T(M)$ construction. This provides a candidate screening explanation for why GR tests pass (binary pulsars: $S \sim 29,000$) while galactic dynamics ($S \sim 10^{-9}$ at $\rho \sim 10^{-24}$ g/cm³) are deeply unscreened, exhibiting strong scalar effects.

The saturation density $\rho_T \approx 20$ g/cm³ emerges as a candidate macroscopic saturation/proximity scale in the TEP environmental-response hierarchy, not an ambient-density switch. GNSS provides the primary terrestrial timing anchor;

SPARC onset scaling, Milky Way structure, RBH-1 crossover behaviour, compact-object screening, and condensed-matter/EOS considerations then test whether the same scale organizes otherwise disconnected gravitational regularities without retuning. Channel-specific quantities such as ρ_{trans} , R_s , κ_{Cep} , κ_{MSP} , and lensing response coefficients are downstream projections of $S_{\Sigma}(\mathcal{E})$, not independent measurements of ρ_T .

The $\rho^{1/3}$ hierarchy is a consistency relation induced by the $R_T(M)$ construction; it is not, by itself, an independent discriminator of microscopic screening mechanism.

Evidence hierarchy. Within the evidence hierarchy of the TEP series, the GNSS clock analyses (Papers 1–3) provide the primary empirical input for the terrestrial correlation scale; the present paper tests the cross-scale consequences of conditionally identifying that scale with $R_T(M_{\oplus})$.

Keywords: dark matter – gravitation – scalar fields – galaxies: kinematics and dynamics – temporal equivalence principle

1. Introduction: The Dark Matter Problem as a Temporal Structure Problem

The Universal Scaling Anomaly

Dark matter observations across cosmological scales reveal an empirical regularity. From dwarf galaxies ($M \sim 10^9 M_{\odot}$) to massive galaxies ($M \sim 10^{12} M_{\odot}$), the characteristic radius at which Newtonian dynamics fails—the "dark matter onset"—scales approximately as $R \propto M^{1/3}$. This implies a universal critical proximity scale ("critical" here meaning a phenomenological saturation/proximity scale, not a hard density threshold), observationally proxied by $\rho_T \sim M/R^3 \sim \text{constant}$, at which gravitational phenomenology changes.

This scaling appears in multiple independent contexts:

- **Galaxy rotation curves:** SPARC database analysis yields $R_{\text{DM}} \propto M^{\alpha_{\text{SPARC}}}$ with $\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ (stat) ± 0.07 (definition) (Lelli et al. 2016; this work).
- **Ultra-diffuse galaxies:** DF2 and DF4 exhibit dark matter deficiency consistent with saturation radii exceeding tidal radii (van Dokkum et al. 2018, 2019).
- **Milky Way:** The SPARC-calibrated $M^{1/3}$ relation predicts a dark-matter onset at $R_{\text{DM}} \approx 3$ kpc for the Milky Way, consistent with the observed transition from baryonic to dark-matter-dominated rotation at $R \sim 3\text{--}5$ kpc (Gaia Collaboration 2023; Sofue 2020).

The persistence of this $M^{1/3}$ scaling across 6 orders of magnitude in mass (screening hierarchy: gas giants to neutron stars) and 15 orders of magnitude in density suggests a fundamental physical scale, not a coincidence of baryonic feedback or halo assembly.

Reframing Dark Matter: Phantom Mass from Temporal Shear

The Temporal Equivalence Principle (TEP) proposes that gravitational phenomena arise from a conformal time field $\phi(x^\mu)$ that modulates proper time rates. This phenomenology arises from the TEP action (Smawfield 2025a, Paper 0):

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} (\partial\phi)^2 - V(\phi) \right] + S_m[\tilde{g}_{\mu\nu}] \quad (1)$$

where matter couples to the Jordan frame metric $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi$. At late times, $B(\phi)$ is strongly constrained by multi-messenger observations and can be neglected, yielding $\tilde{g}_{\mu\nu} \approx A^2(\phi)g_{\mu\nu}$ with $A(\phi) = \exp(\beta_A\phi/M_{\text{Pl}})$. The environmental screening response suppresses the observed gradient in dense environments, producing effective field saturation at characteristic density ρ_T . In this framework, the "dark matter" problem is reinterpreted as a violation of the *isochrony axiom*—the assumption that clocks at the same gravitational potential tick at the same rate regardless of their spatial separation or the mass distribution's history.

When this axiom is relaxed, mass estimates made through different observable pairs diverge, and "phantom mass" — the apparent mass inferred when temporal structure is analysed as spatial mass — can be measured per channel. In the *dynamical–baryonic* channel studied here (rotation curves), phantom mass is the apparent excess of the dynamical mass $V_{\text{obs}}^2 R/G$ over the baryonic mass: the orbital dynamics respond to the local time gradient, sourcing centripetal support with no additional baryons. In the *lensing* channel (Papers 4, 19), photon paths are conformally invariant and track the gravitating matter alone, so no conformal propagation delay exists and lensing inference reads the true mass — the phantom mass therefore appears only in dynamics, and the dynamical–lensing mass difference is its decisive test. Both measurements arise from spatial structure in proper time—without invoking non-baryonic particles—and are kept distinct throughout.[†]

[†]Note: This geometric "phantom mass" from temporal shear differs from cosmological "phantom energy" (dark energy with $w < -1$). The former arises from spatial gradients in proper time; the latter from exotic equation-of-state matter.

The characteristic scale at which this temporal structure becomes significant is set by the saturation scale ρ_T , where the scalar field ϕ reaches its self-interaction threshold and saturates—forming a stable, localized configuration (a soliton under that interpretation) with radius:

$$R_T = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3}. \quad (2)$$

This $M^{1/3}$ scaling is a direct consequence of the saturation condition; under a soliton interpretation, this corresponds to soliton formation.

The saturation scale parameter $\rho_T \approx 20 \text{ g/cm}^3$ serves as the bulk-matter realization of the abstract environmental screening operator $\mathcal{S}_\Sigma(\mathcal{E})$ defined in the foundational TEP framework. In macroscopic atomic matter, ρ_T acts as an empirical proxy for the geometric saturation of the Temporal Topology, yielding cross-scale predictions without committing to specific chameleon, Vainshtein, or symmetron microphysics.

$$\Sigma_\mu^{\text{obs}} = \mathcal{S}_\Sigma(\mathcal{E}) \nabla_\mu \ln A(\phi), \quad (3)$$

where the environmental state is $\mathcal{E} = \{\rho, \Phi/c^2, \nabla\rho, \nabla\Phi, \text{compactness}, R_T(M), \text{proximity}, T, z, \text{boundary geometry, coherence volume}\}$. This single operator unifies density screening, compactness screening, proximity screening, thermodynamic/environmental-state response, lensing/cosmological covariance screening, and wide-binary environmental screening across the entire TEP corpus.

Box 1.1: Joint Constraint Architecture of ρ_T and $\mathcal{S}_\Sigma(\mathcal{E})$

ρ_T is the shared macroscopic saturation/proximity scale. Other papers in the TEP programme do not all measure ρ_T directly; they measure domain-specific projections of the same environmental operator $\mathcal{S}_\Sigma(\mathcal{E})$:

- **Clock covariance lengths** (GNSS, Paper 1–3)
- **Galactic onset radii** (SPARC, Milky Way, this work)
- **Response coefficients** (Cepheids, pulsars, Papers 10–11)
- **Lensing residuals** (strong-lens time delays, Paper 4)
- **Weak-field recovery scales** (wide binaries, Paper 13)
- **Compactness-dependent residuals** (LLR, Paper 17)
- **Cosmological temporal-shear amplitudes** (hi_class, C0, TH, Papers 18, 26, 27)

Each measurement channel has a response coefficient κ_X , an environmental screening projection, and a domain-specific observable functional. This architecture protects the programme from the criticism that different channels report different numerical values: they are *observable response coefficients*, not bare scalar couplings, and each channel measures a different projection of $\mathcal{S}_\Sigma(\mathcal{E})$.

The Multi-Scale Consistency Strategy

As summarized in Table 1, this paper tests ρ_T through a convergent multi-scale approach:

Table 1: Cross-Scale Consistency of ρ_T

Scale	System	Mass Range	Density Range	Key Observable	Result
Terrestrial	GNSS Clocks	$M_\oplus \sim 6 \times 10^{27} \text{ g}$	$\rho_\oplus \sim 5.5 \text{ g/cm}^3$	$L_c \approx 4200 \text{ km}$	$\rho_T \approx 20 \text{ g/cm}^3$
Galactic	SPARC Galaxies	$10^9\text{--}10^{12} M_\odot$	$\rho \sim 10^{-24} \text{ g/cm}^3$	$R_{\text{DM}} \propto M^{\alpha_{\text{SPARC}}}$	$\alpha_{\text{SPARC}} = 0.355 \pm 0.043 \text{ (stat)} \pm 0.07 \text{ (def.)}$
Local	Milky Way	$M_{\text{bar}} \sim 6 \times 10^{10} M_\odot$	$\rho \sim 10^{-24} \text{ g/cm}^3$	Inner mass-discrepancy onset	$R_{\text{DM}} \approx 3 \text{ kpc}$ (observed $\sim 3\text{--}5 \text{ kpc}$)

Scale	System	Mass Range	Density Range	Key Observable	Result
Screening	26 Objects	$\sim 10^{27}\text{--}10^{33}$ g	$\sim 10^0\text{--}10^{14}$ g/cm ³	$S \propto \rho^{\beta_{\text{scr}}}$	$\beta_{\text{scr}} = 0.334$ ($R^2 = 0.99995$)

Note: The convergence of constraints across 18 orders of magnitude in mass (Earth to galaxy, full range) motivates ρ_T as a candidate saturation scale.

Paper Structure

The remainder of this paper is organized as follows:

- **Section 2:** Terrestrial calibration via GNSS atomic clock correlations, establishing $L_c \approx 4200$ km $\rightarrow \rho_T \approx 20$ g/cm³.
- **Section 3:** Galactic consistency test via SPARC rotation curves, confirming $M^{1/3}$ scaling.
- **Section 4:** Screening hierarchy analysis, providing a candidate screening account for why precision GR tests remain satisfied despite scalar-field structure.
- **Section 5:** Physical context for ρ_T , including a dimensional identity bridge preserving WEP, a condensed-matter EOS coincidence, and open derivation targets.
- **Section 6:** Universal Scaling as a Cross-Scale Consistency Test, synthesizing constraints across all regimes.
- **Section 7:** Milky Way dark-matter onset at $R_{\text{DM}} \approx 3$ kpc.
- **Section 8:** Discussion of dark matter as phantom mass and cosmological implications.
- **Section 9:** Conclusion and astrophysical applications (including RBH-1, companion paper).

This multi-scale convergence identifies $\rho_T \approx 20$ g/cm³ as a candidate GNSS-anchored saturation parameter (with explicit systematic uncertainty), enabling sharply constrained astrophysical applications such as the RBH-1 runaway black hole candidate (Smawfield 2025h, Paper 7).

Evidence hierarchy. Within the evidence hierarchy of the TEP series, the GNSS clock analyses (Papers 1–3) provide the primary empirical input for the terrestrial correlation scale; the present paper tests the cross-scale consequences of conditionally identifying that scale with $R_T(M_{\oplus})$. It is a cross-scale normalization test, not an independent primary proof of the TEP framework.

2. Terrestrial Calibration: GNSS Atomic Clock Correlations

Distance-Structured Correlations in Atomic Clocks

The first empirical constraint on ρ_T comes from terrestrial atomic clock networks. Multi-center analysis of GNSS (Global Navigation Satellite System) clock products reveals distance-structured correlations with a characteristic length scale $L_c \approx 4200$ km for Earth's mass. Operationally, L_c denotes the best-fit exponential decay length recovered by fitting clock-residual covariance versus inter-station separation (see Appendix A). Recent analysis of GNSS products (Smawfield 2025b, Paper 1; Smawfield 2025c, Paper 2) reports this correlation across multiple processing centers.

In this corpus, the terrestrial scale L_c is operationally identified with the projected covariance scale associated with the geometric Temporal Topology scale $R_T(M_{\oplus})$, not with the in-medium Compton wavelength $\lambda_C(\rho)$ of a candidate completion. The empirical claim is that distance-structured clock correlations decay on the geometric scale $R_T(M)$ of the source mass, rather than on a Yukawa range set by local ambient density.

These correlations exhibit seven independent signatures:

- **Distance structure:** Correlation amplitude scales with inter-station separation.
- **Directional anisotropy:** Stronger correlations along Earth's orbital velocity vector.
- **Orbital velocity coupling:** Annual modulation consistent with $v_{\oplus} \approx 30$ km/s.
- **Annual-phase direction recovery:** RA = 6°, Dec = +4°, about 6° from Earth's aphelion velocity tangent; the directed CMB-apex template is anti-phase.
- **Multi-center consistency:** CODE, IGS, and ESA products show consistent characteristic scales and directional structure.
- **25-year stability:** Correlation structure persists across 1999–2024 (Smawfield 2025c, Paper 2).
- **Raw RINEX validation:** Pattern survives in raw-RINEX Single Point Positioning (SPP) analysis, constraining processing artifacts (Smawfield 2025d, Paper 3).

Operational Calibration of ρ_T from L_c

The characteristic length L_c is used as the observed projected covariance scale associated with $R_T(M_{\oplus})$. For a spherical mass distribution, the saturation radius is:

$$R_T = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3}. \quad (4)$$

Solving for ρ_T :

$$\rho_T = \frac{3M}{4\pi R_T^3}. \quad (5)$$

Substituting $M_\oplus \approx 6 \times 10^{27}$ g and $L_c \approx 4200$ km $\approx 4.2 \times 10^8$ cm:

$$\rho_T = \frac{3 \times (6 \times 10^{27} \text{ g})}{4\pi \times (4.2 \times 10^8 \text{ cm})^3} \approx 20 \text{ g/cm}^3. \quad (6)$$

Systematic Uncertainties

The primary uncertainty is the length scale determination. Four distinct contributions are reported: (i) the 25-year CODE per-measurement statistical uncertainty ($4,201 \pm 1,967$ km, $\pm 47\%$; Paper 2); (ii) the inter-center systematic spread (3,330–4,549 km, $\sim \pm 12\%$; Paper 1), with the caveat that centers share largely overlapping underlying data so this spread is not an independent ensemble; (iii) the epoch/window instability established by the executed held-out discriminator (Paper 14 Step 3.1): the identical estimator on the identical product returns λ from 1,050 to 1,935 km across quarterly windows, while matched-day independent chains agree at $\sim 30\%$ (2,744 vs. 1,933 km — the CNES value landing within 4% of the pooled 1,862 km — same-product fallback days excluded by epoch-aligned detrended-residual comparison) — the dominant calibration systematic is therefore the ~ 30 – 40% epoch/window scatter, not the inter-center figure; and (iv) the $\mathcal{O}(1)$ prefactor in the transfer sketch (Section 2), treated as a model-defining choice. The adopted operational value is $L_c = 4,200$ km with ± 500 km ($\pm 12\%$) reflecting the inter-center range, plus the epoch/window systematic carried separately. Since $\rho_T \propto L_c^{-3}$, the 30–40% window scatter propagates to a factor ≈ 2 – 3 in ρ_T : the operational value $\rho_T = 20$ g/cm³ then carries an effective systematic band of roughly 8–70 g/cm³, within which the MGEX held-out scale ($\lambda \approx 1.4$ – 1.9×10^3 km, ρ_T up to $\sim 5 \times 10^2$ g/cm³ if adopted at face value) is recorded as an unresolved epoch-scale tension pending longer MGEX baselines. The $\pm 47\%$ per-measurement statistical uncertainty from the single-center CODE fit is retained as a conservative upper bound on measurement-to-measurement scatter. The operational mapping from L_c to the projected covariance scale associated with $R_T(M_\oplus)$, together with the use of total Earth mass, are model-defining choices rather than additional empirical uncertainties.

Propagating the length-scale uncertainty alone:

$$\rho_T = 20 \pm 7 \text{ g/cm}^3 \text{ (35\% inter-center)}, \quad \rho_T \in [8, 70] \text{ g/cm}^3 \text{ (epoch/window systematic)}. \quad (7)$$

This range lies in the same condensed-matter density regime where electron degeneracy becomes dynamically relevant to the equation of state (Section 5.3) and provides the calibration scale for all subsequent tests. The key point is that the $M^{1/3}$ structural form is independent of ρ_T ; only the normalization changes. Even at the extremes ($L_c = 3,700$ km $\rightarrow \rho_T \approx 30$ g/cm³, or $L_c = 4,700$ km $\rightarrow \rho_T \approx 14$ g/cm³), the cross-scale consistency tests remain viable because the same ρ_T value must describe all systems simultaneously. The wider epoch/window band does not alter this structural conclusion — it re-prices the normalization confidence: cross-scale checks that pass at $\rho_T = 20$ g/cm³ must be re-evaluated under the widened prior, and the Level 1/2/3 claim hierarchy (Section 8) is designed to absorb exactly this class of anchor-level hit.

Discrimination from Geophysical Signals

A critical objection to long-range clock correlations is the presence of tropospheric and ionospheric delays. However, spectral power analysis reveals distinct scale separation:

- **Tropospheric covariance:** Decays at ~ 100 – 500 km (weather systems)
- **Ionospheric covariance:** Decays at ~ 500 – 2000 km (TEC structures)
- **Scalar field signature:** Persists as a covariance floor at ~ 4200 km

The 4200 km feature shows no counterpart in ionospheric or tropospheric delay products. Furthermore, the signal recovers an annual phase about 6° from Earth's aphelion velocity tangent (companion analysis); the CMB-apex template is anti-phase and is not

identified as a rest frame. Power spectral analysis of the GPS-only clean baseline confirms this as a persistent background covariance floor, not a geophysical artifact; multi-constellation validation shows consistent signatures.

Null Tests and Robustness

The GNSS correlation pattern survives multiple null tests:

- **Geophysical coupling:** Correlations persist after removing tidal, atmospheric, and ionospheric signals.
- **Orbital mechanics:** Pattern is independent of satellite constellation geometry.
- **Processing artifacts:** Raw RINEX analysis (Paper 3) constrains "black box" concerns.
- **Temporal stability:** 25-year consistency (Paper 2) disfavors transient instrumental effects.

The convergence of multi-center, multi-decade, and raw-data analyses identifies $L_c \approx 4200$ km as the canonical terrestrial screening length from the 25-year GNSS baseline (Papers 1–2), independent of theoretical interpretation. All cross-regime calibration and forward models in this paper adopt that value. Paper 14 provides an independent MGEX held-out verification on a ~ 1 yr combined-clock span (recovered scale $\approx 1,396 \pm 90$ km, $R^2 \approx 0.49$), confirming signal presence on a shorter baseline and different product; it is cited as verification only, not as the dimensional scale for ρ_T extrapolation. The scale tension between that figure and the 4,200 km anchor has since been tested directly (Paper 14 extension, Step 3.1): on strictly matched, verified-independent days the CODE and CNES processing chains return $\lambda = 2,744$ km and 1,933 km respectively ($\sim 30\%$ offset, the CNES value within 4% of the pooled 1,862 km), while quarterly windows of the same product scatter between 1,050 and 1,935 km — the held-out discrepancy is dominated by epoch/sampling instability of the short-span estimate, not by a product-dependent physical scale, and the independent-center agreement bounds the processing contribution to $\sim 30\%$ (Appendix B.8).

Phenomenological Transfer Sketch: Mapping the Static Profile to Clock Covariance

A central question is how the static Earth-sourced Temporal Topology profile produces the dynamic observable recovered in GNSS clock products: a spatial covariance length $L_c \approx 4200$ km. The calibration does not require GNSS satellites to orbit within a saturation shell, nor does it posit a literal shell at $R_T(M_\oplus)$. Instead, $R_T(M_\oplus)$ defines the geometric scale controlling the projected covariance kernel of the Earth-sourced exterior clock-transport field. In this sense, L_c is not an arbitrary fitted length: it is the observed network projection of the same mass-dependent scale used in the cross-regime $R_T \propto M^{1/3}$ construction. This subsection gives an order-of-magnitude argument — a transfer sketch, not a derivation — for this identification, together with the assumptions that a full demonstration would have to close (Appendix B.2).

What $R_T(M_\oplus)$ is, and is not

$R_T(M) = (3M/4\pi\rho_T)^{1/3}$ is the radius a body of mass M would have at uniform density ρ_T . For Earth this is ≈ 4146 km. It is a *scaling variable*, not a location: because Earth lies in the partial-transition regime, $R_T(M_\oplus)$ is used as a geometric scaling variable rather than as the radius of a literal material shell (Appendix B.2). The question the sketch addresses is what spatial scale governs the *exterior* field of such a partially screened source.

The exterior profile scale

In the static exterior problem, the available length scales are the source radius $R_\oplus$, the geometric saturation scale $R_T(M_\oplus)$, and the ambient-medium Compton wavelength $\lambda_C(\rho_{\text{amb}})$ (very long in the low-density exterior ($T^{(m)} \approx 0$)). Under the corpus's screening ontology the transition in the screening function $S(\rho)$ is *gradual*, spanning $\Delta\rho/\rho_T \sim \mathcal{O}(1)$ — a continuous Temporal Topology with no thin shells (Paper 0, A4). For such a gradual transition, the exterior field of the partially screened source carries no structure sharper than the source scale itself, and the characteristic scale over which the effective source charge accumulates is $R_T(M)$: radii enclosing mean density above the transition range contribute suppressed source charge, radii beyond contribute unsuppressed charge, and the changeover in the *integrated* source — not in any local shell — occurs over a radial span of order R_T . (Note that a *steep* transition would instead produce thin-shell behaviour with structure on scales $\ll R_T$; the gradual ontology is therefore not incidental to this argument but required by it.) Both $R_\oplus = 6371$ km and $R_T(M_\oplus) = 4146$ km lie within a factor ~ 1.5 , so at sketch precision the exterior profile varies on the scale of a few thousand kilometres, with $R_T(M)$ — not $R_\oplus$ — as the mass-scaling variable the cross-scale law requires.

From a static profile to a time-series covariance

A deterministic static field produces constant clock offsets, which clock-bias estimation absorbs; by itself it generates no time-series covariance. The sketch therefore requires one explicit assumption:

Modulated-sampling assumption. The residual variance analysed in Papers 1–3 arises from temporal modulation of each station's sampling of the static exterior profile — diurnal rotation of the ground network through the profile's anisotropy, satellite orbital motion, and tropospheric/orientation geometry that multiplies the local conformal rate $A(\vec{\phi}(\mathbf{x}, t))$. Two clocks then acquire *correlated* time variance to the extent that their modulated samples traverse the same large-scale field structure.

Writing $\delta\tau_i(t) \approx \tau_0(\beta_A/M_{\text{Pl}})\bar{\phi}(\mathbf{x}_i(t))$ for the modulated residual, the equal-time network covariance $\langle\delta\tau_A\delta\tau_B\rangle_t$ inherits its baseline dependence from the spatial structure of $\bar{\phi}$: stations separated by d small compared with the profile's variation scale sample nearly the same field excursion and correlate strongly; stations separated by d large compared with that scale decorrelate. The decay length of the network covariance is then set by the exterior profile scale,

$$L_c \sim \mathcal{O}(1) \times R_T(M_\oplus), \quad (8)$$

with the $\mathcal{O}(1)$ factor determined by the unspecified completion (profile shape, anisotropy spectrum, sampling geometry). This is the entire quantitative content of the sketch: a coincidence of scales with an undetermined order-unity prefactor, *not* a prediction of $L_c = R_T$ to any stated precision.

Implications for the calibration

This sketch binds the clock observable (L_c) to the geometric scaling variable (R_T) at the level of dimensional analysis under two explicit assumptions: the gradual-transition ontology and the modulated-sampling mechanism. It does not prove that the GNSS correlation is caused by the TEP scalar — that remains the empirical hypothesis tested by null-model exclusion (Papers 1–3) — and it does not by itself discriminate $R_T(M)$ from $R_\oplus$ as the controlling scale, since the two differ by only a factor ~ 1.5 for Earth. What *would* discriminate them is the mass-scaling test: $R_T \propto M^{1/3}$ while a source-radius scale does not follow that law across the astrophysical systems of Sections 3–7. The single-potential numerical demonstration that would replace this sketch with a derivation — an explicit exterior profile and sampling kernel under a specified completion, with the quartic normalization held at the corpus self-coupling value $\lambda = 7.5 \times 10^{-66}$ (the scalar-potential coefficient, not the covariance length denoted λ in this section) and ρ_T fixed by the calibrated L_c — remains the open target recorded in Appendix B.2.

For detailed methodology and validation, see Appendix A and Papers 1–3.

3. The SPARC Galaxy Analysis: Phantom Mass as Unscreened Time-Field

Galaxies can be understood as "baryon-anchored" solitons—where the same scalar field structure calibrated from terrestrial GNSS data is pinned to a baryonic mass concentration. This section extends the test to galactic scales, where the $M^{1/3}$ scaling is expected to govern the transition between screened (Newtonian) and unscreened (phantom mass) regimes. The SPARC database provides an ideal testbed: 175 disk galaxies with high-quality rotation curves and well-constrained baryonic mass distributions. The theoretical basis for the phantom mass interpretation is developed in detail in Smawfield (2025e), which demonstrates how differential proper-time accumulation in gravitational lensing can produce apparent dark matter phenomenology.

Phantom Mass Interpretation

In the TEP framework, the apparent "dark matter" observed in galaxy rotation curves is identified as *Phantom Mass*—defined as the apparent mass derived from temporal shear under the assumption of isochrony (Smawfield 2025e). It is not a particle species but a geometric manifestation of the unscreened proper-time field. A critical distinction must be emphasized: "Phantom Mass" is an *observational inference artifact* (analogous to a refractive index effect), not a "ghost field" in the quantum field theory sense. Ghost fields imply negative kinetic energy and Hamiltonian instability; phantom mass carries no such pathology. It arises simply from modeling a bi-metric spacetime with a single-metric prior—the "missing mass" is the unmodeled temporal shear contribution.

The transition radius R_{trans} marks the boundary between two regimes:

- Inside R_{trans} : The enclosed mean baryonic density exceeds the critical screening density ($\bar{\rho} > \rho_{\text{trans}}$). The time-field is screened, and gravity follows Newtonian predictions based on visible baryonic matter.
- Outside R_{trans} : The enclosed mean density drops below ρ_{trans} . The time-field becomes unscreened, and the refractive proper-time gradient produces an apparent gravitational excess—the "phantom mass" conventionally attributed to dark matter.

If this interpretation is correct, the radius at which rotation curves diverge from Newtonian prediction should scale as $R_{\text{DM}} \propto M_{\text{bar}}^{1/3}$. While the fundamental core saturation occurs at the proximity scale observationally proxied by $\rho_T \approx 20 \text{ g/cm}^3$, the halo transition is governed by a much lower proximity scale, proxied by ρ_{trans} , characteristic of the diffuse field tail.

Invariant: ρ_T denotes the universal saturation density of the scalar sector that fixes the compact soliton scale via $R_T \propto (M/\rho_T)^{1/3}$. By contrast, ρ_{trans} is an *emergent* screening threshold for the onset of halo phenomenology in diffuse environments; it is not treated as a second fundamental constant and can depend on coupling, geometry, and baryonic structure.

Theoretical Expectation from ρ_T

The TEP scaling law $R_T \propto M^{1/3}$, anchored by the GNSS calibration $\rho_T \approx 20 \text{ g/cm}^3$, has a direct consequence for galactic structure. In the TEP framework, the dark-matter onset radius R_{DM} is identified with the radius at which the enclosed mean baryonic density drops below the emergent halo threshold ρ_{trans} . Because ρ_{trans} is an emergent threshold intended to be tied to the same scalar

potential that fixes ρ_T — a linkage whose derivation is the open target of the Formal Link — dimensional analysis yields the same mass scaling:

$$R_{\text{DM}} \propto \left(\frac{M_{\text{bar}}}{\rho_{\text{trans}}} \right)^{1/3} \propto M_{\text{bar}}^{1/3}. \quad (9)$$

Thus the TEP framework *expects* $\alpha = 1/3$ for the DM onset scaling. This is a theoretical consequence of the same saturation scale that fixes the terrestrial correlation length, not a new free parameter. The SPARC analysis below tests whether the empirical exponent α_{SPARC} is consistent with this expectation.

The analysis was performed *post hoc*: the SPARC data were examined after the TEP framework was established, so the comparison is a consistency check rather than a blind prediction. A measured exponent $\alpha_{\text{SPARC}} \neq 1/3$ (outside measurement uncertainty) would challenge the framework, while agreement supports it.

Methodology

The analysis proceeds as follows for each of the 175 SPARC galaxies:

1. Calculate the total baryonic mass: $M_{\text{bar}} = M_* + 1.33M_{\text{HI}}$, where $M_* = (M/L)_{3.6\mu} \times L_{3.6}$ with $(M/L)_{3.6\mu} = 0.5 M_{\odot}/L_{\odot}$ for disk populations.
2. Compute the expected Newtonian rotation velocity: $V_{\text{bar}}^2 = V_{\text{gas}}^2 + (M/L)_{\text{disk}} V_{\text{disk}}^2 + (M/L)_{\text{bulge}} V_{\text{bulge}}^2$.
3. Identify the mass discrepancy onset radius R_{DM} : the first radius where $V_{\text{obs}}/V_{\text{bar}} > 1.3$. Radial bins with nonpositive or unreliable baryonic velocity ($V_{\text{bar}} \leq 5 \text{ km s}^{-1}$, including $V_{\text{bar}}^2 \leq 0$ from negative model contributions) are excluded from the ratio; galaxies whose ratio never exceeds the threshold contribute no R_{DM} and are dropped from the fit — a selection, disclosed here — leaving 167 of 175 systems at the fiducial threshold.
4. Fit the power-law relation $R_{\text{DM}} = k \cdot M_{\text{bar}}^{\alpha}$ across the full sample.

Results

Of the 175 SPARC galaxies, 167 yield valid mass discrepancy onset radii spanning five decades in baryonic mass (10^7 – $10^{12} M_{\odot}$).

Robustness to Onset Definition

To address concerns that the "onset radius" is a noise-sensitive functional, a bootstrap analysis (1000 resamples) was performed across multiple definition criteria. The scaling exponent α_{SPARC} remains consistent with the TEP expectation ($\alpha_{\text{SPARC}} = 1/3$) across a wide range of velocity ratio thresholds:

Definition (Threshold)	Fitted Exponent (α_{SPARC})	Consistency with 1/3
Loose ($V_{\text{obs}}/V_{\text{bar}} > 1.1$)	0.274 ± 0.036	1.6σ
Fiducial ($V_{\text{obs}}/V_{\text{bar}} > 1.3$)	0.372 ± 0.041	1.0σ
Strict ($V_{\text{obs}}/V_{\text{bar}} > 1.5$)	0.413 ± 0.045	1.8σ

While individual definitions shift the normalization, the slope consistently clusters around the $M^{1/3}$ expectation. An alternative definition based on acceleration thresholds ($a < a_0$) yields a steeper slope ($\alpha_{\text{SPARC}} \approx 0.5$). This estimator is the $M^{1/3}$ vs. $M^{1/2}$ (TEP vs. MOND-like) discrimination in miniature; the fact that the verdict is currently estimator-dependent is noted as an open systematic. Because the acceleration-threshold method is degenerate with the MOND acceleration scale itself, the kinematic divergence test is retained as the fiducial estimator. This preference is theoretically motivated: TEP models phantom mass as a geometric effect driven by temporal shear (a density/proximity threshold) rather than an instantaneous acceleration threshold, so a purely kinematic divergence estimator is intrinsically less biased toward MOND-like phenomenologies.

$$R_{\text{DM}} = k \cdot M_{\text{bar}}^{\alpha_{\text{SPARC}}} \quad (10)$$

$$\alpha_{\text{SPARC}} = 0.355 \pm 0.043 \text{ (stat)} \pm 0.07 \text{ (definition)} \quad (11)$$

The bootstrap-marginalized exponent deviates from the TEP expectation of $\alpha_{\text{SPARC}} = 1/3 = 0.333$ by approximately 0.3σ under the combined uncertainty—consistent within typical statistical tolerance. The correlation coefficient $r \approx 0.6$ indicates a significant relationship. Importantly, this result is robust: individual threshold choices yield exponents ranging from 0.28 to 0.42, but the ensemble average converges near $1/3$.

Size-Scaling Confound Test

The $M^{1/3}$ onset exponent is not by itself discriminating: for approximately self-similar exponential disks, a fixed velocity-ratio crossing occurs at a nearly fixed multiple of the disk scale length R_d , so the galaxy mass–size relation can produce the same exponent without any density threshold. The confound is tested directly on the same SPARC sample (pipeline step 8). The in-sample mass–size relation is $R_d \propto M_{\text{bar}}^{0.322 \pm 0.017}$ ($r = 0.821$, $n = 175$), which alone predicts an onset exponent ≈ 0.32 — within the stated uncertainty of the measured 0.355 ± 0.043 . Comparing the two candidate invariants per galaxy — the size ratio R_{DM}/R_d and the enclosed mean baryonic density $\bar{\rho}(< R_{\text{DM}}) = 3M_{\text{bar}}(< R_{\text{DM}})/(4\pi R_{\text{DM}}^3)$, with $M_{\text{bar}}(< R)$ estimated spherically as $V_{\text{bar}}(R)^2 R/G$ — the size invariant is mass-independent at the fiducial threshold (slope 0.056 ± 0.044 , $p = 0.21$; median $R_{\text{DM}}/R_d = 0.64$), while the enclosed density retains a significant mass correlation (slope 0.190 ± 0.068 , $p = 5.7 \times 10^{-3}$). The pattern is threshold-dependent: at $V_{\text{obs}}/V_{\text{bar}} > 1.2$ the size null wins decisively (density slope 0.302 , $p = 5.4 \times 10^{-6}$), while at 1.4 the two are comparable (0.086 vs 0.084).

The onset scaling is therefore consistent with, but not by itself discriminating evidence for, the density-threshold mechanism: a fixed fraction of the disk scale length produces the same exponent on this sample. The discriminating weight of the SPARC test accordingly rests on the two channels the size-scaling null cannot supply — the residual correlations with baryonic content (step 7) and the open normalization linkage of the Formal Link, which the confound makes the sharper falsification condition: if ρ_{trans} cannot be derived from the scalar-sector parameters (λ, β_A) , or if a forward-modelled amplitude fails, the onset-scaling result contributes no independent support.

Phantom-Mass Amplitude and Profile Test

The onset radius alone does not test the mechanism — any threshold criterion produces an onset. The discriminating content of the phantom-mass claim lies in the amplitude and the radial profile, both of which are evaluated directly on the same SPARC sample (pipeline step 9). For a conformal coupling of charge β_A , an unscreened scalar channel contributes a centripetal term of order $2\beta_A^2$ times the Newtonian one, giving the plateau ratio $V_{\text{obs}}/V_{\text{bar}} = \sqrt{1 + 2\beta_A^2 \mathcal{S}_{\text{eff}}}$, which for the bare coupling $|\beta_A| = 1$ at full saturation is $\sqrt{3} \approx 1.73$. Measured on the outer third of each curve ($n = 164$ galaxies with ≥ 6 valid points and $V_{\text{bar}} > 5 \text{ km s}^{-1}$), the median plateau ratio is 1.98 (16–84th percentile 1.38 – 2.49) — order-consistent with the bare-coupling saturation value but systematically above it: the implied effective response is $\mathcal{S}_{\text{eff}} = 1.46$ at the median, and 71% of galaxies require $\mathcal{S}_{\text{eff}} > 1$. The plateau amplitude is therefore quantified rather than asserted: it sits within a factor of order unity of the bare-coupling prediction and the residual excess is precisely the response-normalization demand already flagged in the Formal Link — the amplitude test converts that open derivation into a measured gap of factor ~ 1.5 – 2 rather than an unconstrained parameter.

The radial profile supplies the second discriminator. Defining the enhancement $E(R) = V_{\text{obs}}^2/V_{\text{bar}}^2 - 1 = M_{\text{ph}}(R)/M_{\text{bar}}(< R)$, the phantom fraction is required by the transition picture to grow outward (screened inside, unscreened outside), whereas a constant coupling rescaling of the baryonic potential predicts $d \log E/d \log R = 0$. The measured profile rises: median slope 0.46 across 161 galaxies with sufficient baseline, positive in 133 (Wilcoxon signed-rank $p = 1.0 \times 10^{-17}$). The rising phantom fraction is the expected morphology of a baryon-anchored screening transition and is inconsistent with a global multiplicative rescaling; it is not, by itself, separable from acceleration-threshold phenomenologies, which predict the same qualitative rise. What the two tests jointly establish is that the SPARC sample carries both an amplitude-level and a profile-level constraint on the mechanism — the onset scaling is no longer the sole empirical content of the claim.

Cross-Regime Consistency

The significance of this result emerges when combined with the Temporal Topology screening analysis (Section 4). The complete screening hierarchy spans ~ 15 orders of magnitude in density—from gas giants ($\rho \sim 1 \text{ g/cm}^3$) to binary pulsars ($\rho \sim 10^{14} \text{ g/cm}^3$)—while the fitted regression uses the 11 dense objects ($\rho > \rho_T$) spanning ~ 12 orders of magnitude ($\sim 10^2$ – 10^{14} g/cm^3). The consistency of this exponent across such a range is striking, though the high R^2 is partly a consequence of the definitions used (see Box 4.1). The result is consistent with the hypothesis that the $M^{1/3}$ scaling reflects a genuine feature of the gravitational sector, though systematic uncertainties in rotation curve fitting and baryonic modeling remain.

Combined with the SPARC galaxy scaling ($\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ (stat) ± 0.07 (definition)), the RBH-1 crossover consistency (within $\sim 25\%$ combined uncertainty), and the Milky Way dark-matter onset ($R_{\text{DM}} \approx 3 \text{ kpc}$, observed ~ 3 – 5 kpc), these results suggest a universal density-limited structure may span 18 orders of magnitude in mass—from Earth to galaxy.

Physical Interpretation

The observed $M^{1/3}$ scaling admits a natural interpretation in terms of density-limited screening. If the screening transition occurs at a characteristic density ρ_{trans} , dimensional analysis requires:

$$R_{\text{trans}} \sim \left(\frac{M}{\rho_{\text{trans}}} \right)^{1/3}. \quad (12)$$

Fixing $\alpha_{\text{SPARC}} = 1/3$ and fitting only the normalization yields $k \approx 7.9 \times 10^{-4} \text{ kpc}/M_{\odot}^{1/3}$. This normalization corresponds to an emergent screening density $\rho_{\text{trans}} \approx 0.5 M_{\odot}/\text{pc}^3$ ($\sim 3 \times 10^{-23} \text{ g/cm}^3$) recovered from the fitted normalization k via Eq.~(12). A completion is conjectured to derive ρ_{trans} from the scalar-sector normalization (the quartic coupling λ and the coupling β_A), but this remains an open target (see Formal Link below). This value is within an order of magnitude of typical disk densities at the optical radius ($0.01\text{--}0.1 M_{\odot}/\text{pc}^3$), providing a physically reasonable anchor for the screening mechanism.

Note: This galactic transition density ρ_{trans} is distinct from the fundamental core saturation scale $\rho_T \approx 20 \text{ g/cm}^3$ calibrated in Section 2 and cross-checked in Section 5. The core density ρ_T governs the compact object size (soliton surface), while ρ_{trans} governs the onset of the diffuse halo effect (screening radius). Both scales follow the same $M^{1/3}$ structural form, consistent with the environment-dependent nature of the theory across 15 orders of magnitude in density.

Formal Link: From Core Saturation to Galactic Transition

The galactic transition density ρ_{trans} is intended to emerge from the same scalar field profile that fixes the core saturation scale ρ_T ; at present it is an independent empirical normalization recovered from the fitted k , and the derivation sketched below is the stated open target rather than an established result. In the TEP framework (Box 6.5), the scalar field obeys

$$\nabla^2 \phi = V'(\phi) + \alpha(\phi)\rho, \quad (13)$$

with effective mass $m_{\text{eff}}^2(\rho) = V''(\phi_{\text{min}})$. The field profile transitions from saturated ($\rho \sim \rho_T$, R_T scale) to screened ($\rho \gg \rho_T$, Compton wavelength $\lambda_C \ll R_T$) to unscreened ($\rho \ll \rho_T$, $\lambda_C \gg R_T$). The galactic halo onset occurs where the enclosed mean baryonic density $\bar{\rho}(< R) = 3M_{\text{bar}}(< R)/(4\pi R^3)$ drops to the value ρ_{trans} at which the scalar-field-induced gravitational acceleration becomes comparable to the baryonic contribution — an enclosed-mean-density criterion, consistent with the form of Eq.~(12); a local-density formulation would require a specified field/density profile and is not used here. Because the field profile is governed by the same scalar sector — the quartic normalization λ and coupling β_A — that underlies the compact-object saturation, ρ_{trans} is expected to be a derived consequence of these parameters rather than a free fit; pending that derivation it is an independent empirical input. The SPARC analysis measures the structural exponent α_{SPARC} and normalization k ; the corresponding ρ_{trans} is then recovered from k via Eq.~(12) as a consistency check.

Connection to the MOND Acceleration Scale

The fitted screening density implies a characteristic transition acceleration. At the transition radius, the gravitational acceleration is:

$$g_{\text{TEP}} = \frac{GM}{R_{\text{trans}}^2} \approx 5 \times 10^{-10} \text{ m/s}^2.$$

This is within a factor of 4 of the MOND acceleration scale $a_0 \approx 1.2 \times 10^{-10} \text{ m/s}^2$. The near-coincidence suggests that the TEP screening mechanism and the empirical MOND phenomenology may share a common origin: both describe the transition from screened (Newtonian) to unscreened (modified) gravity at a characteristic acceleration scale set by cosmological boundary conditions.

Connection to the Radial Acceleration Relation

The SPARC database is the source of the celebrated Radial Acceleration Relation (RAR), which shows a tight empirical coupling between observed acceleration g_{obs} and baryonic acceleration g_{bar} (McGaugh et al. 2016). The RAR has been interpreted as evidence for modified gravity (MOND) or as a consequence of dark matter halo profiles fine-tuned to baryonic distributions.

The TEP framework offers a third interpretation: the RAR may emerge from the environment-dependent screening of the proper-time field. At high accelerations (high densities), screening is complete and $g_{\text{obs}} \approx g_{\text{bar}}$. At low accelerations (low densities), the unscreened time-field gradient contributes additional apparent gravity, producing the characteristic upturn in the RAR. The $M^{1/3}$ scaling of the transition radius is the spatial manifestation of this acceleration-dependent screening.

A fixed-normalization test applies the existing alternative low-gradient kinetic realization directly to the SPARC mass models (pipeline steps 10–13). Its response $P_X(\xi) = k\sqrt{\xi} + 2\xi$, with $\xi = |X|/\Lambda_X^4$, differs from the unit-term baseline $1 + 2\xi$ of Paper 0 §7 and yields the derived flat-tail and baryonic Tully–Fisher limits. Step 10 fixes $k = 4\sqrt{2} g_t/a_0 \simeq 16.03$ using the adopted empirical $a_0 = 1.2 \times 10^{-10} \text{ m/s}^2$. No galaxy-specific parameter is adjusted in this comparison: the normalization is prespecified from that empirical scale, not independently derived from the wide-binary/Solar crossover. The algebraic flux law $\bar{u}P_X(\bar{u}^2/2) = g_{\text{bar}}/g_t$ yields the pointwise prediction $g_{\text{obs}} = g_{\text{bar}} + 2g_t\bar{u}$. Evaluated on all 3,389 measured radii across 175 galaxies ($\Upsilon_d = 0.5$, $\Upsilon_b = 0.7$), this curve tracks the RAR across four decades in g_{bar} (rms log-residual 0.219 dex, against 0.281

dex for the unit-term baseline and 0.199 dex for the empirical MOND interpolating function), with a uniform $\approx 20\%$ normalization offset (median -0.086 dex) rather than a shape-dependent failure; a single free normalization prefers $a_{\text{eff}} \approx 0.52 a_0$ ($k \approx 31$). The same fixed sector predicts the onset-radius scaling $R_{\text{onset}} = \sqrt{GM_b/a_{\text{eff}}}$: over the 54 galaxies bracketing the crossing, the fixed-slope-1/2 intercept reproduces the predicted normalization to 0.022 dex (rms 0.277 dex). The baryonic Tully–Fisher outer-edge evaluation returns median $v_{\text{obs}}/v_{\text{pred}} = 0.910$ (rms 0.093 dex) over 135 galaxies — consistent with the same normalization offset in the same $\bar{u}$ regime and with mass-to-light systematics. A field-value coupling $K(\phi) = e^{c\psi}$ improves the RAR match monotonically (rms $\rightarrow 0.199$ dex at $c = 3$) but is excluded by a sign tension with the wide-binary pair channel, which requires the opposite sign, so the monotone form is not adopted. The residual offset is attributed to the identified corrections — disk geometry, Υ systematics, and ambient-field vertex factors — rather than treated as a free refit.




Figure 1: Representative SPARC rotation curves. Observed velocities diverge from baryonic predictions near the TEP transition radius R_{trans} . Shaded regions indicate the screened (Newtonian) zone. The agreement between R_{trans} and the observed onset R_{DM} supports the screening interpretation.

Linking RBH-1 to Galactic Halos

The SPARC analysis identifies a cross-scale consistency between the RBH-1 wake phenomenon and galactic dark matter halos:

- RBH-1 represents a "naked saturation"—a Temporal Topology saturation not anchored to a baryonic host, propagating freely through the intergalactic medium and inducing star formation via its time-dilation wake.
- Galactic halos represent "baryon-anchored solitons"—the same temporal field structure, but centered on and shaped by the baryonic mass distribution of the host galaxy.

Both phenomena are consistent with the same $M^{1/3}$ scaling law, calibrated from terrestrial GNSS observations. The universality of this scaling across planetary, stellar, galactic, and cosmological mass scales constitutes a central theoretical consequence of the Temporal Equivalence Principle.

Residual Analysis: Discriminating Baryonic Feedback vs Field Theory

The RMS scatter of 0.48 dex around the $M^{1/3}$ relation indicates substantial galaxy-to-galaxy variation. A critical test distinguishes whether this scatter arises from baryonic feedback processes (Standard Model) or from the intrinsic field theory mechanism (TEP): correlate the residuals with baryonic properties versus screening proxies.

Methodology

For each of the 167 SPARC galaxies with valid mass discrepancy radii, the residual from the $M^{1/3}$ expectation is calculated as:

$$\text{Residual} = \log_{10}(R_{\text{DM}}/R_{\text{exp}}) \quad (14)$$

where $R_{\text{exp}} = k \cdot M_{\text{bar}}^{1/3}$ with $k = 7.86 \times 10^{-4} \text{ kpc}/M_{\odot}^{1/3}$. These residuals are then correlated with:

- *Baryonic properties.* Gas fraction (f_{gas}), surface brightness (Σ), and inclination, treated as proxies for feedback efficiency, star formation history, and observational systematics.
- *Screening proxy.* Central density (ρ_{central}), treated as a proxy for local screening strength independent of R_{DM} .

Results

The residual correlations reveal a clear pattern:

Property	Correlation (r)	P-value	Interpretation
Gas Fraction	−0.164	0.034	Weak, marginally significant
Surface Brightness	−0.129	0.096	Weak, not significant
Inclination	+0.297	0.0001	Moderate (observational systematic)
Central Density	−0.108	0.164	Weak, not significant

Interpretation

The maximum baryonic correlation is $|r| = 0.297$ (inclination), which likely reflects observational systematics in rotation curve deprojection rather than physical feedback. The gas fraction and surface brightness—the primary tracers of baryonic feedback efficiency—show weak correlations ($|r| < 0.2$) that are either marginally significant or non-significant.

The central density (screening proxy) shows no significant correlation ($r = -0.108$, $p = 0.16$). The weak correlation with central surface density is consistent with (though does not by itself prove) a geometric origin: if baryonic feedback were the dominant driver of the scatter in this particular estimator, stronger covariance with surface brightness would be expected. The present results remain compatible with a substantial contribution from measurement uncertainty and geometric effects.

Summary: the residuals do not show strong correlations with gas fraction or surface brightness, while inclination shows a moderate correlation that is plausibly attributable to rotation-curve deprojection systematics. Central density does not show a statistically significant correlation in this analysis.




Figure 2: Residual analysis of the SPARC $M^{1/3}$ scaling. (a) Scaling relation colored by residual. (b-e) Correlations with baryonic properties and screening proxies. (f) Gaussian residual distribution ($\sigma = 0.48$ dex). (g) Correlations are weak in amplitude ($|r| < 0.3$); inclination is statistically significant but plausibly reflects deprojection systematics rather than baryonic feedback physics.

Scatter Quantification and Comparative Analysis

The observed scatter of $\sigma = 0.48$ dex around the $M^{1/3}$ relation is substantial (corresponding to a factor of ~ 3 variation). Quantifying the origin of this scatter is essential for distinguishing the TEP signal from competing hypotheses.

Intrinsic vs. Measurement Scatter

Standard error budgeting suggests that a significant fraction of this scatter is observational. Distance uncertainties in the SPARC sample typically range from 10–20%, which propagate to a ~ 0.1 – 0.2 dex uncertainty in R_{DM} . Inclination corrections and non-circular motions contribute additional noise. However, the residual scatter (~ 0.3 dex) likely reflects intrinsic variation in the galaxy population.

Comparison with MOND and LCDM+Feedback

MOND: The Modified Newtonian Dynamics framework typically yields a tighter scatter (~ 0.13 dex) in the Radial Acceleration Relation (RAR). This is expected because MOND modifies gravity based on instantaneous local acceleration, a continuous variable. In contrast, this analysis extracts a single discrete parameter (R_{DM}) from each rotation curve, a method inherently more sensitive to local irregularities and noise than the integrated RAR analysis. The larger scatter in R_{DM} does not invalidate the mean scaling; rather, it indicates that the "onset radius" is a noisy estimator of the underlying screening transition.

LCDM + Feedback (Null Hypothesis): The standard cosmological model relies on baryonic feedback to explain rotation curve diversity. In this scenario, the "dark matter onset" is not a fundamental scale but an emergent property of halo assembly and feedback history. Hydrodynamic simulations (e.g., NIHAO, FIRE) typically predict a scaling of $R_{DM} \propto M^\alpha$ with $\alpha \approx 0.3$ – 0.4 , broadly

consistent with observation. However, they offer no account of why the galactic relation shares its structural exponent with the compact-object scaling. The fitted normalization corresponds to $\rho_{\text{trans}} \approx 0.5 M_{\odot}/\text{pc}^3$ ($\sim 3 \times 10^{-23} \text{ g/cm}^3$), twenty-four orders of magnitude below ρ_T : what is presently demonstrated is a shared structural exponent between two independently normalized scales, not a normalization link, and whether ρ_{trans} follows from the same scalar-sector normalization that underlies the ρ_T saturation law is the open derivation target recorded in the Formal Link.

Distinctive Cross-Scale Linkage

While MOND provides a tighter fit to galaxy kinematics alone, it offers no explanation for the GNSS clock correlations ($L_c \approx 4200$ km). LCDM+Feedback can accommodate the galaxy scaling but treats the normalization as a free parameter, offering no predictive link to other scales. TEP is distinctive in that it poses the normalization of the galactic relation ($k \approx 7.9 \times 10^{-4} \text{ kpc}/M_{\odot}^{1/3}$, equivalently ρ_{trans}) as a quantity to be derived from the same GNSS-anchored scalar sector — a stated falsification condition rather than an established link: the measured normalization is an independent empirical input pending that derivation, within the stated systematic uncertainty.

Reproducibility

Full analysis code, data products, and methodology documentation are available in the public repository:

- *Repository*. github.com/matthewsmawfield/TEP-UCD
- *SPARC Scaling Analysis*. `scripts/steps/step_4_sparc_analysis.py`
- *SPARC Residual Analysis*. `scripts/steps/step_7_sparc_residuals.py`
- *Input Data*. `data/sparc/` (SPARC database tables)

Methodological Details

The "onset radius" R_{DM} is defined as the *first radial bin* where $V_{\text{obs}}/V_{\text{bar}} > 1.3$ (fiducial threshold). Specifically:

1. For each galaxy, the rotation curve is evaluated at the native SPARC radial sampling (not interpolated).
2. The velocity ratio $V_{\text{obs}}/V_{\text{bar}}$ is computed at each radius.
3. R_{DM} is the radius of the first bin exceeding the threshold.
4. Galaxies with non-monotonic or noisy $V_{\text{obs}}/V_{\text{bar}}$ profiles are retained; the "first crossing" definition is robust to subsequent fluctuations.
5. Galaxies that never exceed the threshold (8 of 175) are excluded from the fit.

The headline value $\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ is the threshold-marginalized bootstrap estimate, obtained by resampling galaxies while marginalizing over the accepted onset definitions. The fiducial ($V_{\text{obs}}/V_{\text{bar}} > 1.3$) definition alone gives 0.372 ± 0.041 ; the difference is retained as onset-definition systematic uncertainty. Varying the threshold from 1.1 to 1.5 yields exponents from 0.28 to 0.42, bracketing the headline estimate.

4. Temporal Topology Screening Analysis

If a scalar field permeates spacetime, why has it not been detected in precision gravitational experiments? This section answers that question. The key insight is that the scalar field's influence depends on the geometric overlap of topological charge cores: in environments where charge cores are tightly packed, the field is suppressed—a phenomenon called Temporal Topology screening. Rather than invoking discrete thin-shell boundaries, screening operates via the continuous spatial profile of the scalar field (Temporal Topology). The tight geometric packing in deep potential wells suppresses the local field gradient (Temporal Shear), ensuring short-range Temporal Shear suppression while leaving the field light cosmologically. General Relativity is recovered in the regimes where it has been tested most stringently.

Screening in TEP is represented at the theory level by the environmental operator $S_{\Sigma}(\mathcal{E})$. Quantities such as ρ_T , $R_T(M)$, $S_{\oplus}(r)$, compactness Φ/c^2 , local stellar density, geometric coherence length, and channel-specific response coefficients are domain-specific projections of $\mathcal{E}$, not independent screening mechanisms and not interchangeable universal thresholds. Each is an observational transfer model that parameterizes the same underlying operator in a regime-appropriate form.

The screening factor S quantifies this suppression. It is defined as the ratio of the Temporal Topology saturation radius (where the scalar field saturates) to the physical radius of the object:

$$S = \frac{R_T}{R_{\text{phys}}}. \quad (15)$$

The quantity S is used here as a geometric proxy for how deeply the baryonic source is embedded within a saturated soliton configuration. When $S \gg 1$, the physical object occupies a small region relative to the saturated field scale; in this regime the phenomenology assumes that non-linear response in the scalar sector suppresses gradients in the dense interior, recovering GR to high precision in local dynamics. When $S \sim 1$, the system lies near the transition between strongly screened and weakly screened behavior. When $S \ll 1$, the Temporal Topology saturation scale is smaller than the object and the saturated region does not envelop the full baryonic configuration; in this regime the model does not assume strong Temporal Topology suppression a priori, and constraints must be assessed case-by-case.

$$\alpha_{\text{PPN}}^{\text{eff}} = \alpha_0 F(S, \rho, \Phi, \nabla\phi), \quad F \rightarrow 0 \quad (16)$$

in the screened or source-charge-suppressed limit. The geometric factor S is not itself the observable PPN coupling. Precision tests constrain $\alpha_{\text{PPN}}^{\text{eff}}$; S is used here only as a proxy for the nonlinear suppression regime.

The near-transition regime $S \sim 0.4\text{--}0.7$ is resolved by the corpus's channel-projection structure (Paper 0). The screening response is not a single function of S : clock-comparison and transport observables probe the amplitude projection S_A , while orbital and geodesic observables probe the source-charge/shear projection S_Σ , which is suppressed more strongly for the pairwise configurations used in ephemeris tests. The same geometric S therefore places different channels on different sides of their respective thresholds: Earth's $S \approx 0.66$ admits a detectable clock-covariance signal (C_A covariance channel, the GNSS calibration of Section 2) while the LLR, Cassini, and MESSENGER orbital observables at $S = 0.42\text{--}0.65$ remain GR-consistent through the S_Σ projection. The reading of sub-unity S is uniform throughout this paper: $S < 1$ denotes that the saturation region does not envelop the object, i.e. the unscreened side of the transition; the observable amplitude in any given channel is then set by that channel's projection, not by S alone.

4.1 The White Dwarf Stress Test

White Dwarfs are the ideal stress test for two reasons. First, their structure is determined by quantum mechanics (electron degeneracy pressure), not thermal physics, so their mass-radius relation is calculable from first principles. Second, their mass-radius scaling runs in the opposite direction to the saturation scaling (see Figure 3):

- White Dwarf (Chandrasekhar): $R_{WD} \propto M^{-1/3}$ — heavier stars are smaller
- Soliton (TEP): $R_{sol} \propto M^{+1/3}$ — heavier fields are larger




Figure 3: The "Scissors" Effect. The defining visual signature is the divergence of scales: as mass increases, the white dwarf physical radius shrinks ($R_{WD} \propto M^{-1/3}$) while the saturation radius grows ($R_{sol} \propto M^{1/3}$). This divergence drives the screening factor $S = R_T/R_{phys}$ to large values ($S \gg 1$), recovering General Relativity within the star.

These opposing scalings create a "scissors" pattern. As mass increases, the saturation radius grows while the physical radius shrinks—the screening factor diverges. This is the signature of Temporal Topology flattening.

For Sirius B ($M \approx 1.02M_\odot$, $\rho \approx 2.4 \times 10^6 \text{ g/cm}^3$):

$$R_{phys} \approx 5,800 \text{ km} \quad (\text{observed}) \quad (17)$$

$$R_{sol} \approx 289,000 \text{ km} \quad (\text{TEP expectation}) \quad (18)$$

$$\text{Screening Factor} = \frac{R_T}{R_{phys}} \approx 50\times. \quad (19)$$

With a mean density $\rho \approx 2.4 \times 10^6 \text{ g/cm}^3$ far exceeding the saturation threshold $\rho_T \approx 20 \text{ g/cm}^3$, the saturation field extends 50 times beyond the physical surface of the star. The dense baryonic matter is deeply embedded within the scalar field's saturation core, consistent with strong screening under the stated model assumptions. Within the star, GR dynamics are recovered; the Keplerian mass measured from binary orbital motion is the true baryonic mass.

4.2 The Empirical Screening Law

Extending this analysis across 26 astrophysical objects (planets, brown dwarfs, main sequence stars, white dwarfs, neutron stars, and binary pulsars) yields a compact summary of how the defined screening factor varies with density. Restricting the regression fit to the 11 dense objects ($\rho > \rho_T$, excluding planets and black holes) gives:

$$S \propto \rho^{\beta_{scr}}, \quad \beta_{scr} = 0.334 \quad (R^2 = 0.99995). \quad (20)$$

The exponent $\beta_{scr} = 0.334$ is statistically indistinguishable from $1/3$. This density-dependent screening law is a projection of the source-charge screening operator $\mathcal{S}_\Sigma(\mathcal{E})$ (Paper 0 \S7) onto the UCD density regime; it is not an independent screening operator. Two sector assignments are stated explicitly. The geometric diagnostic $S = R_T/R_{phys}$ tracks the shear-sector $\mathcal{S}_\Sigma$ regime — density is the licensed stand-in for the ambient-steepness criterion inside dense macroscopic matter (Paper 0 \S8) — while the scale ρ_T that sets R_T is itself calibrated through the clock-amplitude projection $S_A = \min[1, (\bar{\rho}/\rho_T)^{1/3}]$ (Paper 0 \S3). Because each object's density is evaluated at its physical radius, $\bar{\rho} = 3M/(4\pi R_{phys}^3)$, the diagnostic carries an enclosed-acceleration equivalent: the criterion $S \sim 1$ corresponds to $GM/R_{phys}^2 \propto \bar{\rho} R_{phys}$, so the density and acceleration readings of the hierarchy agree object-by-object. Under the stated definitions, this is the expected scaling: if $R_T \propto M^{1/3}$ and $R_{phys} \propto (M/\rho)^{1/3}$, then:

$$S = \frac{R_T}{R_{phys}} = \frac{M^{1/3}}{M^{1/3}/\rho^{1/3}} = \rho^{1/3}. \quad (21)$$

Interpretation of the Screening Law

It is important to recognize that the regression $S \propto \rho^{0.334}$ is primarily a consistency check, not independent evidence for the scaling law. Since $S \equiv R_T/R_{phys}$ and the radii are defined by mass-density relations, the slope $\sim 1/3$ is algebraically expected given the model assumptions.

The value of this analysis is not to prove the scaling 'ab initio', but to demonstrate that a *single* saturation scale ρ_T yields a consistent screening hierarchy across 15 orders of magnitude in density ($R^2 \approx 1$) without requiring regime-dependent adjustments. The high R^2 confirms internal consistency; it does not constitute independent confirmation of TEP.

Furthermore, the extreme screening factors observed in binary pulsars ($S \sim 29,000$) are consistent with a strongly non-linear suppression mechanism in the scalar sector. The empirical screening hierarchy (S vs ρ) derives from the canonical Temporal

Topology mechanism (Paper 0), where screening operates via the continuous spatial profile of the scalar field governed by non-linear superposition of field gradients (Temporal Shear). Box 6.5 (Section 6) sketches a candidate route from the canonical action to soliton-like saturation, showing how the interplay between the kinetic term and a candidate saturation potential could generate the characteristic $M^{1/3}$ scaling and screening behavior. The detailed dynamical derivation is not required for the empirical hierarchy used here.

The empirical screening law is a direct consequence of the $M^{1/3}$ soliton scaling, providing cross-regime consistency under a single ρ_T rather than independent confirmation (see Note on Interpretation above).

4.3 Complete Screening Hierarchy




Figure 4: The Screening Hierarchy. Screening factor S plotted against density for 26 astrophysical objects. The empirical power law $S \propto \rho^{0.334}$ explains why dense objects (binary pulsars) are screened ($S \gg 1$) while diffuse systems (galaxies) are weakly screened ($S \ll 1$).

The complete hierarchy of screening factors across all object classes is visualized in Figure 4 and tabulated below.

Object Class	Density (g/cm ³)	Screening	Physical Meaning
Gas Giants	0.7 – 1.6	0.3 – 0.4×	R_T smaller than physical radius; saturated region does not envelop the object — the unscreened side of the transition, with channel-projected response (clock channels more active than orbital channels)
Main Sequence Stars	0.6 – 57	0.3 – 1.3×	Mixed regime; scalar and baryonic scales comparable
Rocky Planets (Earth)	3.3 – 5.5	0.56 – 0.66×	Saturation scale comparable to object radius; GNSS probes this boundary
Brown Dwarfs	~100	~1.7×	Just above ρ_T ; screening onset begins
White Dwarfs	$\sim 4 \times 10^5 - 2.5 \times 10^6$	27 – 50×	Baryonic star embedded within saturation scale; Keplerian mass = baryonic mass
Neutron Stars	10^{14}	26,800×	Scalar contribution less than 0.004%; pure GR dynamics
Binary Pulsars	10^{14}	29,500×	GR verified to 0.2%; Nobel Prize 1993

Table 4.1: Complete screening hierarchy across astrophysical object classes. The screening factor increases monotonically with density, consistent with a Temporal Topology screening hierarchy under the stated definitions. Binary pulsars provide the strongest validation, with GR verified to 0.2% precision at screening factors exceeding 29,500×

4.4 Precision GR Tests: Explained, Not Violated

A critical question for any modified gravity theory is: why do precision tests of General Relativity show no deviation? Temporal Topology provides the answer: the suppression of Temporal Shear (vanishing field gradient) reduces the screened PPN scalar charge $\alpha_{\text{PPN}}^{\text{eff}}$ where the ambient landscape is steep. The five most precise tests of GR all occur in regimes where screening is operative:

Test	Observable	Precision	Screening Factor	Status
Lunar Laser Ranging	Nordtvedt effect	10^{-13}	$0.56\times$	GR-consistent in the orbital ($\mathcal{S}_\Sigma$) projection; candidate residual reported by Paper 17 pending ephemeris refits
Cassini Conjunction	Shapiro delay	2×10^{-5}	$0.42\times$	Scalar sub-dominant; consistent with GR
MESSENGER	Perihelion precession	3×10^{-4}	$0.65\times$	GR-consistent in the orbital projection
Hulse-Taylor Pulsar	GW emission	0.2%	$29,500\times$	Completely screened
Double Pulsar	7 PPN tests	0.05%	$26,400\times$	Completely screened

Table 4.2: Precision GR tests and their screening factors. All tests are consistent with GR; TEP explains this via environment-dependent screening rather than requiring the scalar sector to be absent.

The Hulse-Taylor binary pulsar is particularly significant. Its orbital decay matches the GR prediction for gravitational wave emission to 0.2% precision—a result honored with the 1993 Nobel Prize (Taylor & Weisberg 1982). At a density of $\sim 10^{14}$ g/cm³, the screening factor reaches $29,500\times$. The scalar field contributes less than 0.003% to the orbital dynamics. This is not a violation of GR tests; it is a requirement of consistent screening.

4.5 Earth as the Calibration Anchor

The screening hierarchy reveals why Earth is the natural calibration point—and why GNSS clocks can detect what neutron stars cannot.

At $\rho \approx 5.5$ g/cm³, Earth sits just below the critical proximity scale, proxied by $\rho_T \approx 20$ g/cm³. The screening factor is $0.66\times$, meaning the saturation radius (4,200 km) is comparable to the physical radius (6,371 km). Earth occupies the narrow window where:

- **The scalar field is not screened:** Unlike neutron stars ($S = 26,800\times$), Earth's density is low enough that the soliton extends to observable scales.
- **The transition is close:** Unlike gas giants ($S \approx 0.3$), where the saturation core sits deep inside the object and well below the channel-projection boundary, Earth's $S \approx 0.66$ places it near the transition, where clock-comparison (S_A) channels retain a measurable response.

GNSS satellites orbit at $\sim 20,000$ km altitude, sampling clock transport through the exterior field region associated with the terrestrial Temporal Topology profile. The measured covariance length $L_c \approx 4200$ km is interpreted as a characteristic correlation/saturation scale of the Earth-sourced field, not as the orbital radius of the satellite network. The observed clock correlation length is therefore not an arbitrary parameter—it is the characteristic scale where the scalar field's gradient becomes steep enough to produce measurable timing correlations.

Within this phenomenology, the same saturation scale is expected to govern both terrestrial clock correlations and compact-object structure. The terrestrial scale provides an empirical calibration point, and the RBH-1 crossover provides an astrophysical consistency test under the same parameter choice.

4.6 The Critical Proximity Scale

The empirical data suggest a critical proximity scale, observationally proxied by the density $\rho_T \approx 20$ g/cm³, corresponding to approximately $4\times$ Earth's mean density. This value has coincidence-level significance: it lies near the onset of electron degeneracy, where the equation of state transitions from thermal to quantum pressure support. This coincidence motivates treating the 10–30 g/cm³ regime as a useful condensed-matter reference scale, while the mechanism setting ρ_T remains an open derivation target.

$\rho_T \approx 20$ g/cm³ is a saturation scale of the temporal-field topology, not a local on/off switch. Systems approach the GR-like limit when the observable shear/source-charge sector is suppressed, $\mathcal{S}_\Sigma \rightarrow 0$, which depends on source structure, environment, and boundary conditions, not $\rho > \rho_T$ alone. The empirical screening hierarchy $S \propto \rho^{0.334}$ (§4.2) is a smooth power-law across 15 orders of magnitude in density, not a step function at ρ_T .

Box 4.1: Summary — Three Cross-Regime Consistency Tests

The screening analysis provides three cross-regime consistency tests of the TEP framework under a single ρ_T :

1. **The RBH-1 Crossover:** The saturation radius is consistent with the Schwarzschild radius at $M \approx 10^7 M_\odot$ within combined uncertainties ($R_T/R_S \approx 1.3$). This provides a tightly constrained cross-regime consistency check under the fixed ρ_T calibration.
2. **The Screening Exponent:** The empirical scaling $S \propto \rho^{\beta_{\text{scr}}}$ with $\beta_{\text{scr}} = 0.334$ is fitted to the 11 dense objects ($\rho > \rho_T$) spanning ~ 12 orders of magnitude in density, with $R^2 = 0.99995$. The exponent $1/3$ is a direct consequence of the $M^{1/3}$ saturation scaling. The high R^2 partly reflects the definitional structure of S (the $\rho^{1/3}$ dependence is algebraically favoured once $R_T(M)$ enters the construction); the substantive content is the absence of large deviations, not the correlation coefficient — see Appendix B.4.
3. **GR Test Consistency:** All five precision tests of General Relativity are consistent with screening factors that suppress scalar contributions below current measurement limits.

The TEP framework does not violate established physics. It extends the metric structure to include a conformal sector that becomes observable only in specific density regimes—such as those probed by GNSS atomic clocks and cosmological soliton-scale systems.

5. Physical Constraints on the Proximity Scale

A natural question arises: why does the saturation scale take the observational proxy value $\rho_T \approx 20 \text{ g/cm}^3$? This section argues that the underlying proximity scale is not treated as an arbitrary fit parameter: it is empirically calibrated, while additional coincidence-level reference scales and open derivation targets are recorded for completeness. The GNSS coherence length L_c is a derived quantity; the fundamental parameter is the proximity scale itself, of which density is an observable proxy. In the canonical TEP framework, the full screening state of any system is determined by the environmental operator $S_\Sigma(E)$; ρ_T is the density projection of that operator, not the complete screening description.

5.1 ρ_T as the Conjectured Fundamental Parameter, L_c as Derived

The saturation scaling law $R_T \propto M^{1/3}$ —where a soliton interpretation is one candidate microscopic realization—implies a constant characteristic density:

$$\rho_T = \frac{M}{\frac{4}{3}\pi R_T^3} = \text{constant}. \quad (22)$$

For any object of mass M , the saturation radius is determined by:

$$R_T(M) = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3}. \quad (23)$$

The terrestrial coherence length $L_c \approx 4200 \text{ km}$ is therefore not an independently tuned galaxy-scale parameter; operationally, it is the observed GNSS covariance scale used to calibrate the projected Earth-scale realization of $R_T(M_\oplus)$. This relation is inverted in Section 2 to calibrate ρ_T from the measured L_c for Earth:

$$L_{c,\oplus} = \left(\frac{3M_\oplus}{4\pi\rho_T} \right)^{1/3} \approx 4200 \text{ km} \quad \text{for } \rho_T \approx 19\text{--}20 \text{ g/cm}^3. \quad (24)$$

The question thus reduces to: what physical considerations constrain ρ_T ?

5.2 The Dimensional Identity Bridge

A natural concern is whether the proximity of $\rho_T \approx 20 \text{ g/cm}^3$ to the density of terrestrial condensed matter implies a composition-dependent coupling that would violate the Weak Equivalence Principle. Under universal conformal coupling, the scalar sector is sourced by the bulk trace $T = -\rho$, independent of microscopic composition. The saturation scale ρ_T must therefore be a property of the vacuum potential $V(\phi)$, not an emergent property of the local matter.

The electron Fermi wavelength does not cause the scalar field to saturate. If it did, the field would require a direct, particle-specific coupling to the electron number density n_e , making the field exert a different force on iron than on hydrogen at the same bulk density. This would introduce a gross WEP violation, already constrained by experiments such as the MICROSCOPE satellite.

Instead, ρ_T is treated as a fundamental constant of the scalar sector — the geometric reference density of the saturation-radius law — rather than an emergent property of the local matter. The recorded coincidence is, in natural units,

$$\rho_T \approx \alpha^{4/3} m_e^4 \quad (\text{a density coincidence, not a potential-scale identification}). \quad (25)$$

Because this is a property of the scalar sector itself, the dimensional identity does not introduce composition-dependent coupling; WEP protection then follows from the universal matter coupling and the screened effective scalar charge. The field universally couples to bulk ρ and begins its continuous screening transition when local ρ approaches this ρ_T . The potential sector's own normalization is a separate quantity — the quartic coupling λ (equivalently $\Lambda_V = \lambda^{1/4} M_{\text{Pl}} \approx 1.3 \times 10^2$ GeV on the Cassini-compatible branch) — and is not to be identified with ρ_T (Appendix C.3).

The bulk density of Thomas-Fermi condensed matter is governed by Coulomb packing. The volume per atom scales with the Bohr radius $a_0 = (\alpha m_e)^{-1}$. In the Thomas-Fermi model, the effective atomic radius is $R_{\text{TF}} \approx a_0 Z^{-1/3}$, giving a volume $V \approx \frac{4\pi}{3} a_0^3 Z^{-1}$ per atom. With nucleon mass $M \approx A m_p$:

$$\rho_{\text{CM}} \approx \frac{3}{4\pi} A Z m_p m_e^3 \alpha^3. \quad (26)$$

For representative rock-forming elements ($A/Z \approx 2$), the product AZ ranges from $\sim 10^2$ (oxygen, $Z = 8$) to $\sim 10^3$ (iron, $Z = 26$). Evaluating with $m_p/m_e \approx 1836$ and $\alpha \approx 1/137.036$ gives $\rho_{\text{CM}} \sim 10^2\text{--}10^3$ g/cm³, one to two orders of magnitude above the calibrated $\rho_T \approx 20$ g/cm³. The Thomas-Fermi Coulomb-packing estimate therefore does not independently converge on the saturation scale; it situates ordinary condensed matter in a denser regime. The relevant condensed-matter reference that lies near ρ_T is the electron-degeneracy onset threshold (Section 5.3).

5.3 The Electron Degeneracy Threshold

Coincidence-level significance attaches to $\rho_T \approx 20$ g/cm³: this density lies in the condensed-matter regime where quantum pressure begins to become dynamically relevant. The crossover from thermal and electrostatic support to degeneracy-dominated support is gradual, with full degeneracy domination occurring only at much higher (white-dwarf) densities.

Density Regime	Dominant Physics	Examples
$\rho < 1$ g/cm ³	Gas pressure, thermal	Planets, main-sequence stars
$\rho \sim 1\text{--}10$ g/cm ³	Coulomb/thermal dominated solids and liquids	Rocky planets
$\rho \sim 10\text{--}10^2$ g/cm ³	Degeneracy onset (crossover)	Earth's core, brown dwarfs
$\rho \gtrsim 10^4$ g/cm ³	Electron degeneracy dominated	White dwarfs

Table 5.1: Equation of state transitions by density. The saturation density $\rho_T \approx 20$ g/cm³ lies near the onset of electron degeneracy.

This coincidence provides a useful condensed-matter reference scale for where quantum pressure becomes dynamically relevant to the equation of state. Under the universal conformal-coupling benchmark used throughout this corpus, the scalar sector is sourced primarily by the non-relativistic mass-density term, with pressure corrections entering only as a small relativistic correction in the condensed-matter regime. Electron degeneracy therefore does not, by itself, provide a mechanism that sets ρ_T , but it motivates the $\sim 10\text{--}30$ g/cm³ window as a natural coincidence-level cross-check.

5.4 Open Derivation Target: From Potential Scale to Transition Width

Box 6.5 adopts the corpus's quartic completion, in which the amplitude sector is normalized by the quartic coupling λ and ρ_T enters as the geometric constant of the saturation-radius law $R_T(M)$. Once ρ_T and the corpus branch of λ are specified, the field equation determines a smooth crossover between Compton-unresolved and Compton-resolved regimes ($\lambda_c(\bar{\rho})$ versus $R_T(M)$). Determining the detailed width and shape of this crossover as a function of ambient density—particularly in the terrestrial transition regime $\rho \sim 1\text{--}30$ g/cm³—requires an explicit numerical profile solution at the specified corpus normalization. This remains a target for future work.

Many-body condensed-matter physics (including the onset of electron degeneracy) governs the baryonic equation of state and provides useful reference densities, but under universal coupling it does not by itself select the saturation scale ρ_T . A complete theoretical account must instead explain both the value of ρ_T and the mapping between the geometric saturation radius $R_T(M)$ and the observable correlation length in clock networks.

While Section 5.3 contextualizes ρ_T within macroscopic condensed-matter physics, establishing its fundamental origin requires examining the scalar coupling scale at the vacuum level.

5.5 Dimensional Analysis: The Scalar Coupling Scale

Expressed in natural units, the saturation density carries the energy-density scale $\rho_T^{1/4} \approx 96.4$ keV for $\rho_T = 20$ g/cm³. This is a fourth-root reading of a density, not the scalar potential's normalization: the potential coefficient is the quartic coupling $\lambda = 7.5 \times 10^{-66}$, equivalently $\Lambda_V \equiv \lambda^{1/4} M_{\text{Pl}} \approx 1.3 \times 10^2$ GeV (Box 6.5, Appendix C; step 14). The kinetic completion carries its own distinct scale, $\Lambda_X = \sqrt{M_{\text{Pl}} H_0} \approx 1.9$ meV, which sets the Temporal-Shear threshold and plays no role in the amplitude-sector density argument.

$$\rho_T \approx \alpha^{4/3} \frac{m_e^4 c^3}{\hbar^3} \approx 22.4 \text{ g/cm}^3. \quad (27)$$

This correspondence is recorded as a numerical identity: the exponent $4/3$ (equivalently $\rho_T^{1/4} \sim \alpha^{1/3} m_e c^2$) is not derived within the present corpus and must be treated as an open target for theoretical work, not as an independent support.

5.6 Status: Phenomenological Constraint with Theoretical Foundation

The present analysis treats $\rho_T \approx 20$ g/cm³ as an empirically calibrated parameter, while recording additional coincidence-level reference scales and open derivation targets:

1. **Condensed-matter EOS coincidence:** ρ_T lies near the density scale where electron degeneracy becomes dynamically relevant to the equation of state
2. **Numerical identity:** $\rho_T \approx \alpha^{4/3} m_e^4 c^3 / \hbar^3$, with the $4/3$ exponent an explicit open target
3. **Open derivation:** a first-principles account must explain why the saturation scale ρ_T and the operational identification of L_c as the projected covariance scale associated with $R_T(M_\oplus)$ govern the observable clock-correlation envelope

Its use in quantum or topological extensions as a microscopic cutoff or defect-overlap scale is a conjectural extrapolation of the macroscopic saturation proxy, not an independent consequence of the present paper.

The key point for the present manuscript is that ρ_T is not treated as a free fit parameter: it is GNSS-anchored by clock correlations (Section 2) and then carried, without retuning, into the cross-scale tests that follow.

Box 5.1: Summary — The Saturation Scale: One Calibration, Two Coincidences, One Open Derivation

Operationally, ρ_T is calibrated from the measured GNSS coherence length L_c by treating L_c as the observed projected covariance scale associated with $R_T(M_\oplus)$, and then inverting $\rho_T = 3M_\oplus / (4\pi L_c^3)$. The framework's conjecture—left as an open derivation target—is that ρ_T is the fundamental saturation parameter and that L_c follows as a derived geometric scale once ρ_T is fixed. Two additional numerical correspondences are noted, but they remain coincidence-level until a first-principles derivation exists:

- Condensed-matter EOS coincidence (electron degeneracy onset)
- Numerical identity $\rho_T \approx \alpha^{4/3} m_e^4 c^3 / \hbar^3$ (exponent underived)

In the present phenomenological calibration, the terrestrial coherence length is treated as the observed projected covariance realization of the Earth-scale $R_T(M_\oplus)$ geometry; deriving this mapping from a specified scalar potential remains the open target. Both the condensed-matter coincidence and the numerical identity are contingent on the transfer-sketch $\mathcal{O}(1)$ prefactor being close to unity; a prefactor of ~ 2 would shift ρ_T to ~ 2.5 g/cm³, evaporating the coincidences. Operationally, GNSS observations provide the calibration by which ρ_T is inferred in the most accessible laboratory: Earth's gravitational field.

6. The Universal Density Constraint

A key test is whether the Temporal Topology saturation radius R_T yields quantitative expectations for otherwise free scales. Under a soliton interpretation, RBH-1 would correspond to a gravitational soliton; however, the saturation radius itself is the primary observable, constrained by the universal density ρ_T rather than introduced as an adjustable parameter.

The Universal Density Hypothesis

The Temporal Topology framework tests the empirical claim that a candidate saturation scale $\rho_T \approx 20 \text{ g/cm}^3$ governs compact-object structure across mass scales. A soliton interpretation provides one candidate microscopic realization of this saturation physics. This hypothesis is testable via two independent windows:

1. **Planetary Scale:** The Earth-scale Temporal Topology parameter $R_T(M_\oplus)$ is calibrated through the observed GNSS projected covariance length ($L_c \approx 4200 \text{ km}$), serving as the empirical anchor.
2. **Cosmological Scale:** The extrapolated Temporal Topology saturation radius for RBH-1 ($M \approx 10^7 M_\odot$) can be compared to the Schwarzschild diameter, placing the object near the crossover mass where horizon and soliton interpretations are maximally degenerate.

The central question is whether a single density parameter, validated cosmologically, remains universal at galactic scales.

The Saturation Scale as Fundamental Parameter

If the scalar sector saturates at a critical energy density ρ_T , dimensional analysis requires a universal mass–radius relation:

$$R_T(M) = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3}. \quad (28)$$

Box 6.1: The Theoretical Priority of ρ_T

A potential misreading of this framework is that "GPS clock noise determines black hole physics." This inverts the logical and historical order of the analysis.

- **Calibration (Terrestrial):** The GNSS analysis (Smawfield 2025b) serves as an empirical anchor. The detection of a correlation length $L_c \approx 4200 \text{ km}$ in atomic clock data provides an empirical estimate of the saturation scale, conditional on treating L_c as the projected covariance scale associated with $R_T(M_\oplus)$.
- **Application (Cosmological):** The resulting $M^{1/3}$ scaling is therefore not an extrapolation of clock noise, but a test of whether this terrestrially-validated constant remains universal at galactic scales.

The alignment of the RBH-1 crossover scale is thus an independent consistency check, not a fitted result.

Box 6.2: Null Hypothesis and Look-Elsewhere Effect

A potential concern is that apparent cross-scale correspondences could arise from post-hoc alignment rather than predictive structure. To address this possibility, the prior probability of simultaneous agreement across multiple reference scales is estimated under an explicit null hypothesis.

Null hypothesis: assume ρ_T is drawn uniformly from the range 10^{-3} – 10^6 g/cm^3 (spanning interstellar gas to nuclear density). The GNSS constraint ($R_T(M_\oplus)$ within 50% of 4200 km) permits ρ_T in the range ~ 6 – 160 g/cm^3 (≈ 1.4 dex). The RBH-1 constraint ($R_T(10^7 M_\odot)$ within 50% of $2R_S$) permits ρ_T in the range ~ 7 – 180 g/cm^3 (≈ 1.4 dex). The overlap is ~ 7 – 160 g/cm^3 (≈ 1.4 dex). Under a uniform logarithmic prior spanning 9 dex, the probability of a randomly drawn ρ_T satisfying both constraints simultaneously is $P \approx 1.4/9 \approx 0.15$ ($\sim 15\%$). While not overwhelmingly small, this probability is reduced further when the condensed-matter coincidence (Section 5.3) is included as a third independent cross-check. Given the GNSS calibration and the condensed-matter coincidence noted in Section 5, the RBH-1 correspondence provides an independent consistency check of the same underlying parameter. The convergence of two physically distinct systems—geodesy and cosmology—on the same density scale across 15 orders of magnitude is not readily explained by independent tuning, but the universality hypothesis remains falsifiable: any future measurement that robustly requires a substantially different ρ_T would exclude this framework.

Independence caveat: the GNSS calibration is currently validated within the author's research program (Smawfield 2025b,c,d). Independent replication by other groups is required before the terrestrial constraint can be treated as established. The cosmological inputs referenced here are derived from external published data (van Dokkum et al. 2025).

The GNSS-derived value $\rho_T \approx 19$ – 20 g/cm^3 (Section 2) lies in the same condensed-matter density regime where electron degeneracy becomes dynamically relevant to the equation of state (Section 5). The central question is whether this same density scale holds for RBH-1.

Box 6.3: Robustness of the GNSS Calibration (Systematics Check)

The terrestrial calibration length $L_c \approx 4200 \text{ km}$ — the GNSS instrument's measured realization of the canonical correlation scale λ_T (Paper 0 §7) — is an extraordinary claim requiring careful exclusion of geodetic systematics. The following tests support its physical origin (full methodology in Smawfield 2025b,c,d):

- **Multi-Center Verification:** The correlation structure persists across independent clock solutions from CODE, ESA, and IGS (1999–2024), disfavoring software-specific processing artifacts (e.g., Bernese vs. GIPSY). The recovered L_c spans $\sim 3,330$ – $4,549$ km across centers (a $\sim 23\%$ spread).
- **Null Tests:** (i) Randomizing satellite epochs destroys the correlation ($r^2 < 0.01$, $N = 1000$ shuffles); (ii) shuffling clock residuals within each satellite eliminates the spatial structure; (iii) replacing real data with white noise yields no coherent scale. All three nulls are satisfied at $> 5\sigma$.
- **Scale Separation:** The 4200 km scale is distinct from tropospheric correlation lengths (~ 100 – 500 km) and orbital period harmonics (half-sidereal, ~ 12 hr). Power spectral analysis shows the 4200 km feature as a persistent background covariance floor in the GPS-only clean baseline, with no counterpart in ionospheric or tropospheric delay products; multi-constellation validation shows consistent signatures.
- **Dataset:** IGS final clock products (CLK files), 30-second sampling, 1999–2024. Preprocessing: removal of satellite/receiver clock offsets, relativistic corrections (Sagnac, gravitational redshift), and reference frame alignment to ITRF2020.

Limitation: The present analysis treats L_c as an empirical calibration parameter. A complete derivation from first principles—linking the empirical L_c to the geometric saturation radius $R_T(M)$ and to a candidate completion's field-profile parameters—remains a target for future theoretical work.

Box 6.4: Derivation of the Scaling Law

The $R \propto M^{1/3}$ scaling is not an ad hoc ansatz but a direct consequence of saturation in the scalar sector. Consider a scalar field ϕ coupled to matter density ρ with a potential $V(\phi)$ that enforces a maximum gradient or energy density.

In the dense limit (compact objects), the scalar field profile saturates to a core of constant effective energy density ρ_T . For a self-gravitating configuration of total mass M , the volume of this saturated core is constrained by mass conservation:

$$M \approx \frac{4}{3}\pi R_T^3 \rho_T. \quad (29)$$

Solving for the radius yields the characteristic scaling:

$$R_T \propto \left(\frac{M}{\rho_T} \right)^{1/3}. \quad (30)$$

This relation describes the boundary of the saturated "soliton" region. Outside this radius, the field decays, recovering Newtonian gravity (Temporal Topology screening). Inside, the field is phase-locked, modifying the effective metric (proper time). The "Universal Scaling" is thus simply the statement that the scalar potential $V(\phi)$ has a maximum capacity to support scalar gradients before saturating at ρ_T .

Box 6.5: Soliton Formation from the Canonical Action

To move beyond phenomenology, a candidate route from the canonical Temporal Topology action (Paper 0) to soliton-like saturation is sketched. The action above contains a canonical kinetic term $-\frac{1}{2}(\partial\phi)^2$ and a candidate saturation potential $V(\phi)$ that would prevent gradient divergence. The interplay between these terms can generate soliton-like saturated configurations with characteristic radius $R_T \propto M^{1/3}$.

1. Field Equation and Equilibrium

For a static, spherically symmetric configuration, the scalar field equation is:

$$\nabla^2 \phi = V'(\phi) + \alpha(\phi)\rho \quad (31)$$

where $\alpha(\phi) \equiv d \ln A / d\phi$ and ρ is the ambient matter density. In the dense limit, the effective potential $V_{\text{eff}}(\phi; \rho) = V(\phi) + [A(\phi) - 1]\rho$ develops a minimum at $\phi_{\text{min}}(\rho)$ with effective mass $m_{\text{eff}}^2(\rho) = V_{\text{eff}}''(\phi_{\text{min}})$.

2. Saturation Mechanism

Under the corpus coupling $\beta_A = -1$ the matter term enters with negative slope ($\alpha\rho = -\rho/M_{\text{Pl}}$), so a minimum of V_{eff} exists only on the confining branch $V' > 0$. Writing $u = \phi/M_{\text{Pl}}$, the corpus's own weak-field branch is the quartic $V(u) = (\lambda/4) M_{\text{Pl}}^4 u^4$ — the member of the confining family $V \propto u^m$ selected by the corpus amplitude law $S_A \propto \rho^{1/3} = \rho^{1/(m-1)}$, i.e. $m = 4$ (Appendix C.2). The equilibrium condition $V'_{\text{eff}} = 0$ reads $\lambda M_{\text{Pl}}^4 u_{\text{min}}^3 = \rho e^{-u_{\text{min}}}$, giving $u_{\text{min}} \approx (\rho/\lambda M_{\text{Pl}}^4)^{1/3}$ for $u_{\text{min}} \ll 1$: the field is

elevated inside matter and approaches zero in the ambient, in the direction required by the corpus sign convention. (The inverse-power branch $V \propto \phi^{-n}$ is inadmissible under $\beta_A = -1$: both slopes are then negative and no minimum exists; its implied $\phi_{\min} \propto \rho^{-1/(n+1)}$ would in any case run opposite to the required direction.) The scalar energy density in a saturated core is:

$$\rho_{\text{eff}} \sim V(u_{\min}) + \frac{1}{2} M_{\text{Pl}}^2 (\nabla u)^2 \simeq \frac{1}{4} \rho u_{\min} \ll \rho. \quad (32)$$

The saturated configuration is therefore a field-geometry statement, not a competing mass density: at the corpus normalization $\lambda = 7.5 \times 10^{-66}$ (the scalar self-coupling, Appendix C) the core energy density is $\sim 3 \times 10^{-8} \text{ g/cm}^3$ at terrestrial densities, and the equilibrium $u_{\min}(\rho_T) \approx 6.9 \times 10^{-9}$ remains perturbative. What saturates at ρ_T is the clock-amplitude response $S_A = \min[1, u_{\min}(\bar{\rho})/u_{\min}(\rho_T)]$ through the geometric radius R_T ; the order-unity self-quenching crossing $u_{\min} \sim 1$ (where e^{-u} suppresses the conformal source) lies at $\rho_{\text{sat}} \sim \lambda M_{\text{Pl}}^4 \approx 6 \times 10^{25} \text{ g/cm}^3$, inside the temporal-well domain of Paper 28 (step 14).

3. Temporal Topology Screening

The effective mass $m_{\text{eff}}^2(\rho) = V''_{\text{eff}}(u_{\min})/M_{\text{Pl}}^2 = [3\lambda M_{\text{Pl}}^4 u_{\min}^2 + \rho e^{-u_{\min}}]/M_{\text{Pl}}^2$ grows with density as $m_{\text{eff}} \propto \lambda^{1/6} \rho^{1/3}$, so the Compton wavelength $\lambda_C = 1/m_{\text{eff}}$ contracts through denser interiors: a body is Compton-resolved once $\lambda_c(\bar{\rho}) \lesssim R_T(M)$, the crossover that for Earth sits at $\sim 4 \text{ g/cm}^3$ (step 14). Inside such a body the field rests near its local equilibrium $u_{\min}(\rho)$ and the spatial variation is confined to a boundary layer of width $\sim \lambda_C$, where the kinetic response $P_{,X}$ controls the gradient amplitude. This creates the characteristic flattening of the scalar field spatial profile—Temporal Topology screening—that reconciles precision local tests with cosmological dynamics. In the saturation regime ($\rho \sim \rho_T$), the Temporal Topology saturation radius scales as $R_T = (3M/4\pi\rho_T)^{1/3}$, yielding the observed $M^{1/3}$ dependence. At still higher densities — where the equilibrium approaches order unity ($\rho \sim \rho_{\text{sat}} \approx 6 \times 10^{25} \text{ g/cm}^3$) — the effective source slope $\rho A_{,u} = -\rho e^{-u_{\min}}$ is exponentially suppressed, so the scalar charge of deep-interior matter quenches as well: that is the temporal-well regime, not the terrestrial one.

4. Coupling and Stability

Matter couples to the Jordan frame metric $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi$ with $A(\phi) = \exp(\beta_A\phi/M_{\text{Pl}})$. Stability is ensured because the canonical kinetic term has the correct sign for a physical scalar field, and the theory satisfies the Null Energy Condition for physically realizing solutions.

Testing the Density Constraint at Cosmological Scales

The saturation scale ρ_T determined from terrestrial clocks is now tested against RBH-1. Using the revised best-fit mass estimate of $M \approx 2 \times 10^7 M_\odot$ (van Dokkum et al. 2025), the predicted Temporal Topology saturation radius is:

$$R_T = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3} \approx 7.8 \times 10^7 \text{ km}. \quad (33)$$

The predicted Temporal Topology saturation radius can be compared directly to the Schwarzschild radius for this mass:

Schwarzschild radius: $R_S = 2GM/c^2 \approx 5.9 \times 10^7 \text{ km}$

Predicted Temporal Topology saturation radius: $R_T = (3M/4\pi\rho_T)^{1/3} \approx 7.8 \times 10^7 \text{ km}$

Ratio: $R_T/R_S \approx 1.3$

This order-unity correspondence ($R_T \approx 1.3R_S$) arises because ρ_T lies near the horizon-formation density threshold in the underlying scalar sector. Within the combined uncertainties from M and ρ_T , which propagate to $\sim 25\%$ uncertainty in R_T , the two scales remain consistent at the factor-of-few level. This places RBH-1 near a crossover regime where horizon and saturated-soliton interpretations can be observationally degenerate.

This geometric match is the primary prediction of the scaling law. While the velocity discontinuity ($\Delta v \sim 650 \text{ km/s}$) provides a secondary constraint on the *amplitude* of the screening (see Smawfield 2025h, Paper 7), the *scale* of the object is set fundamentally by the saturation scale ρ_T . The correspondence suggests that what is conventionally identified as the event horizon scale in GR may correspond to the saturation boundary of the scalar field in the TEP framework.

In the TEP interpretation, what is conventionally called a "black hole" at this mass scale is modeled as a saturated soliton core, i.e., a region where the conformal time-field approaches a maximum gradient set by ρ_T . In this phenomenology, the characteristic radius is set by the saturation scale rather than by horizon formation. The observed central dimming is attributed to extreme time dilation (strong redshifting) rather than causal disconnection.

A further structural implication follows from the different mass scalings. The soliton relation predicts $R_T \propto M^{1/3}$, whereas the Schwarzschild radius scales as $R_S \propto M$. These opposing scalings intersect at a unique crossover mass $M_\times$ where the two radii coincide (see Figure D.1). Empirically, the terrestrial calibration places this crossover near $M_\times \approx 10^7 M_\odot$ —precisely where RBH-1 resides. Objects near $M_\times$ are expected to be maximally degenerate between horizon and soliton interpretations, making RBH-1 an unusually diagnostic system.

The identification is thereby restricted to the crossover neighbourhood: it is a statement about $M \approx M_\times$, not a general reinterpretation of compact objects. The directional implication is explicit. For $M \ll M_\times$ the weak-field construction gives $R_T \gg R_S$ — a $10 M_\odot$ object would carry R_T of order 10^6 km against $R_S \approx 30$ km — so soliton-realized compact objects at stellar masses would appear as extended structures and are incompatible with observed merger phenomenology. For $M \gg M_\times$ the saturation radius retreats far inside the horizon (Appendix E computes $S \sim 0.03$ for M87*), and the identification ceases to be degenerate. Two resolutions are admitted, and the corpus requires one of them: either soliton-realized compact objects exist only in the crossover band, or the weak-field $R_T(M)$ construction fails at horizon compactness and must be replaced by the relativistic strong-field treatment of Paper 28, within which the $M^{1/3}$ ansatz is not guaranteed to survive. The crossover statement is therefore conditional on weak-field validity at horizon-compactness — an assumption stated rather than assumed — while the degeneracy at $M_\times$ itself remains a legitimate and diagnostic coincidence that feeds the RBH-1 analysis of Paper 7.

Status of the Scaling Law

The relation $R \propto M^{1/3}$ is a density-limited scaling expected for compact, self-bound field configurations whose cores approach a finite saturation scale. Such behavior is familiar in non-topological solitons and bosonic compact objects (Coleman 1985; Seidel & Suen 1991; Gleiser 1994; Hui et al. 2017).

The central claim is not that terrestrial clocks "predict" black hole sizes, but that a fundamental density constant ρ_T appears across scales. The saturation scale is determined empirically from Earth—the most accessible high-precision gravitational laboratory—and then tested for consistency at cosmological scales. The present analysis should be read as a phenomenological identification of an unexpected regularity: the same density constraint that governs clock correlations in Earth's gravitational field also governs the characteristic size of a $10^7 M_\odot$ compact object.

Dependency Structure and Constraint Isolation

A critical requirement for the robustness of this framework is to demonstrate that the apparent convergence of ρ_T is not a result of circular reasoning. Specifically, it must be determined whether the "independent" tests are truly distinct or if they all implicitly rely on the same prior assumptions.

To resolve this, a constraint-isolation analysis is performed, isolating each constraint to determine what value of ρ_T it yields without input from the others.

Test 1: Condensed-Matter Density Coincidence Isolation (Dropping GNSS)

If all GNSS data are discarded, condensed-matter equation-of-state transition scales (Section 5.3) provide a coincidence-level density estimate.

Result: Condensed-matter transition considerations yield densities in the $\sim 10\text{--}30$ g/cm³ range, broadly consistent with the GNSS-derived value.

Test 2: Terrestrial Isolation (External Calibration Check)

If the condensed-matter coincidence is discarded and reliance is placed *solely* on the observed GNSS correlation length $L_c \approx 4200$ km for Earth's mass ($M_\oplus$), what density is derived?

$$\rho_T(\text{GNSS}) = \frac{3M_\oplus}{4\pi L_c^3} \approx 20 \pm 7 \text{ g/cm}^3 \text{ (inter-center), } \in [8, 70] \text{ g/cm}^3 \text{ (epoch/window)}. \quad (34)$$

Result: Terrestrial data independently recovers the same density scale, consistent with the condensed-matter coincidence estimate but derived from a system 51 orders of magnitude more massive. The quoted band carries the epoch/window systematic established by the executed held-out discriminator (Appendix B.8); the identification of L_c with $R_T(M_\oplus)$ remains conditional on that systematic not being resolved in favour of the lower MGEX scale.

Test 3: Cosmological Isolation (Dropping Local Constraints)

If both condensed-matter considerations and terrestrial inputs are discarded and the density required to place the RBH-1 Temporal Topology saturation radius at the horizon scale (the crossover condition) is computed, the result is:

$$\rho_T(\text{RBH-1}) = \frac{3M_{\text{RBH-1}}}{4\pi(1.3R_S)^3} \approx 15\text{--}30 \text{ g/cm}^3. \quad (35)$$

Result: The cosmological constraint points to the same narrow density window.

Conclusion on Circularity

The convergence is not a trivial post-hoc fit. Multiple physically distinct systems—the Earth (geodesy) and galactic black holes (cosmology)—are mutually consistent with the same saturation scale $\rho_T \approx 20 \text{ g/cm}^3$. The framework remains falsifiable: any future measurement that robustly requires a substantially different ρ_T would exclude this universal-density soliton model as formulated here.

7. Milky Way Test: Dark Matter Onset

Expected Onset Radius

The Milky Way provides a local test of the galactic $M^{1/3}$ scaling law established in Section 3. Unlike external galaxies where distance uncertainties dominate, the Milky Way offers high-precision kinematic data. This section tests whether the SPARC-calibrated relation predicts the radius at which the mass discrepancy first becomes significant in the Milky Way.

Using the SPARC-calibrated normalization ($k \approx 7.9 \times 10^{-4} \text{ kpc}/M_\odot^{1/3}$) and the Milky Way's total baryonic mass estimated at $M_{\text{bar}} \approx 6 \times 10^{10} M_\odot$ (Bland-Hawthorn & Gerhard 2016), the expected dark-matter onset radius is:

$$R_{\text{DM}} = k \cdot M_{\text{bar}}^{1/3} \approx 7.9 \times 10^{-4} \cdot (6 \times 10^{10})^{1/3} \text{ kpc} \approx 3.1 \text{ kpc}. \quad (36)$$

This radius marks the scale where the baryonic rotation curve (bulge + disk) first falls below the observed circular velocity, signaling the onset of the phantom mass effect. It is the same quantity R_{DM} that is measured directly in the SPARC ensemble; the outer Keplerian decline at larger radii is a separate phenomenon governed by the full halo profile, not by the onset scaling alone.

Gaia DR3 and Local Observations

Gaia Data Release 3 (Gaia Collaboration 2023) provides precise proper motions and radial velocities for millions of stars, enabling construction of the Milky Way rotation curve to greater distances than previously possible. Recent analyses (e.g., Jiao et al. 2023) also report evidence for a Keplerian-like decline beginning near $R \approx 19 \text{ kpc}$; this outer feature traces the halo edge and is distinct from the inner onset scale tested here.

Key observational features relevant to the R_{DM} prediction:

- **Baryonic dominance ($R \lesssim 3 \text{ kpc}$):** The bulge and inner disk dominate; the rotation curve rises steeply and is well described by visible matter alone (Sofue 2020).
- **Dark-matter onset ($R \sim 3\text{--}5 \text{ kpc}$):** Beyond the peak of the baryonic contribution, the observed circular velocity ($v_{\text{circ}} \approx 220 \text{ km/s}$) stays flat while the baryonic model predicts a decline. This is the Milky Way's equivalent of the SPARC R_{DM} — the radius where the mass discrepancy first becomes significant.
- **Flat regime ($5 \lesssim R \lesssim 15 \text{ kpc}$):** The rotation curve remains approximately flat, indicating substantial phantom mass contribution.
- **Keplerian decline ($R \gtrsim 19 \text{ kpc}$):** At large radii the enclosed mass growth slows and v_{circ} begins to decline. This outer feature is not predicted by the onset scaling $R_{\text{DM}} \propto M^{1/3}$.

Uncertainty Analysis and Error Budget

Parameter	Value	Uncertainty	Impact on R_{DM}
MW Baryonic Mass	$6.0 \times 10^{10} M_\odot$	$\pm 15\%$ (0.9×10^{10})	$\pm 5\%$ (scaling $M^{1/3}$)
SPARC Constant (k)	7.9×10^{-4}	$\pm 4\%$ (fit error)	$\pm 4\%$
Model Systematic	-	$\pm 10\%$ (geometry)	$\pm 10\%$
Total Expected R_{DM}	3.1 kpc	$\pm 0.4 \text{ kpc}$ ($\sim 14\%$)	-

Parameter	Value	Uncertainty	Impact on R_{DM}
Observed Onset	$\sim 3\text{--}5$ kpc	± 1 kpc	-

The expected value (3.1 ± 0.4 kpc) and the observed dark-matter onset scale ($\sim 3\text{--}5$ kpc) are consistent within the combined uncertainty. Because the SPARC-calibrated normalization k is the sole empirical input to this prediction, the Milky Way test is a SPARC-internal consistency check rather than a GNSS cross-scale test, consistent with the Numerology Firewall classification (Section 8).

Comparison with Dark Matter Halo Models

NFW Halo vs. TEP Soliton

- **NFW Halo (CDM):** In standard CDM, the inner rotation curve profile depends on the halo concentration c and virial mass M_{vir} , both of which are free parameters. The radius where baryons cease to dominate is not predicted a priori; it emerges from the halo assembly history and can vary significantly between cosmological simulations.
- **TEP Soliton:** The $M^{1/3}$ scaling yields the onset radius directly from the baryonic mass, with no Milky Way-specific tuning. The normalization is fixed by the SPARC ensemble ($k \approx 7.9 \times 10^{-4} \text{ kpc}/M_{\odot}^{1/3}$) and gives $R_{\text{DM}} \approx 3$ kpc for the Milky Way. The agreement with the observed transition at $\sim 3\text{--}5$ kpc supports the universality of the scaling law.

Future Refinements

Ongoing Gaia data releases will improve the precision of the inner rotation curve, potentially constraining the exact baryonic-to-dark-matter transition radius to $\sim 10\%$ accuracy. Combined with improved Milky Way mass estimates from satellite kinematics and gravitational lensing, this could provide a sub-10% test of the $M^{1/3}$ onset scaling in the Milky Way.

8. Discussion: The Nature of the Dark Sector

Synthesis of Multi-Scale Evidence

The central result of this work is the identification of a single density scale, $\rho_T \approx 20 \text{ g/cm}^3$, that organizes gravitational anomalies across 18 orders of magnitude in mass (Earth to galaxy). This scale, originally calibrated from terrestrial GNSS atomic clock correlations ($L_c \approx 4200$ km), is consistent with:

- The dark matter onset radius in spiral galaxies ($R_{\text{DM}} \propto M^{1/3}$).
- The dark-matter onset radius of the Milky Way ($R_{\text{DM}} \approx 3$ kpc, observed $\sim 3\text{--}5$ kpc).
- The screening hierarchy in binary pulsars vs. galaxies ($S \propto \rho^{1/3}$).

The convergence of these mutually reinforcing constraints suggests that ρ_T is not merely a fitting parameter for a specific system, but a candidate universal parameter of the effective description of the dark sector, within the stated uncertainties.

Claim Hierarchy and Falsification Scope

The empirical content of this work can be read at three distinct levels. This separation is critical for interpreting the consequences of future measurements.

- **Level 1 (Empirical regularities):** The existence of approximate $M^{1/3}$ scaling features in rotation-curve onset radii, and the density-ordered screening hierarchy summarized by $S = R_T/R_{\text{phys}}$.
- **Level 2 (Universal-density soliton model):** The hypothesis that a single saturation scale ρ_T organizes these regularities via $R_T(M) = (3M/4\pi\rho_T)^{1/3}$, with GNSS providing an empirical calibration (subject to independent replication) and cross-regime consistency tests (SPARC ensemble, Milky Way, RBH-1 crossover).
- **Level 3 (TEP microphysics):** The full dynamical realization in which temporal shear and a conformal time field $A(\phi) = \exp(\beta_A \phi/M_{\text{Pl}})$ generate the effective soliton/screening phenomenology. Failures at Level 2 primarily constrain the soliton implementation and/or calibration mapping, without necessarily excluding the broader TEP framework.

Numerology Firewall: Which Agreements Were Predicted?

The elegance of cross-scale agreement is vulnerable to the accusation of "curve-fitting across scales." This section provides a candid assessment of which constraints were predicted *a priori* and which were identified *post hoc*.

Comparison	Before/after GNSS calibration?	Free parameters?	Independent?	Weight
Earth GNSS L_c	Calibration anchor	yes/no (fit to correlation)	no — defines scale	Calibration

Comparison	Before/after GNSS calibration?	Free parameters?	Independent?	Weight
SPARC $M^{1/3}$ scaling	After	Yes (normalization k)	Medium	Moderate
DF2/DF4 UDGs	After	Yes (extrapolation)	Weak	Illustrative
Milky Way inner mass-discrepancy onset	After	No new normalization; uses SPARC-calibrated k	Medium	Local consistency check
Wide binaries (if tested)	Before/after	Yes	Stronger (if replicated)	High if replicated

Honesty Principle: The Value of ρ_T

The value of ρ_T should be treated as a candidate organizing scale until at least one held-out mass regime is predicted *before* analysis and recovered without refitting. Current cross-scale consistency is motivating but not discriminating; the $\rho^{1/3}$ hierarchy is a consistency relation induced by the $R_T(M)$ construction, not an independent discriminator of microscopic mechanism.

The per-constellation MGEX analysis (Paper 14 extension) provides a concrete pre-registered discriminator: if GPS, Galileo, and GLONASS individually recover $\sim 4,200$ km while the combined product compresses to the measured $\sim 1,900$ km scale (pooled MGEX: 1,862 km), the MGEX discrepancy is resolved as a projection artifact. If per-constellation analysis also yields $\sim 1,900$ km, the $L_c \leftrightarrow R_T(M_\oplus)$ identification fails as calibrated, and the Level 1/2/3 claim hierarchy absorbs the hit exactly as designed. Writing this test down before running it converts the corpus's largest latent vulnerability into its sharpest falsifiable statement.

Dark Matter as Phantom Mass

In the TEP framework, "dark matter" is reinterpreted not as a particle species, but as "phantom mass"—an apparent excess inferred when a geometry with temporal shear (two metrics) is analyzed under the assumption of isochrony (single metric). The saturation scale ρ_T represents the scale at which the conformal factor $A(\phi) = \exp(\beta_A \phi / M_{Pl})$ enters a non-linear, self-supporting phase (the soliton).

This explains the phenomenology of "cores" in dark matter halos. In standard CDM simulations, halos tend to form cusps ($\rho \propto r^{-1}$), which are often in tension with observations of flat cores in dwarf galaxies (the core-cusp problem). In the soliton picture, the core is a distinct physical state—a Bose-Einstein condensate-like configuration of the scalar field—that naturally supports a flat density profile with a characteristic radius R_T .

Comparison with Alternative Frameworks

The TEP framework shares phenomenological features with both Modified Newtonian Dynamics (MOND) and Fuzzy Dark Matter (FDM) but is distinguished by its screening mechanism and cross-scale predictive power.

Feature	MOND	Fuzzy DM (FDM)	TEP (Soliton)
Physical Basis	Modified Inertia/Gravity	Ultralight Boson ($m \sim 10^{-22}$ eV)	Conformal Time Field Saturation
Fundamental Scale	Acceleration $a_0 \approx 1.2 \times 10^{-10}$ m/s ²	Particle Mass m_ψ	Proximity Scale ξ_T (obs. proxy $\rho_T \approx 20$ g/cm ³)
Galactic Scaling	Fits RAR (a_0 is free parameter)	$M_c \propto M_h^{1/3}$ (Core-Halo)	$R \propto M^{1/3}$ (derived from ρ_T)
Screening	External Field Effect (EFE)	None (requires tuning)	Temporal Topology ($S \propto \rho^{1/3}$)
Solar System	Recovered via interpolation function	Unsuppressed (tension)	Screened ($S \sim 0.6$ for Earth, $> 30k$ for Pulsars)
Key Prediction	Exact RAR	Solitonic Cores	GNSS Correlation Length + Screening Hierarchy

TEP vs. MOND

MOND provides an excellent fit to galaxy rotation curves but offers no explanation for the GNSS clock correlations. The critical distinction lies in the triggering mechanism: MOND modifies kinematics at a specific *acceleration* scale ($a_0 \approx 1.2 \times 10^{-10}$ m/s²), which is a derived, environment-independent parameter that does not predict where screening will occur in dense systems. By contrast, TEP is triggered by a *density/proximity* scale ($\rho_T \approx 20$ g/cm³), enabling it to explain anomalous dynamics in environments where MOND either makes no prediction or fails—such as ultra-diffuse galaxies (where the low surface brightness yields a different

acceleration at fixed density) and the Milky Way core (where the high baryonic density should suppress MOND-like effects but does not in the data).

TEP recovers MOND-like phenomenology ($a_0 \sim G\Sigma_c$) as an emergent property of the screening transition, while correctly predicting the environment-dependent suppression required for precision GR tests.

TEP vs. FDM

Standard FDM predicts solitonic cores but fundamentally struggles to recover the correct structural scales for both dwarf galaxies and the Milky Way simultaneously, requiring contradictory particle mass bounds. TEP naturally and uniquely links the core scale directly to the ambient saturation scale, generating the correct cross-scale hierarchy anchored directly by GNSS empirical measurements.

Response-Coefficient Hierarchy

A natural criticism of the multi-channel TEP programme is that different papers report different numerical coefficients. This is expected, and it is a feature rather than a flaw. In the TEP framework, each measurement channel carries an *observable response coefficient* κ_X that absorbs stellar physics, environment, and projection geometry, not the bare microscopic conformal coupling β_A .

Paper 11 (TEP-H0) independently calibrates the weakly screened galactic-disk response via Cepheid period–luminosity residuals. The Paper 0 benchmark normalization is $\kappa_{\text{Cep}}^{\text{bench}} \sim 10^6$, while the empirical channel returns $0.33\text{--}0.45 \times 10^6$ (dimensionless, in the appropriate observable basis) — a factor 2–3 benchmark/empirical gap carried as an unresolved normalization target (Paper 0, issue-class 0-17), attributable to host-screening normalization or additional period-structure in the transfer function. Paper 10 (TEP-COS) measures the globular-cluster pulsar response $\tilde{\kappa}_{\text{MSP}} \approx 3 \times 10^4$ (full response coefficient, `step_44_kappa_msp_prior.json`) after dense-cluster geometric suppression; the mapping to Paper 10's effective channel coefficient is $\kappa_{\text{MSP}} = \tilde{\kappa}_{\text{MSP}} \mathcal{S}_{\text{cluster}} \approx 0.05$, i.e. a cluster suppression factor $\mathcal{S}_{\text{cluster}} \approx 1.7 \times 10^{-6}$ applied to the full response — the same coefficient quoted under the un-tilde symbol. The two values are consistent with the bare TEP geometric-factor estimate ($\sim 10^6\text{--}10^7$) only after environmental transfer factors are included; they do not assert a direct one-to-one equality of raw channel coefficients.

This hierarchy protects the programme from the criticism: "*Why does each paper have a different coefficient?*" The answer is that they are *observable response coefficients*, not bare scalar couplings. Each channel measures a different projection of $S_\Sigma(\mathcal{E})$:

Channel	Response coefficient	Environment	Projection
GNSS clocks	$L_c \approx 4200$ km	Planetary crust/mantle	Radial covariance scale
Cepheids (Paper 11)	$\kappa_{\text{Cep}} = 0.33\text{--}0.45 \times 10^6$ empirical; benchmark $\sim 10^6$	Galactic disk (weakly screened)	Period–luminosity modulation
Pulsars (Paper 10)	$\tilde{\kappa}_{\text{MSP}} \sim 10^4$ full; $\kappa_{\text{MSP}} \approx 0.05$ effective	Globular cluster (strongly screened)	Spin-down excess (suppressed)
Wide binaries (Paper 13)	$\alpha_{\text{sat}} \approx 0.366$	Galactic halo (unscreened)	Velocity-profile saturation

The cross-paper comparison is therefore through a shared clock-response structure, not raw equality of coefficients. This preserves the internal consistency of the programme while acknowledging that each domain probes a different facet of the same environmental operator.

Resolving Galactic Anomalies

The Radial Acceleration Relation (RAR)

The SPARC analysis (Section 3) confirms that the onset of mass discrepancies follows the baryonic mass distribution, a key feature of the Radial Acceleration Relation (McGaugh et al. 2016). In TEP, this coupling is a natural consequence of the assumed sourcing: the scalar field is sourced by the trace of the energy-momentum tensor (T_μ^μ), so the "dark matter" halo is directly anchored to the baryonic mass. This recovers the MOND-like phenomenology of the RAR without modifying inertia or requiring new force laws, through the non-linear response of the scalar sector.

Ultra-Diffuse Galaxies (DF2/DF4)

The "dark matter free" galaxies NGC 1052-DF2 and DF4 pose a challenge to theories where dark matter and baryons are dynamically coupled. TEP offers an illustrative mechanism — "soliton stripping" — stated here as a qualitative possibility pending a dynamical treatment: the scalar field is not a separable substance, so "stripping" denotes the disruption and slow re-equilibration of the baryon-anchored field configuration during a high-velocity encounter (as proposed by van Dokkum et al. 2022), with the timescale set by the effective mass m_{eff} and the encounter velocity rather than by material removal. The remaining baryonic component would appear

devoid of dark matter until the field configuration re-equilibrates. A quantitative version requires the relaxation timescale of the disrupted profile to be computed; until then the mechanism is registered as illustrative, not as a demonstrated channel.

Systematic Uncertainties

The primary uncertainty in the global fit remains the GNSS calibration length L_c . Three distinct contributions comprise this uncertainty: (i) the 25-year CODE per-measurement statistical uncertainty ($4,201 \pm 1,967$ km, $\pm 47\%$; Paper 2); (ii) the inter-center systematic spread ($3,330$ – $4,549$ km, $\sim \pm 12\%$; Paper 1), with the caveat that centers share largely overlapping underlying data so this spread is not an independent ensemble; and (iii) the $\mathcal{O}(1)$ prefactor in the transfer sketch (Section 2), treated as a model-defining choice. The adopted operational value is $L_c = 4,200$ km with ± 500 km ($\pm 12\%$) reflecting the inter-center range. The $\pm 47\%$ per-measurement statistical uncertainty from the single-center CODE fit is retained as a conservative upper bound on measurement-to-measurement scatter, but the $\pm 12\%$ inter-center spread is preferred as the operational systematic because it reflects the actual range of scales recovered by independent processing pipelines applied to the same underlying data, and the consensus value ($4,200$ km) lies within that range. Propagating this alone gives $\rho_T = 20 \pm 7$ g/cm³; the epoch/window instability established by the executed held-out discriminator (Paper 14 Step 3.1; quarterly-window scatter of $1,050$ – $1,935$ km on an identical product) is the dominant calibration systematic and widens the effective band to roughly 8 – 70 g/cm³ (Appendix B.8). The $M^{1/3}$ structural form is independent of ρ_T ; only the normalization changes. Even the widened band is far smaller than the dynamic range over which the model is tested, preserving the falsifiability of the scaling law.

Predictions for High-Redshift JWST Observations

The TEP framework makes concrete predictions for the "Little Red Dots"—compact, massive galaxies at $z > 5$ discovered by JWST. These objects are a critical testing ground because they are compact enough for scalar-envelope effects to dominate their apparent structure, yet massive enough for JWST to resolve the predicted kpc-scale transition region.

Qualitative Predictions for High-Redshift JWST Observations

For a target galaxy with stellar mass $M_* = 10^{10} M_\odot$ (typical of LRDs), the TEP model makes the following qualitative predictions:

1. Soliton Envelope Scale

The fundamental core saturation density $\rho_T \approx 20$ g/cm³ governs the compact soliton surface, but the observable galactic transition is set by the diffuse screening density $\rho_{\text{trans}} \approx 3 \times 10^{-23}$ g/cm³ (fitted from SPARC, Section 4). The predicted emission extent therefore scales as $R_{\text{trans}} \propto M^{1/3}$ at the transition density, not at ρ_T . Precise numerical predictions for high-redshift galaxies require epoch-specific calibration of ρ_{trans} ; until then, the model predicts that resolved gas tracers (e.g., H α) should reveal more extended kinematic structure than expected from a purely stellar compact core.

2. Velocity Dispersion

The dynamical support speed is set by the soliton potential at the transition radius: $\sigma_v^2 \sim GM/R_{\text{trans}}$. Because $R_{\text{trans}} \propto M^{1/3}$, the velocity dispersion scales as $\sigma_v \propto M^{1/3}$. This predicts characteristic velocity dispersions intermediate between the broad-line AGN regime (> 1000 km/s) and the low-mass dwarf regime (< 50 km/s). Precise normalization awaits epoch-specific calibration of ρ_{trans} .

Discriminant: Soliton vs. Supermassive Black Hole

A leading alternative explanation is that LRDs are reddened Active Galactic Nuclei (AGNs) powered by supermassive black holes ($M_{\text{BH}} \sim 10^7$ – $10^8 M_\odot$).

- **AGN Hypothesis:** Emission should be point-like or confined to the nuclear region (< 1 kpc). Broad lines reflect BLR physics ($v > 1000$ km/s).
- **TEP Soliton Hypothesis:** Emission should be extended over the galactic transition scale (kpc-scale for typical LRD masses, pending epoch-specific calibration of ρ_{trans}). Broad lines reflect the deep gravitational potential of the soliton well (~ 100 – 200 km/s).

Discriminant: Spatially resolve the H α emission. Extended kpc-scale emission with velocity dispersions ~ 100 – 200 km/s would disfavor a purely nuclear AGN interpretation, while strongly compact emission with $v > 1000$ km/s would favor an AGN-dominated explanation. This constitutes a concrete observational discriminator for this particular soliton-interpretation channel.

Implications for Paper 7 (RBH-1)

The GNSS-calibrated ρ_T identified here provides the working normalization for the analysis of RBH-1 in the companion paper (Smafield 2025h). By adopting $\rho_T \approx 20$ g/cm³ (with stated uncertainty), the soliton interpretation for RBH-1 becomes tightly constrained. The wake structure, velocity jump, and thermal emission properties can then be compared against the interaction of the wake with a soliton of characteristic radius $R \approx 7.8 \times 10^7$ km. Deviations would primarily constrain the object-specific soliton

interpretation and/or the assumed mapping between ρ_T and the effective radius in this environment, rather than the broader TEP framework.

Cross-Domain Constraint Network

Tier	Role	Channels	Relation to ρ_T
A	Direct / quasi-direct calibration	GNSS, SPARC/MW, RBH-1, condensed-matter EOS	Enters ρ_T normalization directly
B	Screening / response-transfer test	Pulsars (10), Cepheids (11), wide binaries (13), LLR (17), Earth flyby (15)	Constrains $S_\Sigma(\mathcal{E})$ and transfer functions, not independent ρ_T
C	Phantom Mass / conformal transport	Lensing (4), multiply-imaged SNe (19), JWST (12)	Tests the mechanism; not a ρ_T calibration
D	Orthogonal architecture test	J0437 (16), hi_class (18), C0 (26), TH (27), QF (23)	Tests deep TEP structure; does not calibrate ρ_T

Only Tier A contributes to the present ρ_T normalization. Tiers B–D constrain the environmental operator, response hierarchy, and theoretical viability of the same TEP architecture.

The TEP programme now spans more than twenty papers across terrestrial, galactic, compact-object, lensing, and cosmological domains. It is essential to distinguish which of these constrain ρ_T directly and which constrain the broader TEP architecture. The following tiered classification makes this dependence explicit and prevents the misreading that every paper independently confirms the same density number.

Tier A — Direct or Quasi-Direct ρ_T Constraints

These channels calibrate the macroscopic saturation scale most directly:

- **GNSS λ_T (Papers 1–3):** The primary empirical anchor. The 4,200 km correlation length is conditionally identified with the projected covariance scale associated with $R_T(M_\oplus)$, up to an order-unity transfer factor, yielding the operational estimate $\rho_T \approx 20 \text{ g/cm}^3$.
- **SPARC/Milky Way onset scaling (this work):** Quasi-direct scaling consistency. The $M^{1/3}$ scaling of dark-matter onset radii across 175 spiral galaxies is consistent with the saturation-radius construction, but does not independently calibrate ρ_T . The Milky Way test uses the SPARC-calibrated normalization and is therefore not an independent measurement.
- **RBH-1 crossover (Paper 7):** The runaway-black-hole candidate tests the extrapolation of $R_T(M)$ to $M \sim 10^7 M_\odot$, with wake structure providing an object-specific consistency test.
- **Condensed-matter EOS coincidence (this work, Section 5):** The proximity of $\rho_T \approx 20 \text{ g/cm}^3$ to warm-dense-matter / degeneracy-onset scales provides a coincidence-level physical plausibility check, not an independent empirical calibration.

Tier B — Screening and Response-Transfer Constraints

These channels do not independently calibrate ρ_T , but they constrain the environmental operator $S_\Sigma(\mathcal{E})$ and the transfer functions that connect ρ_T to observables:

- **Globular-cluster pulsars (Paper 10):** The suppressed density scaling of spin-down excess ($\Gamma \approx 0.39 \text{ dex/dex}$, vs. Newtonian 0.75) constrains how dense environments screen the temporal-field response. The observable coefficient $\tilde{\kappa}_{\text{MSP}}$ is a downstream projection, not a direct ρ_T measurement.
- **Cepheids (Paper 11):** The Cepheid clock-bias response — $0.33\text{--}0.45 \times 10^6$ empirically, against the $\sim 10^6$ benchmark normalization — in galactic disks bridges the weakly screened regime. It agrees with $\tilde{\kappa}_{\text{MSP}}$ only after environmental transfer factors are applied, demonstrating that response coefficients are channel-specific projections.
- **Wide binaries (Paper 13):** Gaia DR3 reveals a weak-field screening transition at a few thousand AU with environmental ordering (high- $|Z|$ vs. midplane). This is an independent *morphology* test of Temporal Shear recovery, not an independent ρ_T calibration: the paper explicitly states that imported ρ_T and g_{TEP} enter only as cross-scale consistency checks.
- **Lunar Laser Ranging (Paper 17):** Reports a compactness-dependent residual channel at high significance, but remains at the "candidate residual" level pending integrator-level ephemeris refits. Classified as a *guardrail*: it suggests the local screened regime may still contain residual-channel structure, but does not yet provide a completed ρ_T likelihood term.
- **Earth flyby (Paper 15):** Uses the UCD $\rho_T \approx 20 \text{ g/cm}^3$ as a *shared input*, not an independent validation. Valuable as a downstream posterior predictive test: if the terrestrial screening profile is correct, flyby anomalies should organize under the restricted trajectory model.

Tier C — Phantom Mass / Conformal Transport Tests

These channels test the *mechanism* by which TEP generates apparent mass discrepancies, without directly calibrating ρ_T :

- **Gravitational lensing (Paper 4):** The Phantom Mass framework shows that the Temporal Shear acts on matter trajectories while lensing tracks the gravitating matter, so the theory's dark-sector signature is the dynamical–lensing mass difference. This is *mechanistic support* for the TEP interpretation of the dark sector, not a ρ_T calibration.
- **Multiply-imaged supernovae (Paper 19):** SN Refsdal provides directional evidence from blind model residuals for conformal time-transport in the lensing sector. The amplitude kernel remains phenomenological; included as a *sign test* for Phantom Mass, pending derivation of the full transfer kernel.
- **JWST high-redshift galaxies (Paper 12):** Tests potential-depth response at $z > 5$ using an external response prior. The signal is not an independent ρ_T measurement; it probes whether the same conformal architecture organizes high-redshift compact galaxies.

Tier D — Orthogonal TEP-Architecture Tests

These papers do not calibrate ρ_T at all, but they test deep structural claims of the TEP framework:

- **J0437 scintillation (Paper 16):** Reports non-zero phase-closure signal in millisecond pulsar scintillation, rejecting standard scalar-delay ISM models ($\psi = 0$) at high confidence. This tests *non-integrable time transport* / synchronization holonomy, one of the deepest TEP claims. It does not calibrate ρ_T .
- **hi_class cosmology (Paper 18):** Demonstrates that the conformal temporal-transport geometry preserves CMB acoustic structure, sound-horizon behaviour, and stable linear perturbations. A *global consistency constraint*, not a ρ_T term.
- **C0 supernova conformal redshift (Paper 26):** Shows that Pantheon+ distance-redshift data do not uniquely require primitive spatial expansion. Supports the temporal-transport interpretation of cosmological redshift.
- **TH temporal horizon (Paper 27):** The apparent $a_{\text{eff}} \rightarrow 0$ limit becomes a temporal conformal boundary rather than a physical singularity. Provides *theoretical closure* for the late-time cosmology.
- **QF quantum tangent limit (Paper 23):** Shows that the screened quantum tangent limit is consistent with the TEP architecture. A *theoretical compatibility* check, not empirical ρ_T evidence.

Summary: Only Tier A enters the present ρ_T normalization argument directly. Tiers B–D constrain the environmental operator, response hierarchy, and theoretical viability of the same TEP architecture. This tiered dependence is stronger than a flat pile of evidence because it acknowledges exactly where each paper contributes and where it does not.

9. Conclusion

A Universal Organizing Parameter

This paper identifies the saturation scale $\rho_T \approx 20 \text{ g/cm}^3$ as a candidate organizing parameter for gravitational phenomena across planetary, galactic, compact-object, and RBH-scale regimes. The convergence of terrestrial GNSS timing structure, SPARC galaxy onset scaling, Milky Way inner mass-discrepancy structure, and the density-ordered screening hierarchy motivates the hypothesis that ρ_T is a real physical scale in the temporal-field topology, not a system-specific fitting parameter. The screening hierarchy is not treated as an independent discovery of the $1/3$ exponent; rather, it demonstrates that the same ρ_T coherently recovers GR-like behavior in dense systems while permitting active-shear scalar phenomenology in diffuse regimes.

Key Results

The primary findings are:

1. **GNSS Calibration:** Distance-structured correlations in atomic clocks yield $L_e \approx 4200 \text{ km} \rightarrow \rho_T \approx 20 \text{ g/cm}^3$. This calibration exhibits 25-year stability, multi-center consistency, and survives raw RINEX validation.
2. **Galactic Scaling:** SPARC rotation curves yield $\alpha_{\text{SPARC}} = 0.355 \pm 0.043 \text{ (stat)} \pm 0.07 \text{ (definition)}$, consistent with the $M^{1/3}$ expectation within $\sim 0.3\sigma$.
3. **Screening Hierarchy:** 26 objects spanning ~ 15 orders of magnitude in density reveal a consistency relation $S \propto \rho^{1/3}$, algebraically expected from the $R_T(M)$ construction, explaining why GR tests pass (binary pulsars: $S \sim 29,000$) while galactic dynamics are in the active-shear regime ($S \sim 10^{-9}$ at $\rho \sim 10^{-24} \text{ g/cm}^3$).
4. **Milky Way Test:** The SPARC-calibrated $M^{1/3}$ relation predicts a dark-matter onset radius $R_{\text{DM}} \approx 3 \text{ kpc}$ for the Milky Way, consistent with the observed transition from baryonic to dark-matter-dominated rotation at $R \sim 3\text{--}5 \text{ kpc}$, providing a local scale-consistency check.

Dark Matter as Phantom Mass

The TEP framework reinterprets dark-matter observations as violations of the isochrony axiom—the assumption that spatially separated clocks at the same gravitational potential tick at the same rate. The Temporal Shear acts on matter trajectories but not on photon paths (null curves are conformally invariant), so mass inferred from dynamics and mass inferred from lensing diverge in low-acceleration environments—creating "phantom mass" as an inference artifact without invoking non-baryonic particles. This dynamical–lensing mass difference — the phantom mass — is the theory's falsifiable dark-sector signature (Paper 19); persistent dynamical–lensing agreement bounds the shear's share of the dynamical anomaly. The companion closure obligation is registered:

replacing particulate dark matter additionally requires a CDM-free gravitational-metric prediction of galaxy, cluster and CMB lensing, and that common optical-sector closure is not established by the present force-level construction (Paper 0, §8).

This reinterpretation offers a candidate common account of several dark-sector puzzles:

- **Universal scaling:** The $M^{1/3}$ relation follows naturally from the saturation-radius construction.
- **Dark-matter-deficient UDGs:** Field-envelope disruption ("soliton stripping") offers an illustrative explanation for DF2/DF4-like systems, pending computation of the re-equilibration timescale.
- **Core-cusp structure:** Saturated temporal-field cores offer a possible route to flat effective density profiles.
- **Baryonic Tully-Fisher behaviour:** The $M^{1/3}$ scaling may contribute to BTFR-like phenomenology, though a full BTFR derivation is not completed here.

Astrophysical Applications

The GNSS-anchored value $\rho_T \approx 20 \text{ g/cm}^3$ enables specific applications for astrophysical systems. The companion paper (Smawfield 2025h, Paper 7) applies this calibration to the RBH-1 runaway black hole candidate, yielding a saturation radius $R_T \approx 7.8 \times 10^7 \text{ km}$. This provides a testable extrapolation: if the RBH-1 mass is revised by a factor of 3, the expected radius would change by $\sim 44\%$, offering a clear consistency check with observations.

Future applications include:

- **JWST high-redshift galaxies:** "Little Red Dots" and massive quiescent galaxies at $z > 10$ may exhibit soliton signatures.
- **EHT polarimetry:** M87* and Sgr A* polarization patterns could reveal scalar field structure near supermassive compact objects.
- **Gravitational waves:** Binary black hole mergers may produce scalar radiation detectable by LISA.
- **Strong lensing time delays:** The phantom-mass sector can act on time-delay cosmography through the mass-model normalization — the dynamical-lensing mass difference biases the velocity-dispersion prior that breaks the mass-sheet degeneracy, in the direction that elevates the inferred H_0 (Paper 19, Step 58 channel audit); the propagated-delay sector itself is clean at $\lesssim 10^{-7}$ fractional.

Theoretical Implications

The convergence of terrestrial, compact, galactic, and cosmological constraints on a single density scale suggests a fundamental organization principle governing gravitational phenomenology across many orders of magnitude in mass. This organization is formalized in the Temporal Equivalence Principle (Smawfield 2025a, Paper 0), which posits that gravitational phenomena arise from a conformal time field $\phi(x^\mu)$ coupled to matter via the action:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} (\partial\phi)^2 - V(\phi) \right] + S_m[\tilde{g}_{\mu\nu}] \quad (37)$$

where $\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu}$ and $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$ in the late-time conformal limit.

The saturation scale ρ_T emerges as the scale at which the scalar field ϕ reaches its self-interaction threshold, saturating the field. Under a soliton interpretation, this would correspond to soliton formation; regardless of microscopic realization, this provides a natural explanation for the $M^{1/3}$ scaling and the dynamical phantom-mass phenomenon.

Systematic Uncertainties and Robustness

The primary uncertainty is the GNSS length scale determination. Four distinct contributions are reported: (i) the 25-year CODE per-measurement statistical uncertainty ($4,201 \pm 1,967 \text{ km}$, $\pm 47\%$; Paper 2); (ii) the inter-center systematic spread ($3,330\text{--}4,549 \text{ km}$, $\sim \pm 12\%$; Paper 1), with the caveat that centers share underlying data so this spread is not an independent ensemble; (iii) the epoch/window instability established by the executed held-out discriminator (Paper 14 Step 3.1): the identical estimator on the identical product returns λ from 1,050 to 1,935 km across quarterly windows, while matched-day independent chains agree at $\sim 30\%$ ($2,744$ vs. $1,933 \text{ km}$ — the CNES value landing within 4% of the pooled $1,862 \text{ km}$ — same-product fallback days excluded by epoch-aligned detrended-residual comparison) — the dominant calibration systematic is the $\sim 30\text{--}40\%$ epoch/window scatter; and (iv) the $\mathcal{O}(1)$ prefactor in the transfer sketch (Section 2), treated as a model-defining choice. The adopted operational value is $L_e = 4,200 \text{ km}$ with $\pm 500 \text{ km}$ ($\pm 12\%$) reflecting the inter-center range, plus the epoch/window systematic carried separately. Propagating the inter-center spread alone gives $\rho_T = 20 \pm 7 \text{ g/cm}^3$; propagating the dominant epoch/window scatter widens the effective band to roughly $8\text{--}70 \text{ g/cm}^3$, and the MGEX held-out scale (ρ_T up to $\sim 5 \times 10^2 \text{ g/cm}^3$ if adopted at face value) is recorded as an unresolved epoch-scale tension pending longer MGEX baselines. The $M^{1/3}$ structural form is independent of ρ_T ; only the normalization changes. The present evidence consists of one primary terrestrial timing calibration, one galactic onset-scaling consistency test, one

local SPARC-internal Milky Way check, and one algebraic screening-consistency hierarchy. These layers are mutually consistent, but they are not statistically independent measurements of ρ_T .

The robustness of the result is demonstrated by:

- **Multi-center consistency:** CODE, IGS, and ESA yield consistent GNSS correlation structure across independent processing centres.
- **Temporal stability:** 25-year consistency disfavors transient effects.
- **Raw data validation:** RINEX analysis constrains processing artifacts.
- **Cross-scale agreement:** Four tests spanning 18 orders of magnitude in mass.

Future Directions

The identification of $\rho_T \approx 20 \text{ g/cm}^3$ as a candidate cross-scale saturation parameter opens several avenues for future research:

1. **Precision tests:** Improved GNSS analysis (longer baselines, more stations) could tighten L_c to $\sim 5\%$ accuracy.
2. **High-redshift galaxies:** JWST observations of $z > 10$ systems could test whether soliton formation affects early galaxy assembly.
3. **Gravitational wave cosmology:** LISA could detect scalar radiation from binary black hole mergers, providing direct evidence for the time-field sector.
4. **Laboratory tests:** Atomic interferometry experiments could search for temporal gradients at sub-meter scales.

Concluding Remarks

This paper identifies $\rho_T \approx 20 \text{ g/cm}^3$ as a candidate macroscopic saturation/proximity scale in the TEP environmental-response hierarchy. GNSS clock correlations provide the primary terrestrial timing anchor for the Earth-scale covariance length. SPARC onset scaling, Milky Way structure, RBH-1 crossover behaviour, compact-object screening, and condensed-matter/EOS considerations then test whether the same scale organizes otherwise disconnected gravitational regularities without retuning. Channel-specific quantities such as ρ_{trans} , R_s , κ_{Cep} , κ_{MSP} , and lensing response coefficients are not independent measurements of ρ_T , but downstream projections of $S_{\Sigma}(\mathcal{E})$.

This paradigm shift—from dark matter as substance to dark matter as temporal shear—provides a candidate unified organizing framework for the $M^{1/3}$ scaling, screening hierarchy, and dynamical phantom-mass phenomenon, to the extent permitted by lensing–dynamics consistency. The GNSS-anchored value of ρ_T enables specific testable applications, including the RBH-1 case study (Paper 7), positioning the Temporal Equivalence Principle as an empirically grounded candidate alternative to the particle dark matter paradigm in the dynamical sector, with the dynamical–lensing mass comparison (Paper 19) as its decisive open test.

References

- Abedi, J., Dykaar, H., & Afshordi, N. 2017, *Phys. Rev. D*, 96, 082004 (arXiv:1612.00266)
- Allen, M. G., Groves, B. A., Dopita, M. A., Sutherland, R. S., & Kewley, L. J. 2008, *ApJS*, 178, 20 (arXiv:0805.0204)
- Barro, G., et al. 2025, *From "The Cliff" to "Virgil": Mapping the Spectral Diversity of Little Red Dots with JWST/NIRSpec* (arXiv:2512.15853)
- Bland-Hawthorn, J., & Gerhard, O. 2016, *ARA&A*, 54, 529 (Milky Way mass model)
- Campanelli, M., Lousto, C. O., Zlochower, Y., & Merritt, D. 2007, *Phys. Rev. Lett.* (arXiv:gr-qc/0702133)
- Cardoso, V., Hopper, S., Macedo, C. F. B., Palenzuela, C., & Pani, P. 2016, *Phys. Rev. D*, 94, 084031 (arXiv:1608.08637)
- Chen, K., Li, Z., Inayoshi, K., & Ho, L. C. 2025, *ApJ Lett.* (DOI: 10.3847/2041-8213/ae1955; arXiv:2505.22600)
- Colpi, M., & Dotti, M. 2011, *Adv. Astron.*, 2011, 1 (arXiv:0906.4339)
- Colpi, M., Geppert, U., & Page, D. 2000, *ApJ*, 529, L29 (DOI: 10.1086/312448; arXiv:astro-ph/9912066)
- Donato, F., Gentile, G., Salucci, P., et al. 2009, *MNRAS*, 397, 1169 (DOI: 10.1111/j.1365-2966.2009.15004.x; arXiv:0904.4054)
- Delvecchio, I., et al. 2025, *A&A* (DOI: 10.1051/0004-6361/202557164; arXiv:2509.07100)
- Event Horizon Telescope Collaboration. 2019, *ApJ Lett.*, 875, L1 (DOI: 10.3847/2041-8213/ab0ec7; M87* image)
- Event Horizon Telescope Collaboration. 2021, *ApJ Lett.*, 910, L12 (DOI: 10.3847/2041-8213/abe71d; M87* polarization)
- Event Horizon Telescope Collaboration. 2021, *ApJ Lett.*, 910, L13 (DOI: 10.3847/2041-8213/abe71e; EHT Paper VII)
- Event Horizon Telescope Collaboration. 2022, *ApJ Lett.*, 930, L12 (DOI: 10.3847/2041-8213/ac6674; Sgr A* image)
- Heeck, J., et al. 2021, *Phys. Rev. D*, 103, 115004 (arXiv:2009.08463)
- Hinterbichler, K., & Khoury, J. 2010, *Phys. Rev. Lett.*, 104, 231301 (DOI: 10.1103/PhysRevLett.104.231301; arXiv:1001.4525)

- Khoury, J., & Weltman, A. 2004, *Phys. Rev. Lett.*, 93, 171104 (DOI: 10.1103/PhysRevLett.93.171104; arXiv:astro-ph/0309300)
- Hui, L., Ostriker, J. P., Tremaine, S., & Witten, E. 2017, *Phys. Rev. D*, 95, 043541 (DOI: 10.1103/PhysRevD.95.043541; arXiv:1610.08297)
- Komossa, S. 2012, *Recoiling black holes: electromagnetic signatures, candidates, and astrophysical implications* (arXiv:1202.1977)
- Coleman, S. 1985, *Nucl. Phys. B*, 262, 263 (DOI: 10.1016/0550-3213(85)90286-X)
- Gleiser, M. 1994, *Phys. Rev. D*, 49, 2978 (DOI: 10.1103/PhysRevD.49.2978; arXiv:hep-ph/9308279)
- Gaia Collaboration. 2023, *A&A*, 674, A1 (DOI: 10.1051/0004-6361/202243940; Gaia Data Release 3)
- Gong, Y., Papantonopoulos, E., & Yi, Z. 2018, *Eur. Phys. J. C*, 78, 738 (DOI: 10.1140/epjc/s10052-018-6227-9; arXiv:1711.04102)
- Gronke, M., & Oh, S. P. 2018, *MNRAS*, 480, L111 (arXiv:1806.02729)
- Hviding, R. E., et al. 2025, *A&A* (DOI: 10.1051/0004-6361/202555816; arXiv:2506.05459)
- Seidel, E., & Suen, W.-M. 1991, *Phys. Rev. Lett.*, 66, 1659 (DOI: 10.1103/PhysRevLett.66.1659)
- Kusenko, A. 1997, *Phys. Lett. B*, 404, 285 (arXiv:hep-th/9704073)
- Lelli, F., McGaugh, S. S., & Schombert, J. M. 2016, *AJ*, 152, 157 (DOI: 10.3847/0004-6256/152/5/157; SPARC Database)
- McGaugh, S. S., Lelli, F., & Schombert, J. M. 2016, *Phys. Rev. Lett.*, 117, 201101 (DOI: 10.1103/PhysRevLett.117.201101; arXiv:1609.05917)
- McGaugh, S. S., Schombert, J. M., Bothun, G. D., & de Blok, W. J. G. 2000, *ApJ Lett.*, 533, L99 (DOI: 10.1086/312628; arXiv:astro-ph/0003001)
- Ogiya, G., & Nagai, D. 2023, *ApJ Lett.*, 958, L5 (arXiv:2309.09031)
- Poggianti, B. M., et al. 2019, *ApJ*, 874, 140 (arXiv:1810.05164)
- Rybicki, G. B., & Lightman, A. P. 1979, *Radiative Processes in Astrophysics* (Wiley-VCH)
- Sanchez Almeida, J., Montes, M., & Trujillo, I. 2023, *A&A* (DOI: 10.1051/0004-6361/202346430; arXiv:2304.12344)
- Shive, H.-Y., Chiueh, T., & Broadhurst, T. 2014, *Phys. Rev. Lett.*, 113, 261302 (DOI: 10.1103/PhysRevLett.113.261302; arXiv:1407.7762)
- Sofue, Y. 2020, *PASJ*, 72, 81 (DOI: 10.1093/pasj/psaa065; arXiv:2006.15008)
- Sutherland, R. S., & Dopita, M. A. 1993, *ApJS*, 88, 253 (DOI: 10.1086/191823)
- TDCOSMO Collaboration. 2025, *A&A* (DOI: 10.1051/0004-6361/202555801; arXiv:2506.03023)
- van Dokkum, P., et al. 2023, *ApJ Lett.*, 946, L50 (arXiv:2302.04888)
- van Dokkum, P., et al. 2025, *JWST Confirmation of a Runaway Supermassive Black Hole via its Supersonic Bow Shock* (arXiv:2512.04166)
- Taylor, J. H., & Weisberg, J. M. 1982, *ApJ*, 253, 908 (DOI: 10.1086/159690)
- Vainshtein, A. I. 1972, *Phys. Lett. B*, 39, 393 (DOI: 10.1016/0370-2693(72)90147-5)
- Weisberg, J. M., & Taylor, J. H. 2005, *Binary Radio Pulsars*, ASP Conf. Ser., 328, 25 (arXiv:astro-ph/0407149)
- Westerweck, J., et al. 2018, *Phys. Rev. D*, 97, 124037 (arXiv:1712.09966)
- Will, C. M. 2014, *Living Rev. Relativ.*, 17, 4 (DOI: 10.12942/lrr-2014-4; arXiv:1403.7377)
- Williams, J. G., Turyshev, S. G., & Boggs, D. H. 2012, *Class. Quantum Grav.*, 29, 184004 (DOI: 10.1088/0264-9381/29/18/184004; arXiv:1203.2150)
- Younes, G., et al. 2020, *ApJ Lett.*, 896, L42 (DOI: 10.3847/2041-8213/ab9a48; arXiv:2006.02814)
- Younes, G., et al. 2022, *Nature Astronomy*, 7, 339 (arXiv:2210.11518)
- Zhang, H., et al. 2025, *Polarization Images of Solitonic Boson Stars* (arXiv:2508.11992)
- Zumalacárregui, M., & Bellini, E. 2018, *JCAP*, 2018, 054 (DOI: 10.1088/1475-7516/2018/10/054; arXiv:1705.05302; hi_class cosmology solver)
- Pintore, F., et al. 2016, *MNRAS*, 458, 2088 (DOI: 10.1093/mnras/stw449; arXiv:1602.05950)
- Ray, P. S., et al. 2019, *ApJ*, 879, 130 (DOI: 10.3847/1538-4357/ab24d8)
- Tuo, Y. L., et al. 2024, *ApJ*, 966, 80 (DOI: 10.3847/1538-4357/ad2fb6; arXiv:2403.12137)
- Kramer, M., et al. 2006, *Science*, 314, 97 (DOI: 10.1126/science.1132305)
- van Dokkum, P., et al. 2018, *Nature*, 555, 629 (DOI: 10.1038/nature25767)
- van Dokkum, P., et al. 2019, *ApJ Lett.*, 874, L5 (DOI: 10.3847/2041-8213/ab0d92)
- van Dokkum, P., et al. 2022, *Nature*, 605, 435 (DOI: 10.1038/s41586-022-04665-6)
- Jiao, Y., et al. 2023, *A&A*, 678, A208 (DOI: 10.1051/0004-6361/202347513; arXiv:2309.00048)

- Archibald, R. F., et al. 2013, *Nature*, 497, 591 (DOI: 10.1038/nature12159)
- Archibald, R. F., et al. 2017, *ApJ*, 829, L21 (DOI: 10.3847/2041-8205/829/1/L21; arXiv:1608.01007)
- de Graaff, A., et al. 2025, *A&A* (arXiv:2503.01891)
- Dib, R., & Kaspi, V. M. 2014, *ApJ*, 784, 37 (DOI: 10.1088/0004-637X/784/1/37; arXiv:1401.5738)
- Nicolis, A., Rattazzi, R., & Trinchieri, E. 2009, *Phys. Rev. D*, 79, 064036 (DOI: 10.1103/PhysRevD.79.064036; arXiv:0811.2197)
- Olausen, S. A., & Kaspi, V. M. 2014, *ApJS*, 212, 6 (DOI: 10.1088/0067-0049/212/1/6; arXiv:1309.4167)
- Şaşmaz Muş, S., et al. 2014, *MNRAS*, 440, 2916 (DOI: 10.1093/mnras/stu445; arXiv:1402.6054)

TEP Research Series

- Smawfield, M. L. (2026). *Temporal Equivalence Principle: Dynamic Time & Emergent Light Speed*. Preprint v0.15 (Jakarta). Zenodo. DOI: 10.5281/zenodo.16921911 (Paper 0)
- Smawfield, M. L. (2025b). *Global Time Echoes: Distance-Structured Correlations in GNSS Clocks*. Preprint v0.27 (Jaipur). Zenodo. DOI: 10.5281/zenodo.17127229 (Paper 1)
- Smawfield, M. L. (2025c). *Global Time Echoes: 25-Year Analysis of CODE Precise Clock Products*. Preprint v0.20 (Cairo). Zenodo. DOI: 10.5281/zenodo.17517141 (Paper 2)
- Smawfield, M. L. (2025d). *Global Time Echoes: Raw RINEX Consistency Test*. Preprint v0.8 (Kathmandu). Zenodo. DOI: 10.5281/zenodo.17860166 (Paper 3)
- Smawfield, M. L. (2025e). *Temporal-Spatial Coupling in Gravitational Lensing: A Reinterpretation of Dark Matter Observations*. Preprint v0.8 (Tortola). Zenodo. DOI: 10.5281/zenodo.17982540 (Paper 4)
- Smawfield, M. L. (2025f). *Global Time Echoes: Empirical Synthesis*. Preprint v0.7 (Singapore). Zenodo. DOI: 10.5281/zenodo.18004832 (Paper 5)
- Smawfield, M. L. (2025g). *Temporal Topology Saturation Scale: Cross-Scale Consistency of ρ_T* . Preprint v0.8 (New Delhi). Zenodo. DOI: 10.5281/zenodo.18064365 (Paper 6 — this work)
- Smawfield, M. L. (2025h). *The Soliton Wake: Exploring RBH-1 as a Temporal Topology Candidate*. Preprint v0.4 (Blantyre). Zenodo. DOI: 10.5281/zenodo.18059250 (Paper 7)
- Smawfield, M. L. (2025). *Global Time Echoes: Optical-Domain Consistency Test via Satellite Laser Ranging*. Preprint v0.6 (Mombasa). Zenodo. DOI: 10.5281/zenodo.18064581 (Paper 8)
- Smawfield, M. L. (2025). *What Do Precision Tests of General Relativity Actually Measure?*. Preprint v0.8 (Istanbul). Zenodo. DOI: 10.5281/zenodo.18109760 (Paper 9)
- Smawfield, M. L. (2026). *Temporal Equivalence Principle: Suppressed Density Scaling in Globular Cluster Pulsars*. Preprint v0.9 (Caracas). Zenodo. DOI: 10.5281/zenodo.18165798 (Paper 10)
- Smawfield, M. L. (2026). *The Cepheid Bias: Resolving the Hubble Tension*. Preprint v0.10 (Kingston upon Hull). Zenodo. DOI: 10.5281/zenodo.18209702 (Paper 11)
- Smawfield, M. L. (2026). *Temporal Equivalence Principle: A Unified Resolution to the JWST High-Redshift Anomalies*. Preprint v0.7 (Kos). Zenodo. DOI: 10.5281/zenodo.19000827 (Paper 12)
- Smawfield, M. L. (2026). *Temporal Equivalence Principle: Temporal Shear Recovery in Gaia DR3 Wide Binaries*. Preprint v0.6 (Kilifi). Zenodo. DOI: 10.5281/zenodo.19102061 (Paper 13)
- Smawfield, M. L. (2026). *Temporal Equivalence Principle: Native hi_class Conformal Implementation, Linear Perturbation Closure, and CMB Acoustic Peak Preservation*. Preprint v0.7 (Cambridge). Zenodo. DOI: 10.5281/zenodo.20572722 (Paper 18)

Contact Information

Author: Matthew Lukin Smawfield

Affiliation: Independent Researcher

Email: matthew@mlsmawfield.com

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
 Version: v0.8 (New Delhi) · Last updated: 30 September 2026
 Revision note (v0.8): consistency revision aligning screening terminology with Jakarta v0.15.

Appendix A: GNSS Calibration — Summary of Evidence

The claim that terrestrial clock correlations calibrate a fundamental density parameter ($\rho_T \approx 20 \text{ g/cm}^3$) is extraordinary and requires rigorous justification. This appendix summarizes the key validation results from the companion GNSS papers (Smawfield 2025b,c,d) and addresses common concerns about systematics.

A.1 Physical Mechanism

In the TEP framework, the scalar field ϕ modulates proper time via a conformal factor $A(\phi) = \exp(\beta_A \phi / M_{\text{Pl}})$. Near a gravitating mass, the field profile creates a spatial gradient in clock rates. GNSS atomic clocks, distributed across Earth's gravitational field, sample this gradient at different radial positions. The characteristic correlation length $L_c \approx 4200 \text{ km}$ corresponds to the scale where the scalar field's gradient becomes steep enough to produce measurable timing correlations—the saturation radius for Earth's mass.

A.2 Why Not Ionospheric, Tropospheric, or Orbital Systematics?

Scale Separation

The 4200 km correlation length is distinct from known geodetic systematics:

- **Tropospheric correlations:** $\sim 100\text{--}500 \text{ km}$ (weather systems)
- **Ionospheric correlations:** $\sim 500\text{--}2000 \text{ km}$ (TEC structures)
- **Orbital harmonics:** $\sim 12 \text{ hr}$ (half-sidereal period)

Power spectral analysis shows the 4200 km feature as a persistent background covariance floor in the GPS-only clean baseline, with no counterpart in ionospheric or tropospheric delay products; multi-constellation validation shows consistent signatures.

Multi-Center Verification

The correlation structure persists across independent clock solutions from three analysis centers:

Center	λ Median (km)	95% CI (km)	R ² (pooled)
CODE	4,181	1,198–5,918	0.920
IGS Combined	3,763	3,197–4,871	0.968
ESA Final	3,193	2,532–3,984	0.970

Table A.1: Multi-center correlation parameters. The bootstrap-median scale spans 3,193–4,181 km across centres (CV 13.4%); the corresponding exponential best-fit range is 3,330–4,549 km (CV 18.2%, Paper 1). Because centers process largely shared underlying data, this spread is not an independent ensemble average; it is treated as a systematic range rather than a statistical error reduction. λ values are bootstrap medians; R^2 is from the exponential best fit.

Null Tests

Three null tests were performed (all satisfied at $>5\sigma$):

1. **Epoch randomization:** Shuffling satellite epochs destroys the correlation ($r^2 < 0.01$, $N = 1000$ shuffles)
2. **Residual shuffling:** Shuffling clock residuals within each satellite eliminates the spatial structure
3. **White noise injection:** Replacing real data with white noise yields no coherent scale

A.3 25-Year Temporal Stability

The correlation length has been confirmed over a 25-year temporal baseline (1999–2024) using CODE final products. The signal shows:

- Decadal stability (no secular drift)
- Persistent spatial anisotropy
- Strong orbital velocity coupling ($r = -0.86$, $p = 0.002$ for Southern Hemisphere)
- Annual-phase direction recovery near Earth's aphelion velocity tangent; the directed CMB-apex template is anti-phase

A.4 Raw RINEX Validation

To strongly constrain the processing artifact hypothesis, the correlation was detected in raw RINEX observations processed with Single Point Positioning (SPP) using broadcast ephemerides only—no network solutions, no precise orbits, no clock products.

Signature	PPP (Processed)	SPP (Raw)	Status
Exponential decay	$R^2 = 0.92\text{--}0.97$	$R^2 = 0.94$	✓ Confirmed
Directional anisotropy	Detected	Detected	✓ Confirmed
Orbital velocity coupling	$r = -0.86$	Detected	✓ Confirmed
Preferred-direction recovery	RA = 6° , Dec = $+4^\circ$; $\sim 6^\circ$ from aphelion velocity tangent; CMB apex anti-phase	Detected	✓ Confirmed

Table A.2: Raw RINEX validation results. All signatures detected in processed (PPP) data are independently confirmed in raw (SPP) observations, constraining processing artifacts as the origin of the observed signatures.

A.5 What the GNSS Measurement Does and Does Not Claim

What GNSS Measures

The GNSS analysis measures a characteristic correlation length $L_c \approx 4200$ km in atomic clock residuals. This length is operationally interpreted as the projected Temporal Topology covariance scale associated with $R_T(M_\oplus)$, the geometric saturation scale for Earth's mass under the TEP framework.

What GNSS Does Not Claim

The GNSS analysis does not claim to have detected the scalar field directly. The correlation length is an empirical calibration parameter. The physical interpretation (scalar field soliton) is a hypothesis to be tested by cross-regime consistency—which is the purpose of this manuscript.

A.6 Companion Papers

Full methodology, data products, and reproducibility information are available in the companion papers:

- **Smawfield 2025b:** Multi-center validation (CODE, ESA, IGS)
- **Smawfield 2025c:** 25-year longitudinal analysis
- **Smawfield 2025d:** Raw RINEX validation
- **Smawfield 2025f:** Integrated synthesis (TEP-GTE)

Analysis code and data products are available at: github.com/matthewsmawfield/TEP-GNSS

Appendix B: Open Questions and Limitations

The TEP-UCD manuscript makes several strong empirical claims. This appendix discusses active research questions and known limitations. Addressing these is essential for the theory to mature from a consistency argument to a fully predictive framework.

B.1 The GNSS Correlation Mechanism

The operational identification of the terrestrial coherence length $L_c \approx 4200$ km with the projected covariance scale associated with the geometric saturation radius $R_T(M_\oplus) = (3M_\oplus/4\pi\rho_T)^{1/3}$ assumes that the observed exponential decay in clock residual correlations is controlled by the source-mass scale $R_T(M)$, rather than by network geometry, processing, or by the in-medium Compton wavelength $\lambda_C(\rho)$ of any specific candidate completion. This is not guaranteed. Alternative explanations include:

- **Network geometry:** The spatial distribution of GNSS stations is not random; continental clustering could imprint a characteristic scale unrelated to any physical field.
- **Common-mode clock processing:** Inter-analysis-center differences in orbit/clock estimation software may introduce correlated noise with scale-dependent structure.
- **Ephemeris systematics:** Orbital errors propagate into clock solutions with characteristic spatial signatures (e.g., along-track harmonics) that could mimic an exponential envelope.

- **Flicker noise and power-law correlations:** The null-model covariance analysis (TEP-GNSS, Paper 1) formally excludes power-law and constant kernels, but a more sophisticated non-stationary model might evade the current tests.

Mitigation: The null-model covariance analysis (`null_model_covariance.py`) systematically compares the exponential hypothesis against ionospheric dipole, power-law flicker, Gaussian, and Matérn kernels. The exponential model is preferred by AIC/BIC across all three analysis centers. Nevertheless, a dedicated hardware-injected timing perturbation experiment (e.g., coordinated clock offset on selected stations) would provide conclusive causal evidence.

B.2 Earth is Not a Soliton

The narrative operationally identifies L_c with the projected covariance scale associated with $R_T(M_\oplus) = (3M_\oplus/4\pi\rho_T)^{1/3}$. For Earth mass $M_\oplus$, this yields $R_{T,\oplus} \approx 4146$ km and $\rho_T \approx 20$ g/cm³. However, Earth's mean density is $\rho_\oplus \sim 5.5$ g/cm³, well below the quoted saturation scale. The planet therefore sits in the *transition* regime, not the deep-screening regime. The identification of the GNSS correlation scale with $R_T(M_\oplus)$ is phenomenologically motivated but lacks a single-potential $V(\phi)$ demonstration that both are the same parameter.

Mitigation: A complete mitigation requires a dedicated numerical solution of a specified completion (e.g., Box 6.5) in the terrestrial transition regime, with the quartic normalization held at the corpus value and ρ_T fixed by the calibrated L_c , to demonstrate how an observable clock-network correlation length maps onto the geometric scale $R_T(M_\oplus)$. This single-potential demonstration remains an open target.

B.3 The SPARC Exponent is Empirical, Not Uniquely $\rho^{1/3}$

The empirical dark-matter onset exponent $\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ (stat) ± 0.07 (definition) is consistent with $1/3 = 0.333$ at approximately 0.3σ under combined uncertainty. This agreement is encouraging but not uniquely predicted by TEP without showing that the TEP field equation *derives* the $M^{1/3}$ scaling at halo densities. The present manuscript presents the scaling as a consistency check: if ρ_T is universal, then $R_T \propto M^{1/3}$, and the DM onset radius should follow the same organization. A first-principles derivation of $\alpha_{\text{pred}} = 1/3$ from the screened Klein-Gordon equation in the galactic potential remains open.

Mitigation: The theoretical expectation $\alpha = 1/3$ is explicitly derived in Section 3 of this manuscript (see *Theoretical Expectation from ρ_T*). The SPARC analysis is framed as a *post hoc* consistency check, not a blind prediction. The 0.021 offset from $1/3$ corresponds to approximately 0.5σ using the bootstrap uncertainty and is therefore not statistically significant; the larger threshold-dependent shifts are treated as targets for future galactic-field modelling rather than a refutation.

B.4 Screening Hierarchy: Algebraic Tautology?

The screening fit $S \propto \rho^{0.334}$ with $R^2 = 0.99995$ (11 dense objects, $\rho > \rho_T$, excluding planets and black holes) is extraordinarily tight. Because $S \equiv R_T/R_{\text{phys}}$ and both radii are defined by mass-density relations, the slope $\sim 1/3$ is algebraically expected given the model assumptions. The high R^2 confirms internal consistency; it does not constitute independent confirmation of TEP.

Mitigation: The screening audit (`screening_audit.py`) performs object-by-object residual analysis and class-wise jackknife. The slope remains stable to ± 0.0004 under leave-one-out perturbation, and no single object class dominates the fit. The audit confirms that the hierarchy is robust, though the manuscript explicitly notes that it is a consistency check, not independent evidence.

B.5 Cross-Paper Parameter Drift

Multiple distinct exponents appear across the TEP literature, and their notation has been ambiguous:

Quantity	Typical Value	Source	Physical Meaning
α_{SPARC}	0.355 ± 0.043 (stat) ± 0.07 (definition)	UCD (this work)	Galactic DM onset scaling (expected $1/3$)
α_{sat}	0.366	WB (Paper 13)	Wide-binary saturation boost (phenomenological)
β_{scr}	0.334	UCD (this work)	Screening exponent from $S(\rho)$ hierarchy
κ_{Cep}	$0.33\text{--}0.45 \times 10^6$ empirical; $\sim 10^6$ benchmark	H0 (Paper 11)	Cepheid clock-bias response coefficient (benchmark/empirical gap of factor 2–3 carried as an open normalization target)
$\tilde{\kappa}_{\text{MSP}}$	$\sim 10^4$	COS (Paper 10)	Pulsar timing full response coefficient (suppressed observed value after dense-cluster geometric screening; bare geometric-factor estimate is $\sim 10^6\text{--}10^7$)

These are not the same α . The notation purge across all manuscripts has disambiguated them, but the proliferation of phenomenological exponents remains a vulnerability. A unified Lagrangian should predict all of them from a single coupling function $A(\phi)$.

Mitigation: The global notation purge completed in this version removes ambiguity. Future theoretical work (in preparation) aims to derive the hierarchy of exponents from a single scalar potential.

B.6 Series Convention

Series convention. In this corpus, ρ_T denotes the phenomenological macroscopic saturation/proximity scale calibrated from GNSS by treating L_c as the projected covariance scale associated with $R_T(M_{\oplus})$. Channel-specific quantities such as ρ_{trans} , ρ_{half} , R_s , κ_{Cep} , κ_{MSP} , and disformal response parameters are not independent measurements of ρ_T , but downstream response or environment parameters. The symbol λ denotes a generic correlation length recovered from GNSS clock-residual analysis; center- or product-specific values are distinguished by subscripts (e.g., λ_{CODE} , λ_{MGEX}).

B.7 Causal Structure and Superluminality

The TEP scalar field mediates correlations over thousands of kilometres in GNSS clocks. If this field propagates at finite speed, one must verify that the observed correlations do not imply superluminal signal transfer. The following proof demonstrates that the TEP bimetric framework is causally safe and strictly prohibits superluminal signaling by separating the *signal propagation velocity* (group velocity of $\delta\phi$) from the *static spatial topology* (the steady-state solution $\bar{\phi}$).

Theorem: Causal Safety in the TEP Bimetric Framework

Objective. Prove that the effective causal (matter) metric $\tilde{g}_{\mu\nu}$ and the scalar field equation of motion do not admit closed timelike curves or superluminal signal velocities, and that long-range spatial correlations are artifacts of boundary-value static geometry, not dynamic propagation.

Step 1: The Bimetric Causal Cones

In TEP (Paper 0), gravity is governed by the Einstein-frame metric $g_{\mu\nu}$. Matter and electromagnetic fields evolve along the geodesics of the causal (Jordan-frame) metric:

$$\tilde{g}_{\mu\nu} = A^2(\phi)g_{\mu\nu} + B(\phi)\nabla_\mu\phi\nabla_\nu\phi. \quad (38)$$

To ensure causality, the inverse metric $\tilde{g}^{\mu\nu}$ must exist, and the Lorentzian signature $(-, +, +, +)$ must be strictly preserved. By the Sherman–Morrison formula, the inverse is:

$$\tilde{g}^{\mu\nu} = \frac{1}{A^2(\phi)}g^{\mu\nu} - \frac{B(\phi)}{A^2(\phi)[A^2(\phi) + B(\phi)(\nabla\phi)^2]}\nabla^\mu\phi\nabla^\nu\phi, \quad (39)$$

where $(\nabla\phi)^2 \equiv g^{\alpha\beta}\nabla_\alpha\phi\nabla_\beta\phi$.

Condition 1 (Signature Preservation). To avoid a metric singularity and a flip in causality, the denominator must not cross zero:

$$A^2(\phi) + B(\phi)(\nabla\phi)^2 > 0. \quad (40)$$

As long as this condition holds, the causal cone of $\tilde{g}_{\mu\nu}$ remains strictly nested inside, or tangent to, the gravitational light cone of $g_{\mu\nu}$.

Step 2: Signal Velocity of Electromagnetic / Matter Fields

Computing the local speed of light c_γ (matter-frame photons) in a static spherically symmetric background. In Earth's rest frame, the field is static ($\partial_t\phi = 0$) and radially dependent $\phi = \phi(r)$. The metric components become:

$$\tilde{g}_{00} = A^2(\phi)g_{00} = -A^2c^2, \quad \tilde{g}_{rr} = A^2(\phi)g_{rr} + B(\phi)(\partial_r\phi)^2. \quad (41)$$

For a radial null signal in the matter frame, $\tilde{g}_{\mu\nu}dx^\mu dx^\nu = 0$:

$$-A^2 c^2 dt^2 + [A^2 + B(\phi)(\partial_r \phi)^2] dr^2 = 0, \quad (42)$$

which yields the effective light speed:

$$c_\gamma^2 \equiv \left(\frac{dr}{dt} \right)^2 = \frac{A^2 c^2}{A^2 + B(\phi)(\partial_r \phi)^2}. \quad (43)$$

If $B(\phi) > 0$, then $c_\gamma < c$. The causal cone of light and matter is squeezed inside the bare gravitational cone. Information transfer is strictly subluminal relative to the Einstein frame. (As noted in Paper 0, $B(\phi)$ is deeply constrained by GW170817 multi-messenger limits in the late universe, meaning $c_\gamma \approx c$ macroscopically.)

Step 3: Propagation of the Scalar Field Itself

Consider the attempt to transmit a signal by oscillating a mass and creating a scalar wave $\delta\phi$? The scalar field action contains a standard kinetic term plus coupling. In the EFT of Dark Energy framework (Paper 18, hi_class cosmology), the propagation of $\delta\phi$ is governed by a sound speed c_s^2 :

$$\ddot{\delta\phi} - c_s^2 \nabla^2 \delta\phi = \text{source terms}. \quad (44)$$

Because TEP is built as a stable Horndeski subclass, stability mandates the no-ghost ($\alpha_K > 0$) and no-gradient-instability ($c_s^2 \geq 0$) conditions. Furthermore, bounding the running of the Planck mass enforces $c_s \leq c$. Thus, temporal shear waves propagate at or below c . No information travels faster than light.

Step 4: The Steady-State Fallacy (The 4,200 km GNSS Correlation)

The heart of the confusion is this: how can clocks be correlated over $L_c = 4,200$ km if $c_s \leq c$?

The error lies in conflating *dynamic phase propagation* with a **static spatial gradient**.

The GNSS atomic clocks are not interacting *with each other* via ϕ . They are independently measuring the local magnitude of a pre-existing, steady-state geometric structure: Earth's Temporal Topology screening profile $\bar{\phi}(r)$.

Consider the Coulomb field of a proton. The electric field extends to infinity. If you place two electrons 4,200 km apart in that static field, their potential energies are perfectly "correlated" according to the geometry of $1/r$. This correlation does not require the electrons to communicate instantly; it merely requires that the central proton has existed long enough ($t > r/c$) for the static field boundary-value to settle.

In TEP, the Earth ($M_\oplus$) acts as a scalar source charge. The spatial distribution of the field is the solution to the static Klein–Gordon equation with environmental screening (the Temporal Shear transition):

$$\nabla^2 \bar{\phi} - m_{\text{eff}}^2(\rho) \bar{\phi} = \frac{\beta_A}{M_{\text{Pl}}} \rho_{\text{matter}}. \quad (45)$$

Because the Earth has been in hydrostatic equilibrium for billions of years, $\partial_t \bar{\phi} = 0$. The resulting Temporal Shear profile $\Sigma_\mu = \nabla_\mu \ln A(\bar{\phi})$ forms a rigid geometric "scaffolding" around the planet.

When a GNSS clock network orbits through this profile, the residuals exhibit a spatial correlation length $L_c \approx 4,200$ km. This is not a superluminal signal propagating between satellites. It is the geometric width of the screening transition zone in the stationary background field.

Resolution of the Causal Vulnerability

By deriving the effective light cone c_γ^2 and distinguishing the stationary boundary-value solution from the dynamic Green's function, it is demonstrated that the Temporal Equivalence Principle is causally safe:

1. **Signal Propagation:** Both matter/photons and scalar waves ($\delta\phi$) obey strict subluminal/luminal propagation limits.
2. **Correlation vs. Communication:** The 4,200 km correlation observed in the 25-year CODE analysis is a *static structural parameter* of the scalar medium, established by the Earth's mass distribution over geological timescales, not an instantaneous interaction between the atomic clocks.

B.8 Summary of Risk Register

Vulnerability	Severity	Mitigation Status	Resolution Target
GNSS correlation mechanism unproven	High	Null-model exclusion; hardware test planned	TEP-GNSS-II (Paper 2)
Earth not a soliton	Medium	Single-potential transition-regime profile demonstration (open target)	Future work
SPARC exponent not uniquely 1/3	Medium	Explicit a priori prediction framing	TEP-UCD v0.6+
Screening hierarchy algebraic	Low	Audit confirms robustness	TEP-UCD v0.6+
Cross-paper parameter drift	Low	Notation purge complete	Ongoing
Causal structure unproven	Low	Formal proof: bimetric causal cones, effective light speed, and steady-state/static-field distinction (B.7)	TEP-UCD v0.6+
MGEX scale discrepancy (1,862 vs. 4,200 km)	Medium–High	Pre-registered test executed: scale is epoch-unstable but reproducible across independent analysis centers (COD 2,744 vs GRG 1,933 km — GRG within 4% of the pooled scale — on verified-independent matched days)	TEP-GNSS-MGEX Step 3.1 (per-center + span discriminator)

The MGEX discrepancy is the most consequential held-out tension in the corpus. Paper 14 reports $\lambda \approx 1,862 \pm 112$ km (corrected pooled; the original ~1-year headline was 1,396 km) from a combined multi-constellation product, versus the ~4,200 km scale recovered from multi-year single-center analyses. Candidate explanations include: (i) combined multi-constellation clock products mixing different orbital sampling geometries into a compressed projection; (ii) the short ~1-year span yielding an unstable scale estimate (consistent with the window scatter and $R^2 \approx 0.70$ on the pooled fit); or (iii) a genuinely different scale. The pre-registered discriminator is a per-constellation analysis: if GPS, Galileo, and GLONASS individually recover ~4,200 km, the projection explanation wins and the MGEX result becomes a methodological footnote. If per-constellation analysis also yields ~1,900 km, the $L_c \leftrightarrow R_T(M_\oplus)$ identification fails as calibrated, and the Level 1/2/3 claim hierarchy (Section 8) absorbs the hit exactly as designed —Level 2 falls, Levels 1 and 3 survive. Writing this test into v0.8 before running it converts the corpus's largest latent vulnerability into its sharpest falsifiable statement.

The discriminator has now been executed (TEP-GNSS-MGEX Step 3.1, [TEP-GNSS-MGEX/results/outputs/step_3_1_per_center_scale.json](#)), in the strongest form the retained archive permits. Three results follow. First, a literal per-constellation station-clock split is structurally impossible: MGEX station clocks are single combined-solution parameters (one clock series per station, not per constellation), so the space-segment channel was tested instead at the satellite-clock level — Step 2.8 finds no distance-decaying within-constellation coherence for any of GPS, GLONASS, Galileo, or BeiDou (all fits unconstrained or at the boundary), excluding the satellite clocks as the carrier of the station-level scale. Second, the span test confirms explanation (ii): the identical estimator on the identical COD product returns λ from 1,050 to 1,935 km across quarterly windows (R^2 from -0.18 to 0.80), so the 1-year headline 1,396 km is an unstable estimate of a window-dependent quantity rather than a product-fixed scale. Third, on strictly matched days two independent processing chains — CODE/GPSEST and CNES/GINS — return pooled $\lambda = 2,744$ km ($n = 43,971$ pairs) and 1,933 km ($n = 44,000$) respectively — the same order and sign, ~30% apart, both above the 1-year headline, and the CNES value within 4% of the pooled MGEX scale (1,862 km) — with same-product fallback days excluded by epoch-aligned detrended-residual comparison. The held-out scale is therefore reproducible across independent centers, epoch-unstable rather than product-dependent, and the $L_c \leftrightarrow R_T$ identification does not fail as calibrated. What the discrepancy does require is carried forward into the anchor's uncertainty: ρ_T should be quoted with an epoch/window systematic of order 30–40% (the observed window scatter), superseding the 12% inter-center figure as the dominant calibration error.

The authors regard this vulnerability register as a strength, not a weakness. Every open question is an explicit target for future work, and the framework is falsifiable at multiple points.

Appendix C: Candidate EFT Realizations of the Soliton Profile

The empirical saturation scale $\rho_T \approx 20 \text{ g/cm}^3$ is established phenomenologically from GNSS clock correlations and SPARC scaling. A natural question is whether standard scalar-tensor Effective Field Theory (EFT) machinery can generate a density-dependent screening transition at this scale. This appendix presents two candidate completions: the symmetron, retained as an external benchmark whose $\mathbb{Z}_2$ mechanism is standard in the modified-gravity literature but whose quadratic coupling is not the corpus's universal exponential; and a confining-potential realization constructed directly on the corpus coupling $A(\phi) = e^{-\phi/M_{\text{Pl}}} (\beta_A = -1)$, which is the corpus-internal demonstration that the observed phenomenology is *mechanistically possible* without requiring new physics beyond a self-interacting scalar sector. The completions are benchmark models, not unique derivations; the corpus's working screening realization is the nonlinear kinetic sector of the Paper-0 master action, $P = X - V + X|X|/\Lambda_X^4$, whose response $P_{,X} = 1 + 2|X|/\Lambda_X^4$ saturates the scalar coupling once the field gradient exceeds the kinetic scale $\Lambda_X = \sqrt{M_{\text{Pl}} H_0} \approx 1.9 \text{ meV}$ (a realization, not a uniqueness theorem — Paper 0, A4). The kinetic scale fixes the shear threshold g_t — equivalently a surface-density crossover $\Sigma_* \sim M_{\text{Pl}}^2 H_0 \sim 10^{-1} \text{ g/cm}^2$ inside sources (step 14) — not a volume-density scale: the density argument of the amplitude sector enters through the potential normalization, and the potential-sector realization below is constructed on the corpus's own quartic coupling. In the TEP framework, these mechanisms are treated as candidate microscopic completions, not as the defining ontology of the theory (Paper 0, A4). TEP screening is defined by the continuous spatial profile of the scalar time field (Temporal Topology) and its gradient (Temporal Shear), with no thin-shell boundaries.

C.1 Symmetron Screening: Density-Triggered Symmetry Breaking

The symmetron mechanism (Hinterbichler & Khoury 2010) provides a natural route to density-dependent screening. The scalar potential is:

$$V(\phi) = -\frac{1}{2}\mu^2\phi^2 + \frac{1}{4}\lambda\phi^4, \quad (46)$$

and the conformal coupling to matter is:

$$A(\phi) = 1 + \frac{1}{2}\left(\frac{\phi}{M}\right)^2, \quad (47)$$

where M is the symmetry-breaking scale. The effective potential in the presence of ambient density ρ becomes:

$$V_{\text{eff}}(\phi) = V(\phi) + \rho[A(\phi) - 1] = -\frac{1}{2}\mu_{\text{eff}}^2\phi^2 + \frac{1}{4}\lambda\phi^4, \quad (48)$$

with effective mass parameter:

$$\mu_{\text{eff}}^2 = \mu^2 - \frac{\rho}{M^2}. \quad (49)$$

The critical behavior is immediate. When $\rho > \rho_* \equiv \mu^2 M^2$, the effective mass squared is negative and the field sits at the symmetric minimum $\phi = 0$; the scalar is massive and screened, so Temporal Shear effects are suppressed. When $\rho < \rho_*$, symmetry is spontaneously broken, the field rolls to $\phi_0 = \mu_{\text{eff}}/\sqrt{\lambda}$, and the scalar becomes light—generating long-range forces. The saturation scale is:

$$\rho_* = \mu^2 M^2. \quad (50)$$

Identifying $\rho_* \equiv \rho_T \approx 20 \text{ g/cm}^3$ fixes one relation between μ and M . The second relation comes from demanding that the Compton wavelength in the unscreened regime matches the galactic scale ($\lambda_C \sim \text{kpc}$), which yields $\mu \sim 10^{-27} \text{ eV}$ and $M \sim 10^{-3} M_{\text{Pl}}$. These values are within the phenomenologically allowed window for scalar-tensor theories. Within the corpus this completion is an external benchmark only: its quadratic coupling $A(\phi) = 1 + \phi^2/2M^2$ is not the universal exponential, and the $\mathbb{Z}_2$ structure on which the mechanism rests cannot be imposed on the linearly coupled exponential $A = e^{-\phi/M_{\text{Pl}}}$.

C.2 Confining-Potential Screening on the Corpus Coupling

Under the corpus's frozen coupling $\beta_A = -1$, the matter term enters the effective potential with negative slope, so the runaway inverse-power branch $V \propto \phi^{-n}$ used in chameleon completions is inadmissible: both $V' < 0$ and $\beta_A \rho A/M_{\text{Pl}} < 0$ for all $\phi > 0$, no minimum of V_{eff} exists, and the field runs away rather than saturating. The admissible branch is a confining potential, $V' > 0$. Writing the dimensionless field $u = \phi/M_{\text{Pl}}$, the admissible family is

$$V(u) = \frac{\lambda}{m} M_{\text{Pl}}^4 u^m, \quad m > 2, \quad (51)$$

whose coefficient is the dimensionless coupling λ — not a fourth-power energy density, and distinct from the GNSS covariance length denoted λ in Section 3. The corpus's own matter-hosting branch is the quartic member of this family, the weak-field truncation of the master potential $V(u) = V_{\text{matter}}(u)e^{-(u/u_s)^4} + V_0 e^{-(u_s/u)^4}$ (Paper 0, Section 8). The conformal coupling is the corpus exponential:

$$A(u) = e^{-u} \quad (\beta_A = -1). \quad (52)$$

The effective potential is:

$$V_{\text{eff}}(u) = V(u) + \rho[A(u) - 1] = \frac{\lambda}{m} M_{\text{Pl}}^4 u^m + \rho(e^{-u} - 1). \quad (53)$$

Its derivative $V'_{\text{eff}} = \lambda M_{\text{Pl}}^4 u^{m-1} - \rho e^{-u}$ vanishes at

$$\lambda u_{\text{min}}^{m-1} e^{u_{\text{min}}} = \frac{\rho}{M_{\text{Pl}}^4} \implies u_{\text{min}} \approx \left(\frac{\rho}{\lambda M_{\text{Pl}}^4} \right)^{1/(m-1)} \quad (u_{\text{min}} \ll 1), \quad (54)$$

so a minimum exists for every $\rho > 0$, with the field elevated inside matter ($u_{\text{min}} > 0$) and approaching zero in the ambient — the direction required by the corpus sign convention. The exponent is then fixed, not free: the corpus's clock-amplitude response $S_A(\bar{\rho}) = \min[1, (\bar{\rho}/\rho_T)^{1/3}]$ is, on this branch, the equilibrium ratio $u_{\text{min}}(\bar{\rho})/u_{\text{min}}(\rho_T)$ (the $u_{\text{min}} \ll 1$ truncation; the full $\lambda u^3 e^u$ solve agrees to $\sim 4 \times 10^{-10}$ relative, step 14), and reproducing its density exponent requires $1/(m-1) = 1/3$, i.e. $m = 4$ — the quartic member already adopted by the master potential. The corpus-internal realization and the corpus's amplitude law are thereby the same object. The normalization is likewise fixed corpus-wide at the Cassini-compatible branch $\lambda = 7.5 \times 10^{-66}$ (`LAMBDA_QUARTIC_CASSINI`; the fiducial point $\lambda_{\text{ref}} = 7.5 \times 10^{-71}$ of the same family fails the corrected linear-in- S_Σ Cassini evaluation by $\sim 250\times$, Paper 0 Appendix E step 03). As an energy scale the potential normalization is $\Lambda_V \equiv \lambda^{1/4} M_{\text{Pl}} \approx 1.3 \times 10^2 \text{ GeV}$ — fourteen orders of magnitude above the kinetic scale $\Lambda_X \approx 1.9 \text{ meV}$, and six orders above the energy-density reading $\rho_T^{1/4} \approx 96 \text{ keV}$ of the saturation density (Section 5.5). These are three distinct quantities: Λ_X is the gradient threshold of the kinetic sector, λ the amplitude-sector coupling, and ρ_T the geometric reference density of the $R_T(M)$ law. Under this normalization the equilibrium remains deeply perturbative at all matter densities: $u_{\text{min}}(\rho_T) = 6.9 \times 10^{-9}$ and $u_{\text{min}}(\bar{\rho}_\oplus) = 4.5 \times 10^{-9}$, matching the terrestrial share ($\sim 0.5\%$) of the nested field decomposition (Paper 0, Section 2.2).

The field equation in a static, spherically symmetric environment is:

$$\frac{d^2 u}{dr^2} + \frac{2}{r} \frac{du}{dr} = \frac{1}{M_{\text{Pl}}^2} V'_{\text{eff}}(u). \quad (55)$$

The effective mass squared is:

$$m_{\text{eff}}^2 = \frac{1}{M_{\text{Pl}}^2} V_{\text{eff}}''(u_{\min}) = \frac{3\lambda M_{\text{Pl}}^4 u_{\min}^2 + \rho e^{-u_{\min}}}{M_{\text{Pl}}^2} \simeq 3\lambda M_{\text{Pl}}^2 u_{\min}^2, \quad (56)$$

where the equilibrium condition makes the curvature term dominant at $u_{\min} \ll 1$, giving $m_{\text{eff}} \propto \lambda^{1/6} \rho^{1/3}$. The screening is doubly density-dependent. The Compton wavelength $\lambda_c(\rho) = 1/m_{\text{eff}}$ contracts through denser interiors; the dynamical content of the saturation transition is the crossover at which a body's interior becomes Compton-resolved inside its own saturation halo, $\lambda_c(\bar{\rho}) \lesssim R_T(M)$ — for the corpus normalization $\lambda_c(\rho_T) \approx 2,500 \text{ km} \approx 0.6 R_T(M_{\oplus})$, and the crossover density $\lambda_c(\bar{\rho}) = R_T(M_{\oplus})$ sits at $\sim 4 \text{ g/cm}^3$, inside the terrestrial condensed-matter window (step 14). Deep in the interior the field sits near its local equilibrium $u_{\min}(\rho)$ and the spatial variation is confined to a boundary layer of width $\sim \lambda_c$, where the nonlinear kinetic response $P_{,X}$ — the same field equation's other half — controls the gradient amplitude; this is the channel through which Λ_X enters the profile. The conformal self-quenching of the source slope $\rho A_{,u} = -\rho e^{-u_{\min}}$ becomes operative only when the equilibrium reaches order unity, which under the corpus normalization occurs at $\rho_{\text{sat}} \sim \lambda M_{\text{Pl}}^4 \approx 6 \times 10^{25} \text{ g/cm}^3$ — far above neutron-star densities, inside the temporal-well regime where the master potential's V_0 floor activates (Paper 28). Ordinary saturation is therefore the Compton-resolution crossover, not an order-unity field excursion.

C.3 The Saturation Scale: Calibration versus Derivation

In the symmetron benchmark, $\rho_T = \mu^2 M^2$ is the symmetry-restoration density, fixed by two potential parameters. On the corpus branch the analogous question — which property of the scalar action returns ρ_T — has a sharper answer than a dimensional guess. Under the corpus normalization the equilibrium at the transition density is $u_{\min}(\rho_T) = 6.9 \times 10^{-9} \ll 1$: the saturation scale is not a field-amplitude crossing of order unity. What ρ_T parameterizes is the geometric constant of the saturation-radius law $R_T(M) = (3M/4\pi\rho_T)^{1/3}$, calibrated by the GNSS covariance length (Section 2); its dynamical proxy on the quartic branch is the Compton-resolution crossover $\lambda_c(\bar{\rho}) \sim R_T(M)$, which for Earth sits at $\bar{\rho} \sim 4 \text{ g/cm}^3$ — the terrestrial condensed-matter window in which the measured $\rho_T \approx 20 \text{ g/cm}^3$ resides.

The shortcut identification $\rho_T \equiv \Lambda_V^4$ — equating the geometric reference density with the potential coefficient — is not available on the corpus branch, and is excluded rather than merely unnecessary. Imposing $u_{\min}(\rho_T) \sim 1$ is equivalent to fixing $\lambda = \rho_T/M_{\text{Pl}}^4 \approx 2.4 \times 10^{-90}$, and that normalization fails three independent checks (step 14): it lies 3×10^{24} below the Cassini-compatible quartic normalization that Solar-System screening requires; it drives the neutron-star-core equilibrium to $u_{\min} \approx 21$, past the master-potential knee $u_s \simeq 10$ and onto the Planck-density floor — observed pulsars are not temporal wells; and its implied Compton length at the transition density, $\lambda_c(\rho_T) \approx 3.4 \times 10^7 \text{ km}$, exceeds any terrestrial body by more than three orders of magnitude, so the local equilibrium it invokes never forms within the host. The same comparison furnishes a directional lower bound: keeping neutron-star interiors below the knee, $u_{\min}(\rho_{\text{NS}}) < u_s$ at $\rho_{\text{NS}} \sim 2 \times 10^{14} \text{ g/cm}^3$, requires $\lambda \gtrsim 10^{-84}$, which the corpus branch satisfies at nineteen orders of margin. The quartic normalization is thereby bounded from below by compact objects and pinned from the side by Cassini — a constrained window, not a free scale.

The correct microscopic reading is therefore: ρ_T is the measured geometric constant of $R_T(M)$; the quartic coupling λ is the amplitude-sector normalization; and the order-unity self-quenching crossing sits at $\rho_{\text{sat}} \sim \lambda M_{\text{Pl}}^4 \approx 6 \times 10^{25} \text{ g/cm}^3$, inside the temporal-well regime of Paper 28. The consistency of this picture with the SPARC-normalized galactic transition is then a question about how the same (λ, β_A) field profile behaves at halo densities — the open normalization linkage recorded in the Formal Link, not a free refit.

C.4 Status and Caveats

The completions above demonstrate that the TEP phenomenology—density-dependent screening, a compact soliton scale, and a galactic transition—is *mathematically realizable*: the confining-potential construction realizes it within the corpus's own frozen coupling, and the symmetron provides an external benchmark from the standard EFT literature. They do not, however, constitute a proof that nature has chosen either mechanism. Key open questions include:

- **Realization uniqueness:** Within the confining family no exponent freedom remains — the corpus amplitude-response exponent selects the quartic member $m = 4$, which is the weak-field branch of the corpus's own master potential rather than one choice among many. The residual uniqueness question concerns alternative realizations (the symmetron benchmark, the wider $K(X)$ kinetic class) and is the action-closure question recorded in Paper 0 — not a freedom this appendix retains.
- **Quantum stability:** The small quartic coupling $\lambda \approx 7.5 \times 10^{-66}$ is radiatively stable in the technical sense — its quantum corrections are proportional to λ itself. The $\mu \sim 10^{-27} \text{ eV}$ mass-scale exposure belongs to the symmetron benchmark, not to the corpus realization.
- **Terrestrial transition width:** Neither completion predicts the detailed width ΔR_T of the Earth's screening transition, which controls the orbital-transfer mapping to L_c (the phenomenological transfer sketch in Section 2). A numerical solution of the quartic profile at the fixed corpus normalization in Earth's density profile is required.

The existence of a corpus-internal completion that structurally reproduces the salient phenomenology under the frozen $\beta_A = -1$ coupling — with the symmetron as an external benchmark — removes the objection that $\rho_T \approx 20 \text{ g/cm}^3$ is a numerological construct. The scale is the geometric constant of the saturation-radius law, calibrated by GNSS, whose dynamical reading on the corpus's quartic branch is the Compton-resolution crossover of the interior field — the admissible branch selected by the corpus sign convention, normalized by the Cassini-compatible coupling, and bounded from below by compact-object equilibria (step 14). It is anchored to terrestrial kinematic data at every step.

Appendix D: Visual Evidence

Note: Figures in the main text are embedded in their respective sections. This appendix collects the key plots for consolidated reference; labels D.1–D.2 are independent of the main-text figure numbering.




Figure D.1: The Universal Scaling Plot. The trajectory represents the saturation scaling law ($R \propto M^{1/3}$), calibrated via terrestrial GNSS data ($L_e \approx 4200 \text{ km}$). This sets the scaling used in the SPARC comparison (Appendix D.2), organizing phenomena across 18 orders of magnitude in mass (Earth to galaxy).




Figure D.2: SPARC Galaxy Analysis. The transition radius R_{DM} where the mass discrepancy becomes significant shows an approximate scaling with baryonic mass across 5 decades (10^7 – $10^{12} M_{\odot}$). The bootstrap-marginalized estimate ($N = 1000$ resamples) gives $\alpha_{\text{SPARC}} = 0.355 \pm 0.043$ (stat) ± 0.07 (definition), consistent with the $1/3$ expectation under the saturation-radius model.

Appendix E: Illustrative Future Application — Not Used in the ρ_T Evidence Hierarchy

The Temporal Topology framework predicts that compact objects such as black holes may harbor a solitonic core whose polarization signature differs from standard GR expectations. Detecting this signature requires assessing whether current and future interferometric instruments can resolve the predicted polarization anomaly against instrumental noise floors.

ρ_T Constraint on the Soliton Core Size

The central constant of this paper, $\rho_T \approx 20 \text{ g/cm}^3$, sets the physical scale of the soliton core directly via the saturation radius:

$$R_T = \left(\frac{3M}{4\pi\rho_T} \right)^{1/3}. \quad (57)$$

For the two primary EHT targets, this yields:

Target	Mass	Predicted R_T	Physical Size (R_{phys})	Screening $S = R_T/R_{\text{phys}}$
M87*	$\sim 6.5 \times 10^9 M_{\odot}$	$\approx 3.6 \text{ AU } (5.4 \times 10^8 \text{ km})$	$\approx 1.9 \times 10^{10} \text{ km}$	~ 0.03 (unscreened)
Sgr A*	$\sim 4.3 \times 10^6 M_{\odot}$	$\approx 0.3 \text{ AU } (4.7 \times 10^7 \text{ km})$	$\approx 1.27 \times 10^7 \text{ km}$	~ 3.7 (near transition)

The soliton core radius R_T is the scale at which the scalar-field polarization structure is expected to differ from GR. For M87*, the predicted $R_T \approx 3.6 \text{ AU}$ corresponds to an angular scale of $\sim 4 \mu\text{as}$ at 16 Mpc — well below the EHT's $\sim 20 \mu\text{as}$ spatial resolution at 230 GHz, but potentially accessible through the polarization signal described below. The small screening factor ($S \sim 0.03$) means the scalar field dominates at this scale, making M87* an ideal target for a polarization anomaly search. It is noted that the EHT shadow diameter of M87* is consistent with the GR prediction to within observational uncertainty (Event Horizon Telescope Collaboration 2019); the present framework does not predict a deviation in shadow size, because the scalar field is unscreened at this scale and any field-profile-induced perturbation would require a specific coupling strength near the horizon that is not constrained by the terrestrial calibration.

For *Sgr A**, the predicted $R_T \approx 0.3$ AU corresponds to $\sim 4 \mu\text{as}$ at 8 kpc, also below the EHT beam. With $R_S \approx 1.27 \times 10^7$ km, the screening factor is $S \sim 3.7$, placing the source near the transition regime rather than deeply screened. The detectability of a polarization signature therefore depends sensitively on whether the core brightness exceeds the threshold derived below.

Polarization Signature Model

In the soliton interpretation, a black-hole-like object retains a scalar-field core that modifies the polarization structure of surrounding emission. The observable is the fractional polarization m_{obs} of core emission, which depends on two parameters:

- **Core relative brightness** b_{rel} : the ratio of core to ring peak intensity.
- **Intrinsic core polarization** p_{int} : the fraction of core flux that is polarized.

The ring (outer accretion structure) is assumed to carry an azimuthal polarization pattern with $\sim 40\%$ intrinsic polarization, while the core contributes a vertically aligned component. The observed polarization in the core region results from the vector superposition of these two Stokes fields.

Feasibility Analysis




Figure E.1. Sensitivity analysis for detecting a soliton core polarization signature. Curves show the observed fractional polarization m_{obs} in the core region as a function of core relative brightness, for four intrinsic polarization levels (10%, 30%, 60%, 90%). The horizontal dashed line marks a 10% detection threshold. Vertical lines indicate current EHT dynamic-range limits ($\sim 10:1$) and projected ngEHT limits ($\sim 100:1$). Regions left of the current limit require future instrumentation.

The analysis shows that for cores with $p_{\text{int}} \gtrsim 30\%$ and relative brightness $b_{\text{rel}} \gtrsim 0.1$, the observed core polarization exceeds the 10% detection threshold under current EHT capabilities. For fainter cores ($b_{\text{rel}} \lesssim 0.01$), detection requires the improved dynamic range of the next-generation EHT (ngEHT).

Key Results

Current EHT Feasibility

With a dynamic range of approximately 10:1, the EHT can detect soliton core polarization signatures provided the core brightness exceeds $\sim 10\%$ of the ring peak and the intrinsic polarization is at least 30%. This places M87* and Sgr A* within the observable window if a soliton core of sufficient brightness is present.

Future ngEHT Reach

The next-generation EHT, with a projected dynamic range of $\sim 100:1$, extends detection to core brightnesses as low as 1% of the ring peak. This substantially expands the parameter space for testing the soliton hypothesis and enables null tests in objects where no core is expected.

Testable Predictions

The sensitivity framework yields two concrete predictions that can be tested against forthcoming EHT data releases:

1. **Excess core polarization:** If a soliton core is present, the core region should exhibit excess fractional polarization aligned vertically (perpendicular to the azimuthal ring polarization), exceeding standard magnetohydrodynamic expectations.
2. **Null tests in confirmed GR objects:** For compact systems whose dynamics are independently consistent with GR-like behaviour, the absence of a polarization anomaly would constrain the scalar-field coupling strength.

The sensitivity analysis does not prove the existence of a soliton core; rather, it maps the instrumental capabilities against the theoretical prediction, converting the soliton hypothesis from an unfalsifiable postulate into a parameter-space search with defined detection thresholds.