

After Visit Summary

Long-acting injection

STAFF PREVIEW - NOT FINAL

PATIENT	DOB	RECORD NO
Print, Vivitrol	02/12/1977	WQ7CEO
PROVIDER	VISIT DATE	GIVEN BY
Print Ordering Provider	08/14/2026	Print QA, MA

Important: opioid tolerance and overdose risk

Vivitrol blocks the effect of opioids. Your body's tolerance to opioids is now much LOWER than it was before you started.

You are most at risk near the end of the month as the block wears off, if you miss a dose, and after you stop Vivitrol altogether. At those times the amount of an opioid you used to take can cause an overdose or death.

Do not try to override the block by taking a large amount of an opioid. Keep naloxone (Narcan) on hand and make sure the people around you know how to use it.

Call before you come in - (909) 887-6222

Vivitrol is kept refrigerated and has to be taken out and brought to room temperature before it can be given. Please call ahead so your dose is ready when you arrive. If you walk in without calling, expect a wait.

Your treatment today

Aug 14
TODAY

Vivitrol, 380 mg

naltrexone ER

Right hip (ventrogluteal), intramuscular

Lot VIV-PRINT-001, exp 07/2028



Your next injection and provider appointment are on page 2. Please read all pages of this handout.

Clinic information	APPOINTMENTS & QUESTIONS (909) 887-6222 - call before coming in	CLINIC HOURS Monday-Friday, 8:30 AM - 5:00 PM (appointments preferred)	AFTER HOURS For concerns that cannot wait, use urgent care. Call 911 for emergencies.
--------------------	--	---	--

After Visit Summary - Continued

PATIENT
Print, Vivitrol

DOB
02/12/1977

RECORD NO.
WQ7CE0

VISIT DATE
08/14/2026

Your follow-up visits

Your next injection

Friday, September 11, 2026

Due date - call us to schedule or reschedule

With clinic confirmation, you may come in up to 3 days before or 3 days after your due date. Call ahead so we can prepare your medication; an injection due date does not reserve an appointment time.

Your next provider appointment

Date **Tuesday, November 3, 2026**

Time _____ AM / PM

With **Synthetic Appointment Provider for Comprehensive
Follow-up and Medication Management, PMHNP-BC**

Connect _____

Video visit

Please have the front desk confirm these details before you leave.

Why timing matters

The opioid-blocking effect fades toward the end of the month. A late injection leaves a gap where you are no longer protected but your tolerance is still low.

That gap is the highest-risk time. Coming in on your due date closes it.

Caring for your injection site

Soreness or a firm lump at the hip is normal for a few days, and it may ache when you sit or walk at first. A cool pack helps.

Do not rub or massage the site.

Gentle walking is fine and usually helps the ache settle faster than sitting still.

Your next injection goes in the other side, so the same muscle is not used twice in a row.

Injection-site reactions to this medication can become serious. Look at the site once a day for the first two weeks.

Clinic information

APPOINTMENTS &
QUESTIONS
**(909) 887-6222 - call
before coming in**

CLINIC HOURS
**Monday-Friday, 8:30 AM - 5:00 PM
(appointments preferred)**

AFTER HOURS
**For concerns that cannot wait, use urgent
care. Call 911 for emergencies.**

After Visit Summary - Continued

PATIENT	DOB	RECORD NO.	VISIT DATE
Print, Vivitrol	02/12/1977	WQ7CEO	08/14/2026

What you may notice

- Nausea, tiredness, or headache can happen in the first few days and usually settles.
- Carry your naloxone (Narcan) if you have it, and make sure someone close to you knows where it is and how to use it.
- Tell any doctor, dentist, or emergency crew treating you that you are on Vivitrol - it changes which pain medications will work.

Call the clinic at (909) 887-6222 if

- The injection site develops intense pain, a hard lump that keeps growing, large swelling, blisters, an open sore, or a dark scab.
- The injection site is not getting better after two weeks.
- You have nausea, vomiting, stomach pain, unusual tiredness, dark urine, or your eyes or skin look yellow.
- You feel unusually down, or have thoughts of hurting yourself.
- You cannot come within the clinic-confirmed 3-day window, or need to change your visit.
- Anything feels unusual or worries you.

Call 911 now or go to the nearest emergency room if

- You suspect an opioid overdose in yourself or someone else - give naloxone and call 911.
- Trouble breathing, chest tightness, or severe dizziness.
- Swelling of the face, lips, tongue, or throat.

Clinic information	APPOINTMENTS & QUESTIONS (909) 887-6222 - call before coming in	CLINIC HOURS Monday-Friday, 8:30 AM - 5:00 PM (appointments preferred)	AFTER HOURS For concerns that cannot wait, use urgent care. Call 911 for emergencies.
---------------------------	---	---	---