

FEM for 2D problems

Robin boundary conditions

Variational problem (heat 2D): Find $\theta \in V_0$ such that

$$\int_{\Omega} \lambda \cdot \nabla \theta \cdot \nabla \delta \theta \, dA + \int_{\Gamma_R} h \cdot \theta \cdot \delta \theta \, ds = \int_{\Omega} w \cdot \delta \theta \, dA + \int_{\Gamma_R} h \cdot \theta^* \cdot \delta \theta \, ds$$

for all test functions $\delta \theta \in V_0$. The space (set) V_0 contains all "nice" functions $u: \Omega \rightarrow \mathbb{R}$ with $u(x, y) = 0$ for $(x, y) \in \Gamma_D$.

We define

$$a_{\text{int}}(\theta, \delta \theta) = \int_{\Omega} \lambda \cdot \nabla \theta \cdot \nabla \delta \theta \, dA$$

$$a_R(\theta, \delta \theta) = \int_{\Gamma_R} h \cdot \theta \cdot \delta \theta \, ds$$

$$a(\theta, \delta \theta) = a_{\text{int}}(\theta, \delta \theta) + a_R(\theta, \delta \theta)$$

$$b_{\text{int}}(\delta \theta) = \int_{\Omega} w \cdot \delta \theta \, dA$$

$$b_R(\delta \theta) = \int_{\Gamma_R} h \cdot \theta^* \cdot \delta \theta \, ds$$

$$b(\delta \theta) = b_{\text{int}}(\delta \theta) + b_R(\delta \theta)$$




Linear System with Robin BCs

$$\left(\int_0^L \nabla \cdot \nabla ds = \frac{1}{3} \cdot L, \int_0^L \nabla \triangle ds = \frac{1}{6} \cdot L \right)$$


$$\underline{K} = \underline{K}^{int} + \underline{K}^R \quad \text{where } k_{ij}^{int} = a_{int}(\varphi_i, \varphi_j) \quad \text{and } k_{ij}^R = a_R(\varphi_i, \varphi_j) = \int_{\Gamma_R} h \cdot \varphi_i \cdot \varphi_j ds$$

$$\underline{r} = \underline{r}^{int} + \underline{r}^R \quad \text{where } r_i^{int} = b_{int}(\varphi_i) \quad \text{and } r_i^R = b_R(\varphi_i) = \int_{\Gamma_R} h \cdot \theta^* \cdot \varphi_i ds$$

Example



On edge e



$$K_{66}^R = \int_0^{l^e} h \cdot \left(1 - \frac{s}{l^e}\right) \cdot \left(1 - \frac{s}{l^e}\right) ds = \frac{1}{3} \cdot h \cdot l^e$$


$$K_{69}^R = \int_0^{l^e} h \cdot \left(1 - \frac{s}{l^e}\right) \cdot \frac{s}{l^e} ds = \frac{1}{6} \cdot h \cdot l^e$$

$$\Gamma_6^R = \int_0^{l^e} h \cdot \theta^* \cdot \left(1 - \frac{s}{l^e}\right) ds = \frac{1}{2} \cdot h \cdot \theta^* \cdot l^e$$

$$r_9^R = \dots = \frac{1}{2} h \cdot \theta^* \cdot l^e$$

Edge wise computation

Functions on edge e




Contributions of edge e

$$k_{R,ij}^e = a_R^e(N_i^e, N_j^e)$$

$$r_{R,i}^e = b_R^e(N_i^e)$$

} Assembly into global system as before

→ See pile foundation



Edge forms

$$a_R^e(\theta, \delta\theta) = \int_0^{l^e} h \cdot \theta \cdot \delta\theta ds$$

$$b_R^e(\delta\theta) = \int_0^{l^e} h \cdot \theta^* \cdot \delta\theta ds$$

Where $\theta, \delta\theta : [0, l^e] \rightarrow \mathbb{R}$

Integraltafel

Nr.		A	B	C	D	E	F	G
		k		k_1	k	k		$\int j^2 dx$
1	j	jk	$\frac{1}{2}jk$	$\frac{1}{2}j(k_1 + k_2)$	0	$\frac{1}{4}jk$	$\frac{1}{2}jk$	j^2
2		$\frac{1}{2}jk$	$\frac{1}{3}jk$	$\frac{1}{6}j(k_1 + 2k_2)$	$-\frac{1}{6}jk$	0	$\frac{1}{6}jk(1 + \alpha)$	$\frac{1}{3}j^2$
3		$\frac{1}{2}jk$	$\frac{1}{6}jk$	$\frac{1}{6}j(2k_1 + k_2)$	$\frac{1}{6}jk$	$\frac{1}{4}jk$	$\frac{1}{6}jk(1 + \beta)$	$\frac{1}{3}j^2$
4	j_1	$\frac{1}{2}k(j_1 + j_2)$	$\frac{1}{6}k(j_1 + 2j_2)$	$\frac{1}{6}[j_1(2k_1 + k_2) + j_2(k_1 + 2k_2)]$	$\frac{1}{6}k(j_1 - j_2)$	$\frac{1}{4}j_1k$	$\frac{1}{6}k[j_1(1 + \beta) + j_2(1 + \alpha)]$	$\frac{1}{3}(j_1^2 + j_1j_2 + j_2^2)$
5	j	0	$-\frac{1}{6}jk$	$\frac{1}{6}j(k_1 - k_2)$	$\frac{1}{3}jk$	$\frac{1}{4}jk$	$\frac{1}{6}jk(1 - 2\alpha)$	$\frac{1}{3}j^2$
6	j	$\frac{1}{4}jk$	0	$\frac{1}{4}jk_1$	$\frac{1}{4}jk$	$\frac{1}{4}jk$	$\frac{1}{4}jk\beta$	$\frac{1}{4}j^2$
7	$-\frac{j}{2}$	$\frac{1}{4}jk$	$\frac{1}{4}jk$	$\frac{1}{4}jk_2$	$-\frac{1}{4}jk$	$-\frac{1}{8}jk$	$\frac{1}{4}jk\alpha$	$\frac{1}{4}j^2$
8		$\frac{1}{2}jk$	$\frac{1}{4}jk$	$\frac{1}{4}j(k_1 + k_2)$	0	$\frac{1}{8}jk$	$\frac{jk}{12\beta}(3 - 4\alpha^2)$	$\frac{1}{3}j^2$
9		$\frac{1}{2}jk$	$\frac{1}{6}jk(1 + \gamma)$	$\frac{1}{6}j[k_1(1 + \delta) + k_2(1 + \gamma)]$	$\frac{1}{6}jk(1 - 2\gamma)$	$\frac{1}{4}jk\delta$	$\frac{jk}{6\beta\gamma}(2\gamma - \gamma^2 - \alpha^2)$ $\gamma \geq \alpha$	$\frac{1}{3}j^2$
10		$\frac{2}{3}jk$	$\frac{1}{3}jk$	$\frac{1}{3}j(k_1 + k_2)$	0	$\frac{1}{6}jk$	$\frac{1}{3}jk(1 + \alpha\beta)$	$\frac{8}{15}j^2$
11		$\frac{1}{3}jk$	$\frac{1}{6}jk$	$\frac{1}{6}j(k_1 + k_2)$	0	$\frac{1}{12}jk$	$\frac{1}{6}jk(1 - 2\alpha\beta)$	$\frac{1}{5}j^2$
12		$\frac{2}{3}jk$	$\frac{1}{4}jk$	$\frac{1}{12}j(5k_1 + 3k_2)$	$\frac{1}{6}jk$	$\frac{7}{24}jk$	$\frac{1}{12}jk(5 - \alpha - \alpha^2)$	$\frac{8}{15}j^2$