

W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere^{*1}, Daniel Argüeso², Andrea Zonato³, and Jonas Kittner¹

¹ Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany ² Physics Department, University of the Balearic Islands, Palma, Spain ³ Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Editor Name](#) ↗

Submitted: 01 January XXXX

Published: 01 January XXXX

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The Python-based WUDAPT-to-WRF (W2W) package is developed to translate Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model (Skamarock et al., 2021). It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides an improved, more simple, and more efficient procedure to use LCZ information in WRF. Some important changes include a direct manipulation of the geogrid files (without the creation of temporary files), and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

This development of this package is in line with the objectives of WUDAPT, the World Urban Database and Access Portals Tools community project, that aims to 1) acquire and make accessible coherent and consistent information on form and function of urban morphology relevant to climate weather, and environment studies, and 2) provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue, because of three recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the LCZ Generator web application (Demuzere et al., 2021), 2) the availability of a global LCZ map (Demuzere et al., 2022), and 3) WRF versions > 4.3 (Skamarock et al., 2021) are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato & Chen (2021) and Zonato et al. (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here, so as to make the translation of LCZ-based parameters better and simpler.

^{*}corresponding author

Initial data requirements

In order to use the tool, two input files are required:

1. A **geo_em.d0X** (.nc) file for the inner WRF model domain in which one would like to use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool as part of the WRF Preprocessing System (WPS), without additional modifications of the standard procedure.
2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of the `geo_em.d0X.nc` file. There are a number of ways to obtain an LCZ map for your region of interest (ROI):
 - Extract your ROI from the global LCZ map (Demuzere et al., 2022), or the continental-scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et al., 2020) (see also [here](#) for more info).
 - Check if your ROI is already covered by the many LCZ maps available in the [submission table](#) of the LCZ Generator.
 - Use the [LCZ Generator](#) to make your own LCZ map for your ROI. See also [here](#) for more information. When using LCZ maps produced with the LCZ Generator, by default the Gaussian filtered LCZ map is used (corresponding to argument `--lcz-band = 1`).

Note: When using LCZ information from any of the large-scale LCZ maps, please make sure to crop your domain of interest first, in order to avoid memory issues.

Workflow

The goal of the Python-based W2W tool is to obtain an inner WRF domain file (`geo_em.d0X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters relevant for all urban parameterizations embedded in WRF: the single layer urban canopy model (Noah/SLUCM, Kusaka et al. (2001)), the Building Environment Parameterization (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al. (2010)).

To get to that point, a number of sequential steps are required:

Step 1: Remove the default urban land cover

The default urban land cover from MODIS is replaced with the dominant surrounding vegetation category, as done in Li et al. (2020). This procedure affects WRF's parameters `LU_INDEX`, `LANDUSEF` and `GREENFRAC`. `LU_INDEX` is selected as the dominant category from the corresponding argument `--npix-nlc` (default = 45) nearest grid points (excluding ocean, urban and lakes). `GREENFRAC` is calculated as the mean over all grid points with that dominant vegetation category among the `--npix-nlc` nearest points. For each grid point, if `LANDUSEF` had any percentage of urban, it is set to zero and the percentage is added to the dominant vegetation category assigned to that grid point.

Resulting output: **geo_em.d0X_NoUrban.nc**

Step 2: Define the LCZ-based urban extent

LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF

75 resolution. Areas with `FRC_URB2D < 0.2` (corresponding to argument `--frc-threshold`)
 76 are currently considered non-urban. This choice has been made to avoid the use of the
 77 urban schemes in areas where the majority of the landuse is vegetated, since the impact of
 78 the impervious surfaces is low. The `FRC_URB2D` field is also used to mask all other urban
 79 parameter fields, so that their extent is consistent.

80 Resulting output: `geo_em.d0X_LCZ_extent.nc`

81 Step 3: Introduce modal built LCZ classes

82 For each WRF grid cell, the mode of the underlying built LCZ classes is added to `LU_INDEX`
 83 (numbered from 31-41). See [here](#) for more info. Note that the `W2W` routine by default considers
 84 LCZ classes 1-10 as built classes (corresponding to argument `--built-lcz`). In some cases,
 85 also LCZ E (or 15 - Bare rock or paved) can be considered as a built LCZ class, as it might
 86 reflect large asphalt surfaces such as big parking lots or airstrips. In that case, the user must
 87 make sure the `--built-lcz` argument is set appropriately.

88 Step 4: Assign urban canopy parameters

89 Two procedures are followed when assigning the various urban canopy parameters to the LCZ
 90 map and translating this information onto WRF's grid:

91 **Procedure 1: Morphological parameters** are assigned directly to the high-resolution LCZ
 92 map, and are afterwards aggregated to the lower-resolution WRF grid. As a result, the method
 93 produces a unique urban morphology parameter value for each WRF grid cell. This was found
 94 to be more efficient in reproducing urban boundary layer features, especially in the outskirts of
 95 the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#) routine (Varentsov
 96 et al., 2020).

97 Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv`, and
 98 are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014).
 99 Note however that the values of `MH_URB2D_MIN`, `MH_URB2D`, `MH_URB2D_MAX` for
 100 LCZ 7 are set to 4, 5 and 6 m instead of 2, 3 and 4 m, because the minimum building height
 101 that can be assigned to BEP-BEM is 5m if `dz_u = 5m` (standard value) is used.

102 In addition:

- 103 ▪ While `URBPARM_LCZ.TBL` (stored in WRF's `run/` folder) has values on street width
 104 (`SW`), `W2W` derives street width from the mean building height (`MH_URB2D`) and the
 105 Height-to-Width ratio (`H2W`), to have these fields consistent.
- 106 ▪ Building width (`BW`), is derived from $(BLDFR_URB2D / (FRC_URB2D - BLDFR_URB2D))$
 107 * `SW`, these values being available from the look-up table `LCZ_UCP_default.csv`.
- 108 ▪ Plan (`LP_URB2D`), frontal (`LF_URB2D`) and total (`LB_URB2D`) area indices are based
 109 on formulas in Zonato et al. (2020).
- 110 ▪ `HI_URB2D` is obtained by fitting a bounded normal distribution to the minimum
 111 (`MH_URB2D_MIN`), mean (`MH_URB2D`), and maximum (`MH_URB2D_MAX`)
 112 building height, as provided in `LCZ_UCP_default.csv`. The building height
 113 standard deviation is also required, and is approximated as $(MH_URB2D_MAX -$
 114 $MH_URB2D_MIN) / 4$.
- 115 ▪ For computational efficiency, `HI_URB2D` values lower than 5% were set to 0 after
 116 resampling, the remaining `HI_URB2D` percentages are re-scaled to 100%.

117 **Procedure 2:** In line with the former Fortran-based `W2W` procedure, **radiative and thermal**
 118 **parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell (see
 119 *Step 3*). These parameter values are not stored in the NetCDF output, but are read from

120 URBPARAM_LCZ.TBL and assigned automatically to the modal LCZ class when running the
121 model.

122 Step 5: Adjust global attributes

123 In a final step, some global attributes are adjusted in the resulting NetCDF files:

- 124 ■ NBUI_MAX is added as a global attribute, reflecting the maximum amount of
125 HI_URB2D classes that are not 0 across the model domain. This parameter can be
126 used when compiling WRF, to optimize memory storage.
- 127 ■ NUM_LAND_CAT is set to 41, to reflect the addition of 10 (or 11) built LCZ classes.
128 This is not only done for the highest resolution domain file (e.g. d04), but also for **all of**
129 **its lower-resolution parent domain files (e.g. d01, d02, d03)**. As such, make sure
130 these files are also available in the input data directory. In case the parent domain files
131 have NUM_CAT_LAND \neq 41, new parent domain files will be written to your drive
132 with the extension _41.

133 Resulting output: `geo_em.d0X_LCZ_params.nc` (and `geo_em.d0X_41.nc`)

134 Integration in WRF's preprocessing

135 The current tool is designed to work with the `geo_em.d0X` files produced by `geogrid.exe`, which
136 is available in the WRF Preprocessing System (WPS). WPS needs to be at a version >3.8 ,
137 in order to incorporate the urban geometrical parameters in the URB_PARAM matrix (Glotfelty
138 et al., 2013). The user should run `geogrid.exe` using its default settings, which will provide
139 the various `geo_em.d0X.nc` files containing the static data fields. No additional variables
140 are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL` table. The W2W tool
141 (Figure 1) reads the standard `geo_em.d0X.nc` files (for all the domains) and produces the
142 aforementioned `geo_em.d0X_LCZ_params.nc` files. The user should then simply rename
143 these files to the standard name for each of the domains (e.g. rename `geo_em.d01_41.nc`
144 to `geo_em.d01.nc`, `geo_em.d04_LCZ_params.nc` to `geo_em.d04.nc`, ...), which will serve as
145 input to the `metgrid.exe` module (Figure 1).




Figure 1: Modified workflow to set-up and run a WRF simulations including urban parameters derived from LCZs using W2W.

146 Potential use cases

147 The files provided as output by W2W allow a wide range of applications, including - but not
148 limited to - addressing the impact of:

- 149 ▪ urbanization, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_NoUrban.nc`
150 files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- 151 ▪ an improved urban land cover extent description, by running WRF with the default
152 `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_extent.nc` files (similar to for example Bhati
153 & Mohan (2018) and Mallard et al. (2018)).
- 154 ▪ a more detailed (LCZ-based) urban description, by running WRF with the default
155 `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_params.nc` files (see for example Brousse
156 et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019),
157 Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021)
158 and Patel et al. (2022)).

159 Important notes

- 160 ▪ Make sure to set `use_wudapt_lcz=1` (default is 0) and `num_land_cat=41` (default is
161 21) in WRF's `namelist.input` when using the LCZ-based urban canopy parameters.
- 162 ▪ The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv`
163 and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your
164 ROI. If available, please adjust the urban canopy parameters values according to the
165 characteristics of your ROI. A custom csv file can be specified using the `--lcz-ucp p`
166 `ath/to/custom_file.csv` flag.
- 167 ▪ It is advised to use this tool with urban parameterization options BEP or BEP+BEM
168 (`sf_urban_physics = 2` or `3`, respectively). In case you use this tool with the
169 SLUCM model (`sf_urban_physics = 1`), make sure your lowest model level is above
170 the highest building height. If not, `real.exe` will provide the following error message:
171 `ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine`
172 `urban - change ZDC and ZOC.`
- 173 ▪ At the end of running W2W, a note is displayed that indicates the `nbui_max` value,
174 e.g. for the sample data: Set `nbui_max` to 5 during compilation, in order t
175 o optimize memory storage. This is especially relevant for users that work with the
176 BEP or BEP+BEM urban parameterization schemes (`sf_urban_physics = 2` or `3`,
177 respectively). See also `num_urban_nbui` in [WRF's README.namelist](#) for more info.
- 178 ▪ It is advised to use WRF versions > 4.3, that are able to ingest 10 or 11 built classes
179 (corresponding to WUDAPT's LCZs) by default (Skamarock et al., 2021), and WPS
180 versions > 3.8, in order to incorporate the urban geometrical parameters in the `URB_P`
181 `ARAM` matrix (Glotfelty et al., 2013).

182 Acknowledgements

183 We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez
184 Sanchez and Oscar Brousse.

185 References

- 186 Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of
187 heat island and thermal comfort over an urban airshed in India under varying land use/land

- cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient land use producing data tool for mesoscale models? Integration of urban LCZ in WRF over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O., Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley, M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018). WUDAPT: An urban weather, climate, and environmental modeling infrastructure for the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924. <https://doi.org/10.1175/BAMS-D-16-0236.1>
- Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert and crowd-sourced training data to map urban form and functions for the continental US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- Demuzere, M., Kittner, J., Martilli, A., Mills, G., Moede, C., Stewart, I. D., Vliet, J. van, & Bechtel, B. (2022). A global map of Local Climate Zones to support earth system modelling and urban scale environmental science. *Earth System Science Data Discussions*. <https://doi.org/10.5194/essd-2022-92>
- Glotfelty, T., Tewari, M., Sampson, K., Duda, M., Chen, F., & Ching, J. (2013). *NUDAPT 44: How to use NUDAPT dataset in WRF/SLUCM/MLUCM models* (p. 9). National Center for Atmospheric Research. <https://www.yumpu.com/en/document/read/26871494/how-to-use-nudapt-dataset-in-wrf-slucm-mlucm-models>
- Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing detailed urban canopy parameters for mesoscale climate modelling: a comparison between WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*, 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler, W. (2021). Resolving the influence of local flows on urban heat amplification during heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>
- Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Urban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>
- Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X., Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>
- Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to contemporary land-use datasets across the contiguous united states using dynamical downscaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>
- Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WUDAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. <https://www.wudapt.org/wp-content/>

- uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021
- Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parameterisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>
- Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged, Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>
- Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M., & Niyogi, D. (2022). Modelling Large-Scale Heatwave by Incorporating Enhanced Urban Representation. *Journal of Geophysical Research : Atmospheres*, 127, 1–33. <https://doi.org/10.1029/2021JD035316>
- Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*, 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model coupled with an urban canopy parameterization for urban climate simulations—part I. formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>
- Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric Research. <https://doi.org/10.5065/1dfh-6p97>
- Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies. *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>
- Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the ‘local climate zone’ scheme using temperature observations and model simulations [Journal Article]. *International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>
- Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters on a Megacity’s Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>
- Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang, R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018), 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones, Mitigation Strategies, building materials permeability and new buildings drag coefficient*. <http://www.wudapt.org/wudapt-to-wrf/>. https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf
- Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the performance of a novel WUDAPT averaging technique to define urban morphology with mesoscale models. *Urban Climate*, 31(May 2019), 100584. <https://doi.org/10.1016/j.uclim.2020.100584>

285 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., Barlage, M., Zardi, D., & Giovannini,
286 L. (2021). Exploring the effects of rooftop mitigation strategies on urban temperatures
287 and energy consumption. *Journal of Geophysical Research: Atmospheres*, 1–30. [https:](https://doi.org/10.1029/2021JD035002)
288 [//doi.org/10.1029/2021JD035002](https://doi.org/10.1029/2021JD035002)

DRAFT