

W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere^{*1}, Daniel Argüeso², Andrea Zonato³, and Jonas Kittner¹

¹ Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany ² Physics Department, University of the Balearic Islands, Palma, Spain ³ Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Editor Name](#) ↗

Submitted: 01 January XXXX

Published: 01 January XXXX

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

The Python-based WUDAPT-to-WRF (W2W) package is developed to translate Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model (Skamarock et al., 2021). It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides an improved, simpler, and more efficient procedure to use LCZ information in WRF. Some important changes include direct manipulation of the geogrid files without the creation of temporary files, and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

This development of this package is in line with the objectives of WUDAPT, the World Urban Database and Access Portals Tools community project, that aims to 1) acquire and make accessible coherent and consistent information on the form and function of urban morphology relevant to climate weather and environmental studies, and 2) provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue because of three recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the LCZ Generator web application (Demuzere et al., 2021), 2) the availability of a global LCZ map (Demuzere et al., 2022), and 3) WRF versions > 4.3 (Skamarock et al., 2021) can ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato & Chen (2021) and Zonato et al. (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here to translate LCZ-based parameters better and simpler. It differs from its Fortran-based predecessor mainly by 1) using a more up-to-date LCZ-based urban extent, 2)

*corresponding author

38 aggregating the morphological parameters instead of using modal values, and 3) the fact that
39 all processing is done with one (automated) tool, whereas the Fortran-based version required
40 multiple pre-processing steps and manual namelist changes, described in more detail by Martilli
41 et al. (2016).

42 Initial data requirements

43 To use the tool, two input files are required:

- 44 1. A **geo_em.dXX** (.nc) file for the inner WRF model domain in which one would like to
45 use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool
46 as part of the WRF Preprocessing System (WPS), without additional modifications of
47 the standard procedure.
- 48 2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of
49 the `geo_em.dXX.nc` file. There are several ways to obtain an LCZ map for your region
50 of interest (ROI):
 - 51 ■ Extract your ROI from the global LCZ map (Demuzere et al., 2022), or the continental-
52 scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et
53 al., 2020) (see [here](#) for more info).
 - 54 ■ Check if your ROI is already covered by the many LCZ maps available in the [submission](#)
55 [table](#) of the LCZ Generator.
 - 56 ■ Use the [LCZ Generator](#) to make your LCZ map for your ROI. See [here](#) for more informa-
57 tion. When using LCZ maps produced with the LCZ Generator, by default the Gaussian
58 filtered LCZ map is used (corresponding to argument `--lcz-band = 1`).

59 **Note:** When using LCZ information from any of the large-scale LCZ maps, please make sure
60 to crop your domain of interest first to avoid memory issues.

61 Workflow

62 The goal of the Python-based W2W tool is to obtain an inner WRF domain file (`geo_em.dX`
63 `X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters
64 relevant for all urban parameterizations embedded in WRF: the single-layer urban canopy
65 model (Noah/SLUCM, Kusaka et al. (2001)), the Building Environment Parameterization
66 (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al.
67 (2010)).

68 To get to that point, a number of sequential steps are required:

69 Step 1: Remove the default urban land cover

70 The default urban land cover from MODIS is replaced with the dominant surrounding veg-
71 etation category, as done in Li et al. (2020). This procedure affects WRF's parameters
72 `LU_INDEX`, `LANDUSEF`, and `GREENFRAC`. First, an initial number of neighboring pixels
73 (corresponding argument `--npix-area`, default = `--npix-nlc ** 2`) are selected using the
74 k-d tree algorithm, to find the nearest pixels based on the great circle arc length, specifi-
75 cally [the chord length formula](#). Afterward, the `LU_INDEX` is set by sampling the dominant
76 category from the corresponding argument `--npix-nlc` (default = 45) nearest initial grid
77 points (excluding ocean, urban, and lakes). `GREENFRAC` is calculated as the mean over all

78 grid points with that dominant vegetation category among the `--npix-nlc` nearest points.
 79 For each grid point, if LANDUSEF had any percentage of urban, it is set to zero and the
 80 percentage is added to the dominant vegetation category assigned to that grid point.

81 Resulting output: `geo_em.dXX_NoUrban.nc`

82 Step 2: Define the LCZ-based urban extent

83 LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`)
 84 are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF
 85 resolution. Areas with `FRC_URB2D < 0.2` (corresponding to argument `--frc-threshold`)
 86 are currently considered non-urban. This choice has been made to avoid the use of the urban
 87 schemes in areas where the majority of the land use is vegetated since the impact of the
 88 impervious surfaces is low. The `FRC_URB2D` field is also used to mask all other urban
 89 parameter fields so that their extent is consistent.

90 Resulting output: `geo_em.dXX_LCZ_extent.nc`

91 Step 3: Introduce modal built LCZ classes

92 For each WRF grid cell, the mode of the underlying built LCZ classes is added to `LU_INDEX`
 93 (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default considers
 94 LCZ classes 1-10 as built classes (corresponding to argument `--built-lcz`). In some cases,
 95 also LCZ E (or 15 - Bare rock or paved) can be considered as a built LCZ class, as it might
 96 reflect large asphalt surfaces such as big parking lots or airstrips. In that case, the user must
 97 make sure the `--built-lcz` argument is set appropriately.

98 Step 4: Assign urban canopy parameters

99 Two procedures are followed when assigning the various urban canopy parameters to the LCZ
 100 map and translating this information onto WRF's grid:

101 **Procedure 1: Morphological parameters** are assigned directly to the high-resolution LCZ
 102 map, and are afterward aggregated to the lower-resolution WRF grid. As a result, the method
 103 produces a unique urban morphology parameter value for each WRF grid cell. This was found
 104 to be more efficient in reproducing urban boundary layer features, especially in the outskirts of
 105 the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#) routine (Varentsov
 106 et al., 2020).

107 Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv` and
 108 are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014).
 109 Note however that the values of `MH_URB2D_MIN`, `MH_URB2D`, and `MH_URB2D_MAX`
 110 for LCZ 7 are set to 4, 5, and 6 m instead of 2, 3, and 4 m because the minimum building
 111 height that can be assigned to BEP-BEM is 5m if `dz_u = 5m` (standard value) is used.

112 In addition:

- 113 ■ While `URBPARM_LCZ.TBL` (stored in WRF's `run/` folder) has values on street width
 114 (SW), W2W derives street width from the mean building height (`MH_URB2D`) and the
 115 Height-to-Width ratio (H2W), to have these fields consistent.
- 116 ■ Building width (BW), is derived from $(BLDFR_URB2D / (FRC_URB2D - BLDFR_URB2D))$
 117 * SW, these values are available in the look-up table `LCZ_UCP_default.csv`.
- 118 ■ Plan (`LP_URB2D`), frontal (`LF_URB2D`), and total (`LB_URB2D`) area indices are
 119 based on formulas in Zonato et al. (2020).

- HI_URB2D is obtained by fitting a bounded normal distribution to the minimum (MH_URB2D_MIN), mean (MH_URB2D), and maximum (MH_URB2D_MAX) building height, as provided in LCZ_UCP_default.csv. The building height standard deviation is also required and is approximated as $(MH_URB2D_MAX - MH_URB2D_MIN) / 4$.
- For computational efficiency, HI_URB2D values lower than 5% were set to 0 after resampling, and the remaining HI_URB2D percentages are re-scaled to 100%.

Procedure 2: In line with the former Fortran-based W2W procedure, **radiative and thermal parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell (see Step 3). These parameter values are not stored in the NetCDF output but are read from URBPARAM_LCZ.TBL and assigned automatically to the modal LCZ class when running the model.

Step 5: Adjust global attributes

In a final step, some global attributes are adjusted in the resulting NetCDF files:

- NBUI_MAX is added as a global attribute, reflecting the maximum amount of HI_URB2D classes that are not 0 across the model domain. This parameter can be used when compiling WRF to optimize memory storage.
- NUM_LAND_CAT is set to 41, to reflect the addition of 10 or 11 built LCZ classes. This is not only done for the highest resolution domain file (e.g. d04), but also for **all of its lower-resolution parent domain files (e.g. d01, d02, d03)**. As such, make sure these files are also available in the input data directory. In case the parent domain files have $NUM_CAT_LAND \neq 41$, new parent domain files will be written to your drive with the extension _41.

Resulting output: `geo_em.dXX_LCZ_params.nc` (and `geo_em.dXX_41.nc`)

Integration in WRF's preprocessing

The current tool is designed to work with the `geo_em.dXX` files produced by `geogrid.exe`, which is available in the WRF Preprocessing System (WPS). WPS needs to be at a version >3.8, to incorporate the urban geometrical parameters in the URB_PARAM matrix (Glotfelty et al., 2013). The user should run `geogrid.exe` using its default settings, which will provide the various `geo_em.dXX.nc` files containing the static data fields. No additional variables are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL` table. The W2W tool (Figure 1) reads the standard `geo_em.dXX.nc` files (for all the domains) and produces the aforementioned `geo_em.dXX_LCZ_params.nc` files. The user should then rename these files to the standard name for each of the domains (e.g. rename `geo_em.d01_41.nc` to `geo_em.d01.nc` and `geo_em.d04_LCZ_params.nc` to `geo_em.d04.nc`), which will serve as input to the `metgrid.exe` module (Figure 1).




Figure 1: Modified workflow to set up and run a WRF simulation including urban parameters derived from LCZs using W2W.

Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_NoUrban.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover extent description, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019), Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2022)).

Important notes

- Make sure to set `use_wudapt_lcz=1` (default is 0) and `num_land_cat=41` (default is 21) in WRF's `namelist.input` when using the LCZ-based urban canopy parameters.
- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the urban canopy parameters values according to the characteristics of your ROI. A custom csv file can be specified using the `--lcz-ucp path/to/custom_file.csv` flag.
- It is advised to use this tool with urban parameterization options BEP or BEP+BEM (`sf_urban_physics = 2` or `3`, respectively). In case you use this tool with the SLUCM model (`sf_urban_physics = 1`), make sure your lowest model level is above the highest building height. If not, `real.exe` will provide the following error message:

181 ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine
 182 urban - change ZDC and ZOC.
 183 ■ At the end of running W2W, a note is displayed that indicates the nbui_max value, e.g. for
 184 the sample data: Set nbui_max to 5 during compilation, to optimize memory
 185 storage. This is especially relevant for users that work with the BEP or BEP+BEM
 186 urban parameterization schemes (sf_urban_physics = 2 or 3, respectively). See
 187 num_urban_nbui in [WRF's README.namelist](#) for more info.
 188 ■ It is advised to use WRF versions > 4.3, that can ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default (Skamarock et al., 2021), and WPS versions
 189 > 3.8, to incorporate the urban geometrical parameters in the URB_PARAM matrix (Glotfelty et al., 2013).
 190
 191

192 Acknowledgements

193 We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez
 194 Sanchez, and Oscar Brousse.

195 References

- 196 Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of
 197 heat island and thermal comfort over an urban airshed in India under varying land use/land
 198 cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- 199 Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient
 200 land use producing data tool for mesoscale models? Integration of urban LCZ in WRF
 201 over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- 202 Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O.,
 203 Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley,
 204 M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018).
 205 WUDAPT: An urban weather, climate, and environmental modeling infrastructure for
 206 the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924.
 207 <https://doi.org/10.1175/BAMS-D-16-0236.1>
- 208 Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate
 209 zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- 210 Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert
 211 and crowd-sourced training data to map urban form and functions for the continental
 212 US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- 213 Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to
 214 Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- 215
 216 Demuzere, M., Kittner, J., Martilli, A., Mills, G., Moede, C., Stewart, I. D., Vliet, J. van,
 217 & Bechtel, B. (2022). A global map of Local Climate Zones to support earth system
 218 modelling and urban scale environmental science. *Earth System Science Data Discussions*.
 219 <https://doi.org/10.5194/essd-2022-92>
- 220 Glotfelty, T., Tewari, M., Sampson, K., Duda, M., Chen, F., & Ching, J. (2013). *NUDAPT 44:*
 221 *How to use NUDAPT dataset in WRF/SLUCM/MLUCM models* (p. 9). National Center
 222 for Atmospheric Research. <https://www.yumpu.com/en/document/read/26871494/how-to-use-nudapt-dataset-in-wrf-slucm-mlucm-models>
 223

- 224 Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing
225 detailed urban canopy parameters for mesoscale climate modelling: a comparison between
226 WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*,
227 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- 228 Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler,
229 W. (2021). Resolving the influence of local flows on urban heat amplification during
230 heatwaves. *Environmental Research Letters*, 16(6), 064066. [https://doi.org/10.1088/](https://doi.org/10.1088/1748-9326/ac0377)
231 [1748-9326/ac0377](https://doi.org/10.1088/1748-9326/ac0377)
- 232 Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Ur-
233 ban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab
234 Models. *Boundary-Layer Meteorology*, 101(3), 329–358. [https://doi.org/10.1023/A:](https://doi.org/10.1023/A:1019207923078)
235 [1019207923078](https://doi.org/10.1023/A:1019207923078)
- 236 Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X.,
237 Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall
238 Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. [https://doi.org/](https://doi.org/10.1029/2020GL088758)
239 [10.1029/2020GL088758](https://doi.org/10.1029/2020GL088758)
- 240 Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to con-
241 temporary land-use datasets across the contiguous united states using dynamical down-
242 scaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. [https://doi.org/](https://doi.org/10.1175/JAMC-D-17-0328.1)
243 [10.1175/JAMC-D-17-0328.1](https://doi.org/10.1175/JAMC-D-17-0328.1)
- 244 Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WU-*
245 *DAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. [https://www.wudapt.org/wp-content/](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf)
246 [uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf)
247 [pdf](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf)
- 248 Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange paramete-
249 risation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. [https://doi.org/](https://doi.org/10.1023/A:1016099921195)
250 [10.1023/A:1016099921195](https://doi.org/10.1023/A:1016099921195)
- 251 Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into
252 WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged,
253 Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. [https://doi.org/](https://doi.org/10.1007/s00704-019-02881-1)
254 [10.1007/s00704-019-02881-1](https://doi.org/10.1007/s00704-019-02881-1)
- 255 Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M.,
256 & Niyogi, D. (2022). Modelling Large-Scale Heatwave by Incorporating Enhanced Urban
257 Representation. *Journal of Geophysical Research : Atmospheres*, 127, 1–33. [https://doi.](https://doi.org/10.1029/2021JD035316)
258 [org/10.1029/2021JD035316](https://doi.org/10.1029/2021JD035316)
- 259 Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy
260 rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban*
261 *Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- 262 Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved
263 WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*,
264 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- 265 Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model
266 coupled with an urban canopy parameterization for urban climate simulations—part I.
267 formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied*
268 *Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>
- 269 Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W.,
270 Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of*
271 *the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric
272 Research. <https://doi.org/10.5065/1dfh-6p97>

- 273 Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies.
 274 *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>
 275
- 276 Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the ‘local climate
 277 zone’ scheme using temperature observations and model simulations [Journal Article].
 278 *International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>
 279
- 280 Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters
 281 on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>
 282
- 283 Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang,
 284 R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on
 285 WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018),
 286 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- 287 Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones,*
 288 *Mitigation Strategies, building materials permeability and new buildings drag coefficient.*
 289 <http://www.wudapt.org/wudapt-to-wrf/>. [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
 290 [product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
- 291 Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the
 292 performance of a novel WUDAPT averaging technique to define urban morphology with
 293 mesoscale models. *Urban Climate*, 31(May 2019), 100584. <https://doi.org/10.1016/j.uclim.2020.100584>
 294
- 295 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., Barlage, M., Zardi, D., & Giovannini,
 296 L. (2021). Exploring the effects of rooftop mitigation strategies on urban temperatures
 297 and energy consumption. *Journal of Geophysical Research: Atmospheres*, 1–30. <https://doi.org/10.1029/2021JD035002>
 298