Homogeneous train schedules

Matthias Görgens

March 11, 2009

Abstract

Although most railroad schedules remain in use for half a year, they tend to repeat themself each week. Thus one can also repeat their implementations, the vehicle schedulings, each week.

But railroad schedules show even more structure: All the working days of the week are usually identical. This work sets out to investigate how to transport this similarity in relations to their vehicle scheduligs. We define homogeneous vehicle schedulings, establish a measure of partial homogenity and search for ways to efficiently find the most homogeneous vehicle scheduling for a given schedule.

1 Some definitions

Let G = (V, A) be a digraph of connected channels, i.e. railway stations, maintenance shops. Now look at relations $(v, t_v, w, t_w) \in (V, \mathbb{R})^2$ where v, t_v name the station and time of departure, and w, t_v name the station and time of arrival. We call a set of relations $S \subseteq (V, \mathbb{R})^2$ a schedule.

We want to focus our attention to cyclic schedules. A schedule S is cyclic, iff there exists a period $w \in \mathbb{R}^+$, so that for each relation $(v,t_v,w,t_w) \in S$ the relations $\{(v,t_v+nw,w,t_w+nw) \mid n \in \mathbb{Z}\}$ are also in S. For a cyclic schedule S we will identify S with its image under the canonic morphism $\varphi:(V,\mathbb{R})^2 \to (V,\mathbb{R}/w\mathbb{Z})^2$.

In practise schedules repeat every week i.e. w = 7days.

This work will deal with schedules that show more structure: Fix an interval $d:=\frac{w}{n}$ with $n\in\mathbb{N}^+$; in addition to the schedule S introduce the set of trains $T\subseteq\mathcal{P}(S)$ partitioning the relations of the schedule. Relations in each $train\ z\in T$ all share the same arrival and departure channels, also their arrival and departure times only differ by a natural multiply of w. That is

$$z \subseteq [(v, t_v, w, t_w)]_d := \{(v, t_v + nd, w, t_w + nd) \mid n \in \mathbb{Z}\}$$
 (1)