

ADVANCES IN TRANSLATIONAL SCIENCE

Joseph H. Sellin, Section Editor

Translating an Understanding of the Pathogenesis of Hepatic Fibrosis to Novel Therapies

DON C. ROCKEY

Department of Internal Medicine, Medical University of South Carolina, Charleston, South Carolina

The response to injury is one of wound healing and fibrogenesis, which ultimately leads to fibrosis. The fibrogenic response to injury is a generalized one across virtually all organ systems. In the liver, the injury response, typically occurring over a prolonged period of time, leads to cirrhosis (although it should be pointed out that not all patients with liver injury develop cirrhosis). The fact that many different diseases result in cirrhosis suggests a common pathogenesis. The study of hepatic fibrogenesis over the past 2 decades has been remarkably active, leading to a considerable understanding of this process. It clearly has been shown that the hepatic stellate cell is a central component in the fibrogenic process. It also has been recognized that other effector cells are important in the fibrogenic process, including resident fibroblasts, bone marrow-derived cells, fibrocytes, and even perhaps cells derived from epithelial cells (ie, through epithelial to mesenchymal transition). A key aspect of the biology of fibrogenesis is that the fibrogenic process is dynamic; thus, even advanced fibrosis (or cirrhosis) is reversible. Together, an understanding of the cellular basis for liver fibrogenesis, along with multiple aspects of the basic pathogenesis of fibrosis, have highlighted many exciting potential therapeutic opportunities. Thus, although the most effective antifibrotic therapy is simply treatment of the underlying disease, in situations in which this is not possible, specific antifibrotic therapy is likely not only to become feasible, but will soon become a reality. This review highlights the mechanisms underlying fibrogenesis that may be translated into future antifibrotic therapies and to review the current state of clinical development.

Keywords: Cirrhosis; Wound Healing; Fibrogenesis; Scar; Tissue; Injury.

The response to chronic injury is a generalized one, with features common among multiple organ systems. This feature suggests thematically related pathogenic events across organs. In the liver, many different kinds of injury, including viral hepatitis, alcohol, fatty liver, biliary tract disease, iron or copper overload, cystic fibrosis, and others cause fibrogenesis, and subsequently cirrhosis.

Over the past 2 decades, much has been learned about the biology and pathophysiology of fibrosis. Understanding the mechanisms underlying fibrosis has pointed out several poten-

tial therapeutic approaches. Preclinical studies have been particularly informative, and have highlighted many possible therapies. Although therapies that are directed at the underlying disease process, including antiviral therapies for patients with hepatitis B and hepatitis C virus (HCV) infections, have proven to be effective at reducing and/or reversing fibrosis, specific and effective antifibrotic therapy remains elusive. The objective of this review is to emphasize fundamental concepts underlying hepatic fibrogenesis, and to review translational therapeutics.

Fibrogenesis: Pathophysiology

The Fibrogenic Process

A critical aspect of the fibrogenic response is that injury, typically to hepatocytes, stimulates the injury response (Figure 1). Multiple forms of injury, including hepatitis, metabolic disease (ie, in particular the metabolic syndrome), biliary injury, toxins (including alcohol), and heavy metals, cause a variety of complicated and often integrated effects in the liver. For example, viral hepatitis causes activation of T cells, with recruitment of other inflammatory cells, as well as inflammatory mediators, and this leads to the fibrogenic wounding response (Figure 1). Alcohol-mediated hepatocyte injury causes a classic inflammatory lesion, including tumor necrosis factor (TNF), which leads to hepatitis, and a fibrogenic wounding response. It should be emphasized that multiple different cell types play a role in the injury milieu. For example, injury to endothelial cells, either directly or indirectly, causes them to produce abnormal extracellular matrix, which in turn stimulates fibrogenesis by stellate cells.¹

A central event in the hepatic wounding response is enhanced extracellular matrix production, or fibrogenesis (Figure 1). Irrespective of the specific cause of liver injury (in both experimental models and human cirrhosis), the wound process leads to increased synthesis of extracellular matrix. The fibrogenic process is characterized by increases in a multiple matrix components, including the interstitial collagens, basement membrane collagens, proteoglycans, and matrix glycoproteins such as

Abbreviations used in this paper: CTGF, connective tissue growth factor; HCV, hepatitis C virus; MMP, metalloproteinase; NASH, nonalcoholic steatohepatitis; PDGF, platelet-derived growth factor; PPAR, peroxisomal proliferator active receptor; TGF, transforming growth factor; TNF, tumor necrosis factor.

© 2013 by the AGA Institute

1542-3565/\$36.00

<http://dx.doi.org/10.1016/j.cgh.2013.01.005>




Figure 1. Liver injury and fibrogenesis. In the liver, many different types of injuries (ie, chronic hepatitis, ethanol, metabolic disease, biliary tract disease, iron, copper, and so forth) lead to hepatocyte injury, and then typically an inflammatory response. This injury process is complicated, but in aggregate it stimulates a wound healing response, which involves a number of different systems. Paramount in this process often is the recruitment of inflammatory cells. Among other properties, inflammatory cells produce a variety of mediators, cytokines, and other factors, which in turn are responsible for stimulation and/or recruitment of other cells. Key among these other cells include effector cells, highlighted in the figure, and including stellate cells, fibrocytes, fibroblasts, and even fibroblasts derived through epithelial to mesenchymal transition. These effectors produce extracellular matrix proteins (see text) and importantly interact with other cells in the wounding milieu. In addition, it is important to emphasize that many forms of injury lead to activation and transformation of other cells in the liver, such as endothelial and bile duct epithelial cells. Injury to these cells in turn leads to a variety of downstream effects. Each injured endothelial bile duct epithelial cell is capable of stimulating effector cells to produce extracellular matrix constituents. LPS, lipopolysaccharide; NK, natural killer.

laminin and fibronectin²; specific changes in matrix composition are highly similar in all forms of liver injury and hepatic fibrogenesis. Among the most prominent extracellular matrix proteins are the collagens (type I > III > IV), but increases in other matrix proteins also are prominent. It is important to emphasize that the wounding process is a dynamic one that includes aspects of matrix synthesis and deposition as well as degradation.³ This point is exemplified by a robust body of literature data indicating that experimental^{4–6} and clinical fibrosis,^{7–9} and even clinical cirrhosis, is reversible.^{10–17} In one study in patients with chronic hepatitis B infection and cirrhosis,¹⁴ 436 of 651 patients were assigned to receive lamivudine and 215 were assigned to receive placebo; 7.8% of patients receiving lamivudine and 17.7% of those receiving placebo developed hepatocellular carcinoma, spontaneous bacterial peri-

tonitis, bleeding gastroesophageal varices, or died of liver disease ($P = .001$). In addition, the Child–Pugh score increased in 3.4% of the patients receiving lamivudine and in 8.8% of those receiving placebo ($P = .02$). Thus, not only is advanced fibrosis reversible, but resolution of fibrosis also is associated with improved clinical outcomes.

Hepatic Stellate Cells and Their Activation in Fibrogenesis

A key concept in the wounding response is that during the fibrogenic response, there is activation of effector cells. Evidence now supports the presence of a number of effector cells including stellate cells,¹⁸ periportal and pericentral fibroblasts,¹⁹ fibrocytes,²⁰ myofibroblasts, and perhaps fibrogenic




Figure 2. Stellate cell activation. A key pathogenic feature underlying liver fibrosis and cirrhosis is activation of hepatic stellate cells (note that activation of other effector cells is likely to parallel that of stellate cells). The activation process is complex, both in terms of the events that induce activation and the effects of activation. Multiple and varied stimuli participate in the induction and maintenance of activation, including but not limited to cytokines, peptides, and the extracellular matrix itself. Recently, signaling through Toll-like receptor (TLR)4 on stellate cells has been identified as important in activation. Key phenotypic features of activation include production of extracellular matrix, proliferation, up-regulation of smooth muscle proteins, secretion of peptides and cytokines (which have autocrine effects on stellate cells and paracrine effects on other cells such as leukocytes and malignant cells), and up-regulation of various cytokine and peptide receptors. In addition, evidence indicates that stellate cells show several cell fates, highlighted at the *bottom* of the figure, and each of these appear to play a role in the biology of fibrogenesis.

cells derived from hepatocytes through epithelial to mesenchymal transition.^{21,22}

Stellate cells (also known previously as lipocytes, Ito, or perisinusoidal cells), perisinusoidal, pericyte-like cells of mesenchymal origin, have garnered great attention as effectors of the fibrogenic response. In the normal liver, these cells function as a major retinoid reservoir for the body, storing much of the body's vitamin A.^{23,24} Given their pericyte-like appearance, they also may function as regulators of blood flow.²³ Notwithstanding, one of their most notable features occurs after liver injury. In this situation, stellate cells transform from quiescent (normal) to an activated (injured) state; this activation step is a central component to the liver wounding process (Figure 1). The activation process is remarkably complex, with multiple and dynamic features. Phenotypically, it consists of many im-

portant cellular changes; characteristic features include loss of vitamin A, acquisition of stress bundles, and development of prominent rough endoplasmic reticulum (Figure 2). Perhaps the most prominent feature of activation is the striking increase in production and secretion of extracellular matrix proteins, including types I, III, and IV collagens, fibronectin, laminin, proteoglycans, and others.^{2,25} An additional important feature of activation is *de novo* expression of smooth muscle-specific proteins, such as smooth muscle α actin.²⁶ This feature further identifies stellate cells as liver-specific myofibroblasts, a cell type typical of fibrogenesis in all organs.^{27,28}

Although the most prominent features of activation include enhanced extracellular matrix production, and the expression of smooth muscle α actin, activation also is associated with other important cellular phenotypes including enhanced pro-

Table 1. Cytokines and Growth Factors Important in Stellate Cell Fibrogenesis

Profibrogenic	Antifibrogenic
TGF- β	Interferon- γ
TGF- α	Interferon- α
CTGF	Adiponectin
Insulin-like growth factor (I, II) ^a	Endostatin ^a
PDGF ^a	Hepatocyte growth factor
Monocyte chemotactic factor ^a	
Fibroblast growth factor ^a	
Interleukin-1 ^a	
Interleukin-4 ^a	
Interleukin-6 ^a	
Thrombin ^a	
Endothelin-1	
Norepinephrine	
Angiotensin II	
Thrombospondin (1, 2)	
Leptin	
Lipopolysaccharide ^a	

^aAgents whose effect is largely via stimulation of proliferation, or indirect effects on stellate cells.

liferation, release of proinflammatory cytokines,²⁹ release of matrix-degrading enzymes and their inhibitors, and recruitment and activation of other cell types such as hepatocellular cancer and cholangiocarcinoma cells^{30,31} and inflammatory cells.³² When designing therapeutics focused on liver fibrogenesis, it is important to emphasize that each of these features of activation (and fibrogenesis) represent a potential target for therapy. Important elements of the activation process are highlighted later.

Stellate cell fibrogenesis. Multiple factors play key pathogenic roles in stellate cell fibrogenesis. Prominent among these factors are cytokines, small peptides, and the extracellular matrix itself. Transforming growth factor β -1 (TGF- β 1) appears to be the most profibrogenic cytokine in the liver.^{33–35} TGF- β 1 is produced by Kupffer cells, sinusoidal endothelial cells, bile duct epithelial cells, hepatocytes, and stellate cells, and has prominent paracrine/autocrine effects on stellate cells.^{36,37} When TGF- β 1 is overexpressed in the liver, it leads to prominent fibrosis,³³ and when inhibited during experimental liver injury, fibrosis is reduced.³⁸ TGF- β 1 signaling in stellate cells is remarkably complex,³⁹ acting via direct (and to a lesser extent, indirect) pathways to stimulate extracellular matrix production in stellate cells. Although none appears to be as potent as TGF- β 1, a variety of other cytokines and peptides have profibrogenic effects on stellate cells (Table 1), including connective tissue growth factor (CTGF),^{40,41} endothelin-1,⁶ leptin,⁴² angiotensin II,⁴³ and others.

It also should be emphasized that cytokines and growth factors that drive stellate cell proliferation are important in the fibrogenic response because they help expand the total number of fibrogenic (stellate) cells. In essentially all forms of fibrosing liver injury, the number of activated effector cells is increased. Although the major mitogen-driving cellular proliferation appears to be platelet-derived growth factor (PDGF), a variety of other factors appear to be important in stimulation of stellate cell proliferation and include epidermal growth factor, fibroblast growth factor, insulin-like growth factor, thrombin, pro-

tease-activated receptor agonists, monocyte chemotactic factor, insulin-like growth factors, interleukin-6, CTGF, endothelin-1, angiotensin II, and others. Although many of these compounds have isolated proliferative effects (eg, PDGF), others (eg, endothelin-1, angiotensin II, and CTGF) stimulate both proliferation and fibrogenesis.

The vasoactive peptides endothelin-1 and angiotensin II, each of which have pleiotropic cell biological and molecular effects, are notable not only because they have been emphasized in the pathogenesis of hepatic fibrogenesis,^{6,43–45} but also because these compounds have vasoactive properties, and, as such, may be important in the pathogenesis of portal hypertension. This raises the possibility that therapy directed at them could affect both fibrogenesis and portal hypertension. Other biologically active peptides (including unidentified compounds) also may be important in fibrogenesis. For example, dopamine β -hydroxylase-deficient mice, which cannot make norepinephrine, are resistant to fibrogenesis.⁴⁶ Thus, antagonism of these systems is attractive.

A number of cytokines and peptides appear to have anti-activation or antifibrogenic properties toward stellate cells. Although the number of agents is considerably less than the number reported to be profibrogenic and/or stimulate proliferation, included in this group are interferon- γ ,⁴⁷ interferon- α ,⁴⁸ adiponectin,⁴⁹ hepatocyte growth factor,⁵⁰ and possibly stellate cell activation-associated protein.⁵¹

Evolving evidence indicates that the extracellular matrix and the local environment play an important role in modulating stellate cell activation. For example, culture of stellate cells on a basement membrane mimicking the normal basement membrane inhibits stellate cell activation and matrix synthesis,⁵² whereas culture of stellate cells on abnormal substrates such as the extra domain-A isoform of fibronectin leads to increased activation of stellate cells.¹ Further, data suggest that stellate cells sense their surrounding environment.⁵³ For example, it was shown that stellate cells became activated preferentially while exposed to a stiff substrate (compared with a softer substrate), and that this stiffness-dependent activation required adhesion to matrix proteins and the generation of mechanical tension.⁵⁴ It also has been shown that integrins, which link the extracellular matrix to stellate (and other cells), play an important role in transmitting fibrogenic and contractile signals.⁵⁵ Recently, integrin-linked kinase, an integrin-intracellular signaling molecule, has been shown to transmit fibrogenic signals in stellate cells.^{56,57}

It also should be pointed out that fibrogenesis is a dynamic process with elements of extracellular matrix synthesis as well as degradation. During fibrosis progression, not only is there increased expression of extracellular matrix proteins as highlighted earlier, but also metalloproteinases (MMPs), and in particular their tissue inhibitors. Evolving evidence suggests that early in the injury process, increases in expression of MMP-2 and membrane type 1 MMP lead to degradation of normal basement membrane matrix, which appears to facilitate stellate cell activation.^{58–60} In addition, overexpression of the tissue inhibitors of MMP (1 and 2) contributes to the profibrogenic phenotype.⁵⁸ This dynamic interplay of matrix synthesis and degradation is complex, but an attractive therapeutic target. As proof of concept, overexpression of MMP-8 has been shown to lead to partial reversal of fibrosis.⁶¹

Table 2. Liver Diseases in Which Treatment of the Underlying Process May Reverse Fibrosis

Disease	Comments
Hepatitis B	Antiviral treatment improves outcomes
Hepatitis C	Viral eradication improves outcomes
Autoimmune hepatitis	Corticosteroids may improve outcomes
Alcoholic hepatitis	Corticosteroids may improve outcomes
Bile duct obstruction	Biliary decompression improves histology
Hemochromatosis	Iron depletion may improve outcomes
Primary biliary cirrhosis	UDCA, MTX have weak effects
NASH	PPAR ligands have weak effects

MTX, methotrexate; UDCA, ursodeoxycholic acid.

Stellate cell contractility. Activation of stellate cells is accompanied by an increase in expression of proteins characteristic of contractile cells (ie, such as smooth muscle α actin and smooth muscle myosins^{26,62}). Stellate cell contraction has been reported to be mediated by Ca^{++} -dependent and Ca^{++} -independent mechanisms.^{63–66} Stellate cell contraction has a multitude of effects in the injured liver including in perisinusoidal constriction and portal hypertension and also may lead to the collapse and shrunken state of cirrhotic livers.⁴⁵ Stellate cell contractility is likely tied to multiple different systems, including the endothelin, angiotensin, adrenergic, and perhaps other systems.^{44,45,66–71}

Other stellate cell activation phenotypes. Beyond the phenotypes highlighted earlier, during liver injury and activation, stellate cells show a number of important features (Figure 2). For example, apoptosis (ie, programmed cell death) is prominent in stellate cells and appears to be an important mechanism for fibrosis regression.⁵ The data suggest that a balance between cell proliferation and apoptosis is important in determining the dynamics of the total overall stellate cell population in the liver. Based on these data, stimulation of stellate cell apoptosis could be an attractive therapeutic approach.⁷² However, it also has been shown that stellate cell apoptosis may

stimulate stellate cell activation, and thus may not be desirable.⁷³ In addition, stellate cells may undergo senescence⁷⁴ or revert to a normal phenotype.⁷⁵ Recently, autophagy, a catabolic mechanism involving cell degradation of unnecessary or dysfunctional cellular components through the lysosomal pathway, appears to play a role in stellate cell activation.^{76–78} Mice with stellate cell-specific deletion of autophagy-related protein 7, a protein important in mammalian autophagy, had reduced activation after liver injury, leading to reduced fibrosis in vivo.⁷⁸

Approach to Therapy for Fibrosis

It is important to emphasize that the most effective antifibrotic therapies are those that target the primary stimulus to fibrogenesis (Table 2). For example, eradication or inhibition of hepatitis B virus^{7,9} or HCV⁸ leads to reversion of fibrosis and is associated with improved clinical outcomes.^{11,12,14} Fibrosis (and cirrhosis) in patients with autoimmune hepatitis who respond to medical treatment (prednisone or equivalent) is reversible.^{13,17} Fibrosis may improve in patients with alcoholic liver disease who respond to anti-inflammatory therapy such as corticosteroids.^{79,80} Fibrosis reverts in patients with hemochromatosis during iron depletion^{81,82} and after relief of bile duct obstruction.¹⁵ In addition, in patients with nonalcoholic steatohepatitis (NASH), treatment with the peroxisomal proliferator active receptor (PPAR) γ agonist, rosiglitazone, reduced both steatosis and fibrosis.⁸³

Experimental studies have shown that many different interventions are capable of inhibiting (usually preventing) fibrogenesis. Such therapies have been targeted at inhibition of collagen synthesis, matrix deposition, modulation of stellate cell activation, stimulation of matrix degradation, or stimulation of stellate cell death. A number of these preclinical approaches have been transitioned to clinical trials in human beings (Table 3). As of this writing, a specific antifibrotic agent that fits the profile of an ideal agent, one that is potent, safe, orally bioavailable, and inexpensive, is not yet available.

Specific Antifibrotic Therapies

Colchicine is a plant alkaloid that inhibits polymerization of microtubules and has antifibrotic properties in experi-

Table 3. Potential Antifibrotic Therapies Tested in Human Beings

Agent	Disease	Comments	Status
Compounds with anti-inflammatory, antioxidant, or general effects			
Interleukin-10	HCV	Increased viral load	Not suitable for therapy
PPC	EtOH	Minimal if any effect	Not recommended
SAM	EtOH	Minimal if any effect	Not recommended
Silymarin	HCV/ETOH	Further studies pending	
Anti-TNF- α	EtOH	Increased mortality	Likely dangerous
UDCA	Multiple	Modestly effective, safe	May be acceptable (PBC)
Vitamin E	HCV/NASH	Modestly effective, safe	May be acceptable
Pentoxifylline	EtOH	Minimally effective, safe	May be acceptable
Compounds with specific antifibrotic effects			
Colchicine	Miscellaneous	Minimal if any effect	Not recommended
Interferon- γ	HCV	Minimal if any effect	Not suitable for therapy
Farglitazar	NASH	No clear effect	Not suitable for therapy
ARBs	Miscellaneous	Minimal if any effect	May be acceptable

ARB, angiotensin-receptor blocker; PBC, primary biliary cirrhosis; PPC, polyenylphosphatidylcholine; SAM, S-adenosylmethionine; UDCA, ursodeoxycholic acid.

mental animal models.⁸⁴ Although it has been studied in a number of clinical trials,^{85–88} including in primary biliary cirrhosis, alcoholic cirrhosis, as well as in miscellaneous other liver diseases,⁸⁶ evidence supporting its effectiveness remains lacking.

Interleukin-10, an anti-inflammatory and immunomodulatory cytokine, can down-regulate production of proinflammatory cytokines, such as TNF- α , interleukin-1, and interleukin-2 from T cells. When administered to patients with HCV, interleukin-10 reduced hepatic inflammation and fibrosis scores (mean change from 5.0 ± 0.2 to 4.5 ± 0.3 ; $P < .05$). However, serum HCV-RNA levels increased during therapy and thus, this approach has not been pursued.

Several studies have shown that **interferon- γ** has potent inhibitory effects on stellate cells, inhibiting multiple aspects of stellate cell activation including fibrogenesis.^{47,89} A preliminary recent report in patients with chronic hepatitis C infection and fibrosis indicated that a subgroup of patients had an antifibrotic response.⁹⁰ However, a larger randomized study found that interferon- γ failed to have an antifibrotic effect in patients with HCV and advanced fibrosis, presumably because it enrolled patients with advanced cirrhosis and did not treat them for enough time.⁹¹

The **PPAR system** has gained considerable attention in the hepatic fibrogenesis field.^{92–94} PPAR γ in particular is reduced during stellate cell activation, and PPAR γ ligands inhibit activation and synthesis of extracellular matrix.^{92–94} Further, the adipocytokine, adiponectin, appears to have prominent antifibrotic actions, and the PPAR γ effects on stellate cells are, at least in part, adiponectin dependent.⁹⁵ Because of its added putative beneficial role in the metabolic syndrome, adiponectin is an attractive therapeutic target. Given the potential of PPAR γ agonists in the treatment of patients with fibrosis and preliminary studies that showed significant antifibrotic effects of the PPAR γ agonist, farglitazar, in animal models of fibrosis,⁹⁶ a large multicenter randomized trial of farglitazar in patients with HCV was performed.⁹⁷ This well-designed study showed that farglitazar therapy lasting 52 weeks failed to have an effect on stellate cell activation or fibrosis in this population.

Polyenylphosphatidylcholine contains a mixture of polyunsaturated phosphatidylcholines extracted from soybeans. Because of its presumed cytoprotective effect, it has been examined in human beings.⁹⁸ Unfortunately, in a major multicenter, prospective, randomized, double-blind, placebo-controlled trial of 789 alcoholic participants (average alcohol intake, 16 drinks/day) comparing either polyenylphosphatidylcholine or placebo for 2 years, there was no significant improvement in fibrosis. Of note, the majority of subjects reduced their ethanol consumption during the trial (presumably leading to an improvement in fibrosis in the control group).

Silymarin extract, derived from the milk thistle *Silybum marianum* (the major active component of which is silybinin), reduces lipid peroxidation and inhibits fibrogenesis in animal models.^{99–101} In human beings with fibrosis, the compound has had mixed effects.^{102,103} Thus, although silymarin appears to be safe, data supporting its use are lacking and further study is under way in patients with HCV (<http://ClinicalTrials.gov> identifier: NCT00680342) and NASH (<http://ClinicalTrials.gov> identifier: NCT00680407).

Ursodeoxycholic acid binds to hepatocyte membranes and appears to be cytoprotective, thereby reducing inflammation and thus fibrogenesis.¹⁰⁴ The aggregate data suggest that ur-

sodeoxycholic acid may impede progression of fibrosis in primary biliary cirrhosis via effects on bile duct inflammation, particularly if given early in the disease course.^{105,106} In a large randomized controlled trial of ursodeoxycholic acid in patients with NASH over a 2-year course, examining 107 subjects who had paired biopsy data, there was no improvement in fibrosis.¹⁰⁶ In aggregate, ursodeoxycholic acid is safe, and, although expensive, it is my belief that the available data justify its use at least in patients with primary biliary cirrhosis as an antifibrotic.

Vitamin E has gained a great deal of attention as a potential antifibrotic; it appears to be effective in animal models.¹⁰⁷ In human beings, vitamin E has had equivocal effects in patients with HCV¹⁰⁸ and alcoholic hepatitis.^{109,110} In patients with NASH, vitamin E led to reductions in aminotransferase levels, hepatic steatosis, and lobular inflammation, but failed to lead to an improvement in fibrosis.¹¹¹

A number of **herbal medicines** have been shown to have antifibrotic properties in animal models, and, in some, specific mechanisms have been identified.^{112–115} Herbal medicines with putative antiviral, anti-inflammatory, and antifibrotic effects are being used extensively in the Far East in patients with a variety of liver diseases.¹¹⁶ Medications containing herbs of the *Salvia* genus have been popular in particular as antifibrotics.¹¹⁶ Although human trials have suggested effectiveness of specific herbal medicines in some studies,¹¹⁶ data in peer-reviewed Western journals remain lacking. Because it is well appreciated that such herbal medicines may have significant toxicity, including hepatotoxicity,¹¹⁷ these medications should be used with caution.

The use of **anti-TNF- α** compounds in patients with alcoholic hepatitis is predicated on the rationale that TNF- α is up-regulated after alcohol-mediated hepatocellular injury (Figure 1), and thus these compounds should reduce inflammation, and resultant fibrosis. Although early studies suggested an improvement in inflammation,^{118–121} further larger studies have revealed that their use was associated with an increase in the risk of serious infection¹²² and mortality.¹²³ Pentoxifylline appears to reduce TNF- α expression and also may have primary antifibrotic effects.^{124,125} Although data suggest an effect on certain clinical outcomes,^{118,126} definitive evidence of an antifibrotic effect in human beings is lacking.

Malotilate, penicillamine, methotrexate, S-adenosylmethionine, and propylthiouracil all have shown some degree of anti-inflammatory and/or cytoprotective effects (presumably through their antioxidant properties) and, as such, may have an effect on fibrogenesis.^{127–129} However, evidence of an effect on fibrosis is equivocal at best.^{130–138} It is important to emphasize that for many of these human studies, subjects with alcoholic hepatitis and liver injury were examined, and in these studies fibrosis was not typically measured as a specific outcome. Thus, it may not be entirely appropriate to consider these agents as primary antifibrotics, but rather as compounds that could have secondary effects on fibrogenesis as a result of other properties.

Novel Approaches

A number of novel approaches to treat liver fibrosis exist. These include novel mechanisms of targeting the liver, such as the use of small interfering RNA^{139,140} or specific targeting systems.^{69,141} For example, TGF- β is well known to play a central role in the fibrogenic cascade and therefore is an important therapeutic target. Multiple proof-of-concept studies

Table 4. Potential Antifibrotic Targets

Agent or system	Mechanism
Intestinal microbiota/ TLR4	TLR4 on multiple cells types, including stellate cells, activates inflammatory pathways
NRF2	Transcription factor whose downstream target genes play an important role in cellular antioxidant defense
Loxl2	Enzyme catalyzes the first step in the formation of cross-links in collagens and elastin
Adiponectin	244-amino-acid-long polypeptide regulating glucose levels as well as fatty acid breakdown that has direct effects on stellate cell fibrogenesis
Angiostatin/endostatin Endothelin	Endogenous angiogenesis inhibitors 21 amino acid potent vasoconstrictor that also stimulates stellate cell activation

Loxl2, lysyl oxidase homolog 2; NRF2, nuclear factor (erythroid-derived 2)-like 2; TLR4, Toll-like receptor 4.

have shown that its inhibition (through use of specific antibodies that immobilize active TGF- β or receptor antagonists) is likely to be effective in fibrosis.^{38,142,143} However, given its important role in the regulation of cell growth, global inhibition of TGF- β or similar agents that have widespread biological effects such as PDGF or endothelin-1 could be potentially harmful. Thus, it likely will be critical to localize biological effects to fibrogenic effector cells. Early studies have provided proof of concept of this approach for stellate cells.¹⁴⁴

Previous and exciting new pathophysiologic studies have pointed to further translational opportunities to treat fibrosis (Table 4). Given the central role of inflammation in chronic hepatic injury and the ensuing wound-healing process (Figure 1), it follows that bacterial products, particularly lipopolysaccharides, may be important pathogenically. New evidence suggests that the microbiota may be important in the pathogenesis of liver inflammation,¹⁴⁵ fibrosis,¹⁴⁶ and even the development of hepatocellular cancer.¹⁴⁷ In quiescent stellate cells, Toll-like receptor 4 (a major lipopolysaccharide receptor) activation not only up-regulates chemokine secretion (further driving inflammation), but it also down-regulates the TGF- β pseudoreceptor Bambi, which in turn sensitizes stellate cells to TGF- β -induced signaling.¹⁴⁸ In another study, liver injury was associated with early onset of increased intestinal permeability and bacterial translocation that preceded changes in the microbiota.¹⁴⁹ Changes in the microbiota also have been associated with fibrosis progression.¹⁴⁶ As such, manipulation of the intestinal flora may be an innovative approach to antifibrotic therapy.

MicroRNAs have become recognized as being important in gene regulation. Recent evidence has suggested that a number of microRNAs are involved in the pathogenesis of different forms of organ fibrosis,¹⁵⁰ in stellate cell function, and in liver fibrosis,^{151,152} and therefore may represent novel therapeutic targets.

A variety of other systems also are attractive. Among these include those related to collagen synthesis, such as the lysyl oxidase system. Inhibition of this copper-dependent extracellular enzyme that catalyzes lysine-derived cross-links in collagen and elastin could abrogate tissue fibrosis.^{153,154} Angiogenic

pathways appear to be important in fibrosis, including the liver, and thus, interruption of this pathway could be an effective treatment approach. For example, a short peptide derived from endostatin, a naturally occurring 20-kilodalton C-terminal fragment derived from type 18 collagen, appeared to have potent antifibrotic activity in skin and pulmonary fibrosis *in vivo*.¹⁵⁵ Nuclear factor (erythroid-derived 2)-like 2, a transcription factor that appears to activate a number of genes involved in oxidative stress response, appears to have protective effects for fibrosis.^{156,157} In addition, compounds such as pirfenidone¹⁵⁸ and 5'-lipoxygenase inhibitors¹⁵⁹ appear to have direct effects on stellate cells and/or *in vivo* effects in hepatic fibrogenesis. Although there has been much interest in manipulating the balance between matrix synthesis and degradation via stimulation of collagen-degrading metalloproteases, or dampening the effect of metalloprotease inhibitors, this area remains largely open. Finally, emerging data suggest that coffee, and one of its ingredients, such as caffeine or another biologically active ingredient(s) (ie, such as cafestol or kahweol), protects against liver fibrosis. This area would be an obvious potential target given the prevalence of coffee consumption in many countries.

Vascular biological systems are intriguing because they potentially could have beneficial effects both for fibrosis and for portal hypertension. Stellate cells express angiotensin II and endothelin receptors and stimulation of these receptors with their cognate ligands leads to prominent stellate cell effects.⁴⁵

Challenges in Developing Antifibrotic Therapy

Currently, a potent and effective antifibrotic drug or agent is not available. This is likely the result of several factors, highlighted later. In addition, to develop a highly effective antifibrotic agent, several key features, as highlighted, are important.

Diagnosis/monitoring of hepatic fibrosis and cirrhosis. Perhaps one of the most difficult challenges in the field of development of antifibrotic medications is monitoring the effectiveness of putative compounds. An ideal test would be one that is noninvasive and simple to perform, yet inexpensive. Currently, liver biopsy is considered the gold standard test for determining the extent and progression of fibrosis.¹⁶⁰ A quantitative measure of collagen content can be made by colorimetric assay of Sirius red in liver tissue or by analytical image quantitation of collagen-containing tissue.⁶ In addition, scoring systems have been developed¹⁶¹⁻¹⁶³ to quantitate fibrosis and to help standardize the interpretation of biopsy specimens among different centers; such systems are most useful for standardization and comparison of fibrosis in studies.

Unfortunately, liver biopsy, although considered the gold standard tool to assess fibrosis, is inexact. Not only is liver biopsy subject to interobserver variability, but sampling error may be important, as evidenced by studies examining liver samples from different regions of the liver.¹⁶⁴ In addition, liver biopsy also is associated with significant potential morbidity, including a significant risk of death.¹⁶⁰ Thus, noninvasive measures that can monitor fibrogenesis would be ideal.¹⁶⁵ Noninvasive tools used to assess fibrosis include radiographic tests,¹⁶⁶ combinations of routine laboratory tests,^{167,168} and specific serum markers.¹⁶⁹ In particular, serum marker panels, including several that use mathematic algorithms,^{167,168,170} have been emphasized. Although some of these may even have predictive

clinical value,^{171,172} they generally have proven to be of limited clinical use.

Finally, the field of molecular imaging is emerging. It is possible that effector cells such as stellate cells may be imaged to more precisely quantitate their activity and/or fibrogenic features.^{173,174}

Cell-specific targeting. As emphasized earlier, it would be ideal to localize therapy to only effector cells. This is particularly important for the targeting systems that have widespread biological effects such as, for example, TGF- β , PDGF, or endothelin-1. TGF- β , in particular, is an attractive target because it appears to be the most potent stimulator of fibrogenesis. However, given its important role in regulation of cell growth, and neoplasia, it is highly likely that its global inhibition would have undesirable effects. A number of studies have provided proof of concept that at least stellate cells can be targeted specifically. By taking advantage of the expression of the mannose 6-phosphate/insulin-like growth factor II receptor on stellate cells, it has been shown elegantly that mannose 6-phosphate-modified albumins conjugated to specific inhibitors or toxins reduced stellate cell-mediated fibrogenesis.^{144,175} Alternatively, it is possible that physical properties of activated stellate cells may be taken advantage of, and that stellate cells could be targeted with specialized liposomes or similar compounds.^{176–178}

Length of therapy. As emphasized earlier, fibrogenesis is a dynamic process that occurs over a period of time; advanced fibrosis typically develops over prolonged periods of time. Thus, it is likely that reversion of fibrosis also would be expected to occur over more prolonged periods of time. Most of the trials examining novel agents have been performed over relatively short periods of time, typically over 6 or 12 months. To see meaningful regression of fibrosis, it is likely that a trial will require longer than 1 year, and perhaps longer than 2 years.

End points. The most appropriate end point for a novel treatment is a signal that the compound has antifibrotic effects. Notwithstanding this point, trials to date have used histologic assessment. This means that it is likely that the agent to be tested must be effective enough to cause a change in histology. It may be more appropriate to use a marker or set of markers that detect a fibrogenic signal. For example, serum markers assessed over time may be acceptable. In addition, some investigators have suggested that an antifibrotic agent should have an effect on clinical outcomes. This would require a prolonged treatment, which would make the likelihood of developing an effective agent difficult.

Summary and Future Directions

The pathogenesis of hepatic fibrogenesis is now better understood than ever before. The central event in fibrogenesis appears to be activation of effector cells, most prominently hepatic stellate cells. Stellate cell activation is characterized by many important features including, prominently, enhanced matrix synthesis and a contractile phenotype. The activation process is complex, leading to multiple potential sites for therapeutic interventions. A further critical concept is that the

fibrogenic lesion, in particular, the extracellular matrix, is a dynamic structure; even advanced fibrosis may be reversible. These data have helped spawn interest in the development of therapeutic antifibrotics. Notwithstanding this discovery, the most effective therapy for hepatic fibrogenesis is removal of the underlying disease process. Although a number of challenges exist, including in the area of cell-specific targeting, fibrosis monitoring, and execution of suitable clinical trials, the prospects for translation of the basic pathophysiology to therapy are bright. As for specific therapy directed primarily at the fibrotic lesion, the most effective therapies most likely will be directed fibrogenic effectors, in most cases hepatic stellate cells. In aggregate, although specific, effective, safe, and inexpensive antifibrotic therapies are not yet currently available, multiple potential targets have been identified, and one or more likely will emerge soon.

Supplementary Material

Note: The first 5 references associated with this article are available below in print. The remaining references accompanying this article are available online only with the electronic version of the article. To access the supplementary material accompanying this article, visit the online version of *Clinical Gastroenterology and Hepatology* at www.cghjournal.org, and at doi:10.1016/j.cgh.2013.01.005.

References

1. Jarnagin WR, Rockey DC, Koteliensky VE, et al. Expression of variant fibronectins in wound healing: cellular source and biological activity of the EIIIA segment in rat hepatic fibrogenesis. *J Cell Biol* 1994;127:2037–2048.
2. Schuppan D. Structure of the extracellular matrix in normal and fibrotic liver: collagens and glycoproteins. *Semin Liver Dis* 1990; 10:1–10.
3. Rockey DC. The cell and molecular biology of hepatic fibrogenesis. Clinical and therapeutic implications. *Clin Liver Dis* 2000; 4:319–355.
4. Rockey DC. Antifibrotic therapy in chronic liver disease. *Clin Gastroenterol Hepatol* 2005;3:95–107.
5. Iredale JP, Benyon RC, Pickering J, et al. Mechanisms of spontaneous resolution of rat liver fibrosis. Hepatic stellate cell apoptosis and reduced hepatic expression of metalloproteinase inhibitors. *J Clin Invest* 1998;102:538–549.

Reprint requests

Address requests for reprints to: Don C. Rockey, MD, Department of Internal Medicine, Medical University of South Carolina, 96 Jonathan Lucas Street, Suite 803, Charleston, South Carolina 29425. e-mail: roockey@musc.edu; fax: (843) 792-5265.

Conflicts of interest

The author discloses the following: The author has received a research grant from Actelion Pharmaceuticals.

Funding

Supported by the National Institutes of Health (R01 DK 57830).

References

6. Rockey DC, Chung JJ. Endothelin antagonism in experimental hepatic fibrosis. Implications for endothelin in the pathogenesis of wound healing. *J Clin Invest* 1996;98:1381–1388.
7. Lai CL, Chien RN, Leung NW, et al. A one-year trial of lamivudine for chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *N Engl J Med* 1998;339:61–68.
8. Poynard T, McHutchison J, Manns M, et al. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology* 2002;122:1303–1313.
9. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med* 2003;348:800–807.
10. Ellis EL, Mann DA. Clinical evidence for the regression of liver fibrosis. *J Hepatol* 2012;56:1171–1180.
11. Bruno S, Crosignani A, Facciotto C, et al. Sustained virologic response prevents the development of esophageal varices in compensated, Child-Pugh class A hepatitis C virus-induced cirrhosis. A 12-year prospective follow-up study. *Hepatology* 2010;51:2069–2076.
12. Mallet V, Gilgenkrantz H, Serpaggi J, et al. Brief communication: the relationship of regression of cirrhosis to outcome in chronic hepatitis C. *Ann Intern Med* 2008;149:399–403.
13. Czaja AJ, Carpenter HA. Decreased fibrosis during corticosteroid therapy of autoimmune hepatitis. *J Hepatol* 2004;40:646–652.
14. Liaw YF, Sung JJ, Chow WC, et al. Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 2004;351:1521–1531.
15. Hammel P, Couvelard A, O'Toole D, et al. Regression of liver fibrosis after biliary drainage in patients with chronic pancreatitis and stenosis of the common bile duct. *N Engl J Med* 2001;344:418–423.
16. Wanless IR, Nakashima E, Sherman M. Regression of human cirrhosis. Morphologic features and the genesis of incomplete septal cirrhosis. *Arch Pathol Lab Med* 2000;124:1599–1607.
17. Dufour JF, DeLellis R, Kaplan MM. Reversibility of hepatic fibrosis in autoimmune hepatitis. *Ann Intern Med* 1997;127:981–985.
18. Rockey DC. Hepatic fibrosis, stellate cells, and portal hypertension. *Clin Liver Dis* 2006;10:459–479, vii–viii.
19. Li Z, Dranoff JA, Chan EP, et al. Transforming growth factor-beta and substrate stiffness regulate portal fibroblast activation in culture. *Hepatology* 2007;46:1246–1256.
20. Kisseleva T, Uchinami H, Feirt N, et al. Bone marrow-derived fibrocytes participate in pathogenesis of liver fibrosis. *J Hepatol* 2006;45:429–438.
21. Kalluri R, Neilson EG. Epithelial-mesenchymal transition and its implications for fibrosis. *J Clin Invest* 2003;112:1776–1784.
22. Kaimori A, Potter J, Kaimori JY, et al. Transforming growth factor-beta1 induces an epithelial-to-mesenchymal transition state in mouse hepatocytes in vitro. *J Biol Chem* 2007;282:22089–22101.
23. Wake K. Perisinusoidal stellate cells (fat-storing cells, interstitial cells, lipocytes), their related structure in and around the liver sinusoids, and vitamin A-storing cells in extrahepatic organs. *Int Rev Cytol* 1980;66:303–353.
24. Blomhoff R, Wake K. Perisinusoidal stellate cells of the liver: important roles in retinol metabolism and fibrosis. *FASEB J* 1991;5:271–277.
25. Maher JJ, McGuire RF. Extracellular matrix gene expression increases preferentially in rat lipocytes and sinusoidal endothelial cells during hepatic fibrosis in vivo. *J Clin Invest* 1990;86:1641–1648.
26. Rockey DC, Boyles JK, Gabbiani G, et al. Rat hepatic lipocytes express smooth muscle actin upon activation in vivo and in culture. *J Submicrosc Cytol Pathol* 1992;24:193–203.
27. Hinz B. Masters and servants of the force: the role of matrix adhesions in myofibroblast force perception and transmission. *Eur J Cell Biol* 2006;85:175–181.
28. Gabbiani G. The myofibroblast in wound healing and fibrocontractive diseases. *J Pathol* 2003;200:500–503.
29. Aleffi S, Petrai I, Bertolani C, et al. Upregulation of proinflammatory and proangiogenic cytokines by leptin in human hepatic stellate cells. *Hepatology* 2005;42:1339–1348.
30. Santamato A, Fransvea E, Dituri F, et al. Hepatic stellate cells stimulate HCC cell migration via laminin-5 production. *Clin Sci (Lond)* 2011;121:159–168.
31. Gentilini A, Rombouts K, Galastri S, et al. Role of the stromal-derived factor-1 (SDF-1)-CXCR4 axis in the interaction between hepatic stellate cells and cholangiocarcinoma. *J Hepatol* 2012;57:813–820.
32. Bonacchi A, Petrai I, Defranco RM, et al. The chemokine CCL21 modulates lymphocyte recruitment and fibrosis in chronic hepatitis C. *Gastroenterology* 2003;125:1060–1076.
33. Sanderson N, Factor V, Nagy P, et al. Hepatic expression of mature transforming growth factor beta 1 in transgenic mice results in multiple tissue lesions. *Proc Natl Acad Sci U S A* 1995;92:2572–2576.
34. Hellerbrand C, Stefanovic B, Giordano F, et al. The role of TGFbeta1 in initiating hepatic stellate cell activation in vivo. *J Hepatol* 1999;30:77–87.
35. Schnabl B, Kweon YO, Frederick JP, et al. The role of Smad3 in mediating mouse hepatic stellate cell activation. *Hepatology* 2001;34:89–100.
36. Friedman SL, Maher JJ, Bissell DM. Mechanisms and therapy of hepatic fibrosis: report of the AASLD Single Topic Basic Research Conference. *Hepatology* 2000;32:1403–1408.
37. Bissell DM, Wang SS, Jamagin WR, et al. Cell-specific expression of transforming growth factor-beta in rat liver. Evidence for autocrine regulation of hepatocyte proliferation. *J Clin Invest* 1995;96:447–455.
38. Yata Y, Gotwals P, Koteliensky V, et al. Dose-dependent inhibition of hepatic fibrosis in mice by a TGF-beta soluble receptor: implications for antifibrotic therapy. *Hepatology* 2002;35:1022–1030.
39. Khimji AK, Shao R, Rockey DC. Divergent transforming growth factor-beta signaling in hepatic stellate cells after liver injury: functional effects on ECE-1 regulation. *Am J Pathol* 2008;173:716–727.
40. Huang G, Brigstock DR. Regulation of hepatic stellate cells by connective tissue growth factor. *Front Biosci* 2012;17:2495–2507.
41. Tong Z, Chen R, Alt DS, et al. Susceptibility to liver fibrosis in mice expressing a connective tissue growth factor transgene in hepatocytes. *Hepatology* 2009;50:939–947.
42. Saxena NK, Saliba G, Floyd JJ, et al. Leptin induces increased alpha2(I) collagen gene expression in cultured rat hepatic stellate cells. *J Cell Biochem* 2003;89:311–320.
43. Bataller R, Schwabe RF, Choi YH, et al. NADPH oxidase signal transduces angiotensin II in hepatic stellate cells and is critical in hepatic fibrosis. *J Clin Invest* 2003;112:1383–1394.
44. Bataller R, Gábele E, Parsons CJ, et al. Systemic infusion of angiotensin II exacerbates liver fibrosis in bile duct-ligated rats. *Hepatology* 2005;41:1046–1055.
45. Rockey DC. Vascular mediators in the injured liver. *Hepatology* 2003;37:4–12.
46. Oben JA, Roskams T, Yang S, et al. Hepatic fibrogenesis requires sympathetic neurotransmitters. *Gut* 2004;53:438–445.
47. Rockey DC, Chung JJ. Interferon gamma inhibits lipocyte activation and extracellular matrix mRNA expression during experimental liver injury: implications for treatment of hepatic fibrosis. *J Invest Med* 1994;42:660–670.
48. Inagaki Y, Nemoto T, Kushida M, et al. Interferon alfa down-

- regulates collagen gene transcription and suppresses experimental hepatic fibrosis in mice. *Hepatology* 2003;38:890–899.
49. Kamada Y, Tamura S, Kiso S, et al. Enhanced carbon tetrachloride-induced liver fibrosis in mice lacking adiponectin. *Gastroenterology* 2003;125:1796–1807.
 50. Ueki T, Kaneda Y, Tsutsui H, et al. Hepatocyte growth factor gene therapy of liver cirrhosis in rats. *Nat Med* 1999;5:226–230.
 51. Kawada N, Kristensen DB, Asahina K, et al. Characterization of a stellate cell activation-associated protein (STAP) with peroxidase activity found in rat hepatic stellate cells. *J Biol Chem* 2001;276:25318–25323.
 52. Friedman SL, Roll FJ, Boyles J, et al. Maintenance of differentiated phenotype of cultured rat hepatic lipocytes by basement membrane matrix. *J Biol Chem* 1989;264:10756–10762.
 53. Gaca MDA, Georges P, Jammey P, et al. Matrix compliance determines hepatic stellate cell phenotype. *Hepatology* 2003;38:776A.
 54. Olsen AL, Bloomer SA, Chan EP, et al. Hepatic stellate cells require a stiff environment for myofibroblastic differentiation. *Am J Physiol Gastrointest Liver Physiol* 2011;301:G110–G118.
 55. Racine-Samson L, Rockey DC, Bissell DM. The role of $\alpha 1 \beta 1$ integrin in wound contraction. A quantitative analysis of liver myofibroblasts in vivo and in primary culture. *J Biol Chem* 1997;272:30911–30917.
 56. Shafiei MS, Rockey DC. The role of integrin-linked kinase in liver wound healing. *J Biol Chem* 2006;281:24863–24872.
 57. Shafiei MS, Rockey DC. The function of integrin-linked kinase in normal and activated stellate cells: implications for fibrogenesis in wound healing. *Lab Invest* 2012;92:305–316.
 58. Arthur MJ. Fibrogenesis II. Metalloproteinases and their inhibitors in liver fibrosis. *Am J Physiol Gastrointest Liver Physiol* 2000;279:G245–G249.
 59. Han YP, Zhou L, Wang J, et al. Essential role of matrix metalloproteinases in interleukin-1-induced myofibroblastic activation of hepatic stellate cell in collagen. *J Biol Chem* 2004;279:4820–4828.
 60. Han YP. Matrix metalloproteinases, the pros and cons, in liver fibrosis. *J Gastroenterol Hepatol* 2006;21(Suppl 3):S88–S91.
 61. Garcia-Bañuelos J, Siller-Lopez F, Miranda A, et al. Cirrhotic rat livers with extensive fibrosis can be safely transduced with clinical-grade adenoviral vectors. Evidence of cirrhosis reversal. *Gene Ther* 2002;9:127–134.
 62. Xu G, Rockey DC. Regulation of smooth muscle myosin heavy chain isoforms in hepatic myofibroblasts during liver injury. *Hepatology* 1999;30:492A.
 63. Pinzani M, Failli P, Ruocco C, et al. Fat-storing cells as liver-specific pericytes. Spatial dynamics of agonist-stimulated intracellular calcium transients. *J Clin Invest* 1992;90:642–646.
 64. Rockey DC, Housset CN, Friedman SL. Activation-dependent contractility of rat hepatic lipocytes in culture and in vivo. *J Clin Invest* 1993;92:1795–1804.
 65. Iizuka M, Murata T, Hori M, et al. Increased contractility of hepatic stellate cells in cirrhosis is mediated by enhanced Ca^{2+} -dependent and Ca^{2+} -sensitization pathways. *Am J Physiol Gastrointest Liver Physiol* 2011;300:G1010–G1021.
 66. Xu H, Zhou Y, Lu C, et al. Salvianolic acid B lowers portal pressure in cirrhotic rats and attenuates contraction of rat hepatic stellate cells by inhibiting RhoA signaling pathway. *Lab Invest* 2012;92:1738–1748.
 67. Trebicka J, Hennenberg M, Schulze Pröbsting A, et al. Role of $\beta 3$ -adrenoceptors for intrahepatic resistance and portal hypertension in liver cirrhosis. *Hepatology* 2009;50:1924–1935.
 68. Zou WL, Yang Z, Zang YJ, et al. Inhibitory effects of prostaglandin E1 on activation of hepatic stellate cells in rabbits with schistosomiasis. *Hepatobiliary Pancreat Dis Int* 2007;6:176–181.
 69. van Beuge MM, Prakash J, Lacombe M, et al. Reduction of fibrogenesis by selective delivery of a Rho kinase inhibitor to hepatic stellate cells in mice. *J Pharmacol Exp Ther* 2011;337:628–635.
 70. Watanabe A, Sohail MA, Gautam S, et al. Adenine induces differentiation of rat hepatic stellate cells. *Dig Dis Sci* 2012;57:2371–2378.
 71. Li J, Kuruba R, Wilson A, et al. Inhibition of endothelin-1-mediated contraction of hepatic stellate cells by FXR ligand. *PLoS One* 2010;5:e13955.
 72. Wright MC, Issa R, Smart DE, et al. Gliotoxin stimulates the apoptosis of human and rat hepatic stellate cells and enhances the resolution of liver fibrosis in rats. *Gastroenterology* 2001;121:685–698.
 73. Canbay A, Taimr P, Torok N, et al. Apoptotic body engulfment by a human stellate cell line is profibrogenic. *Lab Invest* 2003;83:655–663.
 74. Krizhanovsky V, Yon M, Dickins RA, et al. Senescence of activated stellate cells limits liver fibrosis. *Cell* 2008;134:657–667.
 75. Deleve LD, Wang X, Guo Y. Sinusoidal endothelial cells prevent rat stellate cell activation and promote reversion to quiescence. *Hepatology* 2008;48:920–930.
 76. Thoen LF, Guimarães EL, Grunsven LA. Autophagy: a new player in hepatic stellate cell activation. *Autophagy* 2012;8:126–128.
 77. Thoen LF, Guimarães EL, Dollé L, et al. A role for autophagy during hepatic stellate cell activation. *J Hepatol* 2011;55:1353–1360.
 78. Hernández-Gea V, Friedman SL. Autophagy fuels tissue fibrogenesis. *Autophagy* 2012;8:849–850.
 79. Ramond MJ, Poynard T, Rueff B, et al. A randomized trial of prednisolone in patients with severe alcoholic hepatitis. *N Engl J Med* 1992;326:507–512.
 80. Spahr L, Rubbia-Brandt L, Pugin J, et al. Rapid changes in alcoholic hepatitis histology under steroids: correlation with soluble intercellular adhesion molecule-1 in hepatic venous blood. *J Hepatol* 2001;35:582–589.
 81. Powell LW, Kerr JF. Reversal of “cirrhosis” in idiopathic haemochromatosis following long-term intensive venesection therapy. *Australas Ann Med* 1970;19:54–57.
 82. Blumberg RS, Chopra S, Ibrahim R, et al. Primary hepatocellular carcinoma in idiopathic hemochromatosis after reversal of cirrhosis. *Gastroenterology* 1988;95:1399–1402.
 83. Neuschwander-Tetri BA, Brunt EM, Wehmeier KR, et al. Improved nonalcoholic steatohepatitis after 48 weeks of treatment with the PPAR- γ ligand rosiglitazone. *Hepatology* 2003;38:1008–1017.
 84. Rodríguez L, Cerbón-Ambríz J, Muñoz ML. Effects of colchicine and colchicine in a biochemical model of liver injury and fibrosis. *Arch Med Res* 1998;29:109–116.
 85. Kaplan MM, Alling DW, Zimmerman HJ, et al. A prospective trial of colchicine for primary biliary cirrhosis. *N Engl J Med* 1986;315:1448–1454.
 86. Kershenobich D, Vargas F, Garcia-Tsao G, et al. Colchicine in the treatment of cirrhosis of the liver. *N Engl J Med* 1988;318:1709–1713.
 87. Rambaldi A, Gluud C. Colchicine for alcoholic and non-alcoholic liver fibrosis or cirrhosis. *Liver* 2001;21:129–136.
 88. Morgan TR, Weiss DG, Nemchauskas B, et al. Colchicine treatment of alcoholic cirrhosis: a randomized, placebo-controlled clinical trial of patient survival. *Gastroenterology* 2005;128:882–890.
 89. Rockey DC, Maher JJ, Jarnagin WR, et al. Inhibition of rat hepatic

- lipocyte activation in culture by interferon-gamma. *Hepatology* 1992;16:776–784.
90. Muir AJ, Sylvestre PB, Rockey DC. Interferon gamma-1b for the treatment of fibrosis in chronic hepatitis C infection. *J Viral Hepat* 2006;13:322–328.
 91. Rockey DC. Current and future anti-fibrotic therapies for chronic liver disease. *Clin Liver Dis* 2008;12:939–962, xi.
 92. Miyahara T, Schrum L, Rippe R, et al. Peroxisome proliferator-activated receptors and hepatic stellate cell activation. *J Biol Chem* 2000;275:35715–35722.
 93. Hazra S, Xiong S, Wang J, et al. Peroxisome proliferator-activated receptor gamma induces a phenotypic switch from activated to quiescent hepatic stellate cells. *J Biol Chem* 2004;279:11392–11401.
 94. Yang L, Chan CC, Kwon OS, et al. Regulation of peroxisome proliferator-activated receptor-gamma in liver fibrosis. *Am J Physiol Gastrointest Liver Physiol* 2006;291:G902–G911.
 95. Shafiei MS, Shetty S, Scherer PE, et al. Adiponectin regulation of stellate cell activation via PPAR γ -dependent and -independent mechanisms. *Am J Pathol* 2011;178:2690–2699.
 96. Yang L, Stimpson SA, Chen L, et al. Effectiveness of the PPAR γ agonist, GW570, in liver fibrosis. *Inflamm Res* 2010;59:1061–1071.
 97. McHutchison J, Goodman Z, Patel K, et al. Farglitazar lacks antifibrotic activity in patients with chronic hepatitis C infection. *Gastroenterology* 2010;138:1365–1373e1–2.
 98. Lieber CS, Weiss DG, Groszmann R II, et al. Veterans Affairs Cooperative Study of polyenylphosphatidylcholine in alcoholic liver disease. *Alcohol Clin Exp Res* 2003;27:1765–1772.
 99. Boigk G, Stroedter L, Herbst H, et al. Silymarin retards collagen accumulation in early and advanced biliary fibrosis secondary to complete bile duct obliteration in rats. *Hepatology* 1997;26:643–649.
 100. Jia JD, Bauer M, Cho JJ, et al. Antifibrotic effect of silymarin in rat secondary biliary fibrosis is mediated by downregulation of procollagen alpha1(I) and TIMP-1. *J Hepatol* 2001;35:392–398.
 101. Lieber CS, Leo MA, Cao Q, et al. Silymarin retards the progression of alcohol-induced hepatic fibrosis in baboons. *J Clin Gastroenterol* 2003;37:336–339.
 102. Ferenci P, Dragosics B, Dittrich H, et al. Randomized controlled trial of silymarin treatment in patients with cirrhosis of the liver. *J Hepatol* 1989;9:105–113.
 103. Parés A, Planas R, Torres M, et al. Effects of silymarin in alcoholic patients with cirrhosis of the liver: results of a controlled, double-blind, randomized and multicenter trial. *J Hepatol* 1998;28:615–621.
 104. Nava-Ocampo AA, Suster S, Muriel P. Effect of colchicine and ursodeoxycholic acid on hepatocyte and erythrocyte membranes and liver histology in experimentally induced carbon tetrachloride cirrhosis in rats. *Eur J Clin Invest* 1997;27:77–84.
 105. Poupon RE, Lindor KD, Parés A, et al. Combined analysis of the effect of treatment with ursodeoxycholic acid on histologic progression in primary biliary cirrhosis. *J Hepatol* 2003;39:12–16.
 106. Lindor KD, Kowdley KV, Heathcote EJ, et al. Ursodeoxycholic acid for treatment of nonalcoholic steatohepatitis: results of a randomized trial. *Hepatology* 2004;39:770–778.
 107. Brown KE, Poulos JE, Li L, et al. Effect of vitamin E supplementation on hepatic fibrogenesis in chronic dietary iron overload. *Am J Physiol* 1997;272:G116–G123.
 108. Houghlum K, Venkataramani A, Lyche K, et al. A pilot study of the effects of d-alpha-tocopherol on hepatic stellate cell activation in chronic hepatitis C. *Gastroenterology* 1997;113:1069–1073.
 109. Mezey E, Potter JJ, Rennie-Tankersley L, et al. A randomized placebo controlled trial of vitamin E for alcoholic hepatitis. *J Hepatol* 2004;40:40–46.
 110. Stewart S, Prince M, Bassendine M, et al. A trial of antioxidant therapy alone or with corticosteroids in acute alcoholic hepatitis. *J Hepatol* 2002;36(Suppl):16.
 111. Sanyal AJ, Chalasani N, Kowdley KV, et al. Pioglitazone, vitamin E, or placebo for nonalcoholic steatohepatitis. *N Engl J Med* 2010;362:1675–1685.
 112. Sakaida I, Tsuchiya M, Kawaguchi K, et al. Herbal medicine Inchin-ko-to (TJ-135) prevents liver fibrosis and enzyme-altered lesions in rat liver cirrhosis induced by a choline-deficient L-amino acid-defined diet. *J Hepatol* 2003;38:762–769.
 113. Shimizu I, Ma YR, Mizobuchi Y, et al. Effects of Sho-saiko-to, a Japanese herbal medicine, on hepatic fibrosis in rats. *Hepatology* 1999;29:149–160.
 114. Zhang XL, Liu L, Jiang HQ. *Salvia miltiorrhiza* monomer IH764-3 induces hepatic stellate cell apoptosis via caspase-3 activation. *World J Gastroenterol* 2002;8:515–519.
 115. Chen ZX, Zhang SJ, Lao SX, et al. He Jie Tang in the treatment of chronic hepatitis B patients. *World J Gastroenterol* 2005;11:6638–6643.
 116. Wang BE. Treatment of chronic liver diseases with traditional Chinese medicine. *J Gastroenterol Hepatol* 2000;15(Suppl):E67–E70.
 117. Stedman C. Herbal hepatotoxicity. *Semin Liver Dis* 2002;22:195–206.
 118. Akriviadis E, Botla R, Briggs W, et al. Pentoxifylline improves short-term survival in severe acute alcoholic hepatitis: a double-blind, placebo-controlled trial. *Gastroenterology* 2000;119:1637–1648.
 119. Spahr L, Rubbia-Brandt L, Frossard JL, et al. Combination of steroids with infliximab or placebo in severe alcoholic hepatitis: a randomized controlled pilot study. *J Hepatol* 2002;37:448–455.
 120. Tilg H, Jalan R, Kaser A, et al. Anti-tumor necrosis factor-alpha monoclonal antibody therapy in severe alcoholic hepatitis. *J Hepatol* 2003;38:419–425.
 121. Menon KV, Stadheim L, Kamath PS, et al. A pilot study of the safety and tolerability of etanercept in patients with alcoholic hepatitis. *Am J Gastroenterol* 2004;99:255–260.
 122. Naveau S, Chollet-Martin S, Dharancy S, et al. A double-blind randomized controlled trial of infliximab associated with prednisolone in acute alcoholic hepatitis. *Hepatology* 2004;39:1390–1397.
 123. Boetticher NC, Peine CJ, Kwo P, et al. A randomized, double-blinded, placebo-controlled multicenter trial of etanercept in the treatment of alcoholic hepatitis. *Gastroenterology* 2008;135:1953–1960.
 124. Hernández E, Bucio L, Souza V, et al. Pentoxifylline downregulates alpha (I) collagen expression by the inhibition of I κ B α degradation in liver stellate cells. *Cell Biol Toxicol* 2008;24:303–314.
 125. Zhang D, Jiang H, Wang Y, et al. Pentoxifylline inhibits hepatic stellate cells proliferation via the Raf/ERK pathway. *APMIS* 2012;120:572–581.
 126. Sidhu SS, Goyal O, Singla M, et al. Pentoxifylline in severe alcoholic hepatitis: a prospective, randomised trial. *J Assoc Physicians India* 2012;60:20–22.
 127. Schaff Z, Lapis K, Szende B, et al. The effect of d-penicillamine on CCl $_4$ -induced experimental liver cirrhosis. *Exp Pathol* 1991;43:111–120.
 128. Lu SC, Tsukamoto H, Mato JM. Role of abnormal methionine metabolism in alcoholic liver injury. *Alcohol* 2002;27:155–162.
 129. Aithal GP, Haugk B, Das S, et al. Monitoring methotrexate-induced hepatic fibrosis in patients with psoriasis: are serial liver biopsies justified? *Aliment Pharmacol Ther* 2004;19:391–399.
 130. Anonymous. The results of a randomized double blind con-

- trolled trial evaluating malotilate in primary biliary cirrhosis. A European Multicentre Study Group. *J Hepatol* 1993;17:227–235.
131. Bodenheimer HC Jr, Schaffner F, Sternlieb I, et al. A prospective clinical trial of D-penicillamine in the treatment of primary biliary cirrhosis. *Hepatology* 1985;5:1139–1142.
 132. Dickson ER, Fleming TR, Wiesner RH, et al. Trial of penicillamine in advanced primary biliary cirrhosis. *N Engl J Med* 1985;312:1011–1015.
 133. Kaplan MM, DeLellis RA, Wolfe HJ. Sustained biochemical and histologic remission of primary biliary cirrhosis in response to medical treatment. *Ann Intern Med* 1997;126:682–688.
 134. Hendrickse MT, Rigney E, Gjafer MH, et al. Low-dose methotrexate is ineffective in primary biliary cirrhosis: long-term results of a placebo-controlled trial. *Gastroenterology* 1999;117:400–407.
 135. Bach N, Bodian C, Bodenheimer H, et al. Methotrexate therapy for primary biliary cirrhosis. *Am J Gastroenterol* 2003;98:187–193.
 136. Kaplan MM, Cheng S, Price LL, et al. A randomized controlled trial of colchicine plus ursodiol versus methotrexate plus ursodiol in primary biliary cirrhosis: ten-year results. *Hepatology* 2004;39:915–923.
 137. Mato JM, Cámara J, Fernández de Paz J, et al. S-adenosylmethionine in alcoholic liver cirrhosis: a randomized, placebo-controlled, double-blind, multicenter clinical trial. *J Hepatol* 1999;30:1081–1089.
 138. Rambaldi A, Gluud C. Meta-analysis of propylthiouracil for alcoholic liver disease—a Cochrane Hepato-Biliary Group Review. *Liver* 2001;21:398–404.
 139. Novobrantseva TI, Akinc A, Borodovsky A, et al. Delivering silence: advancements in developing siRNA therapeutics. *Curr Opin Drug Discov Devel* 2008;11:217–224.
 140. Vaishnaw AK, Gollob J, Gamba-Vitalo C, et al. A status report on RNAi therapeutics. *Silence* 2010;1:14.
 141. Bisht S, Khan MA, Bekhit M, et al. A polymeric nanoparticle formulation of curcumin (NanoCurc™) ameliorates CCl₄-induced hepatic injury and fibrosis through reduction of pro-inflammatory cytokines and stellate cell activation. *Lab Invest* 2011;91:1383–1395.
 142. Isaka Y, Brees DK, Ikegaya K, et al. Gene therapy by skeletal muscle expression of decorin prevents fibrotic disease in rat kidney. *Nat Med* 1996;2:418–423.
 143. George J, Roulot D, Kotliansky VE, et al. In vivo inhibition of rat stellate cell activation by soluble transforming growth factor beta type II receptor: a potential new therapy for hepatic fibrosis. *Proc Natl Acad Sci U S A* 1999;96:12719–12724.
 144. Gonzalo T, Beljaars L, van de Bovenkamp M, et al. Local inhibition of liver fibrosis by specific delivery of a platelet-derived growth factor kinase inhibitor to hepatic stellate cells. *J Pharmacol Exp Ther* 2007;321:856–865.
 145. Seki E, Schnabl B. Role of innate immunity and the microbiota in liver fibrosis: crosstalk between the liver and gut. *J Physiol* 2012;590:447–458.
 146. Gómez-Hurtado I, Santacruz A, Peiró G, et al. Gut microbiota dysbiosis is associated with inflammation and bacterial translocation in mice with CCl₄-induced fibrosis. *PLoS One* 2011;6:e23037.
 147. Dapito DH, Mencin A, Gwak GY, et al. Promotion of hepatocellular carcinoma by the intestinal microbiota and TLR4. *Cancer Cell* 2012;21:504–516.
 148. Seki E, De Minicis S, Osterreicher CH, et al. TLR4 enhances TGF-beta signaling and hepatic fibrosis. *Nat Med* 2007;13:1324–1332.
 149. Fouts DE, Torralba M, Nelson KE, et al. Bacterial translocation and changes in the intestinal microbiome in mouse models of liver disease. *J Hepatol* 2012;56:1283–1292.
 150. Jiang X, Tsitsiou E, Herrick SE, et al. MicroRNAs and the regulation of fibrosis. *FEBS J* 2010;277:2015–2021.
 151. Sekiya Y, Ogawa T, Yoshizato K, et al. Suppression of hepatic stellate cell activation by microRNA-29b. *Biochem Biophys Res Commun* 2011;412:74–79.
 152. Lakner AM, Steuerwald NM, Walling TL, et al. Inhibitory effects of microRNA 19b in hepatic stellate cell-mediated fibrogenesis. *Hepatology* 2012;56:300–310.
 153. Barry-Hamilton V, Spangler R, Marshall D, et al. Allosteric inhibition of lysyl oxidase-like-2 impedes the development of a pathologic microenvironment. *Nat Med* 2010;16:1009–1017.
 154. López B, González A, Hermida N, et al. Role of lysyl oxidase in myocardial fibrosis: from basic science to clinical aspects. *Am J Physiol Heart Circ Physiol* 2010;299:H1–H9.
 155. Yamaguchi Y, Takihara T, Chambers RA, et al. A peptide derived from endostatin ameliorates organ fibrosis. *Sci Transl Med* 2012;4:136ra71.
 156. Xu W, Hellerbrand C, Köhler UA, et al. The Nrf2 transcription factor protects from toxin-induced liver injury and fibrosis. *Lab Invest* 2008;88:1068–1078.
 157. Kim IH, Kim DG, Hao P, et al. Anti-fibrotic effects of L-2-oxothiazolidine-4-carboxylic acid via modulation of nuclear factor erythroid 2-related factor 2 in rats. *BMB Rep* 2012;45:348–353.
 158. Zhao XY, Zeng X, Li XM, et al. Pirfenidone inhibits carbon tetrachloride- and albumin complex-induced liver fibrosis in rodents by preventing activation of hepatic stellate cells. *Clin Exp Pharmacol Physiol* 2009;36:963–968.
 159. Titos E, Clària J, Planagumà A, et al. Inhibition of 5-lipoxygenase induces cell growth arrest and apoptosis in rat Kupffer cells: implications for liver fibrosis. *FASEB J* 2003;17:1745–1747.
 160. Rockey DC, Caldwell SH, Goodman ZD, et al. Liver biopsy. *Hepatology* 2009;49:1017–1044.
 161. Pilette C, Rousselet MC, Bedossa P, et al. Histopathological evaluation of liver fibrosis: quantitative image analysis vs semi-quantitative scores. Comparison with serum markers. *J Hepatol* 1998;28:439–446.
 162. O'Brien MJ, Keating NM, Elderiny S, et al. An assessment of digital image analysis to measure fibrosis in liver biopsy specimens of patients with chronic hepatitis C. *Am J Clin Pathol* 2000;114:712–718.
 163. Bedossa P, Dargère D, Paradis V. Sampling variability of liver fibrosis in chronic hepatitis C. *Hepatology* 2003;38:1449–1457.
 164. Regev A, Berho M, Jeffers LJ, et al. Sampling error and intraobserver variation in liver biopsy in patients with chronic HCV infection. *Am J Gastroenterol* 2002;97:2614–2618.
 165. Rockey DC, Bissell DM. Noninvasive measures of liver fibrosis. *Hepatology* 2006;43:S113–S120.
 166. Colli A, Fraquelli M, Andreoletti M, et al. Severe liver fibrosis or cirrhosis: accuracy of US for detection—analysis of 300 cases. *Radiology* 2003;227:89–94.
 167. Imbert-Bismut F, Ratzin V, Pieroni L, et al. Biochemical markers of liver fibrosis in patients with hepatitis C virus infection: a prospective study. *Lancet* 2001;357:1069–1075.
 168. Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology* 2003;38:518–526.
 169. Callewaert N, Van Vlierberghe H, Van Hecke A, et al. Noninvasive diagnosis of liver cirrhosis using DNA sequencer-based total serum protein glycomics. *Nat Med* 2004;10:429–434.
 170. Parkes J, Guha IN, Roderick P, et al. Performance of serum marker panels for liver fibrosis in chronic hepatitis C. *J Hepatol* 2006;44:462–474.
 171. Mayo MJ, Parkes J, Adams-Huet B, et al. Prediction of clinical outcomes in primary biliary cirrhosis by serum enhanced liver fibrosis assay. *Hepatology* 2008;48:1549–1557.

172. Parkes J, Roderick P, Harris S, et al. Enhanced liver fibrosis test can predict clinical outcomes in patients with chronic liver disease. *Gut* 2010;59:1245–1251.
173. Li F, Song Z, Li Q, et al. Molecular imaging of hepatic stellate cell activity by visualization of hepatic integrin $\alpha v \beta 3$ expression with SPECT in rat. *Hepatology* 2011;54:1020–1030.
174. Wang QB, Han Y, Jiang TT, et al. MR imaging of activated hepatic stellate cells in liver injured by CCl₄ of rats with integrin-targeted ultrasmall superparamagnetic iron oxide. *Eur Radiol* 2011;21:1016–1025.
175. Hagens WI, Beljaars L, Mann DA, et al. Cellular targeting of the apoptosis-inducing compound gliotoxin to fibrotic rat livers. *J Pharmacol Exp Ther* 2008;324:902–910.
176. Park K, Hong SW, Hur W, et al. Target specific systemic delivery of TGF- β siRNA/(PEI-SS)-g-HA complex for the treatment of liver cirrhosis. *Biomaterials* 2011;32:4951–4958.
177. Poelstra K, Schuppan D. Targeted therapy of liver fibrosis/cirrhosis and its complications. *J Hepatol* 2011;55:726–728.
178. Li Q, Yan Z, Li F, et al. The improving effects on hepatic fibrosis of interferon- γ liposomes targeted to hepatic stellate cells. *Nanotechnology* 2012;23:265101.