

Connective tissue growth factor (CTGF/CCN2) in hepatic fibrosis

Amy W. Rachfal^{a,c,d}, David R. Brigstock^{a,b,c,d,*}

^a Department of Surgery, The Ohio State University, Columbus, OH 43212, USA

^b Department of Molecular and Cellular Biochemistry, The Ohio State University, Columbus, OH 43212, USA

^c Molecular, Cellular and Developmental Biology Program, The Ohio State University, Columbus, OH 43212, USA

^d Center for Cell and Vascular Biology, Children's Research Institute, 700 Children's Drive, Columbus, OH 43205, USA

Received 24 December 2002; received in revised form 16 February 2003; accepted 18 February 2003

Abstract

Connective tissue growth factor (CTGF/CCN2) is a highly profibrogenic molecule which is overexpressed in many fibrotic lesions, including those of the liver. CTGF/CCN2 is transcriptionally activated by transforming growth factor-beta (TGF- β) and appears to mediate some of the extracellular matrix (ECM)-inducing properties that have been previously attributed to TGF- β . CTGF/CCN2 and TGF- β stimulate connective tissue cell proliferation and ECM synthesis in vitro and exhibit shared fibrogenic and angiogenic properties in vivo. In fibrotic liver, CTGF/CCN2 mRNA and protein are produced by fibroblasts, myofibroblasts, hepatic stellate cells (HSCs), endothelial cells, and bile duct epithelial cells. CTGF/CCN2 is also produced at high levels in hepatocytes during cytochrome P-450E1-mediated ethanol oxidation. CTGF/CCN2 expression in cultured HSCs is enhanced following their activation or stimulation by TGF- β while exogenous CTGF/CCN2 is able to promote HSC adhesion, proliferation, locomotion, and collagen production. Collectively, these data suggest that during initiating or downstream fibrogenic events in the liver, production of CTGF/CCN2 is regulated primarily by TGF- β in one or more cell types and that CTGF/CCN2 plays important roles in HSC activation and progression of fibrosis. This article reviews the data that support the importance of CTGF/CCN2 in hepatic fibrosis and highlights the concept that CTGF/CCN2 may represent a new therapeutic target in this disease.

© 2003 Elsevier Science B.V. All rights reserved.

Keywords: CTGF/CCN2; TGF- β ; Hepatic stellate cell; Extracellular matrix; Matricellular; CCN

1. Introduction

Connective tissue growth factor (CTGF/CCN2) is a multi-functional matricellular protein that is produced by a variety of cell types and acts via autocrine and paracrine circuits to regulate many cellular functions including growth, proliferation, apoptosis, adhesion, migration, extracellular matrix (ECM) production, and differentiation [1–4]. CTGF/CCN2 appears to play a role in diverse biological processes including angiogenesis, chondrogenesis, embryogenesis, implantation, development, ovarian function, tumorigenesis and fibrosis. CTGF/CCN2 belongs to the CTGF/CYR61/NOV (CCN) family that currently comprises six members

that are 30–40 kDa cysteine-rich mosaic proteins [1–4]. CCN proteins likely exert many of their biological effects by direct binding interactions with components of the ECM and the cell surface.

Of the many functions ascribed to CTGF/CCN2, that of its ability to promote fibrosis has attracted considerable attention and has been the major driving force behind many of the studies published to date [5,6]. This area of focus arose following the demonstration that CTGF/CCN2 was induced by TGF- β (itself highly fibrogenic) and that fibroblasts were targets for CTGF/CCN2 action [7–10]. CTGF/CCN2 regulates proliferation, migration, adhesion and production of ECM molecules in cultured fibroblasts [10–14], while subcutaneous injection of CTGF/CCN2 into neonatal mice causes a rapid increase in the amount of granulation tissue comprising connective tissue cells and abundant ECM [13,15]. Additionally, CTGF/CCN2 was shown to be present and frequently over-expressed and

* Corresponding author. Tel.: +1-614-722-2840; fax: +1-614-722-2007.

E-mail address: brigstod@pediatrics.ohio-state.edu (D.R. Brigstock).

co-expressed with TGF- β in fibrotic skin disorders such as systemic sclerosis, localized skin sclerosis, keloids, scar tissue, eosinic fasciitis, nodular fasciitis, and Dupuytren's contracture [16–21]. CTGF/CCN2 mRNA and/or protein are over-expressed in fibrotic lesions of major organs and tissues including the liver [22–28], kidney [29–37], lung [18,38–40], cardiovascular system [41–43], pancreas [44–46], bowel [47], eye [48–51], and gingiva [52,53]. CTGF/CCN2 is also over-expressed in the stromal compartment of melanoma as well as mammary, pancreatic and fibrohistiocytic tumors that are characterized as having significant connective tissue involvement [54–57]. These studies have led to the notion that CTGF/CCN2 may be a new molecular target for therapeutic intervention in fibrotic diseases [5,6,58,59].

2. The link between TGF- β and CTGF/CCN2

Early studies using fibroblasts established that CTGF/CCN2 is a TGF- β 1-induced immediate early gene [9,10]. Expression of CTGF/CCN2 following TGF- β stimulation occurs within minutes in the absence of de novo protein synthesis, an effect that was initially attributed to a TGF- β response element in the CTGF/CCN2 promoter [60]. However, more recent studies have demonstrated that while this element may be important for basal CTGF/CCN2 expression, TGF- β -induced CTGF/CCN2 expression requires Smad-binding sequences and a novel promoter sequence which is preferentially activated in fibroblasts as compared with epithelial cells [61–63].

The intimate relationship between TGF- β action and CTGF/CCN2 expression has been the basis for examining the role played by CTGF/CCN2 in mediating the biological activities of TGF- β [64]. While in the long run, this may prove to be a somewhat narrow view of the role of CTGF/CCN2, the outcome of these studies has been the demonstration that TGF- β and CTGF/CCN2 share pro-fibrogenic properties whereas anti-inflammatory and immunosuppressive properties are unique to TGF- β . TGF- β 1 and CTGF/CCN2 are both over-expressed in many fibrotic diseases and are fibrogenic in vitro and in vivo [5]. TGF- β -induced collagen production is antagonized by CTGF/CCN2 antibodies or antisense oligonucleotides in normal rat kidney cells (NRK) and human fibroblasts [65]. TGF- β is able to support anchorage-independent growth of NRK cells, a process that is antagonized by CTGF/CCN2 antibodies or antisense oligonucleotides [66]. Additionally, subcutaneous injection of TGF- β into neonatal mice, which causes a rapid increase in the amount of granulation tissue comprising connective tissue cells and abundant ECM, results in enhanced levels of CTGF/CCN2 mRNA in connective tissue fibroblasts but not in

epithelial cells or endothelial cells [13,67]. Finally, injection of CTGF/CCN2 causes a very similar fibrotic reaction as TGF- β and is not mimicked by other growth factors [13,15,67,68]. Collectively, these data support a role for CTGF/CCN2 as a downstream mediator of some of the fibrogenic actions of TGF- β 1, particularly the promotion of fibroblast proliferation and ECM production.

3. Hepatic stellate cells in liver fibrosis

A central event in liver fibrosis is the activation of hepatic stellate cells (HSCs) which represents a transition from quiescent vitamin A-rich cells to vitamin A-deficient, proliferative, fibrogenic and contractile myofibroblasts [69–73]. Thereafter, activated HSCs demonstrate perpetuation which involves changes in cell behavior, proliferation, chemotaxis, fibrogenesis, contractility, matrix degradation, retinoid loss, leukocyte chemotaxis, and cytokine release [71,72]. Collectively, these changes increase ECM accumulation such that HSCs become the principal effector cells of the fibrotic response. HSC activation is also concurrent with a shift from normal low-density basement membrane-like matrix containing non-fibril-forming collagens (types IV and VI) to fibril-forming collagens (types I and III) [71]. Data from several laboratories have demonstrated that CTGF/CCN2 is overexpressed in fibrotic human liver and in animal models of the disease. Importantly, CTGF/CCN2 is produced by HSCs as a function of their activation status and exogenous CTGF/CCN2 is able to modulate HSC function. These aspects are discussed in detail below.

4. TGF- β 1 and hepatic fibrosis

Cytokines play an important role in the development of liver fibrosis. One of the major cytokines associated with hepatic fibrosis and involved in HSC activation and deposition of ECM is TGF- β 1 [74,75]. The evidence for TGF- β 1 having a central role in liver fibrosis is sizable. First, its overexpression has been correlated with the degree of fibrosis in both animal models as well as in human disease [76–82]. Second, TGF- β 1 inhibits matrix degradation by upregulation of TIMP-1 and -2, while stimulating the deposition of ECM by HSCs [81,83,84]. Third, TGF- β 1 modulates the expression of integrin matrix receptors [85]. Fourth, transgenic mice that over-express TGF- β 1 develop acute hepatic fibrosis [86,87]. Finally, blocking TGF- β 1 action with neutralizing antibodies, inhibitors or receptor agonists inhibits animal models of renal or liver fibrosis [88–90].




Fig. 1. CTGF/CCN2 expression in human congenital hepatic fibrosis. Sections of liver from a 9-year-old patient with congenital hepatic fibrosis were treated with (A) sense CTGF/CCN2 mRNA, (B, F) anti-sense CTGF/CCN2 mRNA, (C) normal rabbit IgG, (D, G) affinity-purified rabbit anti-CTGF/CCN2 (81–94) peptide or (E, H) hematoxylin and eosin. CTGF/CCN2 mRNA (purple staining) and protein (brown staining) was localized primarily to bile duct epithelial cells; this staining pattern was observed in 3/3 cases of the disease. A–E, 10 \times ; F–H, 40 \times .

5. CTGF/CCN2 and hepatic fibrosis

5.1. Human disease

Data reported in the last few years have provided compelling evidence that CTGF/CCN2 mRNA and protein levels are correlated with the degree of hepatic fibrosis, irrespective of the type of disease. Ribonuclease protection assay analysis of cirrhotic livers from patients with primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC) or biliary atresia showed that CTGF/CCN2 mRNA in fibrotic livers was elevated 3–5-fold as compared with their normal counterparts [25]. Similarly,




Fig. 2. CTGF/CCN2 production and action in HSC cells. (A) Myofibroblastic HSCs (HSC-T6; Scott Friedman, Mount Sinai Hospital, New York, NY) were grown for 3 days in DMEM containing 0.2% fetal bovine serum until they reached >90% confluence. Cells were left untreated (lane 1) or stimulated for 4 h with 2 ng/ml TGF- β 1 (lane 2) or 1 μ g/ml recombinant CTGF/CCN2 (lane 3). Total RNA from the cells was isolated and subjected to RT-PCR to amplify mRNA for CTGF/CCN2 or GAPDH. (B) Balb/c 3T3 cells were grown to confluence and quiescence in DMEM containing 10% calf serum and then left untreated (lane 1) or stimulated for 4 h with 20 ng/ml TGF- β 1 (lane 2), 200 μ M acetaldehyde (lane 3), 2 mM acetaldehyde (lane 4), 12 mM ethanol (lane 5) or 120 mM ethanol (lane 6). RNA from the cells was isolated and subjected to RT-PCR to amplify mRNA for CTGF/CCN2 or GAPDH. (C) Day 7 primary activated rat HSCs were left untreated or stimulated with 100 ng/ml CTGF/CCN2(247–349) [68,97] for 24 h. α SMA was detected by immunocytochemical staining. (D) Non-tissue culture plastic ELISA plates were coated with 1 μ g/ml human recombinant CTGF/CCN2 and then blocked with 3% BSA. Wells were incubated for 1 h at 37 $^{\circ}$ C with 100 μ l PBS containing 5 \times 10 4 HSC-T6 cells in the presence or absence of 5 μ g/ml heparin or 10 mM EDTA. Adherent cells were fixed and quantified as described [68].

Northern blot analysis showed that, as compared with normal livers, CTGF/CCN2 and TGF- β 1 mRNA expression was 6- to 8-fold higher in cirrhotic livers from patients with chronic viral hepatitis, PBC, PSC, crypto-

genic and alcoholic liver disease (ALD) [24]. Elevated levels of CTGF/CCN2 mRNA were also present in livers from patients with non-alcoholic steatohepatitis (NASH) and the degree of fibrosis was correlated with CTGF/CCN2 protein levels in the ECM [91]. Immunohistochemical staining of normal and hepatitis C livers showed that increased CTGF/CCN2 protein levels were associated with a higher score of fibrosis [26].

In situ hybridization of cirrhotic livers resulting from chronic viral hepatitis, PBC, PSC, cryptogenic, and ALD contained marked CTGF/CCN2 staining in fibroblast and myofibroblast-like cells within the fibrotic portal tracts and fibrous septa [22,24,92]. CTGF/CCN2 mRNA was also observed in HSCs, myofibroblast-like spindle cells around proliferating ductules, a few duct and ductular epithelial cells, inflammatory cells, and sinusoidal and vascular endothelial cells [24,26]. CTGF/CCN2 protein staining was present in portal tracts, the ECM of the fibrous septa, sinusoidal lining and in the proliferating bile ducts in hepatitis C diseased livers [22,26]. Additionally, many cells in fibrotic liver that stained for α -smooth muscle actin (α SMA) were also shown to contain CTGF/CCN2 protein [26]. In congenital hepatic fibrosis, which is marked by profound hyperplasia of the bile ducts, CTGF/CCN2 mRNA and protein are strongly expressed by bile duct epithelial cells (Fig. 1). In ALD, hepatocytes may be an important source of CTGF/CCN2 since it is produced in cultured HepG2 cells downstream of cytochrome P-450E1-mediated ethanol oxidation [92,93]. In livers from patients suffering from idiopathic portal hypertension, periductal mononuclear cells demonstrate enhanced CTGF/CCN2 expression and this was correlated with collagen and elastin deposition and the presence of activated HSCs and pericellular fibrosis [94]. Thus, while its timing of production and mode of action may vary according to the initial hepatic insult and type of damage, CTGF/CCN2 has the potential of being produced by virtually all major cell types in the liver. Moreover, high intra-hepatic levels of CTGF/CCN2 are associated with its entry into the circulation since, as compared with control subjects, serum CTGF/CCN2 levels were higher in patients with biliary atresia and were correlated with the progression of liver fibrosis [28].

5.2. Animal models of hepatic fibrosis

Analysis of mRNA by real-time PCR or reverse transcriptase PCR showed that hepatic CTGF/CCN2 and collagen expression was increased with the degree of fibrosis following CCl₄ treatment or bile duct ligation (BDL) in rats [25,26]. Both CTGF/CCN2 and TGF- β mRNA increased 6 h post CCl₄ administration and levels peaked at 72 h [23]. In the BDL model, TGF- β and CTGF/CCN2 mRNA were increased 4- and 7-fold,

respectively, as compared with normal liver [23]. CTGF/CCN2 mRNA and protein was also upregulated in the obesity and type II diabetes Zucker (fa/fa) rat as compared with the wild type counterpart [91]. In the CCl₄ model, CTGF/CCN2 mRNA was expressed at high levels in HSCs and the protein was localized to fibrous septa and centrilobular regions [23,26]. In contrast, the main site of CTGF/CCN2 mRNA in BDL livers was proliferating bile duct epithelial cells, though the protein was present in the fibrous septa as well as in areas of ductular proliferation [23,26]. TGF- β was only expressed at minimal levels in these cells but was highly expressed in the surrounding HSCs [23].

5.3. CTGF/CCN2 biology in cultured HSCs

In a cultured HSC line, CTGF/CCN2 was localized exclusively to the Golgi apparatus and found to be quantitatively secreted into the medium [95]. Cultured activated primary HSCs were found to contain CTGF/CCN2 by Western blotting [26] and both Northern and Western blots demonstrated CTGF/CCN2 was increasingly expressed during progressive activation of cultured primary rat HSCs [25]. Additionally, both intermediate and activated cultured HSCs were induced by TGF- β 1 to express CTGF/CCN2 mRNA and protein [25]. CTGF/CCN2 mRNA expression was also induced in HSCs treated with VEGF, lipid peroxidation products, acetaldehyde or PDGF-BB [27]. Similarly, CTGF/CCN2 mRNA expression is stimulated by both TGF- β 1 and CTGF/CCN2 in HSC-T6 cells and by TGF- β 1, ethanol, or acetaldehyde in mouse fibroblasts (Fig. 2A, B). Finally, HSCs isolated from fa/fa rats incubated with glucose or insulin showed a TGF- β -independent increase in CTGF/CCN2 mRNA and protein [91]. While these data show that CTGF/CCN2 is produced as a function of HSC activation or under conditions that favor activation, the relative contribution to the activation process of CTGF/CCN2 derived from HSCs as compared with the other cells in the liver remains uncertain. Nonetheless, both autocrine and paracrine pathways of action of CTGF/CCN2 on HSCs seem likely since the cells are responsive to exogenous CTGF/CCN2. For example treatment of rat HSCs with CTGF/CCN2 induces migration and proliferation [27], as well as stimulated expression of type I collagen mRNA [27] and α SMA (Fig. 2C). Moreover when added to CTGF/CCN2-coated ELISA plates, HSC-T6 cells demonstrate heparin-dependent and divalent-cation-dependent adhesion (Fig. 2D), consistent with previously reported data on the interaction of CTGF/CCN2 with heparin-like molecules and integrins on fibroblasts [12,96]. These binding interactions appear to occur principally with the C-terminal 103 residues of CTGF/CCN2 [97] which can stimulate α SMA production in the absence of the rest of the molecule (Fig. 2C and [68]). However, residues 199–




Fig. 3. Pathways of CTGF/CCN2 production and action in hepatic fibrosis. CTGF/CCN2 is produced by most major cell types in the liver in response to diverse stimuli, including those associated with fibrosis and/or HSC activation. As the principal effector cells of the fibrotic response, HSCs appear to be particularly important as autocrine or paracrine targets for CTGF/CCN2, especially in their activated state. CTGF/CCN2 binds to both LRP and integrins on the HSC surface via mechanisms that involve HSPGs. Binding to LRP may facilitate CTGF/CCN2 clearance via endosomal degradation, though additional pathways may also be activated. By analogy to data obtained with fibroblasts, binding of integrins to CTGF/CCN2 likely results in the activation of intracellular signaling cascades that promote proliferation, adhesion, motility and collagen synthesis. As a mosaic protein, CTGF/CCN2 may also be involved in direct synergistic binding interactions with other fibrogenic molecules such as TGF- β or PDGF.

243 of CTGF are also involved in promoting HSC adhesion through the ability of this domain to bind to cell surface low density lipoprotein receptor-related protein (LRP) [98], a multifunctional scavenger, signaling, and adhesion receptor [99]. Upon binding to HSCs, soluble 125 I-labeled CTGF/CCN2 is internalized and degraded in the endosome [95] through a process that may involve LRP [100].

5.4. CTGF/CCN2 transgene expression *in vivo*

While several reports have documented the development of fibrotic lesions after repeated subcutaneous injections of CTGF/CCN2 [13,15,68], there is no published evidence that CTGF/CCN2 is able to drive organ fibrosis *in vivo*. This will likely be obtained from animal models in which CTGF/CCN2 expression or action is ablated or in which the CTGF/CCN2 gene is over-expressed. Short-term (3 days) over-expression of CTGF/CCN2 or TGF- β in mouse livers was recently achieved by infection with replication-deficient recombinant adenovirus [101]. Gene array and Northern blot analysis demonstrated that both CTGF/CCN2 and

TGF- β caused enhanced expression of urokinase receptor, laminin receptor, bone morphogenic protein-1 and the serine proteinase inhibitors Spi 2 and Spi 2.4, whereas TGF- β alone also suppressed immune response genes that were induced by adenoviral infection [101]. While the time-course of these experiments was too short to study disease processes *per se*, these data support an overlapping role for CTGF/CCN2 and TGF- β in stimulating expression of genes *in vivo* that are involved in tissue repair, fibrosis and angiogenesis.

6. Summary and perspectives

The rapid pace of research into CTGF/CCN2 over the last 5–6 years can be attributed largely to studies focusing attention on its relationship to TGF- β and its role in fibrosis. From the standpoint of specific organs, most studies of CTGF/CCN2 expression and action have focused on the liver, kidney and skin and there much anticipation that CTGF/CCN2 may offer a new lead as a therapeutic target in fibrosis of these and other organ systems [5,6,20,31,34]. It remains to be deter-

mined whether the excitement surrounding the possibility that CTGF/CCN2 is an effective target for TGF- β -mediated fibrosis is well founded, but it should be noted that several TGF- β -independent pathways of CTGF/CCN2 action have already been identified [2]. In addition to its role in fibrosis, additional studies need to be undertaken to more carefully establish the role of CTGF/CCN2 in development, differentiation, and normal function of the liver. Since CTGF/CCN2 has also been implicated in liver regeneration [102] and hepatocarcinoma [103], a better understanding is needed of the pathways by which CTGF/CCN2 is overexpressed and whether it exerts distinct or overlapping biological roles in various disease states. Since many hepatic cell types have the potential for CTGF/CCN2 synthesis, complex paracrine networks of CTGF/CCN2 action within the liver are likely in view of its broad interactions with target cells that express cell surface heparin-like molecules and/or integrins [3]. Nonetheless, it is already fairly clear that HSCs are one of the principal target cells for CTGF/CCN2 in liver fibrosis, as summarized in Fig. 3. Additionally, a major role of CTGF/CCN2 may be to act as a binding protein for other bioactive molecules, thereby regulating the activity and bioavailability of both binding partners [104]. As further knowledge of CTGF/CCN2 protein–protein and protein–cell interactions within the liver is gained, its role and mechanisms of action in normal liver function, injury, tumorigenesis, and fibrosis will be clarified.

Acknowledgements

We thank Dr E. Moussad and Dr M. Rageh for technical assistance. D.R. Brigstock is funded by NIH grant AA12817-02.

References

- Brigstock DR. The connective tissue growth factor/cysteine-rich 61/nephroblastoma overexpressed (CCN) family. *Endocr Rev* 1999;20:189–206.
- Moussad EE, Brigstock DR. Connective tissue growth factor: what's in a name. *Mol Genet Metab* 2000;71:276–92.
- Lau LF, Lam SC. The CCN family of angiogenic regulators: the integrin connection. *Exp Cell Res* 1999;248:44–57.
- Perbal B. NOV (nephroblastoma overexpressed) and the CCN family of genes: structural and functional issues. *Mol Pathol* 2001;54:57–79.
- Leask A, Holmes A, Abraham DJ. Connective tissue growth factor: a new and important player in the pathogenesis of fibrosis. *Curr Rheumatol Rep* 2002;4:136–42.
- Blom IE, Goldschmeding R, Leask A. Gene regulation of connective tissue growth factor: new targets for antifibrotic therapy. *Matrix Biol* 2002;21:473.
- Almendral JM, Sommer D, Macdonald-Bravo H, Burckhardt J, Perera J, Bravo R. Complexity of the early genetic response to growth factors in mouse fibroblasts. *Mol Cell Biol* 1988;8:2140–8.
- Ryseck RP, Macdonald-Bravo H, Mattei MG, Bravo R. Structure, mapping, and expression of fisp-12, a growth factor-inducible gene encoding a secreted cysteine-rich protein. *Cell Growth Differ* 1991;2:225–33.
- Brunner A, Chinn J, Neubauer M, Purchio AF. Identification of a gene family regulated by transforming growth factor- β . *DNA Cell Biol* 1991;10:293–300.
- Igarashi A, Okochi H, Bradham DM, Grotendorst GR. Regulation of connective tissue growth factor gene expression in human skin fibroblasts and during wound repair. *Mol Biol Cell* 1993;4:637–45.
- Brigstock DR, Steffen CL, Kim GY, Vegunta RK, Diehl JR, Harding PA. Purification and characterization of novel heparin-binding growth factors in uterine secretory fluids. Identification as heparin-regulated Mr 10000 forms of connective tissue growth factor. *J Biol Chem* 1997;272:20275–82.
- Kireeva ML, Latinkic BV, Kolesnikova TV, et al. Cyr61 and Fisp12 are both ECM-associated signaling molecules: activities, metabolism, and localization during development. *Exp Cell Res* 1997;233:63–77.
- Frazier K, Williams S, Kothapalli D, Klapper H, Grotendorst GR. Stimulation of fibroblast cell growth, matrix production, and granulation tissue formation by connective tissue growth factor. *J Invest Dermatol* 1996;107:404–11.
- Chen CC, Chen N, Lau LF. The angiogenic factors Cyr61 and connective tissue growth factor induce adhesive signaling in primary human skin fibroblasts. *J Biol Chem* 2001;276:10443–52.
- Mori T, Kawara S, Shinozaki M, et al. Role and interaction of connective tissue growth factor with transforming growth factor- β in persistent fibrosis: a mouse fibrosis model. *J Cell Physiol* 1999;181:153–9.
- Igarashi A, Nashiro K, Kikuchi K, et al. Connective tissue growth factor gene expression in tissue sections from localized scleroderma, keloid, and other fibrotic skin disorders. *J Invest Dermatol* 1996;106:729–33.
- Igarashi A, Nashiro K, Kikuchi K, et al. Significant correlation between connective tissue growth factor gene expression and skin sclerosis in tissue sections from patients with systemic sclerosis. *J Invest Dermatol* 1995;105:280–4.
- Sato S, Nagaoka T, Hasegawa M, et al. Serum levels of connective tissue growth factor are elevated in patients with systemic sclerosis: association with extent of skin sclerosis and severity of pulmonary fibrosis. *J Rheumatol* 2000;27:149–54.
- Querfeld C, Sollberg S, Huerkamp C, Eckes B, Krieg T. Pseudoscleroderma associated with lung cancer: correlation of collagen type I and connective tissue growth factor gene expression. *Br J Dermatol* 2000;142:1228–33.
- Denton CP, Abraham DJ. Transforming growth factor- β and connective tissue growth factor: key cytokines in scleroderma pathogenesis. *Curr Opin Rheumatol* 2001;13:505–11.
- Stratton R, Shiwen X, Martini G, et al. Iloprost suppresses connective tissue growth factor production in fibroblasts and in the skin of scleroderma patients. *J Clin Invest* 2001;108:241–50.
- Hayashi N, Kakimura T, Soma Y, et al. Connective tissue growth factor is directly related to liver fibrosis. *Hepatogastroenterology* 2002;49:133–5.
- Sedlacek N, Jia JD, Bauer M, et al. Proliferating bile duct epithelial cells are a major source of connective tissue growth factor in rat biliary fibrosis. *Am J Pathol* 2001;158:1239–44.
- Abou-Shady M, Friess H, Zimmermann A, et al. Connective tissue growth factor in human liver cirrhosis. *Liver* 2000;20:296–304.
- Williams EJ, Gaca MD, Brigstock DR, Arthur MJ, Benyon RC. Increased expression of connective tissue growth factor in

- fibrotic human liver and in activated hepatic stellate cells. *J Hepatol* 2000;32:754–61.
- [26] Paradis V, Dargere D, Vidaud M, et al. Expression of connective tissue growth factor in experimental rat and human liver fibrosis. *Hepatology* 1999;30:968–76.
- [27] Paradis V, Dargere D, Bonvoult F, Vidaud M, Segarini P, Bedossa P. Effects and regulation of connective tissue growth factor on hepatic stellate cells. *Lab Invest* 2002;82:767–74.
- [28] Tamatani T, Kobayashi H, Tezuka K, et al. Establishment of the enzyme-linked immunosorbent assay for connective tissue growth factor (CTGF) and its detection in the sera of biliary atresia. *Biochem Biophys Res Commun* 1998;251:748–52.
- [29] Yokoi H, Sugawara A, Mukoyama M, et al. Role of connective tissue growth factor in profibrotic action of transforming growth factor-beta: a potential target for preventing renal fibrosis. *Am J Kidney Dis* 2001;38:S134–8.
- [30] Yokoi H, Mukoyama M, Sugawara A, et al. Role of connective tissue growth factor in fibronectin expression and tubulointerstitial fibrosis. *Am J Physiol Renal Physiol* 2002;282:F933–42.
- [31] Riser BL, Cortes P. Connective tissue growth factor and its regulation: a new element in diabetic glomerulosclerosis. *Renal Fail* 2001;23:459–70.
- [32] Riser BL, Denichilo M, Cortes P, et al. Regulation of connective tissue growth factor activity in cultured rat mesangial cells and its expression in experimental diabetic glomerulosclerosis. *J Am Soc Nephrol* 2000;11:25–38.
- [33] Wang S, Denichilo M, Brubaker C, Hirschberg R. Connective tissue growth factor in tubulointerstitial injury of diabetic nephropathy. *Kidney Int* 2001;60:96–105.
- [34] Goldschmeding R, Aten J, Ito Y, Blom I, Rabelink T, Weening JJ. Connective tissue growth factor: just another factor in renal fibrosis. *Nephrol Dial Transplant* 2000;15:296–9.
- [35] Clarkson MR, Gupta S, Murphy M, Martin F, Godson C, Brady HR. Connective tissue growth factor: a potential stimulus for glomerulosclerosis and tubulointerstitial fibrosis in progressive renal disease. *Curr Opin Nephrol Hypertens* 1999;8:543–8.
- [36] Ito Y, Aten J, Bende RJ, et al. Expression of connective tissue growth factor in human renal fibrosis. *Kidney Int* 1998;53:853–61.
- [37] Murphy M, Godson C, Cannon S, et al. Suppression subtractive hybridization identifies high glucose levels as a stimulus for expression of connective tissue growth factor and other genes in human mesangial cells. *J Biol Chem* 1999;274:5830–4.
- [38] Lasky JA, Ortiz LA, Tonthat B, et al. Connective tissue growth factor mRNA expression is upregulated in bleomycin-induced lung fibrosis. *Am J Physiol* 1998;275:L365–71.
- [39] Pan LH, Yamauchi K, Uzuki M, et al. Type II alveolar epithelial cells and interstitial fibroblasts express connective tissue growth factor in IPF. *Eur Respir J* 2001;17:1220–7.
- [40] Allen JT, Knight RA, Bloor CA, Spiteri MA. Enhanced insulin-like growth factor binding protein-related protein 2 (connective tissue growth factor) expression in patients with idiopathic pulmonary fibrosis and pulmonary sarcoidosis. *Am J Respir Cell Mol Biol* 1999;21:693–700.
- [41] Oemar BS, Luscher TF. Connective tissue growth factor. Friend or foe. *Arterioscler Thromb Vasc Biol* 1997;17:1483–9.
- [42] Oemar BS, Werner A, Garnier JM, et al. Human connective tissue growth factor is expressed in advanced atherosclerotic lesions. *Circulation* 1997;95:831–9.
- [43] Ohnishi H, Oka T, Kusachi S, et al. Increased expression of connective tissue growth factor in the infarct zone of experimentally induced myocardial infarction in rats. *J Mol Cell Cardiol* 1998;30:2411–22.
- [44] Pan LH, Beppu T, Kurose A, et al. Neoplastic cells and proliferating endothelial cells express connective tissue growth factor (CTGF) in glioblastoma. *Neurol Res* 2002;24:677–83.
- [45] di Mola FF, Friess H, Martignoni ME, et al. Connective tissue growth factor is a regulator for fibrosis in human chronic pancreatitis. *Ann Surg* 1999;230:63–71.
- [46] di Mola FF, Friess H, Riesle E, et al. Connective tissue growth factor is involved in pancreatic repair and tissue remodeling in human and rat acute necrotizing pancreatitis. *Ann Surg* 2002;235:60–7.
- [47] Dammeier J, Brauchle M, Falk W, Grotendorst GR, Werner S. Connective tissue growth factor: a novel regulator of mucosal repair and fibrosis in inflammatory bowel disease. *Int J Biochem Cell Biol* 1998;30:909–22.
- [48] Wunderlich K, Pech M, Eberle AN, Mihatsch M, Flammer J, Meyer P. Expression of connective tissue growth factor (CTGF) mRNA in plaques of human anterior subcapsular cataracts and membranes of posterior capsule opacification. *Curr Eye Res* 2000;21:627–36.
- [49] Wunderlich K, Senn BC, Reiser P, Pech M, Flammer J, Meyer P. Connective tissue growth factor in retrocorneal membranes and corneal scars. *Ophthalmologica* 2000;214:341–6.
- [50] Lee EH, Joo CK. Role of transforming growth factor-beta in transdifferentiation and fibrosis of lens epithelial cells. *Invest Ophthalmol Vis Sci* 1999;40:2025–32.
- [51] Folger PA, Zekaria D, Grotendorst G, Masur SK. Transforming growth factor-beta-stimulated connective tissue growth factor expression during corneal myofibroblast differentiation. *Invest Ophthalmol Vis Sci* 2001;42:2534–41.
- [52] Hong HH, Uzel MI, Duan C, Sheff MC, Trackman PC. Regulation of lysyl oxidase, collagen, and connective tissue growth factor by TGF-beta1 and detection in human gingiva. *Lab Invest* 1999;79:1655–67.
- [53] Uzel MI, Kantarci A, Hong HH, et al. Connective tissue growth factor in drug-induced gingival overgrowth. *J Periodontol* 2001;72:921–31.
- [54] Frazier KS, Grotendorst GR. Expression of connective tissue growth factor mRNA in the fibrous stroma of mammary tumors. *Int J Biochem Cell Biol* 1997;29:153–61.
- [55] Kubo M, Kikuchi K, Nashiro K, et al. Expression of fibrogenic cytokines in desmoplastic malignant melanoma. *Br J Dermatol* 1998;139:192–7.
- [56] Igarashi A, Hayashi N, Nashiro K, Takehara K. Differential expression of connective tissue growth factor gene in cutaneous fibrohistiocytic and vascular tumors. *J Cutaneous Pathol* 1998;25:143–8.
- [57] Wenger C, Ellenrieder V, Alber B, et al. Expression and differential regulation of connective tissue growth factor in pancreatic cancer cells. *Oncogene* 1999;18:1073–80.
- [58] Grotendorst GR. Connective tissue growth factor: a mediator of TGF-beta action on fibroblasts. *Cytokine Growth Factor Rev* 1997;8:171–9.
- [59] Schuppan D, Porov Y. Hepatic fibrosis: from bench to bedside. *J Gastroenterol Hepatol* 2002;17(Suppl. 3):S300–5.
- [60] Grotendorst GR, Okochi H, Hayashi N. A novel transforming growth factor beta response element controls the expression of the connective tissue growth factor gene. *Cell Growth Differ* 1996;7:469–80.
- [61] Holmes A, Sa S, Shiwen X, Black CM, Abraham DJ, Leask A. CTGF and SMADs: maintenance of scleroderma phenotype is independent of SMAD signalling. *J. Biol. Chem.*, 2001;276:10594–601.
- [62] Leask A, Sa S, Holmes A, Shiwen X, Black CM, Abraham DJ. The control of *ccn2* (*ctgf*) gene expression in normal and scleroderma fibroblasts. *Mol Pathol* 2001;54:180–3.
- [63] Leask A, Holmes A, Black CM, Abraham DJ. CTGF gene regulation: requirements for its induction by TGFbeta 2 in fibroblasts. *J Biol Chem* 2003;5:5.
- [64] Ihn H. Pathogenesis of fibrosis: role of TGF-beta and CTGF. *Curr Opin Rheumatol* 2002;14:681–5.

- [65] Duncan MR, Frazier KS, Abramson S, et al. Connective tissue growth factor mediates transforming growth factor beta-induced collagen synthesis: down-regulation by cAMP. *FASEB J* 1999;13:1774–86.
- [66] Kothapalli D, Frazier KS, Welply A, Segarini PR, Grotendorst GR. Transforming growth factor beta induces anchorage-independent growth of NRK fibroblasts via a connective tissue growth factor-dependent signaling pathway. *Cell Growth Differ* 1997;8:61–8.
- [67] Shinozaki M, Kawara S, Hayashi N, Kakinuma T, Igarashi A, Takehara K. Induction of subcutaneous tissue fibrosis in newborn mice by transforming growth factor beta—simultaneous application with basic fibroblast growth factor causes persistent fibrosis [corrected and republished article originally printed in *Biochem. Biophys. Res. Commun.* Aug 18 1997;237(2):292–6]. *Biochem Biophys Res Commun* 1997;240:292–7.
- [68] Ball DK, Moussad EE, Rageh MA, Kemper SA, Brigstock DR. Establishment of a recombinant expression system for connective tissue growth factor (CTGF) that models CTGF processing in utero. *Reproduction* 2003;125:271–84.
- [69] Burt AD. Pathobiology of hepatic stellate cells. *J Gastroenterol* 1999;34:299–304.
- [70] Britton RS, Bacon BR. Intracellular signaling pathways in stellate cell activation. *Alcohol Clin Exp Res* 1999;23:922–5.
- [71] Friedman SL. Stellate cell activation in alcoholic fibrosis—an overview. *Alcohol Clin Exp Res* 1999;23:904–10.
- [72] Friedman SL. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *J Biol Chem* 2000;275:2247–50.
- [73] Eng FJ, Friedman SL. Fibrogenesis I. New insights into hepatic stellate cell activation: the simple becomes complex. *Am J Physiol Gastrointest Liver Physiol* 2000;279:G7–G11.
- [74] Li D, Friedman SL. Liver fibrogenesis and the role of hepatic stellate cells: new insights and prospects for therapy. *J Gastroenterol Hepatol* 1999;14:618–33.
- [75] Gressner AM. Transdifferentiation of hepatic stellate cells (Ito cells) to myofibroblasts: a key event in hepatic fibrogenesis. *Kidney Int Suppl* 1996;54:S39–45.
- [76] Czaja MJ, Weiner FR, Flanders KC, et al. In vitro and in vivo association of transforming growth factor-beta 1 with hepatic fibrosis. *J Cell Biol* 1989;108:2477–82.
- [77] Armendariz-Borunda J, Seyer JM, Kang AH, Raghov R. Regulation of TGF beta gene expression in rat liver intoxicated with carbon tetrachloride. *FASEB J* 1990;4:215–21.
- [78] Nakatsukasa H, Evarts RP, Hsia CC, Thorgeirsson SS. Transforming growth factor-beta 1 and type I procollagen transcripts during regeneration and early fibrosis of rat liver. *Lab Invest* 1990;63:171–80.
- [79] Nakatsukasa H, Nagy P, Evarts RP, Hsia CC, Marsden E, Thorgeirsson SS. Cellular distribution of transforming growth factor-beta 1 and procollagen types I, III and IV transcripts in carbon tetrachloride-induced rat liver fibrosis. *J Clin Invest* 1990;85:1833–43.
- [80] Matsuoka M, Tsukamoto H. Stimulation of hepatic lipocyte collagen production by Kupffer cell-derived transforming growth factor beta: implication for a pathogenetic role in alcoholic liver fibrogenesis. *Hepatology* 1990;11:599–605.
- [81] Castilla A, Prieto J, Fausto N. Transforming growth factors beta 1 and alpha in chronic liver disease. Effects of interferon alfa therapy. *New Engl J Med* 1991;324:933–40.
- [82] Bayer EM, Herr W, Kanzler S, et al. Transforming growth factor-beta1 in autoimmune hepatitis: correlation of liver tissue expression and serum levels with disease activity. *J Hepatol* 1998;28:803–11.
- [83] Edwards DR, Murphy G, Reynolds JJ, et al. Transforming growth factor beta modulates the expression of collagenase and metalloproteinase inhibitor. *EMBO J* 1987;6:1899–904.
- [84] Knittel T, Mehde M, Kobold D, Saile B, Dinter C, Ramadori G. Expression patterns of matrix metalloproteinases and their inhibitors in parenchymal and non-parenchymal cells of rat liver: regulation by TNF-alpha and TGF-beta1. *J Hepatol* 1999;30:48–60.
- [85] Cai T, Lei QY, Wang LY, Zha XL. TGF-beta 1 modulated the expression of alpha 5 beta 1 integrin and integrin-mediated signaling in human hepatocarcinoma cells. *Biochem Biophys Res Commun* 2000;274:519–25.
- [86] Clouthier DE, Comerford SA, Hammer RE. Hepatic fibrosis, glomerulosclerosis, and a lipodystrophy-like syndrome in PEPCK-TGF-beta1 transgenic mice. *J Clin Invest* 1997;100:2697–713.
- [87] Sanderson N, Factor V, Nagy P, et al. Hepatic expression of mature transforming growth factor beta 1 in transgenic mice results in multiple tissue lesions. *Proc Natl Acad Sci USA* 1995;92:2572–6.
- [88] Border WA, Okuda S, Languino LR, Sporn MB, Ruoslahti E. Suppression of experimental glomerulonephritis by antiserum against transforming growth factor beta 1. *Nature* 1990;346:371–4.
- [89] Border WA, Noble NA, Yamamoto T, et al. Natural inhibitor of transforming growth factor-beta protects against scarring in experimental kidney disease. *Nature* 1992;360:361–4.
- [90] George J, Roulot D, Kotliansky VE, Bissell DM. In vivo inhibition of rat stellate cell activation by soluble transforming growth factor beta type II receptor: a potential new therapy for hepatic fibrosis. *Proc Natl Acad Sci USA* 1999;96:12719–24.
- [91] Paradis V, Perlemuter G, Bonvoust F, et al. High glucose and hyperinsulinemia stimulate connective tissue growth factor expression: a potential mechanism involved in progression to fibrosis in nonalcoholic steatohepatitis. *Hepatology* 2001;34:738–44.
- [92] Konishi M, Kajihara M, Kato S, Horie Y, Aiso S, Ishii H. Ethanol enhances expression of connective tissue growth factor (CTGF) in HEPG2 cell line stably expressing cytochrome P-4502E1. *Hepatology* 2001;34:377A.
- [93] Konishi M, Kato S, Kajihara M, et al. Expression of connective tissue growth factor (CTGF) in alcoholic liver fibrosis and in HEPG2 cell line stably expressing cytochrome P-4502E1. *Hepatology* 2002;36:248A.
- [94] Tsuneyama K, Kouda W, Nakanuma Y. Portal and parenchymal alterations of the liver in idiopathic portal hypertension: a histological and immunochemical study. *Pathol Res Pract* 2002;198:597–603.
- [95] Chen Y, Segarini P, Raoufi F, Bradham D, Leask A. Connective tissue growth factor is secreted through the Golgi and is degraded in the endosome. *Exp Cell Res* 2001;271:109–17.
- [96] Chen CC, Chen N, Lau LF. The angiogenic factors Cyr61 and CTGF induce adhesive signaling in primary human skin fibroblasts. *J Biol Chem* 2000;276:10443–52.
- [97] Ball DK, Wilson AW, Kemper SA, Brigstock DR. The heparin-binding 10 kDa fragment of connective tissue growth factor (CTGF) containing module 4 alone stimulates cell adhesion. *J Endocrinol* 2003;176:R1–7.
- [98] Gao R, Brigstock DR. Heparan sulfate proteoglycans and low density lipoprotein receptor-related protein are involved in adhesion of rat activated hepatic stellate cells to CTGF. *Mol Pathol* 2003, in press.
- [99] Herz J, Strickland DK. LRP: a multifunctional scavenger and signaling receptor. *J Clin Invest* 2001;108:779–84.
- [100] Segarini PR, Nesbitt JE, Li D, Hays LG, Yates JR, III, Carmichael DF. The low density lipoprotein receptor-related protein/alpha2-macroglobulin receptor is a receptor for connective tissue growth factor. *J Biol Chem* 2001;276:40659–67.

- [101] Haberberger TC, Kupfer K, Murphy JE. Profiling of genes which are differentially expressed in mouse liver in response to adenoviral vectors and delivered genes. *Gene Ther* 2000;7:903–9.
- [102] Ujike K, Shinji T, Hirasaki S, et al. Kinetics of expression of connective tissue growth factor gene during liver regeneration after partial hepatectomy and D-galactosamine-induced liver injury in rats. *Biochem Biophys Res Commun* 2000;277:448–54.
- [103] Hirasaki S, Koide N, Ujike K, Shinji T, Tsuji T. Expression of Nov, CYR61 and CTGF genes in human hepatocellular carcinoma. *Hepatol Res* 2001;19:294–305.
- [104] Bork P. The modular architecture of a new family of growth regulators related to connective tissue growth factor. *FEBS Lett* 1993;327:125–30.