

New paradigms in the treatment of hepatic cholestasis: From UDCA to FXR, PXR and beyond

Ulrich Beuers^{1,*}, Michael Trauner², Peter Jansen¹, Raoul Poupon³

¹Department of Gastroenterology and Hepatology, Tytgat Institute for Liver and Intestinal Research, Academic Medical Centre University of Amsterdam, Amsterdam, The Netherlands; ²Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Austria; ³UPMC Université Paris 06, INSERM, UMR_S 938, Centre de Recherche Saint-Antoine, AP-HP, Hôpital Saint-Antoine, Service d'Hépatologie, F-75012 Paris, France

Summary

Cholestasis is an impairment of bile formation/flow at the level of the hepatocyte and/or cholangiocyte. The first, and for the moment, most established medical treatment is the natural bile acid (BA) ursodeoxycholic acid (UDCA). This secretagogue improves, e.g. in intrahepatic cholestasis of pregnancy or early stage primary biliary cirrhosis, impaired hepatocellular and cholangiocellular bile formation mainly by complex post-transcriptional mechanisms. The limited efficacy of UDCA in various cholestatic conditions urges for development of novel therapeutic approaches. These include nuclear and membrane receptor agonists and BA derivatives. The nuclear receptors farnesoid X receptor (FXR), retinoid X receptor (RXR), peroxisome proliferator-activated receptor α (PPAR α), and pregnane X receptor (PXR) are transcriptional modifiers of bile formation and at present are under investigation as promising targets for therapeutic interventions in cholestatic disorders. The membrane receptors fibroblast growth factor receptor 4 (FGFR4) and apical sodium BA transporter (ASBT) deserve attention as additional therapeutic targets, as does the potential therapeutic agent norUDCA, a 23-C homologue of UDCA. Here, we provide an overview on established and future promising therapeutic agents and their

potential molecular mechanisms and sites of action in cholestatic diseases.

© 2015 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

Introduction

Bile was first mentioned in the Ebers Papyrus (circa 1550 B.C.) as a useful remedy and purge [1]. Dried black bear's bile rich in ursodeoxycholic acid (UDCA) was recommended in China for treatment of jaundice at times of the Tang dynasty (618–907 A.D.) as documented in the *Tang Materia Medica*, the first state pharmacopoeia worldwide. In the Western hemisphere, bile had been regarded as a major constituent of the human body by the Corpus Hippocraticum and Galen of Pergamon, but was increasingly seen as a useless excrement by the 16th and 17th century (“cloaca sordium et superfluitatum”) [1]. Only during the last centuries, the digestive function of bile was recognized and during the last 30 years the signaling and therapeutic potential of its major constituents, bile acids (BAs), and the (patho-) physiological role of BAs, phospholipids, bicarbonate and other bile constituents were unraveled. Today, bile formation is regarded as a vital secretory process modulated by complex transcriptional and post-transcriptional mechanisms in hepatocytes, cholangiocytes and ileocytes [2–4].

Cholestasis is an impairment of bile formation and flow. It may result from; (i) hepatocellular and/or cholangiocellular secretory defects; or (ii) obstruction of bile ducts by bile duct lesions, stones or tumours, but may also be related to mixed mechanisms in conditions such as primary biliary cirrhosis/choolangitis (PBC) or primary sclerosing cholangitis (PSC). For adequate treatment of cholestasis and cholestatic injury, identification and targeting of the defective hepatocellular and cholangiocellular secretory mechanisms and/or bile duct lesions (or removal of obstructing stones and tumours) is required. Evolving pathophysiological insights in cholestatic disorders, particularly chronic fibrosing cholangiopathies such as PBC, PSC and (other) secondary forms of cholangitis, provide novel opportunities for the development of therapeutic approaches for these disorders. Currently, treatment with UDCA may slow the

Keywords: UDCA; FXR; PXR; Cholestasis.

Received 7 January 2015; received in revised form 16 February 2015; accepted 16 February 2015

* Corresponding author. Address: Department of Gastroenterology & Hepatology, Tytgat Institute for Liver and Intestinal Research, Academic Medical Centre, G4-216, University of Amsterdam, P.O. Box 22600, NL-1100 DD Amsterdam, The Netherlands. Tel.: +31 20 566 24 22; fax: +31 20 566 95 82.

E-mail address: u.h.beuers@amc.uva.nl (U. Beuers).

Abbreviations: AE2, anion exchanger 2; ASBT, apical sodium bile salt transporter; BA, bile acid; CA, cholic acid; CDCA, chenodeoxycholic acid; CFALD, cystic fibrosis-associated liver disease; CFTR, cystic fibrosis transmembrane conductance regulator; DCA, deoxycholic acid; FGF, fibroblast growth factor; FXR, farnesoid X receptor; GR, glucocorticoid receptor; IBD, inflammatory bowel disease; ICP, intrahepatic cholestasis of pregnancy; LCA, lithocholic acid; LPAC, low phospholipid-associated cholelithiasis; MAPK, mitogen-activated protein kinase; NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; PBC, primary biliary cirrhosis; PFIC, progressive familial intrahepatic cholestasis; PKA, protein kinase A; PKC, protein kinase C; PPAR α , peroxisome proliferator-activated receptor α ; PSC, primary sclerosing cholangitis; PXR, pregnane X receptor; UDCA, ursodeoxycholic acid; VDR, vitamin D receptor.



ELSEVIER

Review

progression of chronic cholangiopathies, but has limited or no proven efficacy in various chronic cholestatic disorders and cannot heal them. Immunosuppressive/immunomodulating interventions aiming to minimize immune-mediated damage in immune-mediated/autoimmune cholestatic disorders such as PBC and PSC have disappointed in the past, but new approaches, which are beyond the scope of this review, are at present under consideration for clinical evaluation. The use of specific modifiers of hepatobiliary secretory and cellular protection mechanisms against BA-mediated cytotoxicity may eventually give rise to new classes of disease-modifying drugs. Here, we provide an overview of transcriptional and post-transcriptional modulators of bile formation which may serve as therapeutic agents in the future for the treatment of cholestatic disorders.

UDCA: clinical use

Therapy with natural BAs arose in the 1970s when it was discovered that oral administration of chenodeoxycholic acid (CDCA) induces the dissolution of cholesterol gallstones. However, CDCA induced biliary cirrhosis in some species and was shown to be mildly hepatotoxic and induced dose-dependent diarrhea in humans [5]. Thereafter, UDCA was shown to have similar efficacy in gallstone disease without any side effects [6]. The markedly different behaviour of the two natural BAs was ascertained by numerous experimental studies *in vitro* and *in vivo*.

UDCA was thereafter proposed as a potential therapeutic approach for chronic cholestatic disorders with the following rationale: (a) accumulation of toxic BAs might be at least in part responsible for liver injury in chronic cholestasis; (b) replacement of endogenous BAs by a non-toxic BA (UDCA) could protect the liver and slow down the progression of these disorders. This hypothesis was first tested in PBC [7]. UDCA was shown to provide marked improvement in serum liver tests [7,8]. Placebo-controlled trials showed that UDCA also improves histological features and delays progression to cirrhosis and the time to liver transplantation [9–14]. Today, UDCA therapy is recommended for all patients with PBC provided that they show abnormal serum liver tests [15,16]. The accepted optimal dose is 13–15 mg/kg/day. All patients with PBC do not respond to UDCA in the same way. The transplant-free survival rate among UDCA-treated patients remains significantly lower than that of an age- and gender-matched control population [17], indicating that there is a need for new therapeutic options particularly for patients with a suboptimal biochemical response to UDCA and predictive factors of a poor outcome [18]. Serum bilirubin was shown to be the most potent prognostic marker in PBC as were also serum albumin, prothrombin time and cirrhosis, the traditional prognostic factors in advanced liver disease. More recently, the biochemical response to UDCA has been shown to predict long-term outcomes and, thus, may be applied as a simple selection criterion for clinical trials [18–23].

In PSC, UDCA may lower disease progression but long-term efficacy remains uncertain [24,25]. A placebo-controlled trial using very high doses of UDCA (28–30 mg/kg/day) showed that UDCA was not only ineffective but also harmful in that more patients developed varices or were listed for liver transplantation [26]. Therefore, no evidence-based recommendation can be given for normal doses, but very high-dose regimens should be avoided [16,27].

UDCA therapy has been used for a number of other clinical conditions. Efficacy is regarded as likely in ABCB4 deficiency with progressive familial intrahepatic cholestasis type 3 (PFIC-3) and/or low phospholipid-associated cholelithiasis (LPAC syndrome) and cystic fibrosis-associated liver disease (CFALD). Efficacy is regarded as uncertain in various forms of sclerosing cholangitis, drug-induced liver injury, progressive familial intrahepatic cholestasis type 1 & 2, sarcoidosis hepatitis, prevention of bile duct injury after liver transplantation, and total parenteral nutrition (TPN)-induced cholestasis [16]. In all of these conditions (as in PSC), no clear-cut survival benefit with UDCA has been shown.

UDCA: major molecular mechanisms and sites of action

Dried black bear's bile was recommended more than a thousand years ago for treatment of jaundice at times of the Tang dynasty in China as mentioned above. UDCA may form up to 60% of black bear's total BAs [28] whereas it forms only 1–3% of total BAs in human bile, but is enriched to 40% in bile of patients with PBC and healthy volunteers treated with therapeutic UDCA doses (13–15 mg/kg/day) [29]. UDCA has potent anticholestatic and antiapoptotic properties in conditions of hepatocellular (e.g., intrahepatic cholestasis of pregnancy (ICP)) or cholangiocellular cholestasis (e.g., early PBC).

Early after the first peer-reviewed reports on UDCA in PBC [7,8] it was proposed that UDCA exerts its hepatoprotective effects in cholestatic liver disease mainly by stimulating impaired hepatobiliary secretion [30]. In the 1990's, UDCA was then unraveled as a potent intracellular signaling molecule acting as a Ca^{2+} agonist [31–34] and an activator of protein kinase C (cPKC α) [35–37], mitogen-activated protein kinases (MAPK: Erk1/2, p38^{MAPK}) [38,39] and $\alpha_5\beta_1$ integrins [40,41] in hepatocytes. It was earlier proposed [33,42] and later experimentally proven that UDCA conjugates as potent signaling molecules that might stimulate secretion of hepatocytes (and cholangiocytes [43]) by activating vesicular exocytosis and carrier insertion into their apical membranes resulting in choleretic effects via a dual MAPK- and integrin-dependent mechanism in healthy liver [39,40] and in anticholestatic effects via Ca^{2+} -type II inositol-1,3,4-trisphosphate receptor/cPKC α /PKA-dependent mechanisms in cholestatic liver [44–46]. It remains to be proven if these complex post-transcriptional molecular mechanisms unraveled in experimental animals may explain the choleretic and anticholestatic effects of UDCA in man.

More recently, the 'biliary HCO_3^- umbrella' hypothesis has been introduced as a protective mechanism for hepatocytes and cholangiocytes against the toxic effects of millimolar BA monomers present in bile [47]. This hypothesis indicates that biliary HCO_3^- secretion in humans serves to maintain an alkaline pH near the apical surface of hepatocytes and cholangiocytes to prevent the uncontrolled membrane permeation of protonated glycine-conjugated BAs which have a $\text{pK}_a \geq 4$. Notably, the experimental proof of concept also unraveled that an intact 'biliary HCO_3^- umbrella' is critically dependent on adequate function of the major HCO_3^- exporter, the $\text{Cl}^-/\text{HCO}_3^-$ exchanger AE2, and an intact biliary glycocalyx in human cholangiocytes [48]. Functional impairment of this biliary HCO_3^- umbrella or its regulation would lead to enhanced vulnerability of cholangiocytes and periportal hepatocytes towards the attack of apolar hydrophobic BAs. Notably, UDCA stimulates biliary HCO_3^- secretion under




Fig. 1. Major mechanisms and sites of action of UDCA in cholestatic diseases. (A) UDCA conjugates exert anticholestatic effects and stimulate bile acid (BA) and organic anion secretion by mainly post-transcriptional molecular mechanisms at the level of the hepatocytes (for details see text). UDCA also stimulates biliary HCO_3^- secretion at the level of hepatocytes and cholangiocytes leading to stabilization of a 'biliary HCO_3^- umbrella' [47,48]. Antiapoptotic and anti-inflammatory effects are discussed in the text. (Figure modified from [28], courtesy of G. Paumgartner). (b) A closer look into the 'biliary HCO_3^- umbrella' on the apical membrane of a cholangiocyte [47,48]. The 20–40 nm glycocalyx on the apical membrane may trap HCO_3^- molecules in order to create an alkaline milieu which keeps bile acids in a negatively loaded state (bile salts) and, thereby, prevents carrier-independent invasion of apolar hydrophobic BA into cholangiocytes (and hepatocytes) [47,48]. (Figure modified after Maillette de Buy Wenniger et al. Dig Dis 2015, in press).

experimental conditions as well as in patients with PBC [28,49]. A defective AE2 expression [50,51] and biliary HCO_3^- secretion [49] have been described in patients with PBC. Thus, stabilization of the 'biliary HCO_3^- umbrella' may be a crucial mechanism of action of UDCA in cholestatic liver diseases (Fig. 1).

The original proposal that UDCA may exert anticholestatic effects by removing major hydrophobic BAs such as CDCA or DCA from the circulating BA pool was disproved during short-term UDCA treatment when cholestasis improved, but hydrophobic BA pool sizes remained stable [52]. In contrast to hydrophobic BAs such as CDCA or lithocholic acid (LCA), UDCA also does not markedly affect transport protein expression *in vivo* at the transcriptional level to modulate transport capacity and probably exerts only limited post-transcriptional modification of carrier expression [53].

Antiapoptotic mechanisms [28,54–56] and effects on endoplasmic reticulum stress may contribute to the cytoprotective action of UDCA in cholestatic liver disease as summarized elsewhere [28]. Of note, UDCA has early been described to act as a glucocorticoid receptor (GR) agonist in a ligand-independent way [57–60]. Interaction of UDCA with the GR has been linked to the antiapoptotic effect of UDCA in TGF- β 1-induced liver-cell apoptosis [61]. Thus, antiapoptotic and anti-inflammatory actions of UDCA in the liver and bile ducts may at least in part be secondary to this and the numerous effects described above.

Norursodeoxycholic acid: experimental and clinical effects

24-*nor*ursodeoxycholic acid (*nor*UDCA) is a side chain shortened UDCA derivate which lacks a methylene group resulting in a relative resistance to amidation with taurine or glycine compared with UDCA [62–65]. As a result, *nor*UDCA is passively absorbed from cholangiocytes and undergoes 'cholehepatic shunting' (instead of a full enterohepatic cycle) which generates a HCO_3^- anion resulting in induction of a HCO_3^- -rich hypercholesteris [62,65,66] counteracting intrinsic BA toxicity [67] and reinforcing the 'biliary HCO_3^- umbrella' [47,48]. Cholehepatic shunting may also allow 'ductular targeting' of drugs [62,68]. As a proof of principle, taurine-conjugated *nor*UDCA and bis-*nor*UDCA (resulting from additional side chain shortening) lack cholehepatic shunting properties [69], emphasizing the unique properties of *nor*UDCA. In addition, *nor*UDCA is more hydrophilic and thereby less toxic for hepatocytes and cholangiocytes *in vitro* than its mother compound UDCA [70] which may further help to counteract (intrinsic) biliary toxicity. Notably, neither *nor*UDCA nor UDCA have relevant affinities for dedicated BA receptors such as the farnesoid X receptor (FXR) or G protein-coupled plasma membrane receptor TGR5 [71]. *nor*UDCA (but not "conventional" UDCA or bis-*nor*UDCA) reversed sclerosing cholangitis in the experimental *Mdr2/Abcb4* knockout mouse (*Mdr2/Abcb4*^{-/-}) cholangiopathy model for sclerosing cholangitis, while the mother compound UDCA even aggravated bile infarcts in cholestatic conditions with biliary obstruction [68–70]. Moreover, *nor*UDCA has anti-lipotoxic, anti-proliferative, anti-fibrotic as well as anti-inflammatory effects which may complement stimulation of HCO_3^- secretion with BA detoxification and induction of alternative export via overflow systems at the basolateral membrane [68,71] (Fig. 2). Notably, in a hepatocellular model of cholestasis induced by TLCA, taurine-conjugated *nor*UDCA had anti-cholestatic and anti-apoptotic properties, suggesting that combination of UDCA and *nor*UDCA may be superior to UDCA or *nor*UDCA monotherapy in biliary disorders in which both hepatocyte as well as cholangiocyte dysfunction are involved in the pathophysiology of the disease [72].




Fig. 2. Proposed mechanisms of action of *norUDCA* in the *Mdr2 (Abcb4)*^{-/-} model of sclerosing cholangitis. As hydrophilic bile acid (BA) and by generating a bicarbonate-rich (hyper)choleresis due to cholehepatic shunting, *norUDCA* counteracts intrinsic biliary toxicity resulting from absent phospholipid (PL) secretion with increased free non-micellar bound BAs in this model. Importantly, cholehepatic shunting allows 'ductal targeting' of anti-inflammatory, anti-fibrotic and anti-proliferative effects to injured bile ducts, resulting in 'ductal healing'. *norUDCA* also counteracts lipotoxic fatty acid composition and promotes BA detoxification and elimination via basolateral efflux pumps facilitating their subsequent renal excretion. Modified after [200].

norUDCA is currently undergoing further clinical development in humans. The final results of a double-blind, randomized, European multicenter, placebo-controlled, comparative, exploratory phase II dose-finding trial in the treatment of PSC are expected for 2015 [73]. Future clinical indications, next to PSC, may include PBC and cystic fibrosis-associated liver disease/cholangiopathy where defects of the biliary HCO_3^- umbrella may also be involved [47,48]. Notably, *norUDCA* induces a HCO_3^- -rich choleresis independent of cystic fibrosis transmembrane conductance regulator (CFTR) [69] consistent with the concept of cholehepatic shunting which does not appear to involve active transport processes [62]. Promotion of HCO_3^- -rich bile flow may also beneficially affect sclerosing cholangitis of critically ill patients, non-anastomotic strictures following liver transplantation, or ABCB4 deficiency with progressive familial intrahepatic cholestasis type 3 (PFIC-3) [47,74]. Collectively, these broad and multiple levels of mechanisms make *norUDCA* an attractive therapeutic agent for cholangiopathies [73].

FXR – FGF19: experimental and clinical effects


The discovery of the nuclear hormone receptors in the 1990s was followed by the notion that BAs serve as their ligands [75–79]. This caused a dramatic turn around in BA research. In addition to understanding the role of BAs in regulating their own synthesis and transport, it is now clear that the post-prandial surge of BAs through intestine and liver, and to a lesser degree through adipose tissues, kidney and muscle, triggers signals that prepare the organism for production or storage of energy [80]. Recent studies also show that BA signaling is a major regulator of the circadian rhythm of metabolism [81,82].

Nuclear hormone receptors act as intracellular ligand-activated receptors (Table 1). Cholic acid (CA) and CDCA bind to the FXR

Table 1. Potential future therapeutic agents which modulate bile formation and secretion. See text for potential molecular mechanisms of action and clinical observations.

Pharmacologic agents under study (examples)	Targets	Natural ligands (examples)
Nuclear receptors		
Obeticholic acid PX-102	FXR	Chenodeoxycholic acid Cholic acid
All-trans retinoic acid	RXR	Retinoic acid (Vit. A)
Bezafibrate Fenofibrate Ciprofibrate	PPAR α	Free fatty acids
Budesonide and other corticosteroids	GR PXR	Cortisol
Rifampicin Statins Corticosteroids	PXR	Lithocholic acid
Vitamin D	VDR	1.25-diOH-vitamin D
Membrane receptors		
FGF19 NGM 282	FGFR4	FGF19
Int-777	TGR5	Chenodeoxycholic acid Lithocholic acid
ASBT inhibitors	ASBT	Conjugated bile acids
Bile acid derivatives		
<i>nor</i> UDCA	?	

(NR1H4). FXR, as a heterodimer with the all-trans retinoic acid (ATRA) receptor RXR, binds to the LR-1 DNA motive in the promoter region of target genes [75,83]. FXR target genes include



FGF19 and FGF21 are members of the fibroblast growth factor family [92]. These proteins serve as hormones in inter-organ signaling (FGF19, gut to liver; FGF21, liver to adipose tissues) supporting the actions of FXR and peroxisome proliferator-activated receptor alpha (PPAR α) [93–95]. If after a meal the gallbladder contracts, BAs enter the intestine, are re-absorbed in the ileum where they activate FXR before they enter the portal circulation and are taken up in the liver. In the ileum, FXR activation leads to FGF19 expression. FGF19 enters the portal circulation

For PSC therapeutic possibilities are limited. FXR agonists can be considered but FGF19 induction by these agonists may be a caveat. FGF19 has carcinogenic properties and effects on cholangiocarcinogenesis have to be carefully considered [117]. Nevertheless, ATRA, a ligand of RXR, the heterodimer of FXR, in combination with UDCA, in a short-term small non-randomized trial, has most recently been reported to reduce alkaline phosphatase, serum ALT and BA levels (Assis *et al.*, unpublished).

In addition to FXR and other nuclear hormone receptors, BAs can also signal through a membrane-bound BA-specific receptor

Review

TGR5 (also known as GPBAR1 or M-BAR/BG37) [118]. The most potent endogenous TGR5 activator is LCA followed by DCA [119], while other BAs are less potent. Of note, TGR5 is expressed in various tissues with low or even absent FXR such as spleen, lung or adipose tissue [118,120] with the highest expression in gallbladder and colon [121–123]. In liver, sinusoidal endothelial cells, Kupffer cells and intrahepatic bile ducts express TGR5 (with high expression in human and rat, lower in mouse), while hepatocytes and quiescent stellate cells do not express TGR5 [124–126].

TGR5 activation inhibits pro-inflammatory cytokine production, migration and phagocytic activity of macrophages and Kupffer cells [120,124], in part by suppression of NF- κ B signaling [127] [120,128,129]. Accordingly, mice lacking TGR5 display aggravated liver injury after LPS challenge [129]. TGR5 is also involved in modulation of intestinal inflammation, motility, and improves intestinal barrier function thereby protecting from DSS induced colitis in rodents [130–134]. These findings may be of potential relevance for the gut–liver axis in cholangiopathies such as PSC.

Mice lacking TGR5 have a decreased total BA pool size [121], increased CYP7A1 gene expression [123] and a more hydrophobic BA composition [135] which may be due to the impact of BA-mediated TGR5 activation in inhibiting gallbladder contractility. As such, TGR5 activation by hydrophobic BAs inhibits gallbladder smooth muscle contractility [136]. TGR5^{−/−} mice display prolonged cholestasis, exacerbated inflammatory response and more severe liver injury after partial hepatectomy, dietary BA-challenge or bile duct ligation [135,137]. Importantly, TGR5 polymorphisms in PSC patients may imply a potential role in cholangiocyte pathophysiology [138,139]. Cholangiocytes express TGR5 at the apical membrane/cilia where it may sense the luminal BA concentration and regulate cholangiocellular HCO₃[−]/fluid secretion via CFTR and AE2 [122]. Surprisingly, a highly potent TGR5 agonist (INT-777) failed to induce HCO₃[−] output and bile flow in healthy mice as well as in *Mdr2*^{−/−} model without improvement of bile duct injury [140], findings which could be explained by relatively low TGR5 expression in mouse cholangiocytes. However, mice overexpressing TGR5 showed less liver injury in a mouse model of xenobiotic (DDC)-induced sclerosing cholangitis, while mice lacking TGR5 showed aggravation of inflammation and fibrosis [141]. Collectively, these findings suggest a critical role of TGR5 for liver protection against BA overload, primarily through the control of bile hydrophobicity and cytokine secretion. Conversely, TGR5 deficient mice are protected against lithogenic diet-induced gallstone formation [123] suggesting that inhibition rather than activation of TGR5 may be beneficial for gallstone disease. On the other hand, TGR5 activation of biliary HCO₃[−] secretion could theoretically counteract gallstone formation, at least at the level of bile ducts.

Importantly, activation of TGR5 may also have some undesired off-target effects. For example, bile reflux-induced pancreatitis has been linked to BA-mediated TGR5 activation in mouse pancreas [142]. Moreover, TGR5 activation promoted oxidative stress in astrocytes [143] and activated AKT signaling in cardiomyocytes possibly contributing to cardiac hypertrophy under cholestatic conditions [144]. More recently, TGR5 has also been implied in the pathogenesis of pruritus [145,146]. TGR5-activation by BAs was linked to increased hepatocellular apoptosis [147] through activation of c-Jun N-terminal kinase (JNK) signaling pathways while cholangiocytes seem to evade apoptosis via

TGR5 [148]. Importantly, TGR5 is also highly expressed in gastric and oesophageal adenocarcinoma as well as gallbladder carcinoma where it promotes cell proliferation in response to BA [149]. Such extrahepatic “off-target effects” may need consideration when developing BA receptor ligands as therapeutics in patients with liver disease.

PPAR α : experimental and clinical effects

The PPARs are enriched in tissues with high energy metabolism such as liver (PPAR α , NR1C1), skeletal muscle, heart and gastrointestinal tract (PPAR β/δ , NR1C2) and adipose tissue (PPAR γ , NR1C3) [150]. Fatty acids and their derivatives are the natural ligands for these receptors. By stimulating fatty acid oxidation, PPAR α has anti-steatotic effects. However, PPAR α also has anti-inflammatory actions. It was recently argued that the anti-inflammatory action of PPAR α is based on trans-repression of AP1 and NF- κ B signaling while its metabolic action depends on direct trans-activation of metabolically active target genes. By introducing a mutation in the zinc-finger domain of PPAR α , a DNA-binding deficient PPAR derivative with maintained anti-inflammatory activity but no metabolic activity was recently produced [151].

For the application of PPAR agonists in cholestatic liver disease the notion that the canalicular phospholipid translocator MDR3 is a PPAR α responsive gene is relevant [152,153]. Treatment with PPAR agonists (fibrates) increases MDR3 insertion into the canalicular membrane [153–155]. This stimulates phosphatidylcholine secretion and protects cholangiocytes against bile salt toxicity. This is among the rationales to test fibrates in PBC and PSC. Other actions of PPAR agonists, that underscore their possible therapeutic use, are repression of CYP7A1 and induction of CYP3A4 enzymes that are instrumental for bile salt synthesis and detoxification, respectively [156]. In cholestasis inflammatory and pro-fibrotic genes are activated and it is possible that the anti-inflammatory action of PPAR agonists is equally important [157]. In most trials in which fibrates were tested as treatment of UDCA-refractory PBC, the endpoint has been a decrease of alkaline phosphatase [158]. In a recent meta-analysis on studies comparing patients treated with UDCA plus bezafibrate vs. UDCA alone, combination therapy performed better than monotherapy regarding biochemical parameters but as for symptoms and survival there was no difference (meta-analysis in [159]). In the one study wherein liver stiffness was measured no change was observed [160]. Notably, a significant improvement of pruritus in PBC patients receiving bezafibrate was recently reported [160]. Rigorous testing of histologic endpoints and transplantation-free survival needs to be performed but fibrates may be too weak a PPAR agonists to be successful. Currently stronger PPAR agonists are developed with considerable anti-inflammatory activity [150]. Although primarily developed for the treatment of NASH, these may be promising agents for the treatment of PBC and PSC as well (Table 1).

The pregnane X receptor (PXR): experimental and clinical effects

The PXR (NR1I2) has a critical role in regulating the expression of genes involved in detoxification and metabolism of BA, drugs and other toxins [161]. PXR modulates expression of CYP3A4 and

CYP7A1 [162,163], SULT2A1 [164], UGT1A1, UGT1A3, and UGT1A4 [165,166], MDR1 [167], MRP2 [168], MRP3 [169], and OST β [166]. Cholestatic PXR knockout mice exhibited more hepatic damage than wild-type mice both after bile duct ligation and cholic acid feeding, [170–172]. The potent PXR ligand 5-pregnen-3 β -ol-20-one-16 α -carbonitrile (PCN) reduced (litho-)cholic acid-induced liver injury in wild-type mice, but not in PXR knockout mice [163,172]. Marked upregulation of the basolateral BA efflux transporter MRP3 (ABCC3) may have been crucial in mediating the beneficial effect of PCN [172].

Human PXR agonists include lithocholic acid and a number of drugs including rifampicin, statins, corticosteroids, phenobarbital, and St. John's wort. The antibiotic rifampicin is a potent human PXR activator and an evidence-based treatment for pruritus in cholestatic patients [16]. Rifampicin has been reported to improve serum liver tests in PBC [173,174]. In otherwise healthy gallstone patients, rifampicin induced upregulation of UGT1A1 and MRP2 facilitating bilirubin elimination and increased CYP3A4 expression facilitating detoxification of BAs [53]. Rifampicin also markedly induced CYP3A metabolism in patients with early stage PBC and healthy controls [175]. All these effects were not observed with UDCA indicating that the combined use of UDCA and rifampicin might have synergistic beneficial effects in patients with non-obstructive cholestasis [175]. Rifampicin was reported to be safe in cholestatic liver disease during short-term use for up to two weeks [176]. However, severe hepatotoxicity has been reported in up to 13% of patients with cholestatic disorders after use for more than 4 weeks [174]. Strikingly, rifampicin has been shown most recently to completely reverse severe persistent hepatocellular secretory failure [166] induced by drugs (e.g., clavulanic acid, flucloxacillin, estrogen + progesterone, testosterone, total parenteral nutrition) or transient biliary obstruction (e.g. choledocholithiasis, pancreatic carcinoma) in formerly healthy individuals, an enormous relief for otherwise desperate patients [166]. A prospective, controlled trial on the promising effect of PXR agonists in severe persistent hepatocellular secretory failure is under preparation.

The GR and UDCA: experimental and clinical effects

The use of glucocorticoids to suppress the inflammation in PBC has been always considered as a very attractive approach, but at the cost of serious side effects, especially aggravating osteopenia. Budesonide is a non-halogenated glucocorticoid mainly absorbed in the small intestine. Of an oral dose, 90% is metabolized during the first liver pass in healthy individuals. Compared with prednisolone, GR binding activity of budesonide is 15–20 times higher, so its effect on liver inflammation may be greater. In patients with inflammatory bowel disease and autoimmune hepatitis, oral budesonide has been shown to exert fewer systemic side effects than conventional corticosteroids [177].

Recent studies have shed some light on the complex relationships between GR activation and BAs. Glucocorticoids promote hepatic cholestasis in mice by inhibiting the transcriptional activity of the FXR [178] and increase intestinal apical sodium-dependent bile salt transporter (ASBT) activity [179] while promoting PXR-mediated hydroxylation and sulfation of hydrophobic BAs [180]. Conversely, high BA concentrations might promote SHP-mediated inhibition of GR activation [181]

that could explain some loss of anti-inflammatory effects of glucocorticoids in cholestatic conditions. Taken together these data suggest that classical glucocorticoids may have negative effects in cholestatic conditions.

In contrast to other natural BAs, UDCA is a GR agonist [57–59]. Moreover, combination of budesonide and UDCA have been shown to promote activity of the Cl[−]/HCO₃[−] exchanger AE2, the key transporter involved in alkaline-rich choleresis in humans [182].

Two randomized studies showed the GR (and PXR) agonist budesonide (6–9 mg/day) combined with UDCA to be more effective in improving liver biochemistries and histology than did UDCA alone in patients with stages I–III PBC [183,184]. This was not observed in a study including late stage PBC patients who also developed serious side effects [185]. In an open trial involving non-cirrhotic PBC patients with severe inflammation and biochemical cholestasis not responding to UDCA, the combination of UDCA, budesonide and mycophenolate mofetil – as a corticosteroid-sparing agent – achieved biochemical remission and marked improvement of liver histology [186]. Combination therapy (UDCA + budesonide) is currently being evaluated in comparison to UDCA monotherapy in PBC patients with suboptimal response to UDCA alone in a European multicenter randomized, placebo-controlled trial. Because of its high first-pass hepatic clearance, budesonide should not be given to patients with evidence of cirrhosis and portal hypertension as portosystemic shunting may occur and development of portal vein thrombosis has been described [187].

The vitamin D receptor (VDR): experimental findings

VDR ligands represent potentially attractive agents for pharmacotherapy of autoimmune cholestatic disorders because they may influence several key processes involved in the pathogenesis such as innate and immune activation, BA metabolism and detoxification, bile duct integrity and fibrogenesis. VDR is expressed in almost all immune cells and mediates the immunoregulatory properties of vitamin D. Indeed, vitamin D through VDR interferes directly with T cells by inhibiting the production of T-helper-1 (T_H1) type cytokines, while promoting those of the T_H2 subtype. Furthermore, vitamin D inhibits dendritic cell differentiation resulting also in a decrease in T_H1 cell development. Taken together, these observations indicate that vitamin D through VDR diminishes the effector T cell response suggesting that the vitamin D-VDR axis may be involved in autoimmune diseases [188]. In bile duct epithelial cells, activation of VDR by BAs or vitamin D induces expression of cathelicidin, an anti-microbial peptide known to be protective against bacterial infection. Cathelicidin is known to neutralize the deleterious effects of LPS that accumulates in the biliary tree in fibrosing cholangiopathies [189].

Treatment with VDR agonists stimulates BA detoxification enzymes (such as CYP3A4 and SULT2A1) in the liver and intestine, and protects against lithocholic acid hepatotoxicity [190]. 1,25(OH)₂D₃ was also shown to decrease hepatic Cyp7a1 expression by increasing the expression of Fgf15 in the intestine. Consistently, Cyp7a1 expression was increased in mice lacking VDR when compared to wild-type mice, indicating that intestinal VDR activity controls the basal expression of Cyp7a1 [191].

Review

The anti-fibrotic potential of VDR stimulation has been demonstrated in several models of liver fibrosis, among them the *Mdr2* (*Abcb4*)^{−/−} mice [192,193]. Recently, the vitamin D-VDR axis has been shown to modulate fibrogenesis and hepatic stellate cell activity through a complex mechanism involving epigenetic modifications induced by the SMAD pathway [194].

ASBT inhibitors

The ASBT (SLC10A2) at the luminal surface of ileum enterocytes transports conjugated BA from the gut lumen into the enterocytes [195]. From here BA are secreted into the portal circulation. Under normal physiological conditions ASBT works at maximum capacity since overproduction of BA in the liver leads to increased spill over of BA into the colon [196]. In the colon, BA activate chloride channels and this causes watery diarrhea [197]. Moderate inhibition of ASBT however can have beneficial effects while avoiding diarrhea. ASBT inhibition lowers the intra-mucosal concentration of BA with less activation of FXR, lowered synthesis of FGF19 and unrepressed expression of CYP7A1 in the liver. This causes an enhanced conversion of cholesterol to BA and lowers serum cholesterol. Spill over of BA into the colon will stimulate 7 α -dehydroxylation of BA and stimulate TGR5. This enhances GLP-1 secretion which will increase insulin secretion by the pancreas and improve insulin sensitivity. Anionic exchange resins such as colestevam have a similar effect [198,199]. Thus, ASBT inhibitors are potential drugs for treatment of NAFLD. ASBT inhibitors interrupt the enterohepatic circulation of BA and this may reduce the circulating BA pool, given that the loss exceeds the upregulated BA synthesis. This may be beneficial in cholestatic liver disease but convincing clinical data are currently lacking.

Conclusion

Three decades after the introduction of UDCA as the first anticholestatic agent into clinical practice to treat patients with chronic cholestatic disorders, an enormous progress in the understanding of the molecular pathophysiology of hepatocellular and cholangiocellular cholestasis has led to the development of a variety of novel therapeutic options which are currently under evaluation. Novel immunomodulating approaches are beyond the scope of this review. While UDCA exerts its anticholestatic effects mainly by post-transcriptional mechanisms as a potent intracellular signaling agent and secretagogue, candidates for future combined treatment with UDCA mostly represent transcriptional modulators of secretion and cell protection or membrane receptor agonists. Among these, evaluation of agonists for FXR, GR, and PPAR α in combination with UDCA is far advanced to large scale phase III trials in chronic cholestatic disorders such as PBC. The next line of upcoming therapeutic agents, often in combination with UDCA, for cholestatic disorders includes the 23C-analogue of UDCA, *nor*UDCA, as well as PXR agonists, FGF19 derivatives and ASBT inhibitors. While there was 30 years ago no effective treatment available, promising times for patients with cholestatic disorders and their caring physicians are visible on the horizon.

Financial support

This work was supported by grants from the Deutsche Crohn Colitis Vereinigung (Sektion PSC; to UB) and the Austrian Science Foundation (F3517-B20, to MT).

Conflict of interest

UB signed consultancy agreements (via University of Amsterdam) with Intercept and Novartis and received lecture fees from Falk Foundation, Gilead, Roche.

MT has received research grants from Albireo, Intercept and Falk Pharma, travel support from Falk Foundation and Gilead, and has served as advisor for Albireo, Falk Pharma, Genfit, Intercept and Phenex. MT is listed as co-inventor of patents on the medical use of *nor*UDCA (WO 2006/119803 A1 and WO 2009/013334).

PJ has a consultancy agreement with Shire and has received obeticholic acid from Intercept for basic studies.

RP received fees for an advisory board meeting from Intercept in 2014.

References

- Beuers U, Boyer JL. Bile – A historical review of studies on its form and function. In: Kirsner JB, editor. Gastroenterology in the 20th century. New York: Lea & Febiger; 1994. p. 267–288.
- Trauner M, Meier PJ, Boyer JL. Molecular pathogenesis of cholestasis. N Engl J Med 1998;339:1217–1227.
- Mallette de Buy Wenniger LJ, Oude Elferink RP, Beuers U. Molecular targets for the treatment of fibrosing cholangiopathies. Clin Pharmacol Ther 2012;92:381–387.
- Zollner G, Trauner M. Nuclear receptors as therapeutic targets in cholestatic liver diseases. Br J Pharmacol 2009;156:7–27.
- Hofmann AF, Roda A. Physicochemical properties of bile acids and their relationship to biological properties: an overview of the problem. J Lipid Res 1984;25:1477–1489.
- Makino I, Shinozaki K, Yoshino K, Nakagawa S. Dissolution of cholesterol gallstones by long-term administration of ursodeoxycholic acid. Nippon Shokakibyo Gakkai Zasshi 1975;72:690–702.
- Poupon R, Chretien Y, Poupon RE, Ballet F, Calmus Y, Darnis F. Is ursodeoxycholic acid an effective treatment for primary biliary cirrhosis? Lancet 1987;1:834–836.
- Leuschner U, Fischer H, Kurtz W, Guldutuna S, Hubner K, Hellstern A, et al. Ursodeoxycholic acid in primary biliary cirrhosis: results of a controlled double-blind trial. Gastroenterology 1989;97:1268–1274.
- Poupon RE, Balkau B, Eschwege E, Poupon R. A multicenter, controlled trial of ursodiol for the treatment of primary biliary cirrhosis. UDCA-PBC Study Group. N Engl J Med 1991;324:1548–1554.
- Poupon RE, Balkau B, Guehot J, Heintzmann F. Predictive factors in ursodeoxycholic acid-treated patients with primary biliary cirrhosis: role of serum markers of connective tissue. Hepatology 1994;19:635–640.
- Poupon RE, Lindor KD, Cauch-Dudek K, Dickson ER, Poupon R, Heathcote EJ. Combined analysis of randomized controlled trials of ursodeoxycholic acid in primary biliary cirrhosis. Gastroenterology 1997;113:884–890.
- Poupon RE, Poupon R, Balkau B. Ursodiol for the long-term treatment of primary biliary cirrhosis. The UDCA-PBC Study Group. N Engl J Med 1994;330:1342–1347.
- Poupon RE, Bonnard AM, Chretien Y, Poupon R. Ten-year survival in ursodeoxycholic acid-treated patients with primary biliary cirrhosis. The UDCA-PBC Study Group. Hepatology 1999;29:1668–1671.
- Pares A, Caballeria L, Rodes J, Bruguera M, Rodrigo L, Garcia-Plaza A, et al. Long-term effects of ursodeoxycholic acid in primary biliary cirrhosis: results of a double-blind controlled multicentric trial. UDCA-Cooperative Group from the Spanish Association for the Study of the Liver. J Hepatol 2000;32:561–566.
- Lindor KD, Gershwin ME, Poupon R, Kaplan M, Bergasa NV, Heathcote EJ. Primary biliary cirrhosis. Hepatology 2009;50:291–308.

- [16] EASL Clinical Practice Guidelines: management of cholestatic liver diseases. *J Hepatol* 2009;51:237–267.
- [17] Corpechot C, Carrat F, Bahr A, Chretien Y, Poupon RE, Poupon R. The effect of ursodeoxycholic acid therapy on the natural course of primary biliary cirrhosis. *Gastroenterology* 2005;128:297–303.
- [18] Lammers WJ, van Buuren HR, Hirschfield GM, Janssen HL, Invernizzi P, Mason AL, et al. Levels of alkaline phosphatase and bilirubin are surrogate end points of outcomes of patients with primary biliary cirrhosis: an international follow-up study. *Gastroenterology* 2014;147:1338–1349.
- [19] Pares A, Caballeria L, Rodes J. Excellent long-term survival in patients with primary biliary cirrhosis and biochemical response to ursodeoxycholic acid. *Gastroenterology* 2006;130:715–720.
- [20] Corpechot C, Abenavoli L, Rabahi N, Chretien Y, Andreani T, Johanet C, et al. Biochemical response to ursodeoxycholic acid and long-term prognosis in primary biliary cirrhosis. *Hepatology* 2008;48:871–877.
- [21] Kuiper EM, Hansen BE, de Vries RA, den Ouden-Muller JW, van Ditzhuijsen TJ, Haagsma EB, et al. Improved prognosis of patients with primary biliary cirrhosis that have a biochemical response to ursodeoxycholic acid. *Gastroenterology* 2009;136:1281–1287.
- [22] Poupon R. Primary biliary cirrhosis: a 2010 update. *J Hepatol* 2010;52:745–758.
- [23] Beuers U, Lindor KD. A major step towards effective treatment evaluation in primary biliary cirrhosis. *J Hepatol* 2011;55:1178–1180.
- [24] Shi J, Li Z, Zeng X, Lin Y, Xie WF. Ursodeoxycholic acid in primary sclerosing cholangitis: meta-analysis of randomized controlled trials. *Hepatol Res* 2009;39:865–873.
- [25] Poropat G, Giljaca V, Stimac D, Glud C. Bile acids for primary sclerosing cholangitis. *Cochrane Database Syst Rev* 2011;CD003626.
- [26] Lindor KD, Kowdley KV, Luketic VA, Harrison ME, McCashland T, Befeler AS, et al. High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009;50:808–814.
- [27] Chapman R, Fevery J, Kalloo A, Nagorney DM, Boberg KM, Shneider B, et al. Diagnosis and management of primary sclerosing cholangitis. *Hepatology* 2010;51:660–678.
- [28] Beuers U. Drug insight: mechanisms and sites of action of ursodeoxycholic acid in cholestasis. *Nat Clin Pract Gastroenterol Hepatol* 2006;3:318–328.
- [29] Dilger K, Hohenester S, Winkler-Budenhofen U, Bastiaansen BA, Schaap FG, Rust C, et al. Effect of ursodeoxycholic acid on bile acid profiles and intestinal detoxification machinery in primary biliary cirrhosis and health. *J Hepatol* 2012;57:133–140.
- [30] Kitani K. Hepatoprotective effect of ursodeoxycholate in experimental animals. In: Paumgartner G et al., editors. *Strategies for the treatment of hepatobiliary diseases*. Lancaster: Kluwer; 1990. p. 44–56.
- [31] Bouscarel B, Fromm H, Nussbaum R. Ursodeoxycholate mobilizes intracellular Ca^{2+} and activates phosphorylase a in isolated hepatocytes. *Am J Physiol* 1993;264:G243–251.
- [32] Beuers U, Nathanson MH, Boyer JL. Effects of tauroursodeoxycholic acid on cytosolic Ca^{2+} signals in isolated rat hepatocytes. *Gastroenterology* 1993;104:604–612.
- [33] Beuers U, Nathanson MH, Isles CM, Boyer JL. Tauroursodeoxycholic acid stimulates hepatocellular exocytosis and mobilizes extracellular Ca^{++} mechanisms defective in cholestasis. *J Clin Invest* 1993;92:2984–2993.
- [34] Beuers U, Thiel M, Bardenheuer H, Paumgartner G. Tauroursodeoxycholic acid inhibits the cytosolic Ca^{++} increase in human neutrophils stimulated by formyl-methionyl-leucyl-phenylalanine. *Biochem Biophys Res Commun* 1990;171:1115–1121.
- [35] Bouscarel B, Gettys TW, Fromm H, Dubner H. Ursodeoxycholic acid inhibits glucagon-induced cAMP formation in hamster hepatocytes: a role for PKC. *Am J Physiol* 1995;268:G300–G310.
- [36] Beuers U, Throckmorton DC, Anderson MS, Isles CM, Thasler W, Kullak-Ublick GA, et al. Tauroursodeoxycholic acid activates protein kinase C in isolated rat hepatocytes. *Gastroenterology* 1996;110:1553–1563.
- [37] Stravitz RT, Rao YP, Vlahcevic ZR, Gurley EC, Jarvis WD, Hylemon PB. Hepatocellular protein kinase C activation by bile acids: implications for regulation of cholesterol 7 α -hydroxylase. *Am J Physiol* 1996;271:G293–G303.
- [38] Schliess F, Kurz AK, vom Dahl S, Haussinger D. Mitogen-activated protein kinases mediate the stimulation of bile acid secretion by tauroursodeoxycholate in rat liver. *Gastroenterology* 1997;113:1306–1314.
- [39] Kurz AK, Graf D, Schmitt M, Vom Dahl S, Haussinger D. Tauroursodeoxycholate-induced cholestasis involves p38(MAPK) activation and translocation of the bile salt export pump in rats. *Gastroenterology* 2001;121:407–419.
- [40] Haussinger D, Kurz AK, Wettstein M, Graf D, Vom Dahl S, Schliess F. Involvement of integrins and Src in tauroursodeoxycholate-induced and swelling-induced cholestasis. *Gastroenterology* 2003;124:1476–1487.
- [41] Gohlke H, Schmitz B, Sommerfeld A, Reinehr R, Haussinger D. Alpha5 beta1-integrins are sensors for tauroursodeoxycholic acid in hepatocytes. *Hepatology* 2013;57:1117–1129.
- [42] Haussinger D, Saha N, Hallbrucker C, Lang F, Gerok W. Involvement of microtubules in the swelling-induced stimulation of transcellular taurocholate transport in perfused rat liver. *Biochem J* 1993;291:355–360.
- [43] Marziani M, Francis H, Benedetti A, Ueno Y, Fava G, Venter J, et al. Ca^{2+} -dependent cytoprotective effects of ursodeoxycholic and tauroursodeoxycholic acid on the biliary epithelium in a rat model of cholestasis and loss of bile ducts. *Am J Pathol* 2006;168:398–409.
- [44] Beuers U, Bilzer M, Chittattu A, Kullak-Ublick GA, Keppler D, Paumgartner G, et al. Tauroursodeoxycholic acid inserts the apical conjugate export pump, Mrp2, into canalicular membranes and stimulates organic anion secretion by protein kinase C-dependent mechanisms in cholestatic rat liver. *Hepatology* 2001;33:1206–1216.
- [45] Wimmer R, Hohenester S, Pusl T, Denk GU, Rust C, Beuers U. Tauroursodeoxycholic acid exerts anticholestatic effects by a cooperative cPKC α -PKA-dependent mechanism in rat liver. *Gut* 2008;57:1448–1454.
- [46] Cruz LN, Guerra MT, Kruglov E, Mennone A, Garcia CR, Chen J, et al. Regulation of multidrug resistance-associated protein 2 by calcium signaling in mouse liver. *Hepatology* 2010;52:327–337.
- [47] Beuers U, Hohenester S, de Buy Wenniger LM, Kremer AE, Jansen PLM, Oude Elferink RP. The biliary HCO_3^- umbrella. *Hepatology* 2010;52:1489–1496.
- [48] Hohenester S, Wenniger LM, Paulusma CC, van Vliet SJ, Jefferson DM, Elferink RP, et al. A biliary HCO_3^- umbrella constitutes a protective mechanism against bile acid-induced injury in human cholangiocytes. *Hepatology* 2012;55:173–183.
- [49] Prieto J, Garcia N, Marti-Clement JM, Penuelas I, Richter JA, Medina JF. Assessment of biliary bicarbonate secretion in humans by positron emission tomography. *Gastroenterology* 1999;117:167–172.
- [50] Prieto J, Qian C, Garcia N, Diez J, Medina JF. Abnormal expression of anion exchanger genes in primary biliary cirrhosis. *Gastroenterology* 1993;105:572–578.
- [51] Medina JF, Martinez A, Vazquez JJ, Prieto J. Decreased anion exchanger 2 immunoreactivity in the liver of patients with primary biliary cirrhosis. *Hepatology* 1997;25:12–17.
- [52] Beuers U, Spengler U, Zwiebel FM, Pauletzki J, Fischer S, Paumgartner G. Effect of ursodeoxycholic acid on the kinetics of the major hydrophobic bile acids in health and in chronic cholestatic liver disease. *Hepatology* 1992;15:603–608.
- [53] Marschall HU, Wagner M, Zollner G, Fickert P, Diczfalussy U, Gumhold J, et al. Complementary stimulation of hepatobiliary transport and detoxification systems by rifampicin and ursodeoxycholic acid in humans. *Gastroenterology* 2005;129:476–485.
- [54] Benz C, Angermuller S, Tox U, Kloters-Plachky P, Riedel HD, Sauer P, et al. Effect of tauroursodeoxycholic acid on bile-acid-induced apoptosis and cytolysis in rat hepatocytes. *J Hepatol* 1998;28:99–106.
- [55] Lazaridis KN, Gores GJ, Lindor KD. Ursodeoxycholic acid 'mechanisms of action and clinical use in hepatobiliary disorders'. *J Hepatol* 2001;35:134–146.
- [56] Rodrigues CM, Fan G, Ma X, Kren BT, Steer CJ. A novel role for ursodeoxycholic acid in inhibiting apoptosis by modulating mitochondrial membrane perturbation. *J Clin Invest* 1998;101:2790–2799.
- [57] Tanaka H, Makino I. Ursodeoxycholic acid-dependent activation of the glucocorticoid receptor. *Biochem Biophys Res Commun* 1992;188:942–948.
- [58] Miura T, Ouchida R, Yoshikawa N, Okamoto K, Makino Y, Nakamura T, et al. Functional modulation of the glucocorticoid receptor and suppression of NF- κ B-dependent transcription by ursodeoxycholic acid. *J Biol Chem* 2001;276:47371–47378.
- [59] Takigawa T, Miyazaki H, Kinoshita M, Kawarabayashi N, Nishiyama K, Hatusue K, et al. Glucocorticoid receptor-dependent immunomodulatory effect of ursodeoxycholic acid on liver lymphocytes in mice. *Am J Physiol Gastrointest Liver Physiol* 2013;305:G427–G438.
- [60] Tanaka H, Makino Y, Miura T, Hirano F, Okamoto K, Komura K, et al. Ligand-independent activation of the glucocorticoid receptor by ursodeoxycholic acid. Repression of IFN- γ -induced MHC class II gene expression via a glucocorticoid receptor-dependent pathway. *J Immunol* 1996;156:1601–1608.
- [61] Sola S, Amaral JD, Castro RE, Ramalho RM, Borralho PM, Kren BT, et al. Nuclear translocation of UDCA by the glucocorticoid receptor is required to

Review

- reduce TGF- β 1-induced apoptosis in rat hepatocytes. *Hepatology* 2005;42:925–934.
- [62] Hofmann AF, Hagey LR. Key discoveries in bile acid chemistry and biology and their clinical applications: history of the last eight decades. *J Lipid Res* 2014;55:1553–1595.
 - [63] Hofmann AF, Zakko SF, Lira M, Clerici C, Hagey LR, Lambert KK, et al. Novel biotransformation and physiological properties of norursodeoxycholic acid in humans. *Hepatology* 2005;42:1391–1398.
 - [64] Yeh HZ, Schteingart CD, Hagey LR, Ton-Nu HT, Bolder U, Gavrilkina MA, et al. Effect of side chain length on biotransformation, hepatic transport, and choleretic properties of chenodeoxycholy homologues in the rodent: studies with dinorchenodeoxycholic acid, norchenodeoxycholic acid, and chenodeoxycholic acid. *Hepatology* 1997;26:374–385.
 - [65] Yoon YB, Hagey LR, Hofmann AF, Gurantz D, Michelotti EL, Steinbach JH. Effect of side-chain shortening on the physiologic properties of bile acids: hepatic transport and effect on biliary secretion of 23-nor-ursodeoxycholate in rodents. *Gastroenterology* 1986;90:837–852.
 - [66] Dumont M, Erlinger S, Uchman S. Hypercholeremia induced by ursodeoxycholic acid and 7-ketolithocholic acid in the rat: possible role of bicarbonate transport. *Gastroenterology* 1980;79:82–89.
 - [67] Trauner M, Fickert P, Halilbasic E, Moustafa T. Lessons from the toxic bile concept for the pathogenesis and treatment of cholestatic liver diseases. *Wien Med Wochenschr* 2008;158:542–548.
 - [68] Fickert P, Wagner M, Marschall HU, Fuchsichler A, Zollner G, Tsybrovskyy O, et al. 24-norUrsodeoxycholic acid is superior to ursodeoxycholic acid in the treatment of sclerosing cholangitis in Mdr2 (Abcb4) knockout mice. *Gastroenterology* 2006;130:465–481.
 - [69] Halilbasic E, Fiorotto R, Fickert P, Marschall HU, Moustafa T, Spirl C, et al. Side chain structure determines unique physiologic and therapeutic properties of norursodeoxycholic acid in Mdr2 $^{-/-}$ mice. *Hepatology* 2009;49:1972–1981.
 - [70] Fickert P, Pollheimer MJ, Silbert D, Moustafa T, Halilbasic E, Krones E, et al. Differential effects of norUDCA and UDCA in obstructive cholestasis in mice. *J Hepatol* 2013;58:1201–1208.
 - [71] Moustafa T, Fickert P, Magnes C, Guelly C, Thueringer A, Frank S, et al. Alterations in lipid metabolism mediate inflammation, fibrosis, and proliferation in a mouse model of chronic cholestatic liver injury. *Gastroenterology* 2012;142:e112.
 - [72] Denk GU, Maitz S, Wimmer R, Rust C, Invernizzi P, Ferdinandusse S, et al. Conjugation is essential for the anticholestatic effect of NorUrsodeoxycholic acid in tauroolithocholic acid-induced cholestasis in rat liver. *Hepatology* 2010;52:1758–1768.
 - [73] Trauner M, Halilbasic E, Kazemi-Shirazi L, Kienbacher C, Stauder K, Traussnigg S, et al. Therapeutic role of bile acids and nuclear receptor agonists in fibrosing cholangiopathies. *Dig Dis* 2014;32:631–636.
 - [74] Trauner M, Fickert P, Wagner M. MDR3 (ABCB4) defects: a paradigm for the genetics of adult cholestatic syndromes. *Semin Liver Dis* 2007;27:77–98.
 - [75] Forman BM, Goode E, Chen J, Oro AE, Bradley DJ, Perlmann T, et al. Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* 1995;81:687–693.
 - [76] Seol W, Choi HS, Moore DD. Isolation of proteins that interact specifically with the retinoid X receptor: two novel orphan receptors. *Mol Endocrinol* 1995;9:72–85.
 - [77] Makishima M, Okamoto AY, Repa JJ, Tu H, Learned RM, Luk A, et al. Identification of a nuclear receptor for bile acids. *Science* 1999;284:1362–1365.
 - [78] Baes M, Gulick T, Choi HS, Martinoli MG, Simha D, Moore DD. A new orphan member of the nuclear hormone receptor superfamily that interacts with a subset of retinoic acid response elements. *Mol Cell Biol* 1994;14:1544–1552.
 - [79] Makishima M, Lu TT, Xie W, Whitfield GK, Domoto H, Evans RM, et al. Vitamin D receptor as an intestinal bile acid sensor. *Science* 2002;296:1313–1316.
 - [80] Houten SM, Watanabe M, Auwerx J. Endocrine functions of bile acids. *EMBO J* 2006;25:1419–1425.
 - [81] Le Martelot G, Claudel T, Gatifield D, Schaad O, Kornmann B, Lo Sasso G, et al. REV-ERB α participates in circadian SREBP signaling and bile acid homeostasis. *PLoS Biol* 2009;7:e1000181.
 - [82] Duez H, van der Veen JN, Duhem C, Pourcet B, Touvier T, Fontaine C, et al. Regulation of bile acid synthesis by the nuclear receptor Rev-erb α . *Gastroenterology* 2008;135:689–698.
 - [83] Wang H, Chen J, Hollister K, Sowers LC, Forman BM. Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. *Mol Cell* 1999;3:543–553.
 - [84] Goodwin B, Jones SA, Price RR, Watson MA, McKee DD, Moore LB, et al. A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Mol Cell* 2000;6:517–526.
 - [85] Ananthanarayanan M, Balasubramanian N, Makishima M, Mangelsdorf DJ, Suchy FJ. Human bile salt export pump promoter is transactivated by the farnesoid X receptor/bile acid receptor. *J Biol Chem* 2001;276:28857–28865.
 - [86] Boyer JL, Trauner M, Mennone A, Soroka CJ, Cai SY, Moustafa T, et al. Upregulation of a basolateral FXR-dependent bile acid efflux transporter OST α -OST β in cholestasis in humans and rodents. *Am J Physiol Gastrointest Liver Physiol* 2006;290:G1124–G1130.
 - [87] Claudel T, Staels B, Kuipers F. The Farnesoid X receptor: a molecular link between bile acid and lipid and glucose metabolism. *Arterioscler Thromb Vasc Biol* 2005;25:2020–2030.
 - [88] Inagaki T, Choi M, Moschetta A, Peng L, Cummins CL, McDonald JG, et al. Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. *Cell Metab* 2005;2:217–225.
 - [89] Watanabe M, Houten SM, Wang L, Moschetta A, Mangelsdorf DJ, Heyman RA, et al. Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c. *J Clin Invest* 2004;113:1408–1418.
 - [90] Wang L, Liu J, Saha P, Huang J, Chan L, Spiegelman B, et al. The orphan nuclear receptor SHP regulates PGC-1 α expression and energy production in brown adipocytes. *Cell Metab* 2005;2:227–238.
 - [91] Wang L, Huang J, Saha P, Kulkarni RN, Hu M, Kim Y, et al. Orphan receptor small heterodimer partner is an important mediator of glucose homeostasis. *Mol Endocrinol* 2006;20:2671–2681.
 - [92] Potthoff MJ, Kliewer SA, Mangelsdorf DJ. Endocrine fibroblast growth factors 15/19 and 21: from feast to famine. *Genes Dev* 2012;26:312–324.
 - [93] Cicione C, Degirolamo C, Moschetta A. Emerging role of fibroblast growth factors 15/19 and 21 as metabolic integrators in the liver. *Hepatology* 2012;56:2404–2411.
 - [94] Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, Parameswara V, et al. Endocrine regulation of the fasting response by PPAR α -mediated induction of fibroblast growth factor 21. *Cell Metab* 2007;5:415–425.
 - [95] Owen BM, Mangelsdorf DJ, Kliewer SA. Tissue-specific actions of the metabolic hormones FGF15/19 and FGF21. *Trends Endocrinol Metab* 2015;26:22–29.
 - [96] Bhatnagar S, Damron HA, Hillgartner FB. Fibroblast growth factor-19, a novel factor that inhibits hepatic fatty acid synthesis. *J Biol Chem* 2009;284:10023–10033.
 - [97] Kir S, Beddow SA, Samuel VT, Miller P, Previs SF, Suino-Powell K, et al. FGF19 as a postprandial, insulin-independent activator of hepatic protein and glycogen synthesis. *Science* 2011;331:1621–1624.
 - [98] Potthoff MJ, Boney-Montoya J, Choi M, He T, Sunney NE, Satapati S, et al. FGF15/19 regulates hepatic glucose metabolism by inhibiting the CREB-PGC-1 α pathway. *Cell Metab* 2011;13:729–738.
 - [99] Kim I, Ahn SH, Inagaki T, Choi M, Ito S, Guo GL, et al. Differential regulation of bile acid homeostasis by the farnesoid X receptor in liver and intestine. *J Lipid Res* 2007;48:2664–2672.
 - [100] Modica S, Petruzzelli M, Bellafante E, Murzilli S, Salvatore L, Celli N, et al. Selective activation of nuclear bile acid receptor FXR in the intestine protects mice against cholestasis. *Gastroenterology* 2012;142:355–365.
 - [101] Li S, Hsu DD, Li B, Luo X, Alderson N, Qiao L, et al. Cytoplasmic tyrosine phosphatase Shp2 coordinates hepatic regulation of bile acid and FGF15/19 signaling to repress bile acid synthesis. *Cell Metab* 2014;20:320–332.
 - [102] Badman MK, Pissios P, Kennedy AR, Koukos G, Flier JS, Maratos-Flier E. Hepatic fibroblast growth factor 21 is regulated by PPAR α and is a key mediator of hepatic lipid metabolism in ketotic states. *Cell Metab* 2007;5:426–437.
 - [103] Chau MD, Gao J, Yang Q, Wu Z, Gromada J. Fibroblast growth factor 21 regulates energy metabolism by activating the AMPK-SIRT1-PGC-1 α pathway. *Proc Natl Acad Sci U S A* 2010;107:12553–12558.
 - [104] Markan KR, Naber MC, Ameka MK, Anderregg MD, Mangelsdorf DJ, Kliewer SA, et al. Circulating FGF21 is liver derived and enhances glucose uptake during refeeding and overfeeding. *Diabetes* 2014;63:4057–4063.
 - [105] Potthoff MJ, Inagaki T, Satapati S, Ding X, He T, Goetz R, et al. FGF21 induces PGC-1 α and regulates carbohydrate and fatty acid metabolism during the adaptive starvation response. *Proc Natl Acad Sci U S A* 2009;106:10853–10858.
 - [106] Holland WL, Adams AC, Brozinick JT, Bui HH, Miyauchi Y, Kusminski CM, et al. An FGF21-adiponectin-ceramide axis controls energy expenditure and insulin action in mice. *Cell Metab* 2013;17:790–797.
 - [107] Lin Z, Tian H, Lam KS, Lin S, Hoo RC, Konishi M, et al. Adiponectin mediates the metabolic effects of FGF21 on glucose homeostasis and insulin sensitivity in mice. *Cell Metab* 2013;17:779–789.

- [108] Pellicciari R, Fiorucci S, Camaioni E, Clerici C, Costantino G, Maloney PR, et al. Galpha-ethyl-chenodeoxycholic acid (6-ECDCA), a potent and selective FXR agonist endowed with anticholestatic activity. *J Med Chem* 2002;45:3569–3572.
- [109] Hambruch E, Miyazaki-Anzai S, Hahn U, Matysik S, Boettcher A, Perovic-Ottstadt S, et al. Synthetic farnesoid X receptor agonists induce high-density lipoprotein-mediated transhepatic cholesterol efflux in mice and monkeys and prevent atherosclerosis in cholesteryl ester transfer protein transgenic low-density lipoprotein receptor (–/–) mice. *J Pharmacol Exp Ther* 2012;343:556–567.
- [110] Flatt B, Martin R, Wang TL, Mahaney P, Murphy B, Gu XH, et al. Discovery of XL335 (WAY-362450), a highly potent, selective, and orally active agonist of the farnesoid X receptor (FXR). *J Med Chem* 2009;52:904–907.
- [111] Soisson SM, Parthasarathy G, Adams AD, Sahoo S, Sitlani A, Sparrow C, et al. Identification of a potent synthetic FXR agonist with an unexpected mode of binding and activation. *Proc Natl Acad Sci U S A* 2008;105:5337–5342.
- [112] Bass JY, Caravella JA, Chen L, Creech KL, Deaton DN, Madauss KP, et al. Conformationally constrained farnesoid X receptor (FXR) agonists: heteroaryl replacements of the naphthalene. *Bioorg Med Chem Lett* 2011;21:1206–1213.
- [113] Hirschfield GM, Mason A, Luketic V, Lindor K, Gordon S, Mayo M, et al. Efficacy of obeticholic acid in patients with primary biliary cirrhosis and inadequate response to ursodeoxycholic acid. *Gastroenterology* 2015, [Epub ahead of print].
- [114] Schaap FG, Trauner M, Jansen PL. Bile acid receptors as targets for drug development. *Nat Rev Gastroenterol Hepatol* 2014;11:55–67.
- [115] Beuers U, Kremer AE, Bolier R, Elferink RP. Pruritus in cholestasis: facts and fiction. *Hepatology* 2014;60:399–407.
- [116] Luo J, Ko B, Elliott M, Zhou M, Lindhout DA, Phung V, et al. A nontumorigenic variant of FGF19 treats cholestatic liver diseases. *Sci Transl Med* 2014;6:247a100.
- [117] Nicholes K, Guillet S, Tomlinson E, Hillan K, Wright B, Frantz GD, et al. A mouse model of hepatocellular carcinoma: ectopic expression of fibroblast growth factor 19 in skeletal muscle of transgenic mice. *Am J Pathol* 2002;160:2295–2307.
- [118] Maruyama T, Miyamoto Y, Nakamura T, Tamai Y, Okada H, Sugiyama E, et al. Identification of membrane-type receptor for bile acids (M-BAR). *Biochem Biophys Res Commun* 2002;298:714–719.
- [119] Sato H, Macchiarulo A, Thomas C, Gioiello A, Une M, Hofmann AF, et al. Novel potent and selective bile acid derivatives as TGR5 agonists: biological screening, structure-activity relationships, and molecular modeling studies. *J Med Chem* 2008;51:1831–1841.
- [120] Kawamata Y, Fujii R, Hosoya M, Harada M, Yoshida H, Miwa M, et al. A G protein-coupled receptor responsive to bile acids. *J Biol Chem* 2003;278:9435–9440.
- [121] Maruyama T, Tanaka K, Suzuki J, Miyoshi H, Harada N, Nakamura T, et al. Targeted disruption of G protein-coupled bile acid receptor 1 (Gpbar1/M-BAR) in mice. *J Endocrinol* 2006;191:197–205.
- [122] Keitel V, Cupisti K, Ullmer C, Knoefel WT, Kubitz R, Haussinger D. The membrane-bound bile acid receptor TGR5 is localized in the epithelium of human gallbladders. *Hepatology* 2009;50:861–870.
- [123] Vassileva G, Golovko A, Markowitz L, Abbondanzo SJ, Zeng M, Yang S, et al. Targeted deletion of Gpbar1 protects mice from cholesterol gallstone formation. *Biochem J* 2006;398:423–430.
- [124] Keitel V, Donner M, Winandy S, Kubitz R, Haussinger D. Expression and function of the bile acid receptor TGR5 in Kupffer cells. *Biochem Biophys Res Commun* 2008;372:78–84.
- [125] Keitel V, Reinehr R, Gatsios P, Rupprecht C, Gorg B, Selbach O, et al. The G-protein coupled bile salt receptor TGR5 is expressed in liver sinusoidal endothelial cells. *Hepatology* 2007;45:695–704.
- [126] Keitel V, Ullmer C, Haussinger D. The membrane-bound bile acid receptor TGR5 (Gpbar-1) is localized in the primary cilium of cholangiocytes. *Biol Chem* 2010;391:785–789.
- [127] Wang YD, Chen WD, Yu D, Forman BM, Huang W. The G-Protein-coupled bile acid receptor, Gpbar1 (TGR5), negatively regulates hepatic inflammatory response through antagonizing nuclear factor kappa light-chain enhancer of activated B cells (NF-kappaB) in mice. *Hepatology* 2011;54:1421–1432.
- [128] Pols TW, Nomura M, Harach T, Lo Sasso G, Oosterveer MH, Thomas C, et al. TGR5 activation inhibits atherosclerosis by reducing macrophage inflammation and lipid loading. *Cell Metab* 2011;14:747–757.
- [129] Wang YD, Chen WD, Yu D, Forman BM, Huang W. The G-protein coupled bile acid receptor Gpbar1 (TGR5) negatively regulates hepatic inflammatory response through antagonizing Nuclear Factor kappaB. *Hepatology* 2011;54:1421–1432.
- [130] Cipriani S, Mencarelli A, Chini MG, Distrutti E, Renga B, Bifulco G, et al. The bile acid receptor GPBAR-1 (TGR5) modulates integrity of intestinal barrier and immune response to experimental colitis. *PLoS One* 2011;6:e25637.
- [131] Alemi F, Poole DP, Chiu J, Schoonjans K, Cattaruzza F, Grider JR, et al. The receptor TGR5 mediates the prokinetic actions of intestinal bile acids and is required for normal defecation in mice. *Gastroenterology* 2013;144:145–154.
- [132] Poole DP, Godfrey C, Cattaruzza F, Cottrell GS, Kirkland JG, Pelayo JC, et al. Expression and function of the bile acid receptor GpBAR1 (TGR5) in the murine enteric nervous system. *Neurogastroenterol Motil* 2010;22:814–825.
- [133] Ward JB, Mroz MS, Keely SJ. The bile acid receptor, TGR5, regulates basal and cholinergic-induced secretory responses in rat colon. *Neurogastroenterol Motil* 2013;25:708–711.
- [134] Rajagopal S, Kumar DP, Mahavadi S, Bhattacharya S, Zhou R, Corvera CU, et al. Activation of G protein-coupled bile acid receptor, TGR5, induces smooth muscle relaxation via both Epac- and PKA-mediated inhibition of RhoA/Rho kinase pathway. *Am J Physiol Gastrointest Liver Physiol* 2013;304:G527–G535.
- [135] Pean N, Doignon I, Garcin I, Besnard A, Julien B, Liu B, et al. The receptor TGR5 protects the liver from bile acid overload during liver regeneration in mice. *Hepatology* 2013;58:1451–1460.
- [136] Lavoie B, Balemba OB, Godfrey C, Watson CA, Vassileva G, Corvera CU, et al. Hydrophobic bile salts inhibit gallbladder smooth muscle function via stimulation of GPBAR1 receptors and activation of KATP channels. *J Physiol* 2010;588:3295–3305.
- [137] Doignon I, Julien B, Serriere-Lanneau V, Garcin I, Alonso G, Nicou A, et al. Immediate neuroendocrine signaling after partial hepatectomy through acute portal hyperpressure and cholestasis. *J Hepatol* 2011;54:481–488.
- [138] Hov JR, Keitel V, Laerdahl JK, Spomer L, Ellinghaus E, ElSharawy A, et al. Mutational characterization of the bile acid receptor TGR5 in primary sclerosing cholangitis. *PLoS One* 2010;5:e12403.
- [139] Hov JR, Keitel V, Schrupf E, Haussinger D, Karlsen TH. TGR5 sequence variation in primary sclerosing cholangitis. *Dig Dis* 2011;29:78–84.
- [140] Baghdasaryan A, Claudel T, Gumhold J, Silbert D, Adorini L, Roda A, et al. Dual farnesoid X receptor/TGR5 agonist INT-767 reduces liver injury in the Mdr2–/– (Abcb4–/–) mouse cholangiopathy model by promoting biliary HCO output. *Hepatology* 2011;54:1303–1312.
- [141] Baghdasaryan A, Jha P, Müller M, Agostinelli L, Saccomanno S, Auer N, et al. Protective role of membrane bile acid receptor TGR5 (GPBAR1) in DDC-induced sclerosing cholangitis in mice. *J Hepatol* 2014;60:S197–S198.
- [142] Perides G, Laukkanen JM, Vassileva G, Steer ML. Biliary acute pancreatitis in mice is mediated by the G-protein-coupled cell surface bile acid receptor Gpbar1. *Gastroenterology* 2010;138:715–725.
- [143] Keitel V, Gorg B, Bidmon HJ, Zemstova I, Spomer L, Zilles K, et al. The bile acid receptor TGR5 (Gpbar-1) acts as a neurosteroid receptor in brain. *Glia* 2010;58:1794–1805.
- [144] Desai MS, Shabier Z, Taylor M, Lam F, Thevananthar S, Kusters A, et al. Hypertrophic cardiomyopathy and dysregulation of cardiac energetics in a mouse model of biliary fibrosis. *Hepatology* 2010;51:2097–2107.
- [145] Alemi F, Kwon E, Poole DP, Lieu T, Lyo V, Cattaruzza F, et al. The TGR5 receptor mediates bile acid-induced itch and analgesia. *J Clin Invest* 2013;123:1513–1530.
- [146] Lieu T, Jayaweera G, Zhao P, Poole DP, Jensen D, Grace M, et al. The bile acid receptor TGR5 activates the TRPA1 channel to induce itch in mice. *Gastroenterology* 2014;147:1417–1428.
- [147] Yang JJ, Yoon JH, Myung SJ, Gwak GY, Kim W, Chung GE, et al. Bile acid-induced TGR5-dependent c-Jun-N terminal kinase activation leads to enhanced caspase 8 activation in hepatocytes. *Biochem Biophys Res Commun* 2007;361:156–161.
- [148] Keitel V, Reich M, Sommerfeld A, Kluge S, Kubitz R, Haussinger D. The bile acid receptor TGR5 promotes cholangiocyte proliferation through a cSRC–EGFR–ERK signaling pathway. *J Hepatol* 2013;58:S574–S575.
- [149] Cao W, Tian W, Hong J, Li D, Tavares R, Noble L, et al. Expression of bile acid receptor TGR5 in gastric adenocarcinoma. *Am J Physiol Gastrointest Liver Physiol* 2013;304:G322–G327.
- [150] Pawlak M, Lefebvre P, Staels B. Molecular mechanism of PPARalpha action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J Hepatol* 2015;62:720–733.
- [151] Pawlak M, Bauge E, Bourguet W, De Bosscher K, Lallouy F, Tailleux A, et al. The transrepressive activity of peroxisome proliferator-activated receptor alpha is necessary and sufficient to prevent liver fibrosis in mice. *Hepatology* 2014;60:1593–1606.

Review

- [152] Kok T, Wolters H, Bloks VW, Havinga R, Jansen PL, Staels B, et al. Induction of hepatic ABC transporter expression is part of the PPARalpha-mediated fasting response in the mouse. *Gastroenterology* 2003;124:160–171.
- [153] Kok T, Bloks VW, Wolters H, Havinga R, Jansen PL, Staels B, et al. Peroxisome proliferator-activated receptor alpha (PPARalpha)-mediated regulation of multidrug resistance 2 (Mdr2) expression and function in mice. *Biochem J* 2003;369:539–547.
- [154] Matsumoto T, Miyazaki H, Nakahashi Y, Hirohara J, Seki T, Inoue K, et al. Multidrug resistance3 is in situ detected in the liver of patients with primary biliary cirrhosis, and induced in human hepatoma cells by bezafibrate. *Hepatol Res* 2004;30:125–136.
- [155] Shoda J, Inada Y, Tsuji A, Kusama H, Ueda T, Ikegami T, et al. Bezafibrate stimulates canalicular localization of NBD-labeled PC in HepG2 cells by PPARalpha-mediated redistribution of ABCB4. *J Lipid Res* 2004;45:1813–1825.
- [156] Honda A, Ikegami T, Nakamuta M, Miyazaki T, Iwamoto J, Hirayama T, et al. Anticholestatic effects of bezafibrate in patients with primary biliary cirrhosis treated with ursodeoxycholic acid. *Hepatology* 2013;57:1931–1941.
- [157] Ghonem NS, Assis DN, Boyer JL. On fibrates and cholestasis: a review. *Hepatology* 2015. <http://dx.doi.org/10.1002/hep.27744>. [Epub ahead of print].
- [158] Cuperus FJ, Halilbasic E, Trauner M. Fibrate treatment for primary biliary cirrhosis. *Curr Opin Gastroenterol* 2014;30:279–286.
- [159] Zhang Y, Chen K, Dai W, Xia Y, Wang F, Shen M, et al. Combination therapy of bezafibrate and ursodeoxycholic acid for primary biliary cirrhosis: a meta-analysis. *Hepatol Res* 2015;45:48–58.
- [160] Lens S, Leoz M, Nazal L, Bruguera M, Pares A. Bezafibrate normalizes alkaline phosphatase in primary biliary cirrhosis patients with incomplete response to ursodeoxycholic acid. *Liver Int* 2014;34:197–203.
- [161] Kliewer SA, Willson TM. Regulation of xenobiotic and bile acid metabolism by the nuclear pregnane X receptor. *J Lipid Res* 2002;43:359–364.
- [162] Bertilsson G, Heidrich J, Svensson K, Asman M, Jendeborg L, Sydow-Backman M, et al. Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc Natl Acad Sci U S A* 1998;95:12208–12213.
- [163] Staudinger JL, Goodwin B, Jones SA, Hawkins-Brown D, MacKenzie KI, LaTour A, et al. The nuclear receptor PXR is a lithocholic acid sensor that protects against liver toxicity. *Proc Natl Acad Sci U S A* 2001;98:3369–3374.
- [164] Fang HL, Strom SC, Ellis E, Duanmu Z, Fu J, Duniec-Dmuchowski Z, et al. Positive and negative regulation of human hepatic hydroxysteroid sulfotransferase (SULT2A1) gene transcription by rifampicin: roles of hepatocyte nuclear factor 4alpha and pregnane X receptor. *J Pharmacol Exp Ther* 2007;323:586–598.
- [165] Gardner-Stephen D, Heydel JM, Goyal A, Lu Y, Xie W, Lindblom T, et al. Human PXR variants and their differential effects on the regulation of human UDP-glucuronosyltransferase gene expression. *Drug Metab Dispos* 2004;32:340–347.
- [166] van Dijk R, Kremer AE, Smit W, van den Elzen B, van Gulik T, Gouma D, Lameris JS, et al. Characterization and treatment of persistent hepatocellular secretory failure. *Liver Int* 2014. <http://dx.doi.org/10.1111/liv.12603>. [Epub ahead of print].
- [167] Geick A, Eichelbaum M, Burk O. Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J Biol Chem* 2001;276:14581–14587.
- [168] Kast HR, Goodwin B, Tarr PT, Jones SA, Anisfeld AM, Stoltz CM, et al. Regulation of multidrug resistance-associated protein 2 (ABCC2) by the nuclear receptors pregnane X receptor, farnesoid X-activated receptor, and constitutive androstane receptor. *J Biol Chem* 2002;277:2908–2915.
- [169] Teng S, Jekerle V, Piquette-Miller M. Induction of ABCC3 (MRP3) by pregnane X receptor activators. *Drug Metab Dispos* 2003;31:1296–1299.
- [170] Stedman C, Liddle C, Coulter S, Sonoda J, Alvarez JG, Evans RM, et al. Benefit of farnesoid X receptor inhibition in obstructive cholestasis. *Proc Natl Acad Sci U S A* 2006;103:11323–11328.
- [171] Stedman CA, Liddle C, Coulter SA, Sonoda J, Alvarez JG, Moore DD, et al. Nuclear receptors constitutive androstane receptor and pregnane X receptor ameliorate cholestatic liver injury. *Proc Natl Acad Sci U S A* 2005;102:2063–2068.
- [172] Teng S, Piquette-Miller M. Hepatoprotective role of PXR activation and MRP3 in cholic acid-induced cholestasis. *Br J Pharmacol* 2007;151:367–376.
- [173] Bachs L, Pares A, Elena M, Piera C, Rodes J. Comparison of rifampicin with phenobarbitone for treatment of pruritus in biliary cirrhosis. *Lancet* 1989;1:574–576.
- [174] Bachs L, Pares A, Elena M, Piera C, Rodes J. Effects of long-term rifampicin administration in primary biliary cirrhosis. *Gastroenterology* 1992;102:2077–2080.
- [175] Dilger K, Denk A, Heeg MH, Beuers U. No relevant effect of ursodeoxycholic acid on cytochrome P450 3A metabolism in primary biliary cirrhosis. *Hepatology* 2005;41:595–602.
- [176] Khurana S, Singh P. Rifampin is safe for treatment of pruritus due to chronic cholestasis: a meta-analysis of prospective randomized-controlled trials. *Liver Int* 2006;26:943–948.
- [177] Manns MP, Woynarowski M, Kreisel W, Lurie Y, Rust C, Zuckerman E, et al. Budesonide induces remission more effectively than prednisone in a controlled trial of patients with autoimmune hepatitis. *Gastroenterology* 2010;139:1198–1206.
- [178] Lu Y, Zhang Z, Xiong X, Wang X, Li J, Shi G, et al. Glucocorticoids promote hepatic cholestasis in mice by inhibiting the transcriptional activity of the farnesoid x receptor. *Gastroenterology* 2012;143:1630–1640.
- [179] Jung D, Fantin AC, Scheurer U, Fried M, Kullak-Ublick GA. Human ileal bile acid transporter gene ASBT (SLC10A2) is transactivated by the glucocorticoid receptor. *Gut* 2004;53:78–84.
- [180] Eloranta JJ, Kullak-Ublick GA. Coordinate transcriptional regulation of bile acid homeostasis and drug metabolism. *Arch Biochem Biophys* 2005;433:397–412.
- [181] Eloranta JJ, Jung D, Kullak-Ublick GA. The human Na⁺-taurocholate cotransporting polypeptide gene is activated by glucocorticoid receptor and peroxisome proliferator-activated receptor-gamma coactivator-1alpha, and suppressed by bile acids via a small heterodimer partner-dependent mechanism. *Mol Endocrinol* 2006;20:65–79.
- [182] Arenas F, Hervias I, Uriz M, Joplin R, Prieto J, Medina JF. Combination of ursodeoxycholic acid and glucocorticoids upregulates the AE2 alternate promoter in human liver cells. *J Clin Invest* 2008;118:695–709.
- [183] Leuschner M, Maier KP, Schlichting J, Strahl S, Herrmann G, Dahm HH, et al. Oral budesonide and ursodeoxycholic acid for treatment of primary biliary cirrhosis: results of a prospective double-blind trial. *Gastroenterology* 1999;117:918–925.
- [184] Rautiainen H, Karkkainen P, Karvonen AL, Nurmi H, Pikkarainen P, Nuutinen H, et al. Budesonide combined with UDCA to improve liver histology in primary biliary cirrhosis: a three-year randomized trial. *Hepatology* 2005;41:747–752.
- [185] Angulo P, Jorgensen RA, Keach JC, Dickson ER, Smith C, Lindor KD. Oral budesonide in the treatment of patients with primary biliary cirrhosis with a suboptimal response to ursodeoxycholic acid. *Hepatology* 2000;31:318–323.
- [186] Rabahi N, Chretien Y, Gaouar F, Wendum D, Serfaty L, Chazouilleres O, et al. Triple therapy with ursodeoxycholic acid, budesonide and mycophenolate mofetil in patients with features of severe primary biliary cirrhosis not responding to ursodeoxycholic acid alone. *Gastroenterol Clin Biol* 2010;34:283–287.
- [187] Hempfling W, Grunhage F, Dilger K, Reichel C, Beuers U, Sauerbruch T. Pharmacokinetics and pharmacodynamic action of budesonide in early- and late-stage primary biliary cirrhosis. *Hepatology* 2003;38:196–202.
- [188] Mora JR, Iwata M, von Andrian UH. Vitamin effects on the immune system: vitamins A and D take centre stage. *Nat Rev Immunol* 2008;8:685–698.
- [189] D'Aldebert E, Biyeyeme Bi Mve MJ, Mergey M, Wendum D, Firrincieli D, Coilly A, et al. Bile salts control the antimicrobial peptide cathelicidin through nuclear receptors in the human biliary epithelium. *Gastroenterology* 2009;136:1435–1443.
- [190] Cheng J, Fang ZZ, Kim JH, Krausz KW, Tanaka N, Chiang JY, et al. Intestinal CYP3A4 protects against lithocholic acid-induced hepatotoxicity in intestine-specific VDR-deficient mice. *J Lipid Res* 2014;55:455–465.
- [191] Schmidt DR, Holmstrom SR, Fon Tacer K, Bookout AL, Kliewer SA, Mangelsdorf DJ. Regulation of bile acid synthesis by fat-soluble vitamins A and D. *J Biol Chem* 2010;285:14486–14494.
- [192] Abramovitch S, Dahan-Bachar L, Sharvit E, Weisman Y, Ben Tov A, Brazowski E, Reif S. Vitamin D inhibits proliferation and profibrotic marker expression in hepatic stellate cells and decreases thioacetamide-induced liver fibrosis in rats. *Gut* 2011;60:1728–1737.
- [193] Hochrath K, Stokes CS, Geisel J, Pollheimer MJ, Fickert P, Dooley S, et al. Vitamin D modulates biliary fibrosis in ABCB4-deficient mice. *Hepatol Int* 2014;8:443–452.
- [194] Ding N, Yu RT, Subramaniam N, Sherman MH, Wilson C, Rao R, et al. A vitamin D receptor/SMAD genomic circuit gates hepatic fibrotic response. *Cell* 2013;153:601–613.

- [195] Hagenbuch B, Dawson P. The sodium bile salt cotransport family SLC10. *Pflugers Arch* 2004;447:566–570.
- [196] Walters JR, Tasleem AM, Omer OS, Brydon WG, Dew T, le Roux CW. A new mechanism for bile acid diarrhea: defective feedback inhibition of bile acid biosynthesis. *Clin Gastroenterol Hepatol* 2009;7:1189–1194.
- [197] Mauricio AC, Slawik M, Heitzmann D, von Hahn T, Warth R, Bleich M, et al. Deoxycholic acid (DOC) affects the transport properties of distal colon. *Pflugers Arch* 2000;439:532–540.
- [198] Harach T, Pols TW, Nomura M, Maida A, Watanabe M, Auwerx J, et al. TGR5 potentiates GLP-1 secretion in response to anionic exchange resins. *Sci Rep* 2012;2:430.
- [199] Potthoff MJ, Potts A, He T, Duarte JA, Taussig R, Mangelsdorf DJ, et al. Colesevelam suppresses hepatic glycogenolysis by TGR5-mediated induction of GLP-1 action in DIO mice. *Am J Physiol Gastrointest Liver Physiol* 2013;304:G371–G380.
- [200] Trauner M, Halilbasic E, Baghdasaryan A, Moustafa T, Krones E, Fickert P, et al. Primary sclerosing cholangitis: new approaches to diagnosis, surveillance and treatment. *Dig Dis* 2012;30:39–47.