

REVIEWS IN BASIC AND CLINICAL GASTROENTEROLOGY AND HEPATOLOGY

John P. Lynch and David C. Metz, Section Editors

Pathogenesis of Cholestatic Liver Disease and Therapeutic Approaches

GIDEON M. HIRSCHFIELD,^{*,‡} E. JENNY HEATHCOTE,^{*,‡} and M. ERIC GERSHWIN[§]

^{*}Liver Centre, Toronto Western Hospital, Toronto, Ontario, Canada; [‡]Department of Medicine, University of Toronto, Toronto, Ontario, Canada; and [§]Division of Rheumatology, Allergy and Clinical Immunology, University of California, Davis, California

Cholestatic liver disorders are caused by genetic defects, mechanical aberrations, toxins, or dysregulations in the immune system that damage the bile ducts and cause accumulation of bile and liver tissue damage. They have common clinical manifestations and pathogenic features that include the responses of cholangiocytes and hepatocytes to injury. We review the features of bile acid transport, tissue repair and regulation, apoptosis, vascular supply, immune regulation, and cholangiocytes that are associated with cholestatic liver disorders. We now have a greater understanding of the physiology of cholangiocytes at the cellular and molecular levels, as well as genetic factors, repair pathways, and autoimmunity mechanisms involved in the pathogenesis of disease. These discoveries will hopefully lead to new therapeutic approaches for patients with cholestatic liver disease.

Keywords: Primary Biliary Cirrhosis; Primary Sclerosing Cholangitis; Cholestasis; Ursodeoxycholic Acid.

Cholestatic liver diseases arise from impaired hepatobiliary production and excretion of bile, which cause bile constituents to enter the circulation. In these disorders, injuries to bile ducts or hepatocytes can lead to a range of clinical presentations, from isolated abnormalities in liver biochemistry, to liver failure or hepatobiliary malignancy; congenital, immunologic, structural (obstructive/vascular), and toxic factors can all contribute to disease (Figure 1 and Table 1). In response to injury, mature cholangiocytes and hepatocytes proliferate, which may lead to periductular fibrosis, biliary fibrosis, and cirrhosis. Disease progression and the efficacy of repair depend on etiology and the individual's response to injury. We review mechanisms and biliary pathophysiology of cholestatic liver disease, focusing on primary biliary cirrhosis (PBC) and primary sclerosing cholangitis (PSC).

Cell Biology of Cholestasis

Approximately 5% of cells in the liver are cholangiocytes; these ciliated epithelial cells line the biliary tree,

an intricate network of interconnecting bile ducts that increase in diameter from the ducts of Hering to the extrahepatic bile ducts.^{1,2} Cholangiocytes that line the large interlobular and major ducts are predominantly involved in secretory functions, whereas cholangiocytes that line smaller bile duct branches, cholangioles, and ducts of Hering have roles in inflammatory and proliferative responses.^{3,4} Each cholangiocyte has a primary cilium that extends from the apical plasma membrane into the ductal lumen and regulates mechanosensory, osmosensory, and chemosensory functions.⁵ Cilia detect and signal changes in bile flow and osmolality.⁶

Secretory Function and Bile Acid Transport

Cholangiocyte stimulation through paracrine and endocrine routes leads to secretion of water and biliary alkalinization with a net luminal flux of chloride and bicarbonate.⁷ The process of integrating prosecretory (secretin, glucagon, vasoactive intestinal polypeptide, acetylcholine, bombesin) and antiseecretory (somatostatin, endothelin-1) stimuli involves transmembrane adenylyl cyclases that regulate the concentration of intracellular 3', 5'-cyclic monophosphate.⁸⁻¹⁰ Basolateral (sinusoidal) and canalicular (apical) transport proteins move bile acids from sinusoidal blood into the canaliculus in a highly efficient, regulated enterohepatic circulation.^{11,12} Nuclear receptors regulate transcription of genes that encode proteins involved in hepatobiliary transport systems, bile acid synthesis, bile acid detoxification,¹³ and fibrogenesis.^{14,15} These nuclear receptors include the farnesoid X receptor, pregnane X receptor, vitamin D receptor, and constitutive androstane receptor. Bile acids are internal-

Abbreviations used in this paper: AMA, antimitochondrial antibody; GI, gastrointestinal; IL, interleukin; MADCAM, mucosal addressin cell adhesion molecule; MDR, multi-drug-resistant; NK, natural killer; PBC, primary biliary cirrhosis; PDC-E2, E2 component of the pyruvate dehydrogenase complex; PFIC, progressive familial intrahepatic cholestasis; PSC, primary sclerosing cholangitis; Treg, T regulatory cell.

© 2010 by the AGA Institute

0016-5085/\$36.00

doi:10.1053/j.gastro.2010.09.004




Figure 1. Cholestasis and hepatobiliary injury. Etiologies of cholestatic liver disease related to the site of hepatobiliary insult. (*Left panel*) Bile production is a complex process that involves hepatocytes and cholangiocytes; a number of different bile acid transporters coordinate bile formation (reviewed by Zollner and Trauner¹¹), with various cell surface receptors on cholangiocytes that regulate cholangiocyte secretion and function (reviewed by Glaser et al and Marziani et al^{30,31}). (*Middle panel*) Histologic representation of cholestatic liver disease illustrating bland hepatic cholestasis in drug injury, the small bile duct lymphocytic cholangitis of PBC, ductopenia as illustrated by immunostaining for keratin, the classic large bile duct involvement seen in PSC (obliterative cholangitis), and immunochemical staining showing IgG4-positive plasma cells surrounding a bile duct in IgG4-associated cholangitis (Histology images are courtesy of Dr M Guindi and Dr S Fischer, Toronto General Hospital, Toronto, Canada). (*Right panel*) Cholestasis develops after injury to hepatocytes and small and large bile ducts.

ized across the basolateral membranes of hepatocytes by the Na⁺/taurocholate cotransporter and organic anion transporting proteins (OATP2/OATP1B1); this transporter also mediates the hepatic uptake of many drugs.¹⁶ Genetic polymorphisms in the gene encoding this transporter, *SLCO1B1*, associate with total bilirubin levels in healthy individuals¹⁷ and development of statin-induced myopathy.¹⁸ Active export into bile is mediated by the canalicular bile salt export pump (ABCB11)¹⁹ and the canalicular conjugate export pump (MRP2).²⁰ MRP2 regulates canalicular excretion of organic anions, such as bilirubin. Formation of mixed micelles in bile results from the presence of bile acids, cholesterol, and phosphatidylcholine, and the phospholipid export pump, multi-drug-resistant 3 protein (MDR3), is actively involved in controlling the process.²¹

Cytokines and Receptors

The cholangiocytes in the smaller bile ducts, cholangioles, and ducts of Hering express receptors that allow proliferation in response to liver damage and participation in inflammatory responses.^{22,23} These receptors mediate protection from pathogens (via signals from antimicrobial peptides or the Toll-like receptors 2, 3, 4, and 5, which bind to bacterial molecules, double-stranded RNA, gram-negative bacteria, and lipopolysaccharide²⁴), antigen presentation (HLA molecules and costimulatory molecules^{25–27}), leukocyte recruitment (adhesion molecules such as intercellular adhesion molecule 1, leukocyte factor antigen 3, and CD40, cytokines, and chemokines^{28,29}), and leukocyte apoptosis. Reactive cholangiocytes also produce growth factors such as vascular endothelial

Table 1. The Characteristics of Common Cholestatic Syndromes

	Primary biliary cirrhosis	Primary sclerosing cholangitis	Drug induced cholestasis	Biliary atresia
Demographics	Female predominant; middle age onset; geographic hotspots; co-existent autoimmunity; Asymptomatic screening now most common mode of presentation	Men more than women; onset before 40 years old common; high prevalence of co-existent colitis; Identified commonly through asymptomatic screening in inflammatory bowel disease; may present with cholangitis	10% of drug induced liver injury is cholestatic; more common in elderly; jaundice and pruritus seen	Exclusively neonatal; Isolated in 65%–90%; Presents with persistent neonatal conjugated hyperbilirubinemia >14 days old
Key diagnostics	Anti-mitochondrial antibody (AMA) positive >95%; confirm immunofluorescence with specific AMA assay ^a	Cholangiographic appearance of stricturing and beading of intra- and/or extra-hepatic bile ducts; MRI largely supplanted ERCP as initial diagnostic modality	History of offending agent ^b and exclusion of alternative etiologies	Liver biopsy differentiates obstructive and hepatocellular causes of cholestasis, with 90% sensitivity and specificity; cholangiography if diagnostic doubt (patent biliary tree proximally and distally excludes biliary atresia)
Characteristic pathology	Granulomatous non-suppurative small duct cholangitis, ductopenia, ductal proliferation, interface hepatitis Biopsy is not required routinely for diagnosis or staging	Fibro-obliterative cholangitis, periductal fibrosis and inflammation, absence of bile ducts in some portal tracts and ductal proliferation in other portal tracts Biopsy is not required routinely for diagnosis or staging	Bland cholestasis or cholestatic hepatitis >Biopsy commonly performed to exclude alternative etiologies	Active inflammation with bile duct degeneration, a chronic inflammatory reaction with proliferation of both ductular and glandular elements, and fibrosis Biopsy commonly performed
Overlapping features	Anti-nuclear antibodies present in 50% ^c and possible overlap with autoimmune hepatitis (AIH) in <10%	Autoimmune serology frequently positive; pediatric onset AIH is accompanied by sclerosing cholangitis in 50% of cases; overlap with AIH in adults <10%.	Non-specific serology may be present	Embryonic form in 10%–35% with associated situs inversus or polysplenia/asplenia +/- other congenital anomalies
Natural history and treatment	Progressive biliary cirrhosis, portal hypertension and liver failure	Recurrent cholangitis, progressive biliary cirrhosis, portal hypertension and liver failure; malignancy (hepatobiliary/colorectal)	Usually self limiting but overt ductopenia may be observed	Biliary cirrhosis, liver failure
Treatment	Ursodeoxycholic acid (UDCA) 13–15 mg/kg/day effective in majority	No primary medical intervention presently; liver transplant effective rescue therapy (>50% require transplant within 15 years of symptoms)	No proven therapy other than cessation of drug	No primary medical therapy; Kasai porto-enterostomy and/or liver transplantation highly effective
Disease severity indices	Biochemical response to UDCA predicts disease progression and survival, thus identifying patients at risk of adverse outcome ^d	No early markers of outcome; Mayo PSC score and MELD of use in late disease	“Hy’s rule”: if both drug-induced hepatocellular injury and jaundice occur simultaneously, a mortality of at least 10% can be expected	If bilirubin is less than 2 mg/dL at 3 months post-surgery, then the chance of being transplant-free at 2 years of age is 84%
Key unmet need	Second line therapy for UDCA non-responders	Effective primary treatment; effective screening for cholangiocarcinoma; intervention for transplant recurrence; surrogate end points of disease progression	No effective therapy for progressive injury; predictive tools for preventing exposure to risk-prone patients	Early diagnosis to facilitate early surgery (<60 days old); no effective medical therapy
Diagnostic pitfalls	AMA identified in autoimmune hepatitis, acute liver failure and 0.5% of healthy individuals; isolated elevated alkaline phosphatase values occur in NAFLD	IgG4 associated autoimmune pancreatitis/sclerosing cholangitis responds to steroid therapy; ~10% of patients with PSC have elevated IgG4	Liver injury can present after the cessation of potentially injurious medicines	Differential diagnosis include Alagille syndrome, progressive familial intrahepatic cholestasis, alpha-1-antitrypsin deficiency, and cystic fibrosis

^aThe mitochondrial autoantigens have been cloned, sequenced, and identified as members of the 2-oxo-acid dehydrogenase pathway, including the E2 subunits of pyruvate dehydrogenase (PDC-E2), branched-chain 2-oxo-acid dehydrogenase (BCOADC-E2), and 2-oxo-glutarate dehydrogenase (OGDC-E2).

^bCholestatic liver injury is reported with many agents including, but not exclusively, estrogens, anabolic steroids, chlorpromazine, erythromycin, the oxypenicillins, tamoxifen, macrolides, ticlopidine, terfenadine, terbinafine, nimesulide, irbesartan, fluoroquinolones, cholesterol-lowering ‘statins,’ herbal remedies (greater celandine, glycyrrhizin, chaparral), amoxicillin-clavulanic acid and ibuprofen.

^cHighly specific anti-nuclear antibodies are detected in ~ 50% of patients with PBC (gp210 and sp100).

^dA number of biochemical predictors of treatment response are associated with histological progression and survival in patients treated with UDCA (eg, Barcelona, Paris and Toronto criteria respectively).

growth factor, endothelin-1, platelet-derived growth factor BB, transforming growth factor β 2, and connective tissue growth factor.

Repair and Regulation

Biliary proliferation contributes to the initiation and progression of liver fibrosis.^{30,31} Cholangiocytes normally do not proliferate because they constitutively express the cyclin-dependent kinase inhibitors p27, Bcl2, Bcl-xL, and Mcl-1. In response to liver damage, gastrointestinal (GI) and neuroendocrine hormones, and autocrine and paracrine signaling factors, cholangiocytes proliferate and acquire a neuroendocrine secretory phenotype. In the “ductular reaction,” an expanded population of epithelial cells accumulates at the interface of the biliary tree and hepatocytes, along with proliferation of preexisting ductules, activation of progenitor cells, and the appearance of intermediate hepatocytes. The integrin α v β 6 is up-regulated in proliferating bile duct epithelia and promotes fibrogenesis via adhesion to fibronectin and autocrine/paracrine activation of transforming growth factor β 1.³²

Reactive cholangiocytes express receptors, such as the β 1 and β 2 adrenergic receptors, the M3 acetylcholine receptor, and serotonin 1A and 1B receptors, that bind neurotransmitters. Cholangiocytes can also directly secrete serotonin, which limits the growth of bile ducts through an inhibitory, autocrine loop.³³ Epithelial-mesenchymal transition is a process that might also occur during cholangiocyte proliferation and fibrogenesis.^{34–36} Some epithelia-derived mesenchymal cells undergo a mesenchymal-epithelial transition, reverting to epithelial cells that ultimately become hepatocytes or cholangiocytes. The hedgehog signaling pathway appears to promote and regulate hepatic accumulation of immune cells that interact with cholangiocytes.

Cell Death

An unbalanced equilibrium between cholangiocyte death (via apoptosis or necrosis) and proliferation leads to duct loss and fibrosis. In advanced biliary disease, cholangiocytes lose the ability to proliferate³⁷; in progressive ductopenia, there is more apoptosis than proliferation. Apoptosis contributes to duct loss and is induced by signals such as activation of death receptors, immune-mediated injury, oxidative stress, infections, and toxins. Apoptosis also promotes fibrogenesis, with apoptotic debris contributing to activation of hepatic stellate cells. Cholangiocytes are the primary epithelial source of tumor necrosis factor α in the liver, a proinflammatory mediator that activates caspase cleavage and apoptosis but also mediates cell survival via activation of the transcription factor nuclear factor κ B. In vitro studies have shown that a combination of tumor necrosis factor α , interleukin (IL)-1, IL-6, and interferon gamma inhibits 3', 5'-cyclic monophosphate-dependent ductal secretion. Nuclear factor κ B, which is regulated by I- κ B and its

kinases (IKKs), regulates immune and inflammatory responses; it protects against cytokine-induced death and oxidative damage. Mice with disruptions in IKK1 and IKK2 or IKK1 and NEMO (nuclear factor κ B essential modulator) develop jaundice and a fatal cholangitis, characterized by inflammatory destruction of small portal bile ducts.³⁸ The combined loss of IKK1-specific functions leads to biliary disease, a process proposed to result from changes to the regulation of tight junctions in biliary epithelial cells. Activation of the receptor tumor necrosis factor-related apoptosis-inducing ligand 2 and death receptor 5 induce cell death and mediate cholestatic liver injury.³⁹ Healthy cholangiocytes do not express tumor necrosis factor-related apoptosis-inducing ligand, but diseased cholangiocytes up-regulate it, which might control inflammatory responses by killing leukocytes that express the death receptors.

The Hepatic Lymphoid System

In healthy liver, lymphocytes are scattered throughout the parenchyma and portal tracts and include subpopulations from the innate and adaptive immune systems. Less than 10% of human intrahepatic lymphocytes are B cells; most are CD5⁺ B1 cells that control innate and adaptive immunity.^{40,41} T-cell populations, primarily CD4⁺ helper (Th1 and Th2), CD8⁺ cytotoxic, T-regulatory, and Th17 cells, have important roles in pathogenesis, along with hepatic macrophages. Nonparenchymal liver cells involved in immune system tolerance to liver include resident dendritic cells, liver sinusoidal endothelial cells, Kupffer cells, and hepatic stellate cells. These cells mediate immunosuppression by producing anti-inflammatory cytokines such as IL-10 and transforming growth factor β ; they also express the inhibitor of T-cell activation, programmed cell death 1. IL-10, transforming growth factor β , programmed cell death 1, and cytotoxic T lymphocyte antigen can contribute to the immunosuppressive mechanisms of CD4⁺CD25⁺Foxp3⁺ regulatory T cells, which appear to be converted in the liver from infiltrating naive CD4⁺ T cells and/or effector CD4⁺ T cells.

Immunologic Role of Cholangiocytes

In response to inflammation, cholangiocytes secrete cytokines and chemokines (eg, tumor necrosis factor α , IL-1, or interferon gamma) that recruit and activate immune cells, including T cells, macrophages, and natural killer (NK) cells.^{23,42} Human cholangiocytes constitutively express and secrete chemotactic agents for neutrophils, monocytes, and T cells, including IL-8, IL-6, and MCP-1; under basal conditions, cholangiocytes express low levels of lymphocyte adhesion molecules. In normal livers, some cholangiocytes express HLA class I but not class II molecules. However, cholangiocytes from patients with PBC express HLA class II, but it is not known if costimulatory molecules for T-cell activation are present.

The presence of the adhesion molecule leukocyte factor antigen 3 on the cholangiocyte cell surface allows their interaction with CD2 on cytotoxic T lymphocytes and NK cells. T cells are also activated by CD40, which is expressed on cholangiocytes.⁴³ The CD40-CD40L and leukocyte factor antigen 2 (CD2)/leukocyte factor antigen 3 (CD58) complexes induce production of IL-12. Cholangiocytes also secrete and transport protective immunoglobulins. In bile, immunoglobulin (Ig) A has a role in biliary mucosal immune defense; by preventing the attachment of pathogens or their toxins to the cholangiocyte surface, it protects biliary ducts.⁴⁴ IgA is synthesized by plasma cells around bile ducts and secreted into bile after it binds to the polymeric Ig receptor located on the basolateral membranes of cholangiocytes.

Vascular Supply

Bile ducts are supplied with blood only from hepatic arteries; in contrast to hepatocytes, the biliary epithelium receives blood from a network of capillaries near the intrahepatic bile ducts, the peribiliary vascular plexus, which originate from the terminal branches of the hepatic artery. This specific vascular supply, lacking in canals of Hering and terminal cholangioles, accounts for the prevalent involvement of the interlobular bile ducts in ischemic injury.⁴⁵

Mechanistic Insights From Genetic Studies

From the many genetic insights recognized in both common and uncommon cholestatic liver diseases, it is possible to appreciate a number of important pathophysiologic pathways likely relevant to cholestasis more generally.

Inherited Cholestatic Syndromes

Alagille syndrome (arteriohepatic dysplasia or congenital deficiency in interlobular bile ducts) is characterized by chronic cholestasis, posterior ocular embryotoxon, butterfly-like vertebral arch defects, peripheral pulmonary artery hypoplasia or stenosis, and dysmorphic facial features.⁴⁶ This autosomal dominant multi-system disorder varies in phenotypic expression and is largely associated with mutations in *JAG1*, which encodes a ligand in the Notch signaling pathway. Less than 1% of patients have mutations in the receptor *NOTCH2*. By altering the expression of liver-enriched transcription factors, Notch signaling contributes to biliary tree development during ductal plate remodeling and controls transdifferentiation of hepatoblasts and mature hepatocytes into cholangiocytes.⁴⁷ In the course of cholestatic liver disease more generally, changes in the expression of Jagged1 and Notch have been described. Changes in cell-cell tight junction function and membrane fusion are also associated with cholestasis, leading to changes in the cytoskeletons and tight junctions of hepatocytes. Neonatal

tal ichthyosis-sclerosing cholangitis syndrome, a rare autosomal recessive condition characterized by hypotrichosis of the scalp, scarring alopecia, ichthyosis, and sclerosing cholangitis, has been associated with changes in the gene that encodes the tight junction protein claudin-1,⁴⁸ whereas other even rarer multi-system disorders, including neonatal cholestasis, have been associated with defects in vesical fusion (eg, exocytosis).^{49,50}

The progressive familial intrahepatic cholestasis (PFIC) syndromes 1, 2, and 3 are characterized by abnormal bile formation and associated with penetrant mutations in genes that encode proteins in the hepatocellular transport system.⁵¹ PFIC1 and PFIC2 usually appear early in life (first few months), whereas onset of PFIC3 varies. PFIC1 and PFIC2 are each caused by impaired bile salt secretion; they are caused by defects in *ATP8B1* (which encodes FIC1) and *ABCB11* (which encodes the bile salt export pump protein), respectively. Changes in these 2 proteins are also associated with a recurrent cholestatic syndrome known as benign recurrent intrahepatic cholestasis. *ATP8B1* is a type 4 P-type adenosine triphosphatase acting as a flippase for phosphatidylserine, such that *atp8b1*-deficient mice display an increase in the biliary extraction of cholesterol from the canalicular (apical) membrane of the hepatocyte. The *ABCB11* gene changes in PFIC2 lead to changes in function of an adenosine triphosphate-dependent canalicular bile salt export pump, which is expressed at the hepatocyte canalicular membrane and acts as the major exporter of primary bile acids. As a result, there is decreased biliary bile salt secretion, decreased bile flow, and hepatic accumulation of bile acids. Functional mutations in *ABCB4*, which encodes MDR3, impair biliary phospholipid secretion and result in PFIC3. In healthy liver, MDR3 is present in the canalicular wall and phosphatidylcholine enters bile to complex with bile acids; it acts as a chaperone to prevent toxic membrane injury. Reduced biliary excretion of phosphatidylcholine leads to a high saturation index of biliary cholesterol and impairs protection of the bile duct epithelium from the detergent properties of bile acids. γ -Glutamyltransferase is present in large quantities in the canalicular and bile duct membranes (mostly canalicular); when bile ducts are damaged, γ -glutamyltransferase is released into the bile and subsequently serum. This explains the increase in γ -glutamyltransferase levels in PFIC3 syndromes but not in PFIC1 and PFIC2 syndromes, because bile is especially toxic to the canalicular membrane. In addition to patients with classic PFIC3, individuals with *ABCB4* mutations (and associated altered MDR3 function) are recognized to develop a wide spectrum of cholestatic diseases, including idiopathic adult biliary cirrhosis, low phospholipid-associated cholelithiasis syndrome, transient neonatal cholestasis, intrahepatic cholestasis of pregnancy, and drug-induced cholestasis.²¹

Cholangiocytes are the only epithelial cells in the liver to have cilia, and signals from the cilia regulate cholangiocyte development and function; developmental abnormalities of the portobiliary system, in association with fibrocystic disease of the kidneys, are known as “ciliopathy diseases” or “cholangiociliopathies.”^{52,53} The most frequent hepatic manifestations include congenital hepatic fibrosis (ductal plate malformation, abnormal portal veins, progressive portal fibrosis), Caroli’s disease (saccular or fusiform dilations of the medium- and large-sized intrahepatic bile ducts), and polycystic liver disease (closed cysts that originate from biliary microhamartomas embedded in fibrous tissue and not in continuity with the biliary tree). Examples of disease include autosomal dominant polycystic kidney disease, caused by mutations in *PKD1* or *PKD2*, and autosomal recessive polycystic kidney disease, caused by mutations in *PKHD1*, which encodes fibrocystin. Autosomal dominant polycystic liver disease is characterized by the presence of only liver cysts; disease results from mutations in *PRKCSH*, which encodes hepatocystin (protein kinase C substrate 80K-H), or *Sec63*. Other rare, genetic, multi-system ciliopathies that involve the liver include nephronophthisis (a medullary cystic kidney disease), Bardet-Biedl syndrome, and Meckel-Gruber syndrome.

HLA and Cholestasis

With the association of the *HLA* locus repeatedly the most significant of any genetic observations for PBC,⁵⁴ PSC,⁵⁵ and drug-induced cholestasis,⁵⁶ there is a clear implication that cholestatic liver diseases are in part triggered in individuals with certain *HLA* alleles that are better at presenting peptides that resemble self-peptides to mature T cells. Studies of PBC from northern and southern Europe and North America have associated *HLA-DR8* (*DRB1*08*) with PBC,^{57,58} whereas the only genomic study of patients with PSC to date found that *HLA-B* at chromosome 6p21 was most strongly associated with this disease.⁵⁵ Previous studies of PSC investigated markers of innate immunity near the *HLA* locus. KIR, inhibitory receptors on NK cells, bind *HLA* class I molecules; NK cells of patients with PSC had lower amounts of *HLA-Bw4* and *HLA-C2* (ligands for the KIRs 3DL1 and 2DL1, respectively) compared with controls. These findings indicate that NK cell activity is increased in patients with PSC (decreased inhibition).^{59,60} In addition to the increased activity of NK cells through *HLA*-KIR interactions, genetic evidence indicates a role for the direct activation of NK cells through the *MHC class I chain* (*MIC*)-related genes. *MICA* and *MICB* lie in the *HLA* region; their products activate the receptor NKG2D on NK cells. The alleles *MICA*002* and *MICA*008* reduce the risk of PSC,⁶¹ whereas a Norwegian study associated the alleles *MICA5.1* and *MICB24* with disease.⁶²

Non-HLA Associations

In the genome-wide studies of PBC available to date, which were designed to discover novel non-*HLA* associations, disease was associated with the *IL-12* signaling cascade and the *IL12A* and *IL12RB2* loci in particular. Other robust gene loci include *IRF5* (also associated with systemic lupus erythematosus and inflammatory bowel disease), *MMEL1* (also associated with rheumatoid arthritis), *17q12.21* (also associated with asthma), and *SPIB*.^{54,63,64} Bigger cohorts will inevitably discover or confirm, at an appropriately robust statistical level of certainty, many more risk loci, some of which may already have been discussed (eg, *STAT4* and *CTLA4*). The findings, however, firmly confirm PBC as a classic autoimmune disease. For PSC, associations outside of the *HLA* locus remain more speculative, pending testing of larger cohorts, but include *Glypican 6* as well as loci at chromosomes 2q35 and 3p21, likely reflecting changes to genes encoding the G protein-coupled bile acid receptor 1 and macrophage-stimulating 1, respectively.⁵⁵ Whole genome sequencing of cohorts of patients with cholestatic liver disease will ultimately be required, because present genome-wide association testing will not identify the majority of genetic risk for disease. Such an approach is predicted to identify presently unknown rare mutations within genes encoding proteins with significant biological effect.

PBC as a Model Autoimmune Disease

PBC is considered a model autoimmune disease because of the homogeneity of its phenotype among patients, its predominance in women, and its association with a specific autoantibody (against a mitochondrial protein).^{65–67} It occurs in 0.1% of women older than 40 years of age^{68–71} and is characterized by chronic small bile duct cholangitis. Previous epidemiologic studies have shown that having a first-degree relative with PBC, a history of urinary tract infections, a history of smoking, or use of hormone replacement therapies increased the risk of PBC.^{68,71} Other autoimmune diseases develop in approximately 33% of patients with PBC (for example, scleroderma and Sjögren’s syndrome), as do cholestasis or pruritus during pregnancy.^{68,72–74} CD4⁺ and CD8⁺ T-cell and innate responses are involved in pathogenesis, following environmental insults in genetically susceptible individuals^{75–77} (Figure 2). PBC is characterized by specific loss of tolerance to a ubiquitous mitochondrial antigen, the E2 component of the pyruvate dehydrogenase complex (PDC-E2)^{78–80} (Figure 3). There are several mouse models of PBC^{81–85} that have many genetic and environmental features of human disease; studies of these models indicate a role for regulatory T cells (Tregs) and autoreactive, CD8⁺ T cells in disease induction and modulation. Limitations remain, and the mouse biliary system inevitably differs from that of humans; for exam-




Figure 2. Apoptotic blebs and specificity in PBC. Immunohistochemical analysis of apoptosis-dependent anti-pyruvate dehydrogenase complex (PDC-E2) staining in biliary epithelial cells incubated with AMA-positive serum. PDC-E2 is a ubiquitous protein present in mitochondria of nucleated cells; biliary epithelial cells translocate intact PDC-E2 to apoptotic bodies and create an apoptosome, which is immunoreactive. N, nucleus.

ple, mice do not have polymeric IgA receptors.⁸¹ The role of B cells in pathogenesis is also complex, but selective deletion of B cells reduces the cholangitis in mouse models.^{81,83,84,86–88}

Treatment of patients with PBC is still not disease specific and has not changed for more than 10 years; the standard of care is therapy with the secondary bile acid ursodeoxycholic acid (UDCA).^{89–93} The clinical consensus is that a biochemical response to UDCA delays progression of disease in most patients.^{94–99}

Environmental Factors

The mitochondrial autoantigen PDC-E2 contains lipoic acid; PDC-E2 captures electrons to alternate oxidation and reduction of a disulfide bond within lipoic acid. In genetically susceptible individuals, drugs or chemicals that modify this disulfide bond might lead to loss of tolerance to PDC-E2. This protein is well conserved throughout evolution, so loss of tolerance could also result from an immune response against similar epitopes in bacteria (molecular mimicry).^{100–103} Urinary tract infections are more frequent in patients with PBC than controls, and several bacterial strains have been proposed to lead to immune cross-reactivity with the mitochondrial protein; serologic and molecular analyses have indicated the xenobiotic-metabolizing, gram-negative bacterium *Novosphingobium aromaticivorans* as a good candidate.¹⁰⁴ *N aromaticivorans* was

found in fecal samples from 25% of patients with PBC as well as in controls. PDC-E2 from this bacteria has high homology to human PDC-E2, and antimitochondrial antibodies (AMAs) in serum samples from patients with PBC have a 1000-fold stronger reaction against *N aromaticivorans* PDC-E2 than against *Escherichia coli* PDC-E2.^{105,106} Notably, infection of mice with *N aromaticivorans* induces production of serum AMAs and development of cholangitis.¹⁰⁷

In addition to bacteria, chemical xenobiotic agents might induce PBC. Some halogenated organic compounds can attach to specific epitopes of PDC-E2 and induce production of antibodies that have higher affinities for modified mitochondrial epitopes than for normal PDC-E2. One such agent is 2-octynoic acid, which is present in cosmetic products, including nail polish; in vitro and in vivo data indicate its role in autoimmune biliary disease.^{108,109} AMAs have equal affinities for 2-octynoic acid and lipoic acid. NOD mice immunized with bovine serum albumin–conjugated 2-octynoic acid developed higher titers of AMAs than those injected with only bovine serum albumin and developed histologic features of PBC, including liver granulomas and portal infiltrates enriched in CD8⁺ cells.⁸²

Disease Specificity

It is not clear why patients with PBC develop defects specifically in the biliary system, but they might arise from specific features of cholangiocyte apoptosis.^{76,77} PDC-E2 remains intact and immunogenic during biliary epithelial cell apoptosis; unlike in other apoptotic cells, it is not modified with glutathiol.¹¹⁰ This allows biliary PDC-E2 to be exposed to dendritic cells and its epitopes to be presented to T cells in draining lymph nodes, leading to production of AMAs.^{76,77} AMAs though are detected in patients with other autoimmune and nonautoimmune liver diseases, so loss of tolerance to PDC-E2 alone is not sufficient to cause autoimmune cholangitis.^{111,112} The specific destruction of small biliary epithelial cells in patients with PBC might also arise from differences in expression of trefoils (protease-resistant peptides that are expressed in specific patterns throughout the normal GI tract, and which help protect epithelial cells) between small and large bile ducts. Small bile ducts do not produce intestinal trefoils in response to damage, so they might be more vulnerable than large bile ducts.¹¹³

Gender Disparities

The predominance of PBC in women is unexplained, but believed by some to result from changes in the X chromosome. Genes on the X chromosome maintain levels of physiologic sex hormones and immune responsiveness. Although genome-wide studies have not associated PBC with the X chromosome, age-dependent increases in monosomy X have been reported in the




Figure 3. Pathogenic model of PBC. The mitochondrial antigen, PDC-E2, which is in biliary epithelial cell apoptotic blebs, is presented to immune cells. Immune complexes form, leading to expansion of autoreactive immune cells that target the biliary epithelia and cause chronic inflammation, cholestasis, and fibrosis. Genetic and environmental factors determine risk for this disorder. APC, antigen-presenting cell; MHC, major histocompatibility complex.

peripheral white blood cells of women with PBC. Loss of the X chromosome has been reported to be more common in women with PBC than in women without the disorder.^{114,115} There are little data on epigenetic factors that contribute to cholestatic liver disease, but sex-specific epigenetic factors might be involved.¹¹⁶

Adaptive Responses in PBC

In patients with PBC, the presence of autoantibodies against PDC-E2 is accompanied by a 100-fold increase in hepatic, antigen-specific CD4⁺ T cells and a 10-fold increase in antigen-specific CD8⁺ T cells.^{117–122} These specific T cells are present even when the AMAs are not detected in serum.¹²³ Among patients who carry the *HLA DR4*0101* allele, there is a single epitope (a 163–176 amino acid sequence that encompasses the lipoic acid-binding residue of the inner lipoyl domain of PDC-E2) recognized by autoreactive, proinflammatory, CD4⁺ T cells. Autoreactive CD8⁺ T cells have also been characterized; the HLA class I-restricted epitope, a 159–167 amino acid sequence, is near the epitopes recognized by CD4⁺ T cells and AMAs. As for CD4⁺ autoreactive T cells, the use of tetramer technology has shown a 10-fold higher frequency of PDC-E2 159–167 specific CD8⁺ T cells in diseased liver tissues compared with peripheral blood, particularly in patients with early-stage

disease. Autoreactive CD8⁺ T cells from patients with PBC specifically kill cells that express PDC-E2; they also produce interferon gamma rather than the anti-inflammatory cytokines IL-4 and IL-10.

T-helper cell subpopulations include Tregs and Th17 cells.^{124,125} The proinflammatory, IL-17-secreting subgroup mediate host defense and autoimmunity; reductions in numbers of CD4⁺ CD25^{high} regulatory T cells are likely to contribute to autoimmunity, chronic viral infection, antitumor immunity, and allergy. These T cells are able to suppress inappropriate immune responses and regulate cellular immune responses. Studies in animals have shown that transfer of T cells that lack the CD4⁺ CD25^{high} population into athymic nude mice results in T-cell-mediated autoimmunity. In diseased liver samples, more than 15% of T cells in areas of inflammation are Tregs that express the forkhead family transcriptional regulator box P3 (FoxP3). These Tregs are recruited by specific chemokines; CXCR3 appears to mediate recruitment via the hepatic sinusoidal endothelium, whereas ligands for CCR4 are secreted by dendritic cells that have been recruited to sites of inflammation.¹²⁶ In patients with PBC, FoxP3⁺ Tregs have been found in lymphoid infiltrates that localized to the portal tracts; significantly

lower proportions of CD4⁺ CD25^{high} Tregs were observed in patients and their family members.^{127,128}

The role of Th17 cells in cholestatic liver disease is not yet clear. Increased numbers of IL-17⁺ cells have been reported in liver tissues from patients with PBC and IL-2 receptor α knockout mice, which develop autoimmune biliary disease. Clusters of IL-17⁺ cells were also observed in portal tracts of diseased liver.^{129,130}

Innate Responses in PBC

In addition to T cells, PBC is mediated by other cells in the innate immune system.^{131–134} Patients with PBC have granulomatous inflammation, increased production of polyclonal IgM, hyperresponsiveness to CpG oligodeoxynucleotides, increased numbers of NK cells, and increased responsiveness to cytokines. Bacterial and viral pathogen-associated molecular patterns induce innate immune responses through cell surface binding to Toll-like receptors and via lipopolysaccharide and lipoteichoic acids that are in bile; Toll-like receptors have been found in cultured human biliary epithelial cells. The downstream effects of biliary epithelial cell Toll-like receptor signaling mediate biliary injury and might involve activation of nuclear factor κ B, which regulates transcription of proinflammatory factors (IL-6, tumor necrosis factor α) and leads to chemokines (IL-8, fractalkine/CX3CL1). Fractalkine can exist in membrane-bound and soluble chemotactic forms; it is produced by several types of epithelial cells, regulates cell adhesion, and is a chemoattractant for cells that express its receptor (CX3CR1), such as CD8⁺ and CD4⁺ T cells. In patients with PBC, CX3CL1 expression is up-regulated in injured bile ducts; CD4⁺ and CD8⁺ cells that express CX3CR1 are found in portal tracts and within the biliary epithelial layer of injured bile ducts.¹³⁵

Immune Consequences of Cholestasis

Bile acid homeostasis is regulated to a large extent at a transcriptional level via nuclear receptors that regulate overlapping genes, including not only those related to bile acid synthesis, transport, and detoxification but innate and adaptive immunity as well.¹³ Therefore, during cholestasis, retained bile acids signaling through a myriad of nuclear receptors could conceivably modulate immune responses. One specific example is the farnesoid X receptor, with data supporting a role for the bile acid sensor farnesoid X receptor in regulating the activation of liver NKT cells,¹³⁶ as well as being a negative modulator of nuclear factor κ B-mediated hepatic inflammation.¹³⁷ Other nuclear receptors with similar potential bridging roles between cholestasis and the immune response include the glucocorticoid receptor, peroxisome proliferator-activated receptor, and vitamin D receptor.

Biliary Function

The secretin pathway is an important trophic factor in biliary growth and function,¹³⁸ and in cholestatic

disorders an impaired response to secretin can be shown, which is improved in patients with PBC treated with UDCA.¹³⁹ The impaired ability to induce alkalinization and dilution of canalicular bile upon stimulation by secretin and several neuropeptides in PBC has been associated with altered regulation of cholangiocyte AE2 (SLC4A2; Cl[−]/HCO₃[−] anion exchanger) and NHE (SLC9A3; Na⁺/H⁺ exchanger) function^{140,141} as well as loss of the expression of inositol 1,4,5-triphosphate receptors involved in Ca²⁺-mediated bicarbonate secretion.¹⁴² An impaired generation of a bicarbonate-rich choleresis also modifies normal biliary epithelial responses to microbes, while maintaining an alkaline pH near the apical surface of hepatocytes and cholangiocytes additionally prevents the uncontrolled membrane permeation of protonated, and potentially toxic, glycine-conjugated bile acids.¹⁴³ Therapeutically, the effects of UDCA and dexamethasone on AE2 gene expression have been elucidated and the combination of UDCA and dexamethasone, but not UDCA or dexamethasone alone, increases the expression of liver-enriched alternative messenger RNA isoforms AE2b1 and AE2b2 and enhanced AE2 activity.¹⁴⁴ In mice with widespread AE2 gene disruption,⁸⁵ splenomegaly, increased production of IL-12 p70 and interferon gamma, expanded CD8⁺ T-cell populations, and underrepresented CD4⁺FoxP3⁺/regulatory T cells are seen. Additionally, most of these mice test positive for AMA and one third develop an extensive portal inflammation with CD8⁺ and CD4⁺ T lymphocytes surrounding damaged bile ducts, akin to human PBC. Mutations in the AE2 gene in patients with PBC are not clearly associated with disease onset,¹⁴⁵ and it remains to be seen whether disease progression rather than onset is modulated by biliary transporter function more generally, because presently there is little appreciated about determinants of response to UDCA.¹⁴⁶

PSC: The Black Box of Cholestatic Liver Disease

PSC is a chronic inflammatory large duct cholangiopathy that results in fibrotic strictures and dilatations of the intrahepatic and extrahepatic bile ducts.^{147–149} PSC predominates among men and is frequently associated with inflammatory bowel disease with specific features (rectal sparing, right-sided disease, and backwash ileitis).¹⁵⁰ In these patients, PSC causes cholangitis, secondary biliary cirrhosis, and progression to liver failure. Additionally, there is an ominous increased risk of malignancy, both within the hepatobiliary tree and in the bowel, if associated with inflammatory bowel disease.¹⁵¹ However, little is known about the pathogenesis of PSC (Figure 4). There is an increased prevalence of autoimmunity in these patients and their relatives; although there is evidence to support genetic predisposition to this disease, it is likely that nongenetic factors have a larger role than in PBC. Many characteristics can mimic the




Figure 4. Pathogenic model of PSC. Although there are animal models of PSC, pathogenesis is still poorly characterized in contrast to PBC. Sensitized bile ducts are damaged by different immune cells that are activated in the GI tract and lymph nodes. p-ANCA, perinuclear antineutrophil cytoplasmic antibodies; TBB5, beta-tubulin isotype 5.

histologic and radiologic features of PSC, including autoimmunity, ischemia, infection, and toxins,^{152,153} indicating common pathways to biliary injury that can arise from heterogeneous primary insults. In PSC, the primary injury is not to hepatocytes but to medium- and large-sized bile ducts, with cholangiography showing aptly the resultant intrahepatic and/or extrahepatic biliary disease as localized or multifocal strictures with intervening segments of normal or dilated ducts. Histologically, concentric periductal fibrosis (onion skinning) occurs and progresses to narrowing and obliteration of small bile ducts. Cholangiocytes become activated to express adhesion molecules, inflammatory and profibrogenic cytokines and receptors, and growth factors that stimulate production of extracellular matrix and accumulation and proliferation of periductal myofibroblasts. Drugs that disrupt this fibro-obliterative process are an important goal for new therapies.

Etiologic Factors

The lymphocytes that infiltrate the livers of patients with PSC are primarily T cells comprising CD4⁺ Th-1 phenotype (portal infiltrate) and CD8⁺ T cells (lobular infiltrate).^{42,154–156} There are many $\gamma\delta^+$ T cells; the percentage of $\gamma\delta^+$ T cells in peripheral blood of patients with PSC is greater than that in healthy controls. The infiltrating T cells express the IL-2 receptor and CD45RO, indicating that they have an activated memory phenotype. Although the inflammatory infiltrate in patients with PSC mostly comprises T cells, other cell types, including NK cells, macrophages, B cells, and biliary epithelial cells, are likely to contribute to immunopathogenesis. Innate immune system activation might also be a primary event in PSC; the disease might be triggered or enhanced by bacteria (and pathogen-associated molecular patterns) that enter the portal circulation through an inflamed permeable intestine. IgGs against bil-

iary epithelial cells have been found in the sera of some patients with PSC; the binding of such antibodies to biliary epithelium induces production of proinflammatory cytokines and up-regulation of Toll-like receptors.¹⁵⁷

Infectious triggers continue to cause speculation, with some, albeit weak, supportive data (eg, *Helicobacter pylori*).¹⁵⁸ Sclerosing cholangitis, however, occurs in immunosuppressed patients with cryptosporidial infections, and autoimmune pancreatitis/sclerosing cholangitis has been associated with antibodies against a peptide that has homology to a plasminogen-binding protein of *H pylori*.¹⁵⁹ Patients with PSC often have perinuclear antineutrophil cytoplasmic antibodies, as do patients with other autoimmune liver diseases. In autoimmune liver disorders, perinuclear antineutrophil cytoplasmic antibodies react with β -tubulin isotype 5 (as an autoantigen) and its bacterial precursor protein FtsZ, indicating an abnormal immune response to intestinal microorganisms.¹⁶⁰

Mouse models of PSC are limited; the only promising model has a targeted deletion of the mouse biliary transporter MDR2.^{161,162} A sclerosing cholangiopathy is also encountered in adult patients with cystic fibrosis, raising the question of a role of cystic fibrosis transmembrane regulator variants in PSC.^{163,164} The association between cystic fibrosis transmembrane regulator mutations and human PSC is not clear, but the G-protein–coupled bile acid recep-

tor 1 (also known as TGR5) has been identified as a potential candidate gene in PSC.⁵⁵ This molecule is involved in bile acid–induced fluid secretion in biliary epithelial cells and colocalizes with cystic fibrosis transmembrane regulator and the apical sodium-dependent bile salt uptake transporter, suggesting a functional coupling of TGR5 to bile acid uptake and chloride secretion.¹⁶⁵ Intriguingly, it has also been identified in cholangiocyte cilia, where it might conceivably couple biliary bile acid concentration and composition to ductular bile formation.¹⁶⁶

Adhesion Molecules and PSC

The hepatic vascular bed is a unique low-flow environment, with regulated and leukocyte lineage-specific recruitment, trafficking, and positioning important to the pathogenesis of immune-mediated liver injury. Chemokines and their receptors have specific roles in PSC^{167–172}; memory lymphocytes develop during colitis and express receptors that direct them to the GI tract and liver. Normally, the GI endothelium expresses the adhesion molecule mucosal addressin cell adhesion molecule 1 (MADCAM1), which is absent from other vascular beds, and the chemokine CCL25, which is restricted to the small bowel. In PSC, hepatic inflammation up-regulates hepatic MADCAM1 and CCL25 and increases recruitment of mucosal T cells. Such memory cells recirculate between the liver and GI tract but

Table 2. Therapeutic Interventions in PBC and PSC

	Drug	Type	Relative effectiveness	Comments
PBC	UDCA (13–15 mg/kg/day)	Bile transport	++++	Standard of care
	Budesonide/Prednisone	Immunosuppressive	+	Speculated of use if AIH overlap features
	Azathioprine/Mycophenolate	Immunosuppressive	–	No evidence of benefit
	Mofetil			
	Methotrexate	Immunosuppressive	+/-	Biochemical response in some but randomized data lacking
	Cyclosporine	Immunosuppressive	+	Side effect profile significant
	Colchicine	Cell replication	+	Potential for further study
	Rituximab	Anti B cell	NA	Animal models show B cell depletion deleterious
	INT-747	Farnesoid X receptor agonist/Bile transport	Phase II	Biochemical response in Phase II studies; pruritus as major side effect
	Bezafibrate	PPAR α and ρ ligand	++	Biochemical response but no randomized data
	Fenofibrate	PPAR α ligand	++	Biochemical response but no randomized data
	Penicillamine	Copper chelation	–	No benefit
	Tetrathiomolybdate	Copper chelation	+/-	Single study report
	Combivir	Anti viral	–	Controversial disease mechanism with no overt evidence in favor of retroviral hypothesis
	Moexipril	Angiotensin II synthesis blocker	+	Pilot data
	Tamoxifen	Estrogen blocker	+	Pilot data
PSC	UDCA (variable, including high dose)	Bile transport	+/-	High dose therapy reported as deleterious, potentially due to lithocholic acid elevations
	INT-747	Farnesoid X receptor agonist/Bile transport	NA	No clinical data
	Minocycline	Antibiotic/Immunosuppressive	+/-	Pilot studies
	Prednisone	Immunosuppressive	+	No evidence of benefit overall but subgroups potentially helped, eg, IgG4 elevations
	Vancomycin	Antibiotic	NA	Pilot studies
	Anti-TNF	Immunosuppressive	–	No benefit

can become deleteriously activated by hepatic antigens. Twenty percent of T cells that infiltrate the liver in patients with PSC are $\alpha_4\beta_7^+$ CCR9⁺ memory/effector T cells that produce interferon gamma; these are present at low frequencies in other liver diseases. $\alpha_4\beta_7$ and CCR9 are only colocalized on lymphocytes that are activated in the GI tract by dendritic cells there. GI-associated dendritic cells up-regulate GI homing receptors on B cells via a mechanism that depends on the vitamin A metabolite retinoic acid. Liver dendritic cells do not migrate to the GI tract in experimental models, indicating that the $\alpha_4\beta_7^+$ CCR9⁺ T-cell infiltrate that occurs during PSC is primed in the GI tract.

$\alpha_4\beta_7$ and CCR9 expression is functionally relevant because MADCAM1 and CCL25, which are absent from normal liver, are present on hepatic endothelium in liver diseases associated with colitis; $\alpha_4\beta_7^+$ CCR9⁺ lymphocytes from livers of patients with PSC bind MADCAM1 and respond to CCL25 in adhesion and migration assays. In addition to a specific role for CCL25 in PSC, the CC chemokines CCL21 and CCL28 and the CXC chemokines CXCL9 and CXCL10 are involved in the recruitment of T cells into the portal tract in patients with PBC or PSC; in this regard biliary epithelial cells are recognized to produce many chemokines.

Therapeutic Options Beyond UDCA

Although management of patients with cholestatic liver disease has advanced significantly, UDCA is the only treatment available (Table 2); immunosuppressive agents to date have generally not been effective. Although the efficacy of UDCA is debated, it is largely accepted for most patients with PBC and obstetric cholestasis, whereas the effects of UDCA in patients with PSC are limited, if present at all.^{75,149,173,174} UDCA is a major primary bile acid in some species of bears, whereas in humans it is a minor secondary product formed by β -epimerization of chenodeoxycholic acid in the GI tract. The choleretic effect of UDCA and its ability to make the bile acid pool more hydrophilic accounts for its beneficial properties. UDCA also stabilizes hepatocyte membranes, increases defense against oxidative stress, and inhibits apoptosis.¹⁷⁵

Obeticholic acid is a novel chenodeoxycholic acid derivative; unlike UDCA, it is a farnesoid X receptor agonist.¹⁷⁶ Activation of these nuclear receptors reduces bile acid synthesis, induces phases I and II bile acid hydroxylation and conjugation, stimulates export of alternative bile acids, and limits hepatocellular bile acid import. Phase 2 studies in PBC are ongoing. Clinical trials are also planned for a modification of UDCA; 24-norursodeoxycholic acid is a side chain-modified ursodeoxycholic acid derivative that has therapeutic effects in experimental cholestasis.¹⁷⁷ 24-Norursodeoxycholic acid is resistant to amidation and can undergo cholehepatic shunting and directly stimulate cholangiocyte secretion; these

cause hyperchloresis (mainly HCO₃⁻) that protects the liver from cholestatic injury.¹⁷⁸

Conclusions

Cholestatic liver disease is the consequence of many hepatobiliary insults. A variety of pathways contribute to liver tissue damage and the reparative response, so new therapies might be developed to target pathways that mediate these processes. As for rheumatoid arthritis, multiple sclerosis, and inflammatory bowel disease, new therapeutic approaches based on rational and biological insights from recent research could change dramatically the outcomes for patients with cholestatic liver disease.

References

- Alpini G, McGill JM, Larusso NF. The pathobiology of biliary epithelia. *Hepatology* 2002;35:1256–1268.
- Sirica AE, Nathanson MH, Gores GJ, et al. Pathobiology of biliary epithelia and cholangiocarcinoma: proceedings of the Henry M. and Lillian Stratton Basic Research Single-Topic Conference. *Hepatology* 2008;48:2040–2046.
- Alpini G, Roberts S, Kuntz SM, et al. Morphological, molecular, and functional heterogeneity of cholangiocytes from normal rat liver. *Gastroenterology* 1996;110:1636–1643.
- Alpini G, Glaser S, Robertson W, et al. Large but not small intrahepatic bile ducts are involved in secretin-regulated ductal bile secretion. *Am J Physiol* 1997;272:G1064–G1074.
- Masyuk AI, Masyuk TV, Splinter PL, et al. Cholangiocyte cilia detect changes in luminal fluid flow and transmit them into intracellular Ca²⁺ and cAMP signaling. *Gastroenterology* 2006;131:911–920.
- Masyuk AI, Gradilone SA, Banales JM, et al. Cholangiocyte primary cilia are chemosensory organelles that detect biliary nucleotides via P2Y₁₂ purinergic receptors. *Am J Physiol Gastrointest Liver Physiol* 2008;295:G725–G734.
- Strazzabosco M, Mennone A, Boyer JL. Intracellular pH regulation in isolated rat bile duct epithelial cells. *J Clin Invest* 1991;87:1503–1512.
- Fiorotto R, Spirli C, Fabris L, et al. Ursodeoxycholic acid stimulates cholangiocyte fluid secretion in mice via CFTR-dependent ATP secretion. *Gastroenterology* 2007;133:1603–1613.
- Minagawa N, Nagata J, Shibao K, et al. Cyclic AMP regulates bicarbonate secretion in cholangiocytes through release of ATP into bile. *Gastroenterology* 2007;133:1592–1602.
- Strazzabosco M, Fiorotto R, Melero S, et al. Differentially expressed adenylyl cyclase isoforms mediate secretory functions in cholangiocyte subpopulation. *Hepatology* 2009;50:244–252.
- Zollner G, Trauner M. Mechanisms of cholestasis. *Clin Liver Dis* 2008;12:1–26, vii.
- Boyer JL. It's all about bile. *Hepatology* 2009;49:711–723.
- Wagner M, Zollner G, Trauner M. Nuclear receptor regulation of the adaptive response of bile acid transporters in cholestasis. *Semin Liver Dis* 2010;30:160–177.
- Fiorucci S, Antonelli E, Rizzo G, et al. The nuclear receptor SHP mediates inhibition of hepatic stellate cells by FXR and protects against liver fibrosis. *Gastroenterology* 2004;127:1497–1512.
- Fickert P, Fuchsbichler A, Moustafa T, et al. Farnesoid X receptor critically determines the fibrotic response in mice but is expressed to a low extent in human hepatic stellate cells and periductal myofibroblasts. *Am J Pathol* 2009;175:2392–2405.
- König J, Seithel A, Gradhand U, et al. Pharmacogenomics of human OATP transporters. *Naunyn Schmiedeberg Arch Pharmacol* 2006;372:432–443.

17. Johnson AD, Kavousi M, Smith AV, et al. Genome-wide association meta-analysis for total serum bilirubin levels. *Hum Mol Genet* 2009;18:2700–2710.
18. Link E, Parish S, Armitage J, et al. SLC01B1 variants and statin-induced myopathy—a genomewide study. *N Engl J Med* 2008;359:789–799.
19. Lam P, Soroka CJ, Boyer JL. The bile salt export pump: clinical and experimental aspects of genetic and acquired cholestatic liver disease. *Semin Liver Dis* 2010;30:125–133.
20. Nies AT, Keppler D. The apical conjugate efflux pump ABCB2 (MRP2). *Pflugers Arch* 2007;453:643–659.
21. Davit-Spraul A, Gonzales E, Baussan C, et al. The spectrum of liver diseases related to ABCB4 gene mutations: pathophysiology and clinical aspects. *Semin Liver Dis* 2010;30:134–146.
22. Chen XM, O'Hara SP, LaRusso NF. The immunobiology of cholangiocytes. *Immunol Cell Biol* 2008;86:497–505.
23. Chuang YH, Lan RY, Gershwin ME. The immunopathology of human biliary cell epithelium. *Semin Immunopathol* 2009;31:323–331.
24. Chen XM, O'Hara SP, Nelson JB, et al. Multiple TLRs are expressed in human cholangiocytes and mediate host epithelial defense responses to *Cryptosporidium parvum* via activation of NF-kappaB. *J Immunol* 2005;175:7447–7456.
25. Leon MP, Bassendine MF, Wilson JL, et al. Immunogenicity of biliary epithelium: investigation of antigen presentation to CD4+ T cells. *Hepatology* 1996;24:561–567.
26. Spengler U, Leifeld L, Braunschweiler I, et al. Anomalous expression of costimulatory molecules B7-1, B7-2 and CD28 in primary biliary cirrhosis. *J Hepatol* 1997;26:31–36.
27. Cruickshank SM, Southgate J, Selby PJ, et al. Expression and cytokine regulation of immune recognition elements by normal human biliary epithelial and established liver cell lines in vitro. *J Hepatol* 1998;29:550–558.
28. Leon MP, Bassendine MF, Gibbs P, et al. Immunogenicity of biliary epithelium: study of the adhesive interaction with lymphocytes. *Gastroenterology* 1997;112:968–977.
29. Afford SC, Ahmed-Choudhury J, Randhawa S, et al. CD40 activation-induced, Fas-dependent apoptosis and NF-kappaB/AP-1 signaling in human intrahepatic biliary epithelial cells. *FASEB J* 2001;15:2345–2354.
30. Glaser SS, Gaudio E, Miller T, et al. Cholangiocyte proliferation and liver fibrosis. *Expert Rev Mol Med* 2009;11:e7.
31. Marzioni M, Fava G, Alvaro D, et al. Control of cholangiocyte adaptive responses by visceral hormones and neuropeptides. *Clin Rev Allergy Immunol* 2009;36:13–22.
32. Patsenker E, Popov Y, Stickel F, et al. Inhibition of integrin alphavbeta6 on cholangiocytes blocks transforming growth factor-beta activation and retards biliary fibrosis progression. *Gastroenterology* 2008;135:660–670.
33. Marzioni M, Glaser S, Francis H, et al. Autocrine/paracrine regulation of the growth of the biliary tree by the neuroendocrine hormone serotonin. *Gastroenterology* 2005;128:121–137.
34. Omenetti A, Porrello A, Jung Y, et al. Hedgehog signaling regulates epithelial-mesenchymal transition during biliary fibrosis in rodents and humans. *J Clin Invest* 2008;118:3331–3342.
35. Choi SS, Diehl AM. Epithelial-to-mesenchymal transitions in the liver. *Hepatology* 2009;50:2007–2013.
36. Scholten D, Osterreicher CH, Scholten A, et al. Genetic labeling does not detect epithelial-to-mesenchymal transition (EMT) of cholangiocytes in liver fibrosis in mice. *Gastroenterology* 2010;139:987–998.
37. Kahraman A, Gerken G, Canbay A. Apoptosis in immune-mediated liver diseases. *Dig Dis* 2010;28:144–1449.
38. Luedde T, Heinrichsdorff J, de Lorenzi R, et al. IKK1 and IKK2 cooperate to maintain bile duct integrity in the liver. *Proc Natl Acad Sci U S A* 2008;105:9733–9738.
39. Takeda K, Kojima Y, Ikejima K, et al. Death receptor 5 mediated-apoptosis contributes to cholestatic liver disease. *Proc Natl Acad Sci U S A* 2008;105:10895–10900.
40. Selmi C, Podda M, Gershwin ME. Old and rising stars in the lymphoid liver. *Semin Immunopathol* 2009;31:279–282.
41. Dienes HP, Drebber U. Pathology of immune-mediated liver injury. *Dig Dis* 2010;28:57–62.
42. Borchers AT, Shimoda S, Bowlus C, et al. Lymphocyte recruitment and homing to the liver in primary biliary cirrhosis and primary sclerosing cholangitis. *Semin Immunopathol* 2009;31:309–322.
43. Alabraba EB, Lai V, Boon L, et al. Coculture of human liver macrophages and cholangiocytes leads to CD40-dependent apoptosis and cytokine secretion. *Hepatology* 2008;47:552–562.
44. Mantis NJ, Forbes SJ. Secretory IgA: arresting microbial pathogens at epithelial borders. *Immunol Invest* 2010;39:383–406.
45. Strazzabosco M, Fabris L. Functional anatomy of normal bile ducts. *Anat Rec (Hoboken)* 2008;291:653–660.
46. Kamath BM, Loomes KM, Piccoli DA. Medical management of Alagille syndrome. *J Pediatr Gastroenterol Nutr* 2010;50:580–586.
47. Kodama Y, Hijikata M, Kageyama R, et al. The role of notch signaling in the development of intrahepatic bile ducts. *Gastroenterology* 2004;127:1775–1786.
48. Hadj-Rabia S, Baala L, Vabres P, et al. Claudin-1 gene mutations in neonatal sclerosing cholangitis associated with ichthyosis: a tight junction disease. *Gastroenterology* 2004;127:1386–1390.
49. Gissen P, Johnson CA, Morgan NV, et al. Mutations in VPS33B, encoding a regulator of SNARE-dependent membrane fusion, cause arthrogryposis-renal dysfunction-cholestasis (ARC) syndrome. *Nat Genet* 2004;36:400–404.
50. Cullinane AR, Straatman-Iwanowska A, Zaucker A, et al. Mutations in VIPAR cause an arthrogryposis, renal dysfunction and cholestasis syndrome phenotype with defects in epithelial polarization. *Nat Genet* 2010;42:303–312.
51. Stapelbroek JM, van Erpecum KJ, Klomp LW, et al. Liver disease associated with canalicular transport defects: current and future therapies. *J Hepatol* 2010;52:258–271.
52. Masyuk T, Masyuk A, LaRusso N. Cholangiociliopathies: genetics, molecular mechanisms and potential therapies. *Curr Opin Gastroenterol* 2009;25:265–271.
53. Gunay-Aygun M. Liver and kidney disease in ciliopathies. *Am J Med Genet C Semin Med Genet* 2009;151C:296–306.
54. Hirschfield GM, Liu X, Xu C, et al. Primary biliary cirrhosis associated with HLA, IL12A, and IL12RB2 variants. *N Engl J Med* 2009;360:2544–2555.
55. Karlsen TH, Franke A, Melum E, et al. Genome-wide association analysis in primary sclerosing cholangitis. *Gastroenterology* 2010;138:1102–1111.
56. Daly AK, Donaldson PT, Bhatnagar P, et al. HLA-B*5701 genotype is a major determinant of drug-induced liver injury due to flucloxacillin. *Nat Genet* 2009;41:816–819.
57. Donaldson PT, Baragiotta A, Heneghan MA, et al. HLA class II alleles, genotypes, haplotypes, and amino acids in primary biliary cirrhosis: a large-scale study. *Hepatology* 2006;44:667–674.
58. Invernizzi P, Selmi C, Poli F, et al. Human leukocyte antigen polymorphisms in Italian primary biliary cirrhosis: a multicenter study of 664 patients and 1992 healthy controls. *Hepatology* 2008;48:1906–1912.
59. Karlsen TH, Boberg KM, Olsson M, et al. Particular genetic variants of ligands for natural killer cell receptors may contribute to the HLA associated risk of primary sclerosing cholangitis. *J Hepatol* 2007;46:899–906.

60. Hov JR, Lleo A, Selmi C, et al. Genetic associations in Italian primary sclerosing cholangitis: heterogeneity across Europe defines a critical role for HLA-C. *J Hepatol* 2010;52:712–717.
61. Norris S, Kondeatis E, Collins R, et al. Mapping MHC-encoded susceptibility and resistance in primary sclerosing cholangitis: the role of MICA polymorphism. *Gastroenterology* 2001;120:1475–1482.
62. Wiencke K, Spurland A, Schrupf E, et al. Primary sclerosing cholangitis is associated to an extended B8-DR3 haplotype including particular MICA and MICB alleles. *Hepatology* 2001;34:625–630.
63. Hirschfield GM, Liu X, Han Y, et al. Variants at IRF5-TNPO3, 17q12-21 and MMEL1 are associated with primary biliary cirrhosis. *Nat Genet* 2010;42:655–657.
64. Liu X, Invernizzi P, Lu Y, et al. Genome-wide meta-analyses identify three loci associated with primary biliary cirrhosis. *Nat Genet* 2010;42:658–660.
65. Gershwin ME, Mackay IR, Sturgess A, et al. Identification and specificity of a cDNA encoding the 70 kd mitochondrial antigen recognized in primary biliary cirrhosis. *J Immunol* 1987;138:3525–3531.
66. Gershwin ME, Ansari AA, Mackay IR, et al. Primary biliary cirrhosis: an orchestrated immune response against epithelial cells. *Immunol Rev* 2000;174:210–225.
67. Kaplan MM, Gershwin ME. Primary biliary cirrhosis. *N Engl J Med* 2005;353:1261–1273.
68. Gershwin ME, Selmi C, Worman HJ, et al. Risk factors and comorbidities in primary biliary cirrhosis: a controlled interview-based study of 1032 patients. *Hepatology* 2005;42:1194–1202.
69. James OF, Bhopal R, Howel D, et al. Primary biliary cirrhosis once rare, now common in the United Kingdom? *Hepatology* 1999;30:390–394.
70. Myers RP, Shaheen AA, Fong A, et al. Epidemiology and natural history of primary biliary cirrhosis in a Canadian health region: a population-based study. *Hepatology* 2009;50:1884–1892.
71. Selmi C, Gershwin ME, Lindor KD, et al. Quality of life and everyday activities in patients with primary biliary cirrhosis. *Hepatology* 2007;46:1836–1843.
72. Walker JG, Doniach D, Roitt IM, et al. Serological tests in diagnosis of primary biliary cirrhosis. *Lancet* 1965;1:827–831.
73. Prince MI, Ducker SJ, James OF. Case-control studies of risk factors for primary biliary cirrhosis in two United Kingdom populations. *Gut* 2010;59:508–512.
74. Corpechot C, Chretien Y, Chazouilleres O, et al. Demographic, lifestyle, medical and familial factors associated with primary biliary cirrhosis. *J Hepatol* 2010;53:162–169.
75. Lindor KD, Gershwin ME, Poupon R, et al. Primary biliary cirrhosis. *Hepatology* 2009;50:291–308.
76. Lleo A, Bowlus CL, Yang GX, et al. Biliary apotopes and anti-mitochondrial antibodies activate innate immune responses in primary biliary cirrhosis. *Hepatology* 2010;52:987–998.
77. Lleo A, Selmi C, Invernizzi P, et al. Apotopes and the biliary specificity of primary biliary cirrhosis. *Hepatology* 2009;49:871–879.
78. Yeaman SJ, Fussey SP, Danner DJ, et al. Primary biliary cirrhosis: identification of two major M2 mitochondrial autoantigens. *Lancet* 1988;1:1067–1070.
79. Fussey SP, Guest JR, James OF, et al. Identification and analysis of the major M2 autoantigens in primary biliary cirrhosis. *Proc Natl Acad Sci U S A* 1988;85:8654–8658.
80. Coppel RL, McNeilage LJ, Surh CD, et al. Primary structure of the human M2 mitochondrial autoantigen of primary biliary cirrhosis: dihydrolipoamide acetyltransferase. *Proc Natl Acad Sci U S A* 1988;85:7317–7321.
81. Oertelt S, Lian ZX, Cheng CM, et al. Anti-mitochondrial antibodies and primary biliary cirrhosis in TGF-beta receptor II dominant-negative mice. *J Immunol* 2006;177:1655–1660.
82. Wakabayashi K, Yoshida K, Leung PS, et al. Induction of autoimmune cholangitis in non-obese diabetic (NOD).1101 mice following a chemical xenobiotic immunization. *Clin Exp Immunol* 2009;155:577–586.
83. Wakabayashi K, Lian ZX, Moritoki Y, et al. IL-2 receptor alpha(-/-) mice and the development of primary biliary cirrhosis. *Hepatology* 2006;44:1240–1249.
84. Irie J, Wu Y, Wicker LS, et al. NOD.c3c4 congenic mice develop autoimmune biliary disease that serologically and pathogenetically models human primary biliary cirrhosis. *J Exp Med* 2006;203:1209–1219.
85. Salas JT, Banales JM, Sarvide S, et al. Ae2a,b-deficient mice develop antimitochondrial antibodies and other features resembling primary biliary cirrhosis. *Gastroenterology* 2008;134:1482–1493.
86. Ueno Y, Ambrosini YM, Moritoki Y, et al. Murine models of autoimmune cholangitis. *Curr Opin Gastroenterol* 2010;26:274–279.
87. Yoshida K, Yang GX, Zhang W, et al. Deletion of interleukin-12p40 suppresses autoimmune cholangitis in dominant negative transforming growth factor beta receptor type II mice. *Hepatology* 2009;50:1494–1500.
88. Moritoki Y, Zhang W, Tsuneyama K, et al. B cells suppress the inflammatory response in a mouse model of primary biliary cirrhosis. *Gastroenterology* 2009;136:1037–1047.
89. Poupon RE, Balkau B, Eschwege E, et al. A multicenter, controlled trial of ursodiol for the treatment of primary biliary cirrhosis. UDCA-PBC Study Group. *N Engl J Med* 1991;324:1548–1554.
90. Poupon RE, Poupon R, Balkau B. Ursodiol for the long-term treatment of primary biliary cirrhosis. The UDCA-PBC Study Group. *N Engl J Med* 1994;330:1342–1347.
91. Poupon RE, Lindor KD, Cauch-Dudek K, et al. Combined analysis of randomized controlled trials of ursodeoxycholic acid in primary biliary cirrhosis. *Gastroenterology* 1997;113:884–890.
92. Lindor KD, Thorneau TM, Jorgensen RA, et al. Effects of ursodeoxycholic acid on survival in patients with primary biliary cirrhosis. *Gastroenterology* 1996;110:1515–1518.
93. Heathcote EJ, Cauch-Dudek K, Walker V, et al. The Canadian multicenter double-blind randomized controlled trial of ursodeoxycholic acid in primary biliary cirrhosis. *Hepatology* 1994;19:1149–1156.
94. Kuiper EM, Hansen BE, de Vries RA, et al. Improved prognosis of patients with primary biliary cirrhosis that have a biochemical response to ursodeoxycholic acid. *Gastroenterology* 2009;136:1281–1287.
95. Poupon RE, Lindor KD, Pares A, et al. Combined analysis of the effect of treatment with ursodeoxycholic acid on histologic progression in primary biliary cirrhosis. *J Hepatol* 2003;39:12–16.
96. Corpechot C, Abenavoli L, Rabahi N, et al. Biochemical response to ursodeoxycholic acid and long-term prognosis in primary biliary cirrhosis. *Hepatology* 2008;48:871–877.
97. Huet PM, Vincent C, Deslaurier J, et al. Portal hypertension and primary biliary cirrhosis: effect of long-term ursodeoxycholic acid treatment. *Gastroenterology* 2008;135:1552–1560.
98. Kumagi T, Guindi M, Fischer SE, et al. Baseline ductopenia and treatment response predict long-term histological progression in primary biliary cirrhosis. *Am J Gastroenterol* 2010 May 25 [epub ahead of print].
99. Pares A, Caballeria L, Rodes J. Excellent long-term survival in patients with primary biliary cirrhosis and biochemical response to ursodeoxycholic acid. *Gastroenterology* 2006;130:715–720.
100. Selmi C, Gershwin ME. The role of environmental factors in primary biliary cirrhosis. *Trends Immunol* 2009;30:415–420.
101. McNally RJ, Ducker S, James OF. Are transient environmental agents involved in the cause of primary biliary cirrhosis? Evidence from space-time clustering analysis. *Hepatology* 2009;50:1169–1174.

102. Prince MI, Chetwynd A, Diggle P, et al. The geographical distribution of primary biliary cirrhosis in a well-defined cohort. *Hepatology* 2001;34:1083–1088.
103. Ala A, Stanca CM, Bu-Ghanim M, et al. Increased prevalence of primary biliary cirrhosis near Superfund toxic waste sites. *Hepatology* 2006;43:525–531.
104. Selmi C, Balkwill DL, Invernizzi P, et al. Patients with primary biliary cirrhosis react against a ubiquitous xenobiotic-metabolizing bacterium. *Hepatology* 2003;38:1250–1257.
105. Padgett KA, Selmi C, Kenny TP, et al. Phylogenetic and immunological definition of four lipoylated proteins from *Novosphingobium aromaticivorans*, implications for primary biliary cirrhosis. *J Autoimmun* 2005;24:209–219.
106. Olafsson S, Gudjonsson H, Selmi C, et al. Antimitochondrial antibodies and reactivity to *N. aromaticivorans* proteins in Icelandic patients with primary biliary cirrhosis and their relatives. *Am J Gastroenterol* 2004;99:2143–2146.
107. Mattner J, Savage PB, Leung P, et al. Liver autoimmunity triggered by microbial activation of natural killer T cells. *Cell Host Microbe* 2008;3:304–315.
108. Amano K, Leung PS, Rieger R, et al. Chemical xenobiotics and mitochondrial autoantigens in primary biliary cirrhosis: identification of antibodies against a common environmental, cosmetic, and food additive, 2-octynoic acid. *J Immunol* 2005;174:5874–5883.
109. Rieger R, Leung PS, Jeddeloh MR, et al. Identification of 2-nonyloic acid, a cosmetic component, as a potential trigger of primary biliary cirrhosis. *J Autoimmun* 2006;27:7–16.
110. Odin JA, Huebert RC, Casciola-Rosen L, et al. Bcl-2-dependent oxidation of pyruvate dehydrogenase-E2, a primary biliary cirrhosis autoantigen, during apoptosis. *J Clin Invest* 2001;108:223–232.
111. Leung PS, Rossaro L, Davis PA, et al. Antimitochondrial antibodies in acute liver failure: implications for primary biliary cirrhosis. *Hepatology* 2007;46:1436–1442.
112. Bhat M, Guindi M, Heathcote EJ, et al. Transient development of anti-mitochondrial antibodies accompanies autoimmune hepatitis-sclerosing cholangitis overlap. *Gut* 2009;58:152–153.
113. Kimura Y, Leung PS, Kenny TP, et al. Differential expression of intestinal trefoil factor in biliary epithelial cells of primary biliary cirrhosis. *Hepatology* 2002;36:1227–1235.
114. Invernizzi P, Miozzo M, Battezzati PM, et al. Frequency of monosomy X in women with primary biliary cirrhosis. *Lancet* 2004;363:533–535.
115. Miozzo M, Selmi C, Gentilin B, et al. Preferential X chromosome loss but random inactivation characterize primary biliary cirrhosis. *Hepatology* 2007;46:456–462.
116. Selmi C, Mayo MJ, Bach N, et al. Primary biliary cirrhosis in monozygotic and dizygotic twins: genetics, epigenetics, and environment. *Gastroenterology* 2004;127:485–492.
117. Kita H, Lian ZX, Van de Water J, et al. Identification of HLA-A2-restricted CD8(+) cytotoxic T cell responses in primary biliary cirrhosis: T cell activation is augmented by immune complexes cross-presented by dendritic cells. *J Exp Med* 2002;195:113–123.
118. Shimoda S, Nakamura M, Ishibashi H, et al. HLA DRB4 0101-restricted immunodominant T cell autoepitope of pyruvate dehydrogenase complex in primary biliary cirrhosis: evidence of molecular mimicry in human autoimmune diseases. *J Exp Med* 1995;181:1835–1845.
119. Van de Water J, Ansari A, Prindiville T, et al. Heterogeneity of autoreactive T cell clones specific for the E2 component of the pyruvate dehydrogenase complex in primary biliary cirrhosis. *J Exp Med* 1995;181:723–733.
120. Kita H, Matsumura S, He XS, et al. Quantitative and functional analysis of PDC-E2-specific autoreactive cytotoxic T lymphocytes in primary biliary cirrhosis. *J Clin Invest* 2002;109:1231–1240.
121. Selmi C, Meda F, Kasangian A, et al. Experimental evidence on the immunopathogenesis of primary biliary cirrhosis. *Cell Mol Immunol* 2010;7:1–10.
122. Shimoda S, Van de Water J, Ansari A, et al. Identification and precursor frequency analysis of a common T cell epitope motif in mitochondrial autoantigens in primary biliary cirrhosis. *J Clin Invest* 1998;102:1831–1840.
123. Shimoda S, Miyakawa H, Nakamura M, et al. CD4 T-cell auto-reactivity to the mitochondrial autoantigen PDC-E2 in AMA-negative primary biliary cirrhosis. *J Autoimmun* 2008;31:110–115.
124. Miossec P, Korn T, Kuchroo VK. Interleukin-17 and type 17 helper T cells. *N Engl J Med* 2009;361:888–898.
125. Littman DR, Rudensky AY. Th17 and regulatory T cells in mediating and restraining inflammation. *Cell* 2010;140:845–858.
126. Oo YH, Weston CJ, Lalor PF, et al. Distinct roles for CCR4 and CXCR3 in the recruitment and positioning of regulatory T cells in the inflamed human liver. *J Immunol* 2010;184:2886–2898.
127. Lan RY, Cheng C, Lian ZX, et al. Liver-targeted and peripheral blood alterations of regulatory T cells in primary biliary cirrhosis. *Hepatology* 2006;43:729–737.
128. Sasaki M, Ikeda H, Sawada S, et al. Naturally-occurring regulatory T cells are increased in inflamed portal tracts with cholangiopathy in primary biliary cirrhosis. *J Clin Pathol* 2007;60:1102–1107.
129. Lan RY, Salunga TL, Tsuneyama K, et al. Hepatic IL-17 responses in human and murine primary biliary cirrhosis. *J Autoimmun* 2009;32:43–51.
130. Harada K, Shimoda S, Sato Y, et al. Periductal interleukin-17 production in association with biliary innate immunity contributes to the pathogenesis of cholangiopathy in primary biliary cirrhosis. *Clin Exp Immunol* 2009;157:261–270.
131. Kikuchi K, Lian ZX, Yang GX, et al. Bacterial CpG induces hyper-IgM production in CD27(+) memory B cells in primary biliary cirrhosis. *Gastroenterology* 2005;128:304–312.
132. Mao TK, Lian ZX, Selmi C, et al. Altered monocyte responses to defined TLR ligands in patients with primary biliary cirrhosis. *Hepatology* 2005;42:802–808.
133. Moritoki Y, Lian ZX, Wulff H, et al. AMA production in primary biliary cirrhosis is promoted by the TLR9 ligand CpG and suppressed by potassium channel blockers. *Hepatology* 2007;45:314–322.
134. Selmi C, Lleo A, Pasini S, et al. Innate immunity and primary biliary cirrhosis. *Curr Mol Med* 2009;9:45–51.
135. Shimoda S, Harada K, Niiri H, et al. CX3CL1 (fractalkine): a signpost for biliary inflammation in primary biliary cirrhosis. *Hepatology* 2010;51:567–575.
136. Mencarelli A, Renga B, Migliorati M, et al. The bile acid sensor farnesoid X receptor is a modulator of liver immunity in a rodent model of acute hepatitis. *J Immunol* 2009;183:6657–6666.
137. Wang YD, Chen WD, Wang M, et al. Farnesoid X receptor antagonizes nuclear factor kappaB in hepatic inflammatory response. *Hepatology* 2008;48:1632–1643.
138. Glaser S, Lam IP, Franchitto A, et al. Knockout of secretin receptor reduces large cholangiocyte hyperplasia in mice with extrahepatic cholestasis induced by bile duct ligation. *Hepatology* 2010;52:204–214.
139. Prieto J, Garcia N, Marti-Climent JM, et al. Assessment of biliary bicarbonate secretion in humans by positron emission tomography. *Gastroenterology* 1999;117:167–172.
140. Medina JF, Martinez A, Vazquez JJ, et al. Decreased anion exchanger 2 immunoreactivity in the liver of patients with primary biliary cirrhosis. *Hepatology* 1997;25:12–17.
141. Melero S, Spirli C, Zsembergy A, et al. Defective regulation of cholangiocyte Cl-/HCO3(-) and Na+/H+ exchanger activities in primary biliary cirrhosis. *Hepatology* 2002;35:1513–1521.

142. Shibao K, Hirata K, Robert ME, et al. Loss of inositol 1,4,5-trisphosphate receptors from bile duct epithelia is a common event in cholestasis. *Gastroenterology* 2003;125:1175–1187.
143. Beuers U, Hohenester S, de Buy Wenniger LJ, et al. The biliary HCO₃⁻ umbrella: a unifying hypothesis on pathogenetic and therapeutic aspects of fibrosing cholangiopathies. *Hepatology* 2010 Jun 16 [Epub ahead of print].
144. Arenas F, Hervias I, Uriz M, et al. Combination of ursodeoxycholic acid and glucocorticoids upregulates the AE2 alternate promoter in human liver cells. *J Clin Invest* 2008;118:695–709.
145. Juran BD, Atkinson EJ, Larson JJ, et al. Common genetic variation and haplotypes of the anion exchanger SLC4A2 in primary biliary cirrhosis. *Am J Gastroenterol* 2009;104:1406–1411.
146. Poupon R, Ping C, Chretien Y, et al. Genetic factors of susceptibility and of severity in primary biliary cirrhosis. *J Hepatol* 2008;49:1038–1045.
147. Weismuller TJ, Wedemeyer J, Kubicka S, et al. The challenges in primary sclerosing cholangitis—etiopathogenesis, autoimmunity, management and malignancy. *J Hepatol* 2008;48(Suppl 1):S38–S57.
148. Karlsen TH, Schrumpf E, Boberg KM. Update on primary sclerosing cholangitis. *Dig Liver Dis* 2010;42:390–400.
149. Chapman R, Fevery J, Kalloo A, et al. Diagnosis and management of primary sclerosing cholangitis. *Hepatology* 2010;51:660–678.
150. Loftus EV Jr, Harewood GC, Loftus CG, et al. PSC-IBD: a unique form of inflammatory bowel disease associated with primary sclerosing cholangitis. *Gut* 2005;54:91–96.
151. Claessen MM, Vleggaar FP, Tytgat KM, et al. High lifetime risk of cancer in primary sclerosing cholangitis. *J Hepatol* 2009;50:158–164.
152. Abdalian R, Heathcote EJ. Sclerosing cholangitis: a focus on secondary causes. *Hepatology* 2006;44:1063–1074.
153. Webster GJ, Pereira SP, Chapman RW. Autoimmune pancreatitis/IgG4-associated cholangitis and primary sclerosing cholangitis—overlapping or separate diseases? *J Hepatol* 2009;51:398–402.
154. Ponsioen CY, Kuiper H, Ten Kate FJ, et al. Immunohistochemical analysis of inflammation in primary sclerosing cholangitis. *Eur J Gastroenterol Hepatol* 1999;11:769–774.
155. Hashimoto E, Lindor KD, Homburger HA, et al. Immunohistochemical characterization of hepatic lymphocytes in primary biliary cirrhosis in comparison with primary sclerosing cholangitis and autoimmune chronic active hepatitis. *Mayo Clin Proc* 1993;68:1049–1055.
156. Aron JH, Bowlus CL. The immunobiology of primary sclerosing cholangitis. *Semin Immunopathol* 2009;31:383–397.
157. Xu B, Broome U, Ericzon BG, et al. High frequency of autoantibodies in patients with primary sclerosing cholangitis that bind biliary epithelial cells and induce expression of CD44 and production of interleukin 6. *Gut* 2002;51:120–127.
158. Krasinskas AM, Yao Y, Randhawa P, et al. *Helicobacter pylori* may play a contributory role in the pathogenesis of primary sclerosing cholangitis. *Dig Dis Sci* 2007;52:2265–2270.
159. Frulloni L, Lunardi C, Simone R, et al. Identification of a novel antibody associated with autoimmune pancreatitis. *N Engl J Med* 2009;361:2135–2142.
160. Terjung B, Sohne J, Lechtenberg B, et al. p-ANCA in autoimmune liver disorders recognise human beta-tubulin isotype 5 and cross-react with microbial protein FtsZ. *Gut* 2010;59:808–816.
161. Fickert P, Fuchsbichler A, Wagner M, et al. Regurgitation of bile acids from leaky bile ducts causes sclerosing cholangitis in Mdr2 (Abcb4) knockout mice. *Gastroenterology* 2004;127:261–274.
162. Trauner M, Fickert P, Baghdasarian A, et al. New insights into autoimmune cholangitis through animal models. *Dig Dis* 2010;28:99–104.
163. Henckaerts L, Jaspers M, Van Steenberghe W, et al. Cystic fibrosis transmembrane conductance regulator gene polymorphisms in patients with primary sclerosing cholangitis. *J Hepatol* 2009;50:150–157.
164. Blanco PG, Zaman MM, Junaidi O, et al. Induction of colitis in cfr^{-/-} mice results in bile duct injury. *Am J Physiol Gastrointest Liver Physiol* 2004;287:G491–G496.
165. Keitel V, Cupisti K, Ullmer C, et al. The membrane-bound bile acid receptor TGR5 is localized in the epithelium of human gallbladders. *Hepatology* 2009;50:861–870.
166. Keitel V, Ullmer C, Haussinger D. The membrane-bound bile acid receptor TGR5 (Gpbar-1) is localized in the primary cilium of cholangiocytes. *Biol Chem* 2010;391:785–789.
167. Grant AJ, Lalor PF, Salmi M, et al. Homing of mucosal lymphocytes to the liver in the pathogenesis of hepatic complications of inflammatory bowel disease. *Lancet* 2002;359:150–157.
168. Eksteen B, Grant AJ, Miles A, et al. Hepatic endothelial CCL25 mediates the recruitment of CCR9⁺ gut-homing lymphocytes to the liver in primary sclerosing cholangitis. *J Exp Med* 2004;200:1511–1517.
169. Mora JR, Iwata M, Eksteen B, et al. Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science* 2006;314:1157–1160.
170. Adams DH, Eksteen B, Curbishley SM. Immunology of the gut and liver: a love/hate relationship. *Gut* 2008;57:838–848.
171. Miles A, Liaskou E, Eksteen B, et al. CCL25 and CCL28 promote alpha4 beta7-integrin-dependent adhesion of lymphocytes to MAdCAM-1 under shear flow. *Am J Physiol Gastrointest Liver Physiol* 2008;294:G1257–G1267.
172. Eksteen B, Mora JR, Haughton EL, et al. Gut homing receptors on CD8 T cells are retinoic acid dependent and not maintained by liver dendritic or stellate cells. *Gastroenterology* 2009;137:320–329.
173. Lindor KD, Kowdley KV, Luketic VA, et al. High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009;50:808–814.
174. EASL clinical practice guidelines: management of cholestatic liver diseases. *J Hepatol* 2009;51:237–267.
175. Beuers U. Drug insight: mechanisms and sites of action of ursodeoxycholic acid in cholestasis. *Nat Clin Pract Gastroenterol Hepatol* 2006;3:318–328.
176. Zollner G, Wagner M, Trauner M. Nuclear receptors as drug targets in cholestasis and drug-induced hepatotoxicity. *Pharmacol Ther* 2010;126:228–243.
177. Fickert P, Wagner M, Marschall HU, et al. 24-norUrsodeoxycholic acid is superior to ursodeoxycholic acid in the treatment of sclerosing cholangitis in Mdr2 (Abcb4) knockout mice. *Gastroenterology* 2006;130:465–481.
178. Halilbasic E, Fiorotto R, Fickert P, et al. Side chain structure determines unique physiologic and therapeutic properties of norursodeoxycholic acid in Mdr2^{-/-} mice. *Hepatology* 2009;49:1972–1981.

Received July 28, 2010. Accepted September 7, 2010.

Reprint requests

Address requests for reprints to: M. Eric Gershwin, MD, Division of Rheumatology, Allergy and Clinical Immunology, University of California at Davis School of Medicine, 451 Health Sciences Drive, Suite 6510, Davis, California 95616. e-mail: megershwin@ucdavis.edu; fax: (530) 752-4669.

Conflicts of interest

G.M.H. and E.J.H. have received unrestricted grant support from Axcan Pharma. M.E.G. discloses no conflicts.

Funding

Supported in part by the Canadian Institutes for Health Research (MOP 74621), the Canadian Primary Biliary Cirrhosis Society, and National Institutes of Health grant DK39588.