



## $\beta$ -Catenin pathway is required for TGF- $\beta$ 1 inhibition of PPAR $\gamma$ expression in cultured hepatic stellate cells

Jingjing Qian<sup>b</sup>, Minghui Niu<sup>a</sup>, Xuguang Zhai<sup>a</sup>, Qian Zhou<sup>a</sup>, Yajun Zhou<sup>a,\*</sup>

<sup>a</sup> Department of Biochemistry & Molecular Biology, Medical College, Nantong University, Nantong, Qi xiu road 19, Nantong 226001, Jiangsu, China

<sup>b</sup> Affiliated Hospital of Nantong University, Qi xiu road 19, Nantong 226001, Jiangsu, China

### ARTICLE INFO

#### Article history:

Received 22 March 2012

Received in revised form 4 June 2012

Accepted 5 June 2012

#### Keywords:

$\beta$ -Catenin

Transforming growth factor- $\beta$ 1

Peroxisome proliferator-activated

receptor- $\gamma$

Hepatic stellate cell

### ABSTRACT

Hepatic stellate cell (HSC) activation is a key step in process of liver fibrosis. Transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) is the most powerful mediator of HSC activation and plays a central role in liver fibrosis. Peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) is an important regulator of adipocyte differentiation and has been proposed as a crucial factor for inhibition of HSC activation. The effect of TGF- $\beta$ 1 on PPAR $\gamma$  in HSCs is largely unknown. This study is aimed to examine whether TGF- $\beta$ 1 can influence PPAR $\gamma$  expression, focusing on the role of  $\beta$ -catenin pathway, a key pathway linked to adipogenesis, in TGF- $\beta$ 1 regulation of PPAR $\gamma$  in cultured HSCs. Our results demonstrated that TGF- $\beta$ 1 evidently inhibited PPAR $\gamma$  expression and activity in cultured HSCs, which were mediated through  $\beta$ -catenin pathway. TGF- $\beta$ 1 promoted  $\beta$ -catenin expression and also increased the stability of  $\beta$ -catenin protein through ERK1/2/glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ) axis in cultured HSCs. Moreover, TGF- $\beta$ 1 inhibition of PPAR $\gamma$  expression by  $\beta$ -catenin pathway caused the increase in  $\alpha$ 1(1) collagen and tissue inhibitor of matrix metalloproteinase expression. These results indicated for the first time that TGF- $\beta$ 1 could down-regulate PPAR $\gamma$  expression through  $\beta$ -catenin pathway and subsequently contributed to the increase in  $\alpha$ 1(1) collagen level in cultured HSCs.

© 2012 Elsevier Ltd. All rights reserved.

### 1. Introduction

Liver fibrosis is a wound healing process characterized by over accumulation of extracellular matrix proteins (ECM) especially collagen types I and III. Hepatic stellate cell (HSC) is the most relevant cell type for the development of liver fibrosis. Its activation is the key step in the process of liver fibrogenesis. Activated HSCs are the main ECM-producing cells in the injured liver. Many studies focus on the mechanisms underlying the process of HSC activation. Interestingly, it has been shown that the process of HSC activation is analogous to the reverse transdifferentiation of adipocyte to preadipocyte (fibroblast) [1]. The nuclear receptor peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) is a key regulator of adipocyte differentiation *in vivo* as well as *in vitro* [2] and acts as a transcriptional activator of many adipocyte-specific genes [3].

It has been demonstrated that the level of PPAR $\gamma$  protein is high in quiescent HSCs and its expression and its activity are reduced

during HSC activation *in vitro* and *in vivo* [4,5]. Introduction of exogenous PPAR $\gamma$  is sufficient to reverse the morphology of activated HSCs to the quiescent phenotype with reduced expression of activation markers and with the accumulation of retinyl palmitate [6], a characteristic of quiescent HSCs. Furthermore, PPAR $\gamma$  over-expression *in vivo* strongly attenuated stellate cell activation and liver fibrosis [7]. PPAR $\gamma$  has been proposed as a crucial factor for inhibition of HSC activation [1,6,7].

It has been shown that Wnt/ $\beta$ -catenin pathway is a crucial regulator of adipogenesis [8] and  $\beta$ -catenin is a key effector of the Wnt signaling pathway. The absence of Wnts facilitates the phosphorylation of  $\beta$ -catenin protein by glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ), leading to  $\beta$ -catenin (a primary substrate of GSK-3 $\beta$ ) ubiquitination and degradation [9], whereas Wnts can induce the phosphorylation and inactivation of GSK-3 $\beta$  and subsequent stabilization of  $\beta$ -catenin protein [10]. Importantly, functional Wnt/ $\beta$ -catenin pathway has been demonstrated in HSCs and contributes to HSC activation and liver fibrosis [11].

Transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) is one of the most important cytokines expressed following liver injury and TGF- $\beta$ 1 is considered the most powerful mediator of HSC activation *in vitro* and *in vivo* [12] and [13] and plays a central role in liver fibrosis [14]. Its role in liver fibrosis has been widely studied. However, the effect of TGF- $\beta$ 1 on PPAR $\gamma$  expression is unknown in HSCs. Interestingly, TGF- $\beta$ 1 appears to be associated with  $\beta$ -catenin pathway

**Abbreviations:** TGF- $\beta$ 1, transforming growth factor- $\beta$ 1; PPAR $\gamma$ , peroxisome proliferator-activated receptor- $\gamma$ ; GSK-3 $\beta$ , glycogen synthase kinase-3 $\beta$ ; ECM, extracellular matrix proteins; HSC, hepatic stellate cell; ERK1/2, extracellular regulated protein kinase1/2; TIMP, tissue inhibitor of matrix metalloproteinase.

\* Corresponding author. Tel.: +86 513 85883082; fax: +86 513 85051796.

E-mail address: [zhoyujun64@yahoo.com](mailto:zhoyujun64@yahoo.com) (Y. Zhou).

in human lung fibroblasts [15]. Moreover, TGF- $\beta$ 1 also involves adipocyte differentiation [16].

Based on the observations above, there is a possibility that TGF- $\beta$ 1 could affect PPAR $\gamma$  expression in HSCs and  $\beta$ -catenin pathway might mediate the role of TGF- $\beta$ 1 in PPAR $\gamma$  expression in HSCs. Therefore, this study is aimed to examine whether TGF- $\beta$ 1 could influence PPAR $\gamma$  expression, focusing on the role of  $\beta$ -catenin pathway in TGF- $\beta$ 1 regulation of PPAR $\gamma$  in cultured HSCs.

## 2. Materials and methods

### 2.1. Materials

TGF- $\beta$ 1 and PD98059 (a MEK inhibitor) were purchased from CalBiochem (La Jolla, CA, USA). XAV939 (a potent and specific inhibitor of  $\beta$ -catenin pathway), and all antibodies were purchased from Santa Cruz (Santa Cruz, CA, USA). Lithium chloride (LiCl, an inhibitor of GSK-3 $\beta$  activation) and GW9662 (a specific PPAR $\gamma$  antagonist) were purchased from Sigma (St. Louis, MO, USA).

### 2.2. HSC isolation and culture

HSCs were isolated from Sprague–Dawley rats as we described previously [17]. Semi-confluent HSCs with 4–8 passages were used for experiments. Cells were serum-starved for 24 h in DMEM with 0.4% FBS before treatment with TGF- $\beta$ 1 for 24 h. Animals received humane care and experimental protocols were conducted according to national and local guidelines.

### 2.3. Western blot analysis

Western blot analysis was performed as we described previously [17]. Target proteins were detected by primary antibodies against PPAR $\gamma$  (diluted 1:500),  $\beta$ -catenin (diluted 1:1000), Ser 9 phosphorylated GSK-3 $\beta$  (diluted 1:500), GSK-3 $\beta$  (diluted 1:1000),  $\alpha$ 1(1) collagen (diluted 1:2000),  $\beta$ -actin (diluted 1:2000), respectively, and subsequently by horseradish peroxidase-conjugated secondary antibodies (diluted 1:4000).  $\beta$ -Actin or total GSK-3 $\beta$  was used as the internal control.

### 2.4. RNA isolation and real-time PCR

Total RNA was extracted by using TRI-Reagent (Sigma, St. Louis, USA) according to the manufacturer's instructions and treated with deoxyribonuclease I. Real-time PCR was performed as we described previously [18]. mRNA levels of target genes relative to the endogenous cyclophilin control were calculated as suggested by Schmittgen et al. [19]. The results were expressed as fold changes relative to the control. The primers used in real-time PCR were as follows:

#### Rat PPAR $\gamma$ :

- (forward) 5'-ATTCTGGCCCACTTCGG-3';
- (reverse) 5'-TGGAAGCTGATGCTTTATCCCCA-3'.

#### Rat $\beta$ -catenin:

- (forward) 5'-CCACGACTAGTTCAGCTGCTTGAC-3';
- (reverse) 5'-ACTGCACAAACAGTGAATGGTATT-3'.

#### Rat $\alpha$ 1(1) collagen:

- (forward) 5'-TTCCCTGGACCTAAGGGTACT-3';
- (reverse) 5'-TTGAGCTCCAGCTTCGCC-3'.

#### Rat tissue inhibitor of matrix metalloproteinase-1 (TIMP-1):

- (forward) 5'-CGCAGCGAGGAGTTTCTCAT-3';
- (reverse) 5'-GGCAGTGATGTGCAATTTCC-3'.

#### Rat TIMP-2:

- (forward) 5'-GACACGCTTAGCATCACCCAGA-3';
  - (reverse) 5'-CTGTGACCCAGTCCATCCAGAG-3'.
- Rat cyclophilin:
- (forward) 5'-TGGATGGCAAGCATGTGGTCTTTG-3';
  - (reverse) 5'-CTTCTGCTGGTCTTGCCATTCCT-3'.

### 2.5. Plasmids and transient transfection assays

The PPAR $\gamma$  activity reporter plasmid pPPRE  $\times$  3-TK-Luc contains three consensus PPAR $\gamma$ -responsive elements linked to the herpes virus thymidine kinase promoter and a Photinus luciferase vector (Addgene Inc., Cambridge, MA, USA). Plasmid p $\beta$ catenin encodes dominant-negative  $\beta$ -catenin (a gift from Dr. Jane B. Treppel, Medical Oncology Branch, NCI, NIH, Bethesda, USA).

HSCs in twelve-well plastic plates were transiently transfected with pPPRE  $\times$  3-TK-Luc (1  $\mu$ g DNA/well) plus 30 ng of control vector expressing Renilla luciferase (pRL-TK; Promega, Madison, USA) by using LipofectAMINE reagent (Life Technologies, New York, USA) following manufacturer's instructions. In some experiments, HSCs were cotransfected with pPPRE  $\times$  3-TK-Luc (1  $\mu$ g DNA/well) plus p $\beta$ catenin (1  $\mu$ g DNA/well) or the empty vector and 30 ng Renilla luciferase. The empty vector was used to ensure an equal amount of total DNA in transfection assay. Luciferase activity was quantified fluorimetrically by using the Dual-Luciferase Reporter Assay System (Promega, Madison, USA) and data were expressed as the ratios of Photinus to Renilla luciferase activity for normalization of Photinus luciferase activity. HSCs in 25 cm<sup>2</sup> flask were transfected with 8  $\mu$ g of p $\beta$ catenin or the empty vector per flask.

### 2.6. $\beta$ -catenin and GSK-3 $\beta$ silencing through RNA interference


HSCs in 25 cm<sup>2</sup> flask were transfected with a pool of four short interfering RNAs (siRNAs; 250 pmol) targeted specifically to rat  $\beta$ -catenin or rat GSK-3 $\beta$  using LipofectAMINE reagent (Life Technologies, New York, USA) following manufacturer's instructions. Rat  $\beta$ -catenin siRNA was purchased from Santa Cruz (Santa Cruz, CA, USA). Rat GSK-3 $\beta$  siRNA duplexes were synthesized by Life Technologies Corporation (Shanghai, China) to target the rat GSK-3 $\beta$ : (a) 5'-GATCTGCCATCGAGACATT-3', (b) 5'-CTCAAGAACTGTCAAGTAA-3', (c) 5'-TCAGAAGTCTAGCCTATAT-3', and (d) 5'-ACACTAAAGTCATTGGAAA-3' [20]. As a negative control, cells were transfected with siRNA against GFP (Life Technologies Corporation, Shanghai, China).

### 2.7. Electrophoretic mobility shift assay (EMSA)

Nuclear proteins were extracted as we previously described [21]. The sense oligonucleotide of the putative DNA binding site for NF $\kappa$ B was: 5'-GGGGACTTCCC-3' [22]. EMSA were performed as we previously reported [21]. Briefly, the probe was end-labeled with [<sup>32</sup>P]dATP and 5  $\mu$ g of nuclear extract was incubated with the <sup>32</sup>P-labeled probe for 30 min. For the competition assay, 5  $\mu$ g of nuclear extract was preincubated with 100-fold molar excess of the unlabeled NF $\kappa$ B probe (cold probe) before addition of the labeled probe. For supershift assay, 5  $\mu$ g of nuclear extract was preincubated with 2  $\mu$ g of anti-NF $\kappa$ B (p50) antibody (Santa Cruz, CA, USA) before addition of the labeled probe. The samples were then loaded onto a 5% nondenaturing polyacrylamide gel. The gels were subjected to electrophoresis, drying, and autoradiography.

### 2.8. Statistical analysis

Differences between means were evaluated using an unpaired two-sided Student's *t*-test (*P* < 0.05 considered as significant). Where appropriate, comparisons of multiple treatment conditions



**Fig. 1.** TGF- $\beta$ 1 down-regulates PPAR $\gamma$  expression and trans-activation activity in cultured HSCs. (A and B) Western blot analysis of PPAR $\gamma$  protein level ( $n=3$ ). HSCs were incubated with 3 ng/ml of TGF- $\beta$ 1 for different time periods (A) or with increasing dose of TGF- $\beta$ 1 for 24 h (B).  $\beta$ -Actin was used as internal control. (C) Real-time PCR analysis of PPAR $\gamma$  mRNA level ( $n=3$ ). HSCs were incubated with increasing dose of TGF- $\beta$ 1 for 24 h.  $*P<0.05$  vs cells without TGF- $\beta$ 1. (D) Transfection assay for analysis of PPAR $\gamma$  trans-activation activity ( $n=6$ ). HSCs were transfected with pPPRE  $\times$  3-TK-Luc (1  $\mu$ g/well) plus pRL-TK (30 ng/well) and stimulated with increasing dose of TGF- $\beta$ 1 for 24 h.  $*P<0.05$  vs cells without TGF- $\beta$ 1.

with controls were analyzed by ANOVA with the Dunnett's test for *post hoc* analysis.

### 3. Results

#### 3.1. TGF- $\beta$ 1 treatment reduces PPAR $\gamma$ expression in cultured HSCs


To investigate the effect of TGF- $\beta$ 1 on PPAR $\gamma$  expression, serum-starved HSCs were treated with 3 ng/ml of TGF- $\beta$ 1 for different time periods. The level of PPAR $\gamma$  protein was examined by Western blot analysis. Fig. 1A showed that TGF- $\beta$ 1 markedly reduced PPAR $\gamma$  protein level in a time-dependent manner. Next, HSCs were treated with increasing dose of TGF- $\beta$ 1 for 24 h and the levels of PPAR $\gamma$  protein and mRNA were analyzed by Western blot and real-time PCR, respectively. The results demonstrated that TGF- $\beta$ 1 inhibited PPAR $\gamma$  expression at protein (Fig. 1B) and mRNA (Fig. 1C) levels in a dose-dependent manner.

TGF- $\beta$ 1-induced decrease in PPAR $\gamma$  expression might affect its activity. Therefore, HSCs were transfected with pPPRE  $\times$  3-TK-Luc (containing three consensus PPAR $\gamma$ -responsive elements) and treated with increasing dose of TGF- $\beta$ 1 for 24 h. Luciferase assay indicated that TGF- $\beta$ 1 treatment led to a dose-dependent decrease in the trans-activation activity of PPAR $\gamma$  (Fig. 1D).

These results strongly suggested that TGF- $\beta$ 1 exerted an inhibitory effect on PPAR $\gamma$  expression and trans-activation activity in cultured HSCs.

#### 3.2. TGF- $\beta$ 1 treatment increases $\beta$ -catenin protein level in cultured HSCs

To test the possibility whether  $\beta$ -catenin pathway might involve the effect of TGF- $\beta$ 1 on PPAR $\gamma$  in cultured HSCs, the influence




**Fig. 2.** TGF- $\beta$ 1 increases  $\beta$ -catenin expression in cultured HSCs. (A and B) Western blot analysis of  $\beta$ -catenin protein level ( $n=3$ ). HSCs were incubated with 3 ng/ml of TGF- $\beta$ 1 for different time periods (A) or with increasing dose of TGF- $\beta$ 1 for 24 h (B).  $\beta$ -Actin was used as internal control. (C) Real-time PCR analysis of  $\beta$ -catenin mRNA level ( $n=3$ ). HSCs were incubated with or without 3 ng/ml of TGF- $\beta$ 1 for 24 h.  $*P<0.05$  vs cells without TGF- $\beta$ 1.

of TGF- $\beta$ 1 on  $\beta$ -catenin protein level was examined. HSCs were stimulated with 3 ng/ml of TGF- $\beta$ 1 for different time periods or stimulated with different doses of TGF- $\beta$ 1 for 24 h. Western blot analysis indicated that TGF- $\beta$ 1 induced  $\beta$ -catenin protein level in a time-dependent manner (Fig. 2A) and 3 or 5 ng/ml of TGF- $\beta$ 1 clearly led to an increase in  $\beta$ -catenin protein level (Fig. 2B). To examine the influence of TGF- $\beta$ 1 on  $\beta$ -catenin mRNA, HSCs were stimulated with or without 3 ng/ml of TGF- $\beta$ 1 for 24 h. Real-time PCR analysis demonstrated that 3 ng/ml of TGF- $\beta$ 1 significantly increased  $\beta$ -catenin mRNA level (Fig. 2C). Fig. 2 indicated that TGF- $\beta$ 1 up-regulated  $\beta$ -catenin expression in cultured HSCs.

Extracellular regulated protein kinase1/2 (ERK1/2) is a member of the mitogen-activated protein kinase (MAPK) family. Its activation leads to the phosphorylation (at Ser9) and inactivation of GSK-3 $\beta$ , subsequently inhibiting the degradation of  $\beta$ -catenin protein (a primary substrate of GSK-3 $\beta$ ) in human lung fibroblasts [15]. It has been shown that stimulation of HSCs with 3 ng/ml of TGF- $\beta$ 1 for 20 min is able to induce ERK1/2 activation [23]. Thereby, we examine the effect of TGF- $\beta$ 1-induced activation of ERK1/2 on phosphorylation (at Ser9) of GSK-3 $\beta$  and on  $\beta$ -catenin protein in HSCs. HSCs were pretreated with or without 10  $\mu$ M of PD98059 for 30 min before stimulation with or without TGF- $\beta$ 1 for an additional 20 min (for analysis of Ser 9 phosphorylated GSK-3 $\beta$ ) or 24 h (for analysis of  $\beta$ -catenin protein). Western blot analysis showed that TGF- $\beta$ 1 treatment clearly caused the increase in the levels of Ser 9 phosphorylated GSK-3 $\beta$  (Fig. 3A) and  $\beta$ -catenin protein (Fig. 3B) which were inhibited by pretreatment with 10  $\mu$ M of PD98059, whereas PD98059 alone had no effect on both types of proteins. These results indicated that TGF- $\beta$ 1-induced activation of ERK1/2 resulted in inactivation of GSK-3 $\beta$  and subsequent stabilization of  $\beta$ -catenin protein in cultured HSCs. To further verify the results, HSCs were incubated with 10 mM of LiCl (an inhibitor of GSK-3 $\beta$  activation) for 24 h or the gene expression of GSK-3 $\beta$  was knocked down by transfection of HSCs with siRNA against GSK-3 $\beta$ . Western blot analysis indicated that  $\beta$ -catenin protein level was up-regulated by LiCl treatment (Fig. 3C) or knockdown of the gene expression of GSK-3 $\beta$  (Fig. 3D), which supported that TGF- $\beta$ 1-induced inhibition of GSK-3 $\beta$  activity led to the stabilization of  $\beta$ -catenin protein in HSCs.






**Fig. 3.** TGF-β1-induced activation of ERK1/2 contributes to the phosphorylation of GSK-3β at Ser9 and the accumulation of β-catenin protein in cultured HSCs. (A and B) Western blot analyses of Ser 9 phosphorylated GSK-3β and β-catenin protein levels ( $n=3$ ). HSCs were pretreated with or without 10 μM of PD98059 for 30 min before stimulation with or without 3 ng/ml of TGF-β1 for an additional 20 min (for analysis of Ser 9 phosphorylated GSK-3β) (A) or 24 h (for analysis of β-catenin protein) (B). Total GSK-3β or β-actin was used as internal control. (C and D) Western blot analysis of level of β-catenin protein or total GSK-3β protein ( $n=3$ ). HSCs were treated with or without 10 mM of LiCl for 24 h (C) or transfected with siRNA against GSK-3β or siRNA against GFP (as a negative control) for 24 h (D). β-Actin was used as internal control. (E and F) EMSA of DNA binding activity of NF-κB. Serum-starved HSCs were stimulated with 3 ng/ml of TGF-β1 for 20 min. Nuclear protein was extracted and 5 μg of nuclear protein was incubated with [<sup>32</sup>P]dATP labeled NF-κB probe or nuclear protein from HSCs treated with TGF-β1 was preincubated with 100-fold molar excess of the unlabeled NF-κB probe (cold probe) before addition of labeled NF-κB probe (for competition assay, lane 3) (E). For supershift assay, nuclear protein from HSCs treated with TGF-β1 was preincubated with anti-NF-κB(p50) antibody (2 μg) before addition of labeled NF-κB probe (lane 2) (F). A representative EMSA was shown from three independent experiments.

Taken together, these data indicated that TGF-β1 was linked to β-catenin pathway in cultured HSCs.

Since TGF-β1 can induce ERK1/2 activity [23] and the latter leads to NF-κB activation in HSCs [24], we further investigated the effect of TGF-β1 on NF-κB trans-activation activity by EMSA. As shown in Fig. 3E and F, TGF-β1 induced marked DNA binding activity of NF-κB in cultured HSCs.

### 3.3. β-Catenin pathway is required for TGF-β1 inhibition of PPARγ expression and activity in cultured HSCs

Since TGF-β1 was associated with β-catenin pathway and TGF-β1 exerted evident inhibition of PPARγ expression in HSCs, we next examined whether β-catenin pathway mediated TGF-β1-induced decrease in PPARγ expression and activity. Our previous study showed that 5 μM of XAV939 (a specific inhibitor of β-catenin pathway) was enough to inhibit β-catenin pathway in cultured HSCs (data not shown). HSCs were pretreated with or without 5 μM of XAV939 for 30 min before treatment with or without 3 ng/ml of TGF-β1 for an additional 24 h, the levels of PPARγ protein and mRNA were analyzed by Western blot and real-time PCR, respectively. Fig. 4 shows that TGF-β1 significantly reduced PPARγ protein level (Fig. 4A, upper panel) and mRNA level (Fig. 4A, down panel), which were partially reversed by treatment with XAV939 (Fig. 4A,




**Fig. 4.** β-Catenin pathway mediates TGF-β1-induced inhibition of PPARγ expression and trans-activation activity in cultured HSCs. (A and B) Analyses of PPARγ protein and mRNA levels or β-catenin protein level ( $n=3$ ). HSCs were pretreated with or without 5 μM of XAV939 for 30 min before treatment with or without 3 ng/ml of TGF-β1 for an additional 24 h (A), or HSCs were transfected with siRNA against β-catenin (β-cat siRNA) or siRNA against GFP (as a negative control) before treatment with or without 3 ng/ml of TGF-β1 for 24 h (B). The levels of PPARγ protein and mRNA were analyzed by Western blot and real-time PCR, respectively and β-catenin protein level was analyzed by Western blot. \* $P<0.05$  vs the control without treatment (the first column on the left). \*\* $P<0.05$  vs cells with TGF-β1 alone (the second column on the left). (C and D) Transfection assay for analysis of PPARγ trans-activation activity ( $n=6$ ). HSCs were transfected with pPPRE × 3-TK-Luc (1 μg/well) plus pRL-TK (30 ng per well) and pretreated with or without 5 μM of XAV939 for 30 min before treatment with or without 3 ng/ml of TGF-β1 for an additional 24 h (C), or HSCs were transfected with pPPRE × 3-TK-Luc (1 μg/well), pdn catenin (pdncat, 1 μg/well) or empty plasmid, and pRL-TK (30 ng/well) and treated with or without 3 ng/ml of TGF-β1 for an additional 24 h (D). \* $P<0.05$  vs the control (the first column on the left in C or D). \*\* $P<0.05$  vs cells with TGF-β1 alone (the second column on the left in C).

the second band or the second column on the right). To further verify the results, HSCs were transfected with siRNA against β-catenin gene before treatment with 3 ng/ml of TGF-β1 for 24 h. Western blot (Fig. 4B, upper panel) and real-time PCR (Fig. 4B, down panel) analyses indicated that knockdown of β-catenin expression also relieved TGF-β1 inhibition of PPARγ expression, which was consistent with the results in Fig. 4A.

Next, the role of β-catenin pathway in TGF-β1 inhibition of PPARγ activity was tested. HSCs were transfected with pPPRE × 3-TK-Luc and pretreated with or without 5 μM of XAV939 for 30 min before treatment with or without 3 ng/ml of TGF-β1 for an additional 24 h. Luciferase assay (Fig. 4C) revealed that TGF-β1 treatment resulted in the decline in the trans-activation activity of PPARγ, which was also partially inhibited by XAV939. To further verify the data, HSCs were co-transfected with pPPRE × 3-TK-Luc plus pdn catenin (encoding dominant-negative β-catenin to interrupt β-catenin pathway) or empty vector and stimulated with or without 3 ng/ml of TGF-β1 for 24 h. As shown in Fig. 4D by luciferase assays, pdn catenin partially reversed TGF-β1 inhibition of PPARγ trans-activation activity.


Collectively, these results evidently suggested the involvement of β-catenin pathway in TGF-β1 inhibition of PPARγ expression and trans-activation activity in cultured HSCs.



**Fig. 5.** The inhibitory effect of TGF- $\beta$ 1 on PPAR $\gamma$  by  $\beta$ -catenin pathway leads to the increase in alpha1(1) collagen gene expression in cultured HSCs. (A and B) Analyses of alpha1(1) collagen mRNA and protein levels ( $n=3$ ). HSCs were pretreated with different doses of XAV939 for 30 min before treatment with 3 ng/ml of TGF- $\beta$ 1 for an additional 24 h (A), or HSCs were pretreated with or without 10  $\mu$ M of GW9662 (a specific PPAR $\gamma$  antagonist) and 5  $\mu$ M of XAV939 for 30 min before stimulation with or without 3 ng/ml of TGF- $\beta$ 1 for an additional 24 h (B). The levels of alpha1(1) collagen mRNA and protein were analyzed by real-time PCR and Western blot ( $n=3$ ), respectively. # $P<0.05$  vs the control (the first column on the left in A or B). \* $P<0.05$  vs cells with TGF- $\beta$ 1 alone (the second column on the left in B).  $\xi P<0.05$  vs cells with XAV939 plus TGF- $\beta$ 1 (the third column on the left in B). (C) Analysis of alpha1(1) collagen mRNA level ( $n=3$ ). HSCs were transfected with pdcn catenin (8  $\mu$ g/flask) or empty plasmid and pretreated with or without 10  $\mu$ M of GW9662 for 30 min before stimulation with or without 3 ng/ml of TGF- $\beta$ 1 for an additional 24 h. The levels of alpha1(1) collagen mRNA and protein were analyzed by real-time PCR (left panel) and Western blot (right panel) analyses, respectively ( $n=3$ ). \* $P<0.05$  vs the control (the first column on the left). # $P<0.05$  vs cells with TGF- $\beta$ 1 alone (the second column on the left).  $\xi P<0.05$  vs cells with pdcn catenin plus TGF- $\beta$ 1 (the third column on the left).

#### 3.4. The inhibitory effect of TGF- $\beta$ 1 on PPAR $\gamma$ by $\beta$ -catenin pathway results in the increase of alpha1(1) collagen and TIMP expression in cultured HSCs

Liver fibrosis is characterized by over accumulation of ECM especially collagen types I and III which were mainly produced by the activated HSCs and TGF- $\beta$ 1 is considered the most powerful mediator of HSC activation [12] and [13]. Since TGF- $\beta$ 1 markedly down-regulated PPAR $\gamma$  (a crucial factor for inhibition of HSC activation) expression through  $\beta$ -catenin pathway, it was interesting to examine the influence of TGF- $\beta$ 1-induced inhibition of PPAR $\gamma$  expression by  $\beta$ -catenin pathway on collagen expression in HSCs. First, HSCs were pretreated with different doses of XAV939 for 30 min and then stimulated with TGF- $\beta$ 1 for 24 h for examining the role of TGF- $\beta$ 1-induced  $\beta$ -catenin pathway in alpha1(1) collagen expression. As shown in Fig. 5A, real-time PCR and Western blot analyses demonstrated that XAV939 inhibited TGF- $\beta$ 1-induced increases in levels of alpha1(1) collagen mRNA (upper panel) and protein (down panel). Next, HSCs were pretreated with or without 10  $\mu$ M of GW9662 (a specific PPAR $\gamma$  antagonist) [25] and 5  $\mu$ M of XAV939 for 30 min before stimulation with or without TGF- $\beta$ 1 for an additional 24 h. The levels of alpha1(1) collagen mRNA and protein were analyzed by real-time PCR and Western blot analyses, respectively. Fig. 5B demonstrated that TGF- $\beta$ 1 alone evidently increased the levels of alpha1(1) collagen mRNA and protein as



**Fig. 6.** The inhibitory effect of TGF- $\beta$ 1 on PPAR $\gamma$  by  $\beta$ -catenin pathway leads to the increase in mRNA levels of TIMP-1 and TIMP-2. HSCs were treated as in Fig. 5(B). The mRNA levels of TIMP-1 and TIMP-2 were analyzed by real-time PCR ( $n=3$ ), respectively. # $P<0.05$  vs the respective control without treatment. \* $P<0.05$  vs the respective cells with TGF- $\beta$ 1 alone.  $\xi P<0.05$  vs the respective cells with XAV939 plus TGF- $\beta$ 1.

compared with the control (the first column or the first band on the left). As expected, pretreatment with XAV939 partially eliminated TGF- $\beta$ 1-induced increase in levels of alpha1(1) collagen mRNA and protein (the third column or the third band on the left), which were attenuated by GW9662 (the fourth column or the fourth band on the left), whereas in the presence of TGF- $\beta$ 1, GW9662 alone did not influence the levels of alpha1(1) collagen mRNA and protein (the first column or the first band on the right) as compared with the control with TGF- $\beta$ 1 alone (the second column or the second band on the left).

To further verify the results, HSCs were transfected with or without 8  $\mu$ g of pdcn catenin (per flask) and then pretreated with or without 10  $\mu$ M of GW9662 for 30 min before stimulation with or without TGF- $\beta$ 1 for an additional 24 h. The levels of alpha1(1) collagen mRNA and protein were analyzed by real-time PCR and Western blot analyses, respectively. As shown in Fig. 5C, interruption of  $\beta$ -catenin pathway by pdcn catenin also partially inhibited TGF- $\beta$ 1-induced increase in alpha1(1) collagen mRNA (the third column on the left in left panel) and protein (the third band on the left in right panel) levels, which were attenuated by GW9662 (the fourth column on the left in the left panel and the fourth band on the left in the right panel).

Results in Fig. 5 suggested that TGF- $\beta$ 1-induced decline in PPAR $\gamma$  expression through  $\beta$ -catenin pathway resulted in the increase in alpha1(1) collagen expression in cultured HSCs.

TIMPs exert the inhibitory effects on matrix metalloproteinase activity, leading to inhibition of the ECM degradation. Therefore, we also examined the role of TGF- $\beta$ 1-induced inhibition of PPAR $\gamma$  expression by  $\beta$ -catenin pathway in the expression of TIMP-1 and TIMP-2 at mRNA level. HSCs were treated as in Fig. 5B. Fig. 6 demonstrated that blockade of  $\beta$ -catenin pathway partially inhibited TGF- $\beta$ 1-induced increase in mRNA levels of TIMP-1 and TIMP-2 (the third black or empty column on the left), which were attenuated by GW9662 (the fourth black or empty column on the left). These results were consistent with that in Fig. 5B and C.

#### 4. Discussion

Because quiescent HSC is much like adipocyte and HSC activation is analogous to adipocyte to preadipocyte (fibroblast) transdifferentiation [1], PPAR $\gamma$ , a key regulator of adipocyte differentiation [2], was found to exert a crucial role in inhibition of HSC activation [1,6,7]. TGF- $\beta$ 1 is considered to be a major factor accelerating the progression of organ fibrosis including liver fibrosis [14] and is the most powerful mediator of HSC activation, a key step in the process of liver fibrogenesis, *in vitro* and *in vivo* [12,13]. This study demonstrated that TGF- $\beta$ 1 exerted an

evident inhibitory effect on PPAR $\gamma$  expression and activity in cultured HSCs and TGF- $\beta$ 1-induced  $\beta$ -catenin pathway mediated the effect of TGF- $\beta$ 1 on PPAR $\gamma$  expression and activity. TGF- $\beta$ 1 could up-regulate  $\beta$ -catenin expression at protein and mRNA levels. Furthermore, our results showed that TGF- $\beta$ 1-induced activation of ERK1/2 resulted in the inactivation of GSK-3 $\beta$  and subsequent stabilization of  $\beta$ -catenin protein. The effect of TGF- $\beta$ 1 on  $\beta$ -catenin expression and on  $\beta$ -catenin protein stability led to the decline in PPAR $\gamma$  expression and trans-activation activity, which contributed to the increase in  $\alpha$ 1(1) collagen and TIMP expression in cultured HSCs. These results indicated for the first time that TGF- $\beta$ 1 could down-regulate PPAR $\gamma$  expression and activity through  $\beta$ -catenin pathway in cultured HSCs.

TGF- $\beta$ 1 triggers multiple signaling pathways in HSCs, such as smad pathway [14], ERK1/2 pathway [23], and p38 MAPK pathway [26]. The existence of functional  $\beta$ -catenin pathway has also been demonstrated in HSC and is related to HSC activation and liver fibrosis [11]. We examined the relationship between TGF- $\beta$ 1 and  $\beta$ -catenin pathway in primary cultured HSCs and found that TGF- $\beta$ 1 evidently increased  $\beta$ -catenin protein level, not only by up-regulation of its expression, but also by maintaining its stability. Recently, it was found that TGF- $\beta$ 1 treatment induced  $\beta$ -catenin pathway in airway smooth muscle cells [27] and in fibroblasts [15] and [28], which were in line with our results. Since TGF- $\beta$ 1 is able to induce ERK1/2 pathway [23] and ERK1/2 appears to be associated with the accumulation of  $\beta$ -catenin (a primary substrate of GSK-3 $\beta$ ) in human lung fibroblasts [15], our further study revealed a crosstalk between TGF- $\beta$ 1-induced ERK pathway and TGF- $\beta$ 1-induced  $\beta$ -catenin pathway in cultured HSCs. This result was based on the observations: First, TGF- $\beta$ 1 induced phosphorylation of GSK-3 $\beta$  at Ser9, an inactive state of GSK-3 $\beta$  [15], and [29], which was inhibited by interruption of TGF- $\beta$ 1-induced ERK1/2 pathway. Second, TGF- $\beta$ 1-induced ERK1/2 pathway up-regulated the protein level of  $\beta$ -catenin. Third, treatment with the inhibitor for GSK-3 $\beta$  activity or knockdown of GSK-3 $\beta$  expression increased  $\beta$ -catenin protein level in HSCs. Therefore, TGF- $\beta$ 1 seemed to be able to increase  $\beta$ -catenin protein level through inhibition of GSK-3 $\beta$  activity *via* activation of ERK1/2 pathway in cultured HSCs. Our previous results showed that ERK1/2 mediated leptin down-regulation of PPAR $\gamma$  expression in HSCs [17]. The mediators between ERK1/2 pathway and PPAR $\gamma$  expression were unknown in leptin regulation of PPAR $\gamma$  expression, but  $\beta$ -catenin seemed to be a mediator between ERK1/2 pathway and PPAR $\gamma$  expression in TGF- $\beta$ 1-induced decline in PPAR $\gamma$  expression in HSCs. These results did not exclude other mechanisms by which TGF- $\beta$ 1 regulated  $\beta$ -catenin protein level. Interestingly, when our paper was being prepared, Alhmetshina et al. found that  $\beta$ -catenin mediated TGF- $\beta$ 1-induced dermal fibrosis [30] and revealed that TGF- $\beta$  stimulated  $\beta$ -catenin signaling in a p38-dependent manner by decreasing the expression of the Wnt antagonist Dickkopf-1. Therefore, there is possibility that Dickkopf-1 might also be correlated with TGF- $\beta$ 1-induced decline in PPAR $\gamma$  expression in HSCs.

NF- $\kappa$ B is an essential signaling pathway involved in conveying inflammatory signals and PPAR $\gamma$  exerts anti-inflammatory effects in many cell types, such as smooth muscle cells, endothelial cells, and macrophages [31,32]. Our results also showed TGF- $\beta$ 1-induced increase in NF- $\kappa$ B activity in HSCs. PPAR $\gamma$  appears to inhibit NF- $\kappa$ B activation in skeletal muscle [33]. The relationship between PPAR $\gamma$  and NF- $\kappa$ B in HSCs is worthy of further investigation.

Our study had indicated that  $\beta$ -catenin pathway mediated TGF- $\beta$ 1 inhibition of PPAR $\gamma$  expression and trans-activation activity in cultured HSCs. As PPAR $\gamma$  is a crucial factor for inhibition of HSC activation [1,6,7], these results might therefore partially explain the mechanisms underlying the involvement of  $\beta$ -catenin pathway in HSC activation and liver fibrosis [11].

Although the results in this report demonstrated  $\beta$ -catenin pathway mediated TGF- $\beta$ 1 inhibition of PPAR $\gamma$  expression and activity in HSCs, induction of  $\beta$ -catenin pathway by LiCl was shown to elevate PPAR $\gamma$  expression and activity in colon cancer cells [34]. Our data in present study cannot explain the discrepancy in HSCs and colon cancer cells. The discrepancy might be due to the different types of cells. Increased production of type I collagen is a common hallmark of fibrotic diseases in various organs including liver [14]. Activated HSCs are the principal cells to produce type I collagen in fibrotic liver and contribute to the development of liver fibrosis through TGF- $\beta$ 1-stimulated collagen production [14]. Our results further demonstrated that TGF- $\beta$ 1 down-regulation of PPAR $\gamma$  expression through  $\beta$ -catenin pathway led to the increase in  $\alpha$ 1(1) collagen expression at mRNA and protein levels in cultured HSCs. Moreover, TGF- $\beta$ 1-induced decline in PPAR $\gamma$  expression by  $\beta$ -catenin pathway also up-regulated the mRNA levels of TIMP-1 and TIMP-2 (the inhibitors of matrix metalloproteinase).

TGF- $\beta$ 1 is the most powerful mediator of HSC activation [12,13] and PPAR $\gamma$  is a key inhibitor of HSC activation [1,6,7]. TGF- $\beta$ 1 negative-regulation of the expression and activity of PPAR $\gamma$  in cultured HSCs might extend the mechanisms underlying the role of TGF- $\beta$ 1 in HSC activation and liver fibrosis. It should be emphasized that these results in this report were generated from cultured HSCs, and that they might not necessarily and comprehensively reflect the situation in HSCs *in vivo*. Gong et al. demonstrated that disruption of TGF- $\beta$ 1 pathway significantly attenuated chronic hypoxia-induced down-regulation of PPAR $\gamma$  in mouse lung [35], implying that TGF- $\beta$ 1 seems also to exert an inhibitory effect on PPAR $\gamma$  expression *in vivo*. The mechanisms by which TGF- $\beta$ -induced  $\beta$ -catenin exerts its inhibitory effect on PPAR $\gamma$  expression in HSCs deserves further investigation. Interestingly, ectopic expression of PPAR $\gamma$  can also inhibit TGF- $\beta$ 1 expression in HSCs [6]. This result suggested that there may be a reciprocal relationship between TGF- $\beta$ 1 and PPAR $\gamma$  in HSCs.

## 5. Conclusions

In summary, this report indicated that TGF- $\beta$ 1 down-regulated PPAR $\gamma$  expression and activity through increasing  $\beta$ -catenin expression and stabilizing  $\beta$ -catenin protein in cultured HSCs. The ERK1/2/GSK-3 $\beta$  axis mediated TGF- $\beta$ 1-induced stabilization of  $\beta$ -catenin protein. TGF- $\beta$ 1 inhibition of PPAR $\gamma$  led to increase in the expression of collagen and TIMPs in cultured HSCs. Since TGF- $\beta$ 1 is the most powerful mediator of HSC activation *in vitro* and *in vivo* [12,13] and plays a central role in liver fibrosis [14], the data in this report might provide novel insights into the molecular mechanisms underlying the role of TGF- $\beta$ 1 in HSC activation and liver fibrosis.

## Conflict of interest

The authors have no any conflicts of interest to disclose.

## Acknowledgments

This work was supported by the grant from the National Science Foundation of China (Grant No. 30971117), the project funded by the Priority Academic Program Development of Jiangsu Higher Education Institution, and the project funded by the Natural Science Foundation of Jiangsu Higher Education Institutions of China (Grant No. 11KJB310009). We are grateful to Dr. Jane B. Trepel (Medical Oncology Branch, NCI, NIH, Bethesda, USA) for kindly providing the plasmid p $\beta$ catenin.



## References

- [1] Tsukamoto H. Adipogenic phenotype of hepatic stellate cells. *Alcoholism, Clinical and Experimental Research* 2005;29:1325–35.
- [2] Farmer SR. Transcriptional control of adipocyte formation. *Cell Metabolism* 2006;4:263–73.
- [3] Lehrke M, Lazar MA. The many faces of PPARgamma. *Cell* 2005;123:993–9.
- [4] She H, Xiong S, Hazra S, Tsukamoto H. Adipogenic transcriptional regulation of hepatic stellate cells. *Journal of Biological Chemistry* 2005;280:4959–67.
- [5] Miyahara T, Schrum L, Rippe R, Xiong S, Yee Jr HF, Motomura K, et al. Peroxisome proliferator-activated receptors and hepatic stellate cell activation. *Journal of Biological Chemistry* 2000;275:35715–22.
- [6] Hazra S, Xiong S, Wang J, Rippe RA, Krishna V, Chatterjee K, et al. Peroxisome proliferator-activated receptor  $\gamma$  induces a phenotypic switch from activated to quiescent hepatic stellate cells. *Journal of Biological Chemistry* 2004;279:11392–401.
- [7] Yang L, Chan CC, Kwon OS, Liu S, McGhee J, Stimpson SA, et al. Regulation of peroxisome proliferator-activated receptor-gamma in liver fibrosis. *American Journal of Physiology: Gastrointestinal and Liver Physiology* 2006;291:G902–11.
- [8] Arango NA, Szotek PP, Manganaro TF, Oliva E, Donahoe PK, Teixeira J. Conditional deletion of beta-catenin in the mesenchyme of the developing mouse uterus results in a switch to adipogenesis in the myometrium. *Developmental Biology* 2005;288:276–83.
- [9] Christodoulides C, Lagathu C, Sethi JK, Vidal-Puig A. Adipogenesis and WNT signalling. *Trends in Endocrinology and Metabolism* 2009;20:16–24.
- [10] Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, et al. Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell* 2002;108:837–47.
- [11] Cheng JH, She H, Han YP, Wang J, Xiong S, Asahina K, et al. Wnt antagonism inhibits hepatic stellate cell activation and liver fibrosis. *American Journal of Physiology: Gastrointestinal and Liver Physiology* 2008;294:G39–49.
- [12] Bataller R, Brenner DA. Liver fibrosis. *Journal of Clinical Investigation* 2005;115:209–18.
- [13] Friedman SL. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *Journal of Biological Chemistry* 2000;275:2247–50.
- [14] Inagaki Y, Okazaki I. Emerging insights into transforming growth factor beta Smad signal in hepatic fibrogenesis. *Gut* 2007;56:284–92.
- [15] Caraci F, Gili E, Calafiore M, Failla M, La Rosa C, Crimi N, et al. TGF-beta1 targets the GSK-3beta/beta-catenin pathway via ERK activation in the transition of human lung fibroblasts into myofibroblasts. *Pharmacological Research* 2008;57:274–82.
- [16] Choy L, Skillington J, Derynck R. Roles of autocrine TGF- $\beta$  receptor and Smad signaling in adipocyte differentiation. *Journal of Cell Biology* 2000;149:667–81.
- [17] Zhou Y, Jia X, Qin J, Lu C, Zhu H, Li X, et al. Leptin inhibits PPARgamma gene expression in hepatic stellate cells in the mouse model of liver damage. *Molecular and Cellular Endocrinology* 2010;323:193–200.
- [18] Zhou Y, Jia X, Wang G, Wang X, Liu J. PI-3K/AKT and ERK signaling pathways mediate leptin-induced inhibition of PPARgamma gene expression in primary rat hepatic stellate cells. *Molecular and Cellular Biochemistry* 2009;325:131–9.
- [19] Schmittgen TD, Zakrajsek BA, Mills AG, Gorn V, Singer MJ, Reed MW. Quantitative reverse transcription-polymerase chain reaction to study mRNA decay: comparison of endpoint and real-time methods. *Analytical Biochemistry* 2000;285:194–204.
- [20] Juhaszova M, Zorov DB, Kim SH, Pepe S, Fu Q, Fishbein KW, et al. Glycogen synthase kinase-3beta mediates convergence of protection signaling to inhibit the mitochondrial permeability transition pore. *Journal of Clinical Investigation* 2004;113:1535–49.
- [21] Zhou Y, Jia X, Zhou M, Liu J. Egr-1 is involved in the inhibitory effect of leptin on PPARgamma expression in hepatic stellate cell in vitro. *Life Sciences* 2009;84:544–51.
- [22] Lee KS, Buck M, Houghlum K, Chojkier M. Activation of hepatic stellate cells by TGF alpha and collagen type I is mediated by oxidative stress through c-myc expression. *Journal of Clinical Investigation* 1995;96:2461–8.
- [23] Liu Y, Wen XM, Lui EL, Friedman SL, Cui W, Ho NP, et al. Therapeutic targeting of the PDGF and TGF-beta-signaling pathways in hepatic stellate cells by PTK787/ZK22258. *Laboratory Investigation* 2009;89:1152–60.
- [24] Itagaki T, Shimizu I, Cheng X, Yuan Y, Oshio A, Tamaki K, et al. Opposing effects of oestradiol and progesterone on intracellular pathways and activation processes in the oxidative stress induced activation of cultured rat hepatic stellate cells. *Gut* 2005;54:1782–9.
- [25] Ho TC, Chen SL, Shih SC, Wu JY, Han WH, Cheng HC, et al. Pigment epithelium-derived factor is an intrinsic antifibrosis factor targeting hepatic stellate cells. *American Journal of Pathology* 2010;177:1798–811.
- [26] Ohyama T, Sato K, Kishimoto K, Yamazaki Y, Horiguchi N, Ichikawa T, et al. Azelnidipine is a calcium blocker that attenuates liver fibrosis and may increase antioxidant defence. *British Journal of Pharmacology* 2012;165:1173–87.
- [27] Baarsma HA, Menzen MH, Halayko AJ, Meurs H, Kerstjens HA, Gosens R.  $\beta$ -Catenin signaling is required for TGF- $\beta$ 1-induced extracellular matrix production by airway smooth muscle cells. *American Journal of Physiology: Lung Cellular and Molecular Physiology* 2011;301:L956–65.
- [28] Amini Nik S, Ebrahim RP, Van Dam K, Cassiman JJ, Teijpar S. TGF-beta modulates beta-catenin stability and signaling in mesenchymal proliferations. *Experimental Cell Research* 2007;313:2887–95.
- [29] Rodionova E, Conzelmann M, Maraskovsky E, Hess M, Kirsch M, Giese T, et al. GSK-3 mediates differentiation and activation of proinflammatory dendritic cells. *Blood* 2007;109:1584–92.
- [30] Akhmetshina A, Palumbo K, Dees C, Bergmann C, Venalis P, Zerr P, et al. Activation of canonical Wnt signalling is required for TGF- $\beta$ -mediated fibrosis. *Nature Communications*, in press.
- [31] Staels B, Koenig W, Habib A, Merval R, Lebret M, Torra IP, et al. Activation of human aortic smooth-muscle cells is inhibited by PPARalpha but not by PPARgamma activators. *Nature* 1998;393:790–3.
- [32] Straus DS, Glass CK. Anti-inflammatory actions of PPAR ligands: new insights on cellular and molecular mechanisms. *Trends in Immunology* 2007;28:551–8.
- [33] Remels AH, Langen RC, Gosker HR, Russell AP, Spaapen F, Voncken JW, et al. PPARgamma inhibits NF-kappaB-dependent transcriptional activation in skeletal muscle. *American Journal of Physiology: Endocrinology and Metabolism* 2009;297:E174–83.
- [34] Jansson EA, Are A, Greicius G, Kuo IC, Kelly D, Arulampalam V, et al. The Wnt/beta-catenin signaling pathway targets PPARgamma activity in colon cancer cells. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:1460–5.
- [35] Gong K, Xing D, Li P, Aksut B, Ambalavanan N, Yang Q, et al. Hypoxia induces downregulation of PPAR- $\gamma$  in isolated pulmonary arterial smooth muscle cells and in rat lung via transforming growth factor- $\beta$  signaling. *American Journal of Physiology: Lung Cellular and Molecular Physiology* 2011;301:L899–907.