# Kinetics of Excess CO<sub>2</sub> Output during and after Intensive Exercise

Takahiro YUNOKI, Masahiro HORIUCHI, and Tokuo YANO

Laboratory of Human Movement Sciences, Faculty of Education, Hokkaido University, Kita-ku, Sapporo, 060–0811 Japan

Abstract: In order to clarify the kinetics of excess CO<sub>2</sub> output during and after intensive exercise, six male subjects were each instructed to perform 40-, 60- and 80-s cycle ergometer exercises (282±9 W, 90 rpm). Ventilation and gas exchange parameters were recorded breath-bybreath, and lactate concentration (La) was repeatedly measured with blood samples from a finger tip. The increase in La from the resting value to peak value and the duration of exercise showed a significant linear relationship (r=0.91, p<0.01) passing through zero, indicating that lactic acid was produced at a constant rate in working muscles from the beginning of exercise. However, in contrast to this increase in La, excess VCO2, defined as the difference between  $\dot{V}$ CO<sub>2</sub> and  $\dot{V}$ O<sub>2</sub>, showed a temporary negative value after the start of exercise. Subsequently, excess VCO2 became positive, reaching a peak at 60 s post-exercise, and then decreased down to zero at about 9 min after the end of the 80-s exercise. End-tidal CO2 rose above the pre-exercise level during exercise and at about 3 min post-exercise, and thereafter remained below the pre-exercise level. Excess CO2, calculated by the sum of excess VCO2 from the start of exercise to the 10th min after the end of exercise, was significantly correlated with the increase in La from resting to 10 min post-exercise (r=0.88, p<0.01). These results suggest that although excessive CO2 output (excess CO2) in response to intensive exercise is related to the increase in lactic acid, the time course of excessive CO2 output (excess VCO2) is delayed, relative to the production of lactic acid, and is affected by hyperventilation. [Japanese Journal of Physiology, 49, 139-144, 1999]

Key words: intensive exercise, lactate, excessive CO<sub>2</sub> output, hyperventilation.

It has been reported that as part of the  $CO_2$  produced at the onset of constant load exercise is stored in the body ( $CO_2$  stores), the respiratory exchange ratio (RER) temporarily decreases [1, 2]. This change in  $CO_2$  stores during exercise can be estimated from the change in mixed venous  $CO_2$  pressure ( $P\bar{v}_{CO_2}$ ), because it has been theoretically and experimentally shown that change in  $CO_2$  stores brings about a change in the level of  $P\bar{v}_{CO_2}$  [3–5]. Actually,  $P\bar{v}_{CO_2}$  during constant load exercise is reported to increase for the first 2–3 min and then maintain a constant level [6–8]. Furthermore, it has been confirmed that there is a correspondence between the change in  $CO_2$  stores, estimated from the change in RER, and the change in

 $P\bar{\mathbf{v}}_{\mathrm{CO}_2}$  at that time [8]. Thus,  $\mathrm{CO}_2$  output during constant load exercise is thought to be affected by a change in the  $\mathrm{CO}_2$  stores accompanied by a change in  $\mathrm{CO}_2$  pressure ( $P\mathrm{CO}_2$ ).

In constant load exercise above the anaerobic threshold (AT), the response of O<sub>2</sub> uptake to the load is delayed compared with that below the AT, but the kinetics of CO<sub>2</sub> output indicate no clear change [2, 9]. This constancy in CO<sub>2</sub> output kinetics is considered to be due to the addition of CO<sub>2</sub> produced by buffering lactic acid to CO<sub>2</sub> aerobically produced in the early stage of exercise [9]. Thereafter, Stringer *et al.* [10] carried out detailed measurements of the arterial acid—base balance relative to changes in lactic acid

Received on July 1, 1998; accepted on March 8, 1999

Correspondence should be addressed to: Takahiro Yunoki, Laboratory of Human Movement Sciences, Faculty of Education, Hokkaido University, Nishi 7, Kita 11, Kita-ku, Sapporo, 060–0811 Japan. Tel: +81–11–706–3422, Fax. +81–11–706–5433, E-mail: yunoki@edu.hokudai. ac.jp

during and after 6-min constant load exercises (two intensities between a level above AT and below maximal O<sub>2</sub> uptake).

During relatively heavy constant load exercise, it is thought that lactate concentration in the blood is not always identical with that in the working muscle due to the influence of lactic acid diffusion, and therefore examination of the relevance of lactate and HCO<sub>3</sub> in the working muscle is thought to be necessary. However, previous studies [11-14] have primarily been concerned with measurement during recovery after exercise, since the duration of exercise in previous studies has been short. Furthermore, although they serve as a reference, these studies on acid-base balance have not directly examined the kinetics of CO<sub>2</sub> output, and there has been very little quantitative investigation of intensive exercise [15]. Additionally, it is unthinkable that all of the CO<sub>2</sub> produced by buffering lactic acid during exercise would be expired during intensive exercise if the duration of exercise is short. In this study, therefore, in order to clarify the kinetics of excess CO2 output in response to intensive exercise, we measured pulmonary gas exchange and blood lactate levels during and after intensive exercise, and we examined the correlation between these values.

### **METHODS**

1. Experimental procedures. The subjects were six active males who did not train regularly. Their mean (SD) age, height, and weight were 23.0 (1.9) years, 170.5 (6.1) cm, and 66.1 (2.9) kg, respectively. Each subject was informed of the purpose of the study, the experimental procedure, and the risks associated with the experiments before consent was obtained. Furthermore, they were instructed to refrain from heavy exercise on the day before the experiment, not to take a meal within 3 h before the experiment, and to remain in a resting state for 1 h before the start of the experiment.

Initially, each subject performed an incremental exercise test in order to estimate maximal  $O_2$  uptake  $(\dot{V}O_{2\,\text{max}})$  using a cycle ergometer controlled by an eddy current (Ergometer 232C: Combi) at 60 rpm. Each subject exercised for 4 min at unloaded work after a 5-min rest, and the work was then increased by a ramp pattern (20 W/min) until exhaustion.

On separate days after the incremental exercise test, each subject performed a cycling exercise with maximal effort for 30 s at a load (kp) corresponding to 7.5%×body weight (Wingate test) [16] using a cycle ergometer controlled by an eddy current (Powermax-

VII: Combi). During the exercise, the power output was measured every 5 s from an imposed load and cycling rate (rpm), and the mean power output (mean power) for 30 s of exercise was obtained. A few days later, each subject, after resting on the cycle ergometer for 5 min, exercised at a constant load for 80 s at 90 rpm (282±9 W), corresponding to 50% of the mean power exerted in the Wingate test. After the end of this exercise, the subjects sat on a chair in a resting state for 30 min. In order to assess the process of change in blood lactate concentration during 80-s exercise and the change in kinetics of CO<sub>2</sub> output with different durations of exercise, 40- and 60-s exercises at the same load were also performed on separate days using the same protocol (intensive exercise test).

Data on ventilation ( $\dot{V}E$ ), oxygen uptake ( $\dot{V}O_2$ ), carbon dioxide output (VCO2) and end-tidal carbon dioxide (ET<sub>CO</sub>,) were obtained breath-by-breath using a respiratory gas analyzer (Aeromonitor AE-280S, Minato) during intensive and incremental exercise tests. These data were averaged over a 20-s period. O<sub>2</sub> concentration and CO<sub>2</sub> concentration were measured by a zirconium sensor and infrared absorption analyzer, respectively. These gas analyzers were calibrated by known standard gases.  $\dot{V}E$  was measured by a hot-wire flow meter, and the flow meter was calibrated by flowing room air at a constant rate (2 l/about 1 s) using a 2l calibrator syringe. Blood lactate concentration (La) was measured by an automatic analyzer (1500 sport, YSI) using 25 µl of blood sampled from a finger tip with a capillary tube at rest before exercise, immediately after the end of exercise, and at 5, 10, 20 and 30 min post-exercise. The analyzer was calibrated with a standard liquid (5 mM of lactate).

According to the method of Cerretelli and Di Prampero [15], excessive  $CO_2$  output per unit of time (excess  $\dot{V}CO_2$ ) was calculated by subtracting  $\dot{V}O_2$  from  $\dot{V}CO_2$  during intensive exercise tests. The total of excessive  $CO_2$  output (excess  $CO_2$ ) was calculated by the addition of values of excess  $\dot{V}CO_2$  from the start of exercise until 10 min post-exercise.

**2. Statistical analysis.** Differences in variables obtained in the three intensive exercise tests were analyzed using analysis of variance (ANOVA). A post hoc test (Turkey's test) was then conducted to assess the differences. The strength of the relationship between dependent and independent variables was expressed by a single correlation coefficient of Pearson. The level of significance was p < 0.05. Results are expressed as mean  $\pm$  standard deviation.

## **RESULTS**

Maximal  $O_2$  uptake ( $\dot{V}O_{2\,max}$ ) obtained in the incremental exercise test was  $2.7\pm0.4\,l/min$ . Maximal and mean powers during the Wingate test were  $726\pm54$ 




Fig. 1. Excess  $\dot{V}\text{CO}_2$ ,  $E\text{T}_{\text{CO}_2}$ , and  $\dot{V}\text{E}/\dot{V}\text{CO}_2$  at rest and during and after 80-s intensive exercise. Vertical bars represent the exercise period. Values are means  $\pm$  SD.

and  $564\pm18$  W, respectively. The powers exerted in the 40-, 60- and 80-s intensive exercise tests were similar. The power exerted in the 80-s intensive exercise test corresponded to  $108\pm11.5\%$  of the power ( $268\pm33.5$  W) at exhaustion in the incremental exercise test.

The time courses of excess  $VCO_2$ ,  $ET_{CO_2}$ , and  $\dot{V}E/\dot{V}CO_2$  in the 80-s intensive exercise test are shown in Fig. 1. The plots indicate mean values per 20 s for all subjects. Excess VCO2 showed a peak negative value at 40 s after the start of exercise and then became positive. This positive value showed a peak at 60 s after the end of exercise and then dropped to zero at about 9 min post-exercise.  $ET_{CO_2}$  increased up to 20 s after the end of exercise and then returned to its pre-exercise value at about 3 min post-exercise.  $ET_{CO}$ continued to decrease gradually up to about 5 min postexercise and then became stable over the whole recovery range. The changes in  $\dot{V}$ E/ $\dot{V}$ CO<sub>2</sub> and ET<sub>CO</sub>, showed a relationship with the reflected image. During 40- and 60-s intensive exercise tests, excess VCO2 showed a temporary negative value at 40 s after the start of exercises. Subsequently, excess  $\dot{V}$ CO<sub>2</sub> became positive, reaching a peak at about 60-80 s post-exercise, and then dropped to zero at about 7 and 8 min after the end of the 40- and 60-s exercises, respectively. The changes in  $ET_{CO_2}$  and  $\dot{V}E/\dot{V}CO_2$  during the 40- and 60-s intensive exercise tests were similar to those during the 80-s test.

Figure 2 shows the changes in La in the intensive exercise tests. The plots indicate mean values for all subjects. The values immediately after the end of the




Fig. 2. Changes in blood lactate concentration at rest and during recovery of 40 ( $\blacksquare$ ), 60 ( $\triangle$ ) and 80 s ( $\blacksquare$ ) of intensive exercises. Values are means  $\pm$  SD.




Fig. 3. Relationship between duration of exercise and increase in blood lactate concentrations (difference between blood lactate concentrations at rest and peak) in three intensive exercises.

40-, 60- and 80-s exercises were  $3.49\pm0.82$ ,  $5.30\pm0.38$  and  $6.75\pm1.29$  mM, respectively. Peak La values, which were observed at 5 min post-exercise, were  $4.06\pm0.47$ ,  $6.00\pm0.62$  and  $7.74\pm0.85$  mM, respectively. The increases in La from immediately after the end of exercises to 5 min postexercise were  $0.58\pm0.85$ ,  $0.70\pm0.79$  and  $0.99\pm1.07$  mM, respectively. Figure 3 shows the relationship between the maximal increase in La after the end of intensive exercise (difference between La at rest and peak La:  $\Delta$ La peak) and the duration of exercise. There was a significant correlation between the duration of exercise and  $\Delta$ La peak (r=0.91, p<0.01). Extrapolating this regression line to zero of  $\Delta$ La peak, the estimated time was around zero.

The total excessive  $CO_2$  output (excess  $CO_2$ ), which was the sum of the values of excess  $VCO_2$  from the start of exercise to 10 min post-exercise, and  $\Delta$ La (difference between La at rest and at 10 min post-exercise) were calculated. As shown in Fig. 4, excess  $CO_2$  per body weight (excess  $CO_2/kg$ ) obtained in the intensive exercise tests was significantly related to  $\Delta$ La (r=0.88, p<0.01). On the other hand, the decreases in  $ET_{CO_2}$  (%) at 10 min post-exercise (differences between averaged  $ET_{CO_2}$  during 4–5 min at rest and 9–10 min post-exercise) in the 40-, 60- and 80-s exercise tests were  $-0.51\pm0.20$ ,  $-0.70\pm0.34$  and  $-0.57\pm0.29\%$ , respectively. These differences were not significant.

### DISCUSSION

In this study, 40-, 60- and 80-s exercises were performed in order to clarify the kinetics of excess CO<sub>2</sub> output during and after intensive exercise. The following results were obtained: (1) values of peak La ob-




Fig. 4. Relationship between excess CO<sub>2</sub> per body weight and increase in blood lactate concentration (difference between blood lactate concentrations at rest and at the 10th min after the end of exercise) in intensive exercises.

tained after the end of exercise increased linearly accompanying an increase in the duration of exercise; (2) excess  $\dot{V}\text{CO}_2$  showed a temporary negative value during exercise, then became positive, reaching a peak at 60 s post-exercise, and subsequently decreased down to zero at about 9 min post-exercise in the 80-s exercise test; and (3) excess  $\text{CO}_2$  per body weight (excess  $\text{CO}_2/\text{kg}$ ) increased in proportion to the increase in La ( $\Delta \text{La}$ ).

As lactic acid formed in the working muscles takes time to become uniformly distributed throughout the whole body, the peak value of La is observed at a few minutes after the end of exercise. Although this uniform value seems to indicate a difference between the amount effused out of working muscle and the amount diffused to other tissue and consumed, as the removal of lactic acid is relatively slow, peak La after the end of exercise is considered to reflect the magnitude of lactic acid produced in the working muscles. When these values were plotted as a function of duration of exercise (Fig. 3), it was inferred that lactic acid had been produced at a constant rate in the working muscles from the beginning of exercise.

As the lactic acid produced in working muscles is fully ionized, an equimolar amount of hydrogen ions (H<sup>+</sup>) is dissociated. This dissociated H<sup>+</sup> is then absorbed by the physicochemical action of muscle protein and the breakdown of creatine phosphate, and it is buffered by a leftward shift in the equation of the bicarbonate buffer system [17]:

$$CO_2 + H_2O \rightleftharpoons H^+ + HCO_3^-.$$
 (1)

However, if the CO<sub>2</sub> formed due to this buffering is

not fully eliminated,  $PCO_2$  increases. Similarly, if metabolically produced  $CO_2$  is not temporarily removed from the body,  $CO_2$  is stored in the body, resulting in an increase in  $PCO_2$ . Actually, it has been demonstrated that the femoral venous  $PCO_2$  increased dramatically with maximal cycling exercise [11, 18].

The increased tissue  $PCO_2$  induces an increase in  $P\bar{v}_{CO_2}$  and consequently an increase in  $CO_2$  transport from the tissue to the lung. In this case, if ventilation fitting for the  $CO_2$  transport is conducted and  $CO_2$  is eliminated, it is thought that arterial  $CO_2$  pressure  $(Pa_{CO_2})$  does not change. Based on the results of the kinetics of  $\dot{V} E / \dot{V} CO_2$  (Fig. 1), elimination of  $CO_2$  by the lung is considered to have been insufficient during exercise. As a result,  $ET_{CO_2}$  increased. These increases in  $P\bar{v}_{CO_2}$  and  $ET_{CO_2}$  induce a rightward shift in Eq. 1. Therefore, the decrease in  $HCO_3$  is masked, and consequently excess  $\dot{V}CO_2$  would have shown a temporary negative value.

After an initial increase,  $ET_{CO}$ , showed a decreasing phase. This is because, as shown in the kinetics of  $VE/VCO_2$ , the amount of  $CO_2$  eliminated by ventilation is larger than that transported to the lung. This decrease in ET<sub>CO</sub>, increases the venoarterial PCO<sub>2</sub> difference and could promote the elimination of CO2 from the tissues. This decrease in tissue PCO<sub>2</sub> simultaneously induces a decrease in venous PCO<sub>2</sub> In that case, Eq. 1 shifts leftward, and the excess  $\dot{V}$ CO<sub>2</sub> that had been masked during the period of increase in PCO<sub>2</sub> could appear during the period of decrease in PCO<sub>2</sub>. Thereafter,  $ET_{CO_2}$  fell below the pre-exercise level. This is thought to be due to excessive elimination of CO<sub>2</sub> by continuous hyperventilation. However, as the venous PCO<sub>2</sub> decreases accompanying the decline in CO<sub>2</sub> stores in the tissues, elimination of CO<sub>2</sub> from the tissues soon stops.

There are other factors that are related to the delay of excess CO<sub>2</sub> output from the production of lactic acid. The first is related to the resynthesis of creatine phosphate. The breakdown of creatine phosphate involves absorption of H<sup>+</sup> [17]. Therefore, the influence of the increase in H<sup>+</sup> accompanying the increase in lactic acid on the excess CO<sub>2</sub> output is inhibited during exercise. However, as the reverse reaction occurs after the end of exercise, H<sup>+</sup> increases even if lactic acid does not increase. Consequently, in Eq. 1, the HCO<sub>3</sub><sup>-</sup> decrement due to lactic acid production decreases during exercise and then the volume corresponding to the diminished amount could appear after the end of exercise.

Another factor is diffusion of lactic acid. The bicarbonate buffer system is more dominant in extracellular fluid than in intracellular fluid. If lactic acid in the

muscle flows into the extracellular fluid, the ratio of the amount of lactic acid buffered by the bicarbonate system becomes high. In this study, La increased by  $0.99\pm1.07\,\text{mM}$  from the end point to 5 min after the end of the 80-s intensive exercise. Furthermore, as it has been demonstrated that shifts of strong ions, including those of lactate, between compartments of body fluids induce an increase in blood H<sup>+</sup> concentration [11, 12, 18, 19–21], these factors could be related to the delay of excess  $CO_2$  output.

Excess  $CO_2$ , calculated as the total of excess  $VCO_2$ , was significantly related to  $\Delta La$ . As mentioned above, excess VCO<sub>2</sub> is affected by changes in H<sup>+</sup> accompanying the diffusion of lactic acid into the whole body and the reaction of creatine phosphate and change in PCO<sub>2</sub>. Therefore, these effects must be considered to understand the effect of only diffused lactic acid on excess CO2. However, as lactic acid is diffused into the whole body at about 5 min after the end of exercise, the excess CO<sub>2</sub> obtained is thought to reflect the effect of lactic acid equipoising to the whole body. As creatine phosphate decreases during exercise and recovers at 2-6 min after the end of exercise [22, 23], the effect of creatine phosphate on the excess CO<sub>2</sub> could be counteracted. On the other hand, although  $ET_{CO_2}$  increases during exercise and recovers after the end of exercise, the recovery is below the resting level. Therefore, excess CO2 obtained in the present study could include the amount affected by the change in  $PCO_2$ .

These results suggest that although excessive  $CO_2$  output (excess  $CO_2$ ) in response to intensive exercise is related to the increase in lactic acid, the time course of the excessive  $CO_2$  output (excess  $\dot{V}CO_2$ ) is delayed, relative to the production of lactic acid, and is affected by hyperventilation.

This study was supported by a Grant-in-Aid for Scientific Research (B) from the Ministry of Education, Science, Sports and Culture of Japan (No. 08458013).

# REFERENCES

- Hughson RL and Inman MD: Gas exchange analysis of immediate CO₂ storage at onset of exercise. Respir Physiol 59: 265–278, 1985
- 2. Linnarsson D: Dynamics of pulmonary gas exchange and heart rate changes at start and end of exercise. Acta Physiol Scand 415 (Suppl): 1–68, 1974
- Cherniack NS, Longobardo GS, Staw I, and Heymann M: Dynamics of carbon dioxide stores changes following an alteration in ventilation. J Appl Physiol 21: 785-793, 1966
- Farhi LE: Gas stores of the body. In: Handbook of Physiology, Sect 3, Respiration, ed. Fenn WO and Rahn H, Am Physiol Soc, Washington DC, Vol 1, pp

- 873-885, 1964
- 5. Jones NL and Jurkowski JE: Body carbon dioxide storage capacity in exercise. J Appl Physiol: Respir Environ Exerc Physiol 46: 811–815, 1979
- Edwards RHT, Denison DM, Jones G, Davies CTM, and Campbell EJM: Changes in mixed venous gas tensions at start of exercise in man. J Appl Physiol 32: 165–169, 1972
- Casaburi R, Daly J, Hansen JE, and Effros RM: Abrupt changes in mixed venous blood gas composition after the onset of exercise. J Appl Physiol 67: 1106–1112, 1989
- 8. Yano T: Immediate CO<sub>2</sub> storage capacity at the onset of exercise. Jpn J Physiol 36: 1241–1252, 1986
- Casaburi R, Barstow TJ, Robinson T, and Wasserman K: Influence of work rate on ventilatory and gas exchange kinetics. J Appl Physiol 67: 547–555, 1989
- Stringer W, Casaburi R, and Wasserman K: Acid-base regulation during exercise and recovery in humans. J Appl Physiol 72: 954–961, 1992
- Kowalchuk JM, Heigenhauser GJF, Lindinger MI, Sutton JR, and Jones NL: Factors influencing hydrogen ion concentration in muscle after intense exercise. J Appl Physiol 65: 2080–2089, 1988
- Kowalchuk JM, Heigenhauser GJF, Lindinger MI, Obminski G, Sutton JR, and Jones NL: Role of lungs and inactive muscle in acid-base control after maximal exercise. J Appl Physiol 65: 2090–2096, 1988
- McKenna MJ, Heigenhauser GJF, McKelvie RS, Mac-Dougall JD, and Jones NL: Sprint training enhances ionic regulation during intense exercise in men. J Physiol (Lond) 501: 687–702, 1997
- Osnes JB and Hermansen L: Acid-base balance after maximal exercise of short duration. J Appl Physiol 32: 59–63, 1972
- 15. Cerretelli P and Di Prampero PE: Gas exchange in exercise. In: Handbook of Physiology, Sect 3, The Respiratory System, ed. Farhi LE and Tenney SM, Am Phys-

- iol Soc, Washington DC, Vol 4, pp 297-339, 1987
- Bar-or O: The Wingate anaerobic test: an update on methodology, reliability and validity. Sports Med 4: 381–394, 1987
- Hultman E and Sahlin K: Acid-base balance during exercise. *In*: Exercise and Sport Science Reviews, ed.
  Hutton RS and Miller DI, The Franklin Institute, Philadelphia, Vol 8, pp 41–128, 1980
- Lindinger MI, Heigenhauser GJF, McKelvie RS, and Jones NL: Blood ion regulation during repeated maximal exercise and recovery in humans. Am J Physiol 262: R126–R136, 1992
- Jones NL: [H<sup>+</sup>] control in exercise: concepts and controversies. *In*: International Series on Sport Sciences, Vol 21, Biochemistry of Exercise VII, ed. Taylor AW, Gollnick PD, Green HJ, Ianuzzo CD, Noble EG, Métivier G, and Sutton JR, Human Kinetics Publishers, Inc, Champaign, IL, pp 333–339, 1990
- Heigenhauser GJF, Jones NL, Kowalchuk JM, and Lindinger MI: The role of the physicochemical systems in plasma in acid–base control in exercise. *In*: International Series on Sport Sciences, Vol 21, Biochemistry of Exercise VII, ed. Taylor AW, Gollnick PD, Green HJ, Ianuzzo CD, Noble EG, Métivier G, and Sutton JR, Human Kinetics Publishers, Inc, Champaign, IL, pp 359–374, 1990
- Kowalchuk JM, Heigenhauser GJF, Sutton JR, and Jones NL: Effect of acetazolamide on gas exchange and acid-base control after maximal exercise. J Appl Physiol 72: 278–287, 1992
- Piiper J and Spiller P: Repayment of O₂ debt and resynthesis of high-energy phosphates in gastrocnemius muscle of the dog. J Appl Physiol 28: 657–662, 1970
- Cooke SR, Petersen SR, and Quinney HA: The influence of maximal aerobic power on recovery of skeletal muscle following anaerobic exercise. Eur J Appl Physiol 75: 512–519, 1997