Canagliflozin PBPK/PD model: fit PD

1 Methods: Canagliflozin PBPK/PD model: fit PD

1.1 Parameters and data

4 parameters were optimized: EC50 reduction in RTG (`KI__RTG_E50`), baseline RTG (`KI__RTG_base`), FPG effect on RTG (`KI__RTG_m_fpg`) and maximal relative reduction of the RTG (`KI__RTG_max_inhibition`). The parameters of the adopted fit PK_20261019_103422__858e3 were fixed at their fitted values.

A fit mapping is a curve of a study, the data and the observable of the model it is compared with. The fit has 67 fit mappings of training data with 227 data points, whose residuals enter the objective; and 36 fit mappings of validation data with 174 data points, which were simulated and evaluated with the fitted parameters but not fitted. The fit was set up and run with sbmlsim [1].

1.2 Objective function

The parameters were estimated by minimizing the objective function

$$F(\vec{p}) = \frac{1}{2} \sum_k \sum_i \rho\big(w_{i,k} r_{i,k}(\vec{p})^2\big) \tag{1}$$

where k runs over the fit mappings of the training data and i over their data points, $r_{i,k}$ is the residual, $w_{i,k} = w_k w^p_{i,k}$ the weight of the data point and ρ the loss function, applied to the squared weighted residual $z = w_{i,k} r^2_{i,k}$. The loss function is linear, $\rho(z) = z$, i.e. a standard least squares problem.

The residuals are normalized by the mean $\overline{y}_k$ of the data of their curve, which makes curves of different quantities and magnitudes comparable:

$$r_{i,k} = \frac{y_{i,k} - f_{k(x_{i,k},\vec{p})}}{\overline{y}_k} \tag{2}$$

The weight of curve k is $w_k = \frac{n_k}{N_k}$, the weight n_k of its fit mapping (the weight its fit mapping collection assigns, 1 by default, or, for a collection which uses the weights of its mappings, the weight of the mapping, by default the number of subjects of its data) divided by the number N_k of its data points.

A data point is weighted with the inverse of its coefficient of variation, $w^p_{i,k} = |y_{i,k} \frac{|}{\sigma_{i,k}}$, with $\sigma_{i,k}$ the standard deviation of the data point, or its standard error for data without a standard deviation. An error of zero is missing, a missing error of a curve with errors is replaced by the largest error of the curve, and the data points of a curve without errors have the weight $w^p_{i,k} = 2$, the weight of a coefficient of variation of 0.5.

1.3 Optimization

The optimizer searched the parameters in log10 space within their bounds; bounds, values and intervals are given in the units of the fit parameters, the units of the table of the parameters.

The objective was minimized in 3 runs with differential evolution [2] of SciPy [3] within the bounds of the parameters, with the strategy `best1bin`, a population of 15 times the number of parameters, at most 1000 generations, the relative tolerance 0.01 of convergence, the recombination constant 0.7 and a mutation dithered between 0.5 and 1. The initial population of a run was drawn by latin hypercube sampling [4], with a seed of every run derived from the seed 1234 of the fit. The best member of the final population was polished with the L-BFGS-B method of SciPy.

The convergence of the runs was assessed from the traces of their cost and the waterfall plot of their final costs, and the parameter set with the lowest cost was used.

The models were simulated with libRoadRunner [5] and its integrator CVODE [6] with the relative tolerance 10^{-6} and the absolute tolerances 10^{-12} of an amount, 10^{-12} of a concentration and 10^{-6} of the other states, which for a species in a concentration is multiplied by the initial volume of its compartment. It ran with the variable step size of libRoadRunner, i.e. with an output at every step of the integrator. A data point at the time of a change after the start of a simulation, e.g. a dose or the reset of a collection, was compared with the state before the change, except the first data point of a curve with data at later times, which starts a collection and was compared with the state after the change.

1.4 Goodness of fit

The goodness of fit is given per kind of data (training data and validation data) by metrics over its n data points with the residuals $e_i = y_i - f\left(x_i, \hat{\vec{p}}\right)$ at the fitted parameters $\hat{\vec{p}}$, the mean squared error, its root and the root mean squared error normalized by the mean $\overline{y}_{k(i)}$ of the data of the curve of a point:

$$\mathrm{MSE} = \frac{1}{n}\sum_{i=1}^{n} e_i^2, \quad \mathrm{RMSE} = \sqrt{\mathrm{MSE}}, \quad \mathrm{NRMSE} = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{e_i}{\overline{y}_{k(i)}}\right)^2} \tag{3}$$

The root mean square of the residuals of the objective, i.e. what the fit minimizes, and the coefficient of determination:

$$\mathrm{RMSE}_w = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\rho(w_i r_i^2)}, \quad R^2 = 1 - \frac{\sum_i e_i^2}{\sum_i \left(y_i - \overline{y}\right)^2} \tag{4}$$

The predictions of a non-linear model are no linear regression of the data, so R^2 can be negative: the prediction is worse than the mean of the data. The information criteria of Akaike [7] and Schwarz [8] with the number n_p of fitted parameters are those of a least squares fit with normally distributed residuals, up to an additive constant, so only differences between models fitted to the same data are meaningful:

$$\mathrm{AIC} = n\ln(\mathrm{MSE}) + 2n_p, \quad \mathrm{BIC} = n\ln(\mathrm{MSE}) + n_p\ln(n) \tag{5}$$

The percentage of the data points whose prediction $f_i = f\left(x_i, \hat{\vec{p}}\right)$ lies within twofold of the data is taken over the n_+ data points with positive data $y_i > 0$ and a positive prediction $f_i > 0$; the others have no ratio and are not counted, and a kind of data without such a point has no percentage:

$$P_2 = \frac{100}{n_+}\left|\left\{i : \frac{1}{2} \le \frac{f_i}{y_i} \le 2\right\}\right| \tag{6}$$

The goodness of fit plot shows the predictions against the data on logarithmic axes, with the lines of twofold $f = 2y$ and $f = \frac{y}{2}$. The Bland-Altman plot [9] shows the ratio of prediction and data as $d_i = \log_{10}\left(\frac{f_i}{y_i}\right)$ against the geometric mean $\sqrt{f_i y_i}$ for the data points with positive values, with the bias $\overline{d}$ and the limits of agreement $\overline{d} \pm 1.96 s_d$, with the sample standard deviation s_d of d_i, of the training data; they are read as the fold factors $10^{\overline{d}}$ and $10^{\overline{d} \pm 1.96 s_d}$.

1.5 Identifiability

The identifiability of the parameters was analysed with the profile likelihood [10], [11]. The profile of a parameter is the lowest objective with the parameter fixed:

$$\mathrm{PL}(p_j) = \min_{p_{l \ne j}} F(\vec{p}) \tag{7}$$

At every point of a scan the other parameters were re-optimized from the previous point with the least squares optimizer of SciPy with the relative step 0.05. A scan ran from the optimum in both directions in the space of the optimizer, with steps in decades of the parameter, starting at 0.1. A step which raised the objective by more than 0.2 of the distance between the objective at the optimum and the threshold was reduced by the factor 2 and repeated, down to 0.01, and the step after one which raised it by less than that divided by the factor was enlarged by the factor, up to 1. A scan stopped at the threshold, at the bound of the parameter, at a point which failed or after 50 points.

The confidence intervals are the values at which the profile crosses the threshold of the likelihood ratio test

$$F_{\mathrm{thr}} = F_{\mathrm{min}} + \chi^2_\alpha \frac{\mathrm{df}}{2} \tag{8}$$

with F_{min} the lowest objective of the fit and of all profiles and $\chi^2_\alpha(\mathrm{df})$ the quantile of the χ^2 distribution at the confidence level $\alpha = 0.95$ with df = 1 degrees of freedom, i.e. 1.92 above the lowest objective. The threshold has this confidence level if F is the negative log-likelihood of the data up to a constant, i.e. if the weighted residuals are independent and normally distributed with unit variance; otherwise the level is approximate. A crossing is interpolated between two points of the profile, on the logarithm of a parameter on a logarithmic scale. A parameter is identifiable if its profile reaches the threshold within the scan on both sides of the optimum, i.e. its confidence interval is finite; practically non-identifiable towards the lower bound if its profile reaches the threshold above the optimum but not below it within the scan, which ends at the lower bound or after 50 points; practically non-

identifiable towards the upper bound if its profile reaches the threshold below the optimum but not above it within the scan, which ends at the upper bound or after 50 points; practically non-identifiable if its profile reaches the threshold on neither side within the scans, which end at the bounds or after 50 points, and is not flat; and structurally non-identifiable if its profile is flat on both sides. A side of a profile is flat if it rises by less than 0.05 of the distance between the lowest objective and the threshold, and a profile with a point which failed or did not converge is not classified. The profile likelihood is preferred over the intervals of the Fisher information, its intervals are invariant under a transformation of the parameters and can be asymmetric [12].

The Fisher information is the local alternative to the profiles: it is the Gauss-Newton approximation of the curvature of the objective at the optimum, from the Jacobian J of the weighted residuals of the training data with respect to the parameters $\vec{p}$ in the space of the optimizer, computed by central finite differences with the step $h_j = \varepsilon \max(|p_j|, 1),\ \varepsilon = 10^{-6}$ of the parameter p_j:

$$I = J^\top J, \quad \mathrm{cov} = \sigma^2 I^{-1}, \quad \sigma^2 = \frac{2F}{n - n_p} \tag{9}$$

with the number n of data points of the training data and the number n_p of parameters. The information has rank 4 of 4, the number of its eigenvalues above 10^{-8} of the largest one, and the condition number 377; the covariance of an information of a lower rank is computed with the pseudo-inverse. The Wald intervals are $\hat{p}_j \pm t_{\frac{1+\alpha}{2}, n-n_p} \mathrm{SE}_j$ with the standard errors $\mathrm{SE}_j = \sqrt{\mathrm{cov}_{jj}}$ at the confidence level $\alpha = 0.95$, transformed back into the units of the fit parameters.

1.6 Provenance

The fit PD_20261019_201254__61546 was run on 2026-10-19 18:13:02 UTC with sbmlsim 0.10.0 [1], libRoadRunner 2.10.0 [5], SciPy 1.18.1 [3], NumPy 2.5.3 [13] and Python 3.14.4 (seed 1234). The models and data were those of commit `e2644f489b973f8d28b845906e093e3af1870fe1` of `/home/mkoenig/git/pkdb_models-canagliflozin` their SHA-256 checksums are listed in Table 1. The fit kept the values `ftissue_can` = 1.08931, `Kp_can` = 0.685123, `ftissue_deep_can` = 0.00663007, `Kp_deep_can` = 9.54277, `GU__CANABS_k` = 0.0152774, `GU__F_can_abs` = 0.66566, `LI__CAN2M5_k` = 0.0254215, `LI__CAN2M7_k` = 0.0351472, `LI__CAN2M9_k` = 0.016767, `LI__M5BIEX_k` = 0.0281205, `LI__M7BIEX_k` = 0.0569284, `GU__GLUHYD_k` = 0.00697926, `KI__CANEX_f` = 1.27294, `KI__M5EX_f` = 0.551072 and `KI__M7EX_f` = 0.801255 of the model fixed.

Table 1: SHA-256 checksums of the models and data of the fit PD_20261019_201254__61546, their first 12 characters; the full checksums are in the provenance of the result (optimization_result.json).

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Chen2015/.Chen2015_Fig1.tsv	11309517d87d...
pkdb_models/models/canagliflozin/data/canagliflozin/Chen2015/.Chen2015_Fig2.tsv	ac40152c7c3e...
pkdb_models/models/canagliflozin/data/canagliflozin/Chen2015/.Chen2015_Tab2.tsv	dbf8674d6ca0...
pkdb_models/models/canagliflozin/data/canagliflozin/Chen2015/.Chen2015_Tab2A.tsv	0d692d95ddf4...
pkdb_models/models/canagliflozin/data/canagliflozin/Chen2015/.Chen2015_Tab3.tsv	474fc25c69c2...
pkdb_models/models/canagliflozin/data/canagliflozin/ Chen2015/.Chen2015_TabGroups.tsv	7df0b6e2b74b...
pkdb_models/models/canagliflozin/data/canagliflozin/ Chen2015/.Chen2015_TabInterventions.tsv	caa47749503b...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Fig1.tsv	99cbb98d385c...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Fig2.tsv	17ddd3ee74e9...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Fig3.tsv	a9788d138f8e...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Tab2.tsv	de641741be66...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Tab3.tsv	003083a994f7...
pkdb_models/models/canagliflozin/data/canagliflozin/ Devineni2012/.Devineni2012_Tab3A.tsv	4ab4230cd24c...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2012/.Devineni2012_TabGroups.tsv	fcbd9926a0bc...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2012/.Devineni2012_TabInterventions.tsv	cc994c72c23e...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Fig1.tsv	d3151c60e782...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Fig2.tsv	d54b41c9d9f3...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Tab2.tsv	a6e88864f675...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Tab2A.tsv	3f08200106ae...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Tab3.tsv	f10df8a57a2c...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_Tab3A.tsv	9877e8bf611b...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_TabGroups.tsv	5ca1502e81fc...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2013/.Devineni2013_TabInterventions.tsv	cb41feee3128...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_Fig2.tsv	04b9e40bb1dc...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_Fig3.tsv	ba3c236103c1...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_Tab1.tsv	096cbdc6db9e...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_Tab2.tsv	c5d09da767d6...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_Tab2A.tsv	c6dda93b2bbf...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_TabGroups.tsv	7b29e1f269b4...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2014/.Devineni2014_TabInterventions.tsv	e0063bf738d2...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig2.tsv	c1a175827351...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig3.tsv	088ab9028590...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig4.tsv	b424a9f15e15...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig5.tsv	ddfbd5c57f48...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig6.tsv	5976a606bf1f...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Fig7.tsv	188d2acb983d...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Tab1.tsv	071331a4faa2...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Tab2A.tsv	f4135efe551d...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Tab2B.tsv	292e0ef0a05b...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_Tab3.tsv	1f36409e81fd...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_TabGroups.tsv	f83495e58cf1...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015/.Devineni2015_TabInterventions.tsv	78abd7831e94...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_Fig1.tsv	3d3fbde8968b...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_Tab1.tsv	ec81b99e0ca6...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_Tab1A.tsv	6a4903dd4e4f...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_Tab2.tsv	04b43205a186...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_Tab2A.tsv	844240dc5dc6...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_TabGroups.tsv	4f4c51134a10...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015a/.Devineni2015a_TabInterventions.tsv	4331918feabe...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_Fig2.tsv	8635c6b62220...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_Tab1.tsv	3293450f44b2...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_Tab1A.tsv	e157aa24d566...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_Tab2.tsv	8fd0702a9085...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_TabGroups.tsv	2c50e3e696b0...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015b/.Devineni2015b_TabInterventions.tsv	f13ca85bf9c7...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Fig2.tsv	306cecb5e301...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Fig3.tsv	afda21ad9861...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Fig4.tsv	f008c659cdb0...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Fig5.tsv	5401fabf3340...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab2.tsv	b9496b829298...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab2A.tsv	c5f0a0b8b56c...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab3.tsv	023b75ee6d50...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab3A.tsv	c6090bb491e7...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab5.tsv	a119dc77b0c6...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_Tab5A.tsv	1b57262d6247...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_TabGroups.tsv	4beb338545d4...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015c/.Devineni2015c_TabInterventions.tsv	f37f547ca7ba...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Fig2.tsv	bcda389be253...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Fig3.tsv	e5050775f881...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Fig4.tsv	b757d7034480...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Tab2.tsv	4b7347040170...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Tab2A.tsv	a0342074e785...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Tab3.tsv	71c08387ea36...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Tab3A.tsv	496033a5bd0b...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_Tab4.tsv	dc77a141fa4a...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_TabGroups.tsv	a4556016f856...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015d/.Devineni2015d_TabInterventions.tsv	3b735ad4f8cf...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_Fig1.tsv	1baed35b0680...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_Fig2.tsv	5ad4836b7e41...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_Tab1.tsv	f93e791bfcd7...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_Tab2.tsv	cb889fba805d...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_Tab3.tsv	96ff9591e565...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_TabGroups.tsv	57bd46aaa7a8...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2015e/.Devineni2015e_TabInterventions.tsv	8ece422a882b...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_Fig2.tsv	84245d2ba2bf...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_Fig3.tsv	b9328fee4365...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_Tab2.tsv	5ea0640764b6...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_Tab3.tsv	e02222b8d262...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_Tab3A.tsv	98c55b823920...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_TabGroups.tsv	2dee9dc252e2...
pkdb_models/models/canagliflozin/data/canagliflozin/Devineni2016/.Devineni2016_TabInterventions.tsv	ffda46d473f9...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Fig2.tsv	91ee6b63d891...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Fig3.tsv	8333b049b72c...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Fig4.tsv	a64350254d61...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Fig5.tsv	da49dae531f0...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Fig6.tsv	f85056a3fc84...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Tab2.tsv	6c281cbfdb3b...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_Tab2A.tsv	c606b0fb2fa3...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_TabGroups.tsv	c0595c0aae4f...
pkdb_models/models/canagliflozin/data/canagliflozin/Iijima2015/.Iijima2015_TabInterventions.tsv	9cb67d9d726c...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Fig1.tsv	e5060ccd6762...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Fig2.tsv	88ac0a7f691b...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Fig4.tsv	d26a2084dfda...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Tab3.tsv	d1cbc8c27327...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Tab3A.tsv	23f0511c9c1a...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_Tab4.tsv	1e0705ae5817...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_TabGroups.tsv	1bf0d899848b...
pkdb_models/models/canagliflozin/data/canagliflozin/Inagaki2014/.Inagaki2014_TabInterventions.tsv	89b6e7ded71c...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Fig2.tsv	0131c0f0b1ac...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Fig3.tsv	483cf8f95e92...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Fig4.tsv	bb43f00e42d6...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Tab1.tsv	0dfa5cb6b43f...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Tab2.tsv	442a8708b838...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_Tab3.tsv	d410ffe2a7ff...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_TabGroups.tsv	d345e2dc019f...
pkdb_models/models/canagliflozin/data/canagliflozin/Kinoshita2015/.Kinoshita2015_TabInterventions.tsv	fb9adf4562f1...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Fig3.tsv	cd8a21860345...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab1.tsv	d24d6386dba1...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab1A.tsv	6c0db0ec5eae...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab2.tsv	2fb24299329d...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab3.tsv	95f5dc03d10b...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab4.tsv	aa5a15782060...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab5.tsv	aab8440e3e6e...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab5A.tsv	6bd2a5f313cf...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab6.tsv	1c8f1853c13b...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_Tab6A.tsv	21d10f2b200f...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_TabGroups.tsv	d6a0315899cd...
pkdb_models/models/canagliflozin/data/canagliflozin/Mamidi2014/.Mamidi2014_TabInterventions.tsv	1df63c1b445c...
pkdb_models/models/canagliflozin/data/canagliflozin/Mohamed2019/.Mohamed2019_Fig4.tsv	dab10af787e0...
pkdb_models/models/canagliflozin/data/canagliflozin/Mohamed2019/.Mohamed2019_Tab5.tsv	85e7d382c1ed...
pkdb_models/models/canagliflozin/data/canagliflozin/Mohamed2019/.Mohamed2019_TabGroups.tsv	ed0881ef74b8...
pkdb_models/models/canagliflozin/data/canagliflozin/Mohamed2019/.Mohamed2019_TabInterventions.tsv	41528e011971...
pkdb_models/models/canagliflozin/data/canagliflozin/Murphy2015/.Murphy2015_Fig1.tsv	3cc61caa2896...
pkdb_models/models/canagliflozin/data/canagliflozin/Murphy2015/.Murphy2015_Tab1.tsv	ecaea8f10002...
pkdb_models/models/canagliflozin/data/canagliflozin/Murphy2015/.Murphy2015_Tab2.tsv	5fabd2ff3788...
pkdb_models/models/canagliflozin/data/canagliflozin/Murphy2015/.Murphy2015_TabGroups.tsv	675eac9484fa...
pkdb_models/models/canagliflozin/data/canagliflozin/Murphy2015/.Murphy2015_TabInterventions.tsv	f93a05f0c73c...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2011/.Sha2011_Fig1.tsv	b42af877b45a...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2011/.Sha2011_Fig2.tsv	3bcd3329fb92...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2011/.Sha2011_TabGroups.tsv	e408392ed910...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2011/.Sha2011_TabInterventions.tsv	984b740a83a7...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Fig2.tsv	d3e08cc01af2...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Fig4.tsv	b4933da0e2d6...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Fig5.tsv	637494be2da3...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Fig6.tsv	2fcf7d202af6...

File	SHA-256
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Fig7.tsv	5b7ac8dec783...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Tab2.tsv	6b6e3a4b6df8...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Tab2A.tsv	ddd07c215259...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_Tab3.tsv	60ed6fab26bc...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_TabGroups.tsv	93df6598e18b...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2014/.Sha2014_TabInterventions.tsv	bde1cbe4c5f4...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2015/.Sha2015_Fig2.tsv	f709539e223d...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2015/.Sha2015_Fig3.tsv	76562e87e8de...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2015/.Sha2015_TabGroups.tsv	1c17a90fc63e...
pkdb_models/models/canagliflozin/data/canagliflozin/Sha2015/.Sha2015_TabInterventions.tsv	866a70d1f6a7...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_Fig1.tsv	ab52656d80c1...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_Fig3.tsv	83dbb1d0aa58...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_Fig4.tsv	2faf49857570...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_Fig5.tsv	2d0425030e0b...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_Tab2.tsv	ed117f76c761...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_TabGroups.tsv	ffe3ccbea06e...
pkdb_models/models/canagliflozin/data/canagliflozin/Tamborlane2018/.Tamborlane2018_TabInterventions.tsv	34993d1baf72...
pkdb_models/models/canagliflozin/data/canagliflozin/Wattamwar2020/.Wattamwar2020_Fig4.tsv	d42dadbff52b...
pkdb_models/models/canagliflozin/data/canagliflozin/Wattamwar2020/.Wattamwar2020_Tab8.tsv	a5c6664ba403...
pkdb_models/models/canagliflozin/data/canagliflozin/Wattamwar2020/.Wattamwar2020_TabGroups.tsv	ee43c63765d3...
pkdb_models/models/canagliflozin/data/canagliflozin/Wattamwar2020/.Wattamwar2020_TabInterventions.tsv	da66713c7ae7...
pkdb_models/models/canagliflozin/models/output/models/canagliflozin_body_flat.xml	ed17cbd80655...

2 Results: Canagliflozin PBPK/PD model: fit PD

2.1 Parameters

Table 2: Parameters of the fit PD_20261019_201254__61546, the parameter set PD_20261019_201254__61546_0, cost 169, with their bounds in the optimization. The 95% confidence intervals are those of the profile likelihood, LB and UB an interval which is open towards the lower or the upper bound, and the Wald intervals of the Fisher information for a parameter without a profile. Identifiable is the classification of the profile: yes, no (LB) or no (UB) if the profile does not reach the threshold towards the lower or the upper bound, no if it reaches it on neither side, structural if it is flat.

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
KI__RTG_E50	EC50 reduction in RTG	7.59e-05 [5.86e-05, 9.61e-05]	mmol/l	[1e-06, 0.0002]	yes
KI__RTG_base	baseline RTG	8.93 [8.46, 9.46]	mmol/l	[6, 13]	yes
KI__RTG_m_fpg	FPG effect on RTG	0.667 [0.604, 0.749]	dimensionless	[0.05, 2]	yes
KI__RTG_max_inhibition	maximal relative reduction of the RTG	0.645 [0.624, 0.664]	dimensionless	[0.3, 0.9]	yes

2.2 Data

Table 3: Data of the fit PD_20261019_201254__61546: the number of fit mappings and, in parentheses, of data points per study and kind. The training data enters the cost, the validation data and the outliers are evaluated only.

Study	Training	Validation	Outlier	Total
Chen2015	4 (6)	1 (2)	-	5 (8)
Devineni2012	5 (11)	1 (1)	-	6 (12)
Devineni2013	7 (24)	1 (4)	-	8 (28)
Devineni2014	2 (2)	2 (2)	-	4 (4)
Devineni2015a	6 (18)	-	-	6 (18)
Devineni2015c	2 (5)	6 (15)	-	8 (20)
Devineni2016	4 (20)	-	-	4 (20)
Iijima2015	5 (10)	4 (40)	-	9 (50)
Inagaki2014	2 (6)	2 (6)	-	4 (12)
Sha2011	14 (21)	3 (5)	-	17 (26)
Sha2014	14 (95)	10 (87)	-	24 (182)
Sha2015	2 (9)	-	-	2 (9)
Tamborlane2018	-	6 (12)	-	6 (12)
Total	67 (227)	36 (174)	-	103 (401)

2.3 Goodness of fit

Table 4: Goodness of fit of the fit PD_20261019_201254__61546 per kind of data, the parameter set PD_20261019_201254__61546_0. The cost is defined on the training data only, 2-fold is the percentage of the data points with a positive value and prediction whose prediction is within twofold of the data.

Kind	n	cost	MSE	RMSE	NRMSE	RMSE_w	R^2	2-fold [%]	AIC	BIC
training	227	169	404	20.1	0.322	1.22	0.852	95.1	1370.1	1383.8
validation	174	-	770	27.7	12.3	30.7	0.4	71.3	1164.5	1177.1

2.4 Figures




Figure 1: Cost of the optimizers over their steps, fit PD_20261019_201254__61546. The cost of every optimization run over its steps. Runs which end at the same cost found the same optimum; a run which stops much higher is stuck in a local minimum.




Figure 2: Cost of the optimization runs, ordered, fit PD_20261019_201254__61546. The final cost of the runs, ordered. A flat plateau at the left is the global optimum found repeatedly, which is the evidence that the multistart converged; steps to the right are local minima.




Figure 3: Prediction against the measured data points, per kind of fit mapping, with the agreement of the training data, the lines of twofold and the metrics of the panel, fit PD_20261019_201254__61546. Within twofold of the data are 193 of the 203 training data points (95.1%) and 112 of the 157 validation data points (71.3%); only the data points with a positive value and prediction are counted. Prediction against measurement, one point per data point, with a panel per kind of fit mapping. The points scatter around the diagonal when the model describes the data; a systematic deviation from it is a systematic error of the model. The band is the agreement of the training data, i.e. the same band the Bland-Altman plot draws, here as lines parallel to the diagonal. The dark red dotted lines are twofold, prediction = 2 x measurement and prediction = measurement / 2; the caption states the points between them of every kind. A point which is zero or negative is not drawn on the logarithmic axes and not counted, the title of a panel then states the points drawn of all. The validation panel shows how the fit describes data it was not fitted on, the outlier panel the data the fit dropped. The box of a panel carries its R², its normalized RMSE, the RMSE of the residuals of the cost and the percentage of its points within twofold.




Figure 4: Agreement of prediction and measurement as a ratio, per kind of fit mapping, with the agreement of the training data and the agreement of the panel, fit PD_20261019_201254__61546. The ratio of prediction and measurement over the geometric mean of the two, i.e. the goodness of fit with the diagonal turned into the horizontal. The band is the same, the bias and the limits of agreement `bias ± 1.96 SD` of the training data as fold factors. A bias away from 1 is a systematic over- or underprediction, wide limits are a large scatter, and a trend over the mean is a model which describes the large or the small values better. The box of a panel carries the bias and the SD of its own points as fold factors and the share of them inside the limits of agreement of the training data.




Figure 5: Profile likelihood of every parameter: the cost with the parameter fixed and the other parameters optimized, the threshold of the confidence level and the confidence interval, fit PD_20261019_201254__61546. One profile per parameter. A profile which crosses the dashed threshold on both sides gives a finite confidence interval; one which stays below it up to a bound is practically non-identifiable; a flat profile is structurally non-identifiable, the parameter is compensated by the others.

2.5 Provenance

Table 5: Provenance of the fit PD_20261019_201254__61546: the versions it ran with and the revision of its models and data; the SHA-256 checksums of the models and data are listed in Table 1.

Item	Value
Fit	`PD_20261019_201254__61546`
Created	2026-10-19 18:13:02 UTC
Seed	1234
Python	3.14.4
sbmlsim	0.10.0
libRoadRunner	2.10.0
SciPy	1.18.1
NumPy	2.5.3
Repository	`/home/mkoenig/git/pkdb_models-canagliflozin`
Commit	`e2644f489b973f8d28b845906e093e3af1870fe1`
Uncommitted changes	no

Bibliography

[1] M. König, "sbmlsim: SBML simulation made easy." Zenodo, 2026. doi: 10.5281/zenodo.3597770. Version 0.10.0.

[2] R. Storn and K. Price, "Differential evolution - a simple and efficient heuristic for global optimization over continuous spaces," *Journal of Global Optimization*, vol. 11, pp. 341–359, 1997, doi: 10.1023/A:1008202821328.

[3] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python," *Nature Methods*, vol. 17, pp. 261–272, 2020, doi: 10.1038/s41592-019-0686-2. Version 1.18.1.

[4] M. D. McKay, R. J. Beckman, and W. J. Conover, "A comparison of three methods for selecting values of input variables in the analysis of output from a computer code," *Technometrics*, vol. 21, no. 2, pp. 239–245, 1979, doi: 10.2307/1268522.

[5] C. Welsh, J. Xu, L. Smith, M. König, K. Choi, and H. M. Sauro, "libRoadRunner 2.0: a high performance SBML simulation and analysis library," *Bioinformatics*, vol. 39, no. 1, p. btac770, 2023, doi: 10.1093/bioinformatics/btac770. Version 2.10.0.

[6] A. C. Hindmarsh *et al.*, "SUNDIALS: suite of nonlinear and differential/algebraic equation solvers," *ACM Transactions on Mathematical Software*, vol. 31, no. 3, pp. 363–396, 2005, doi: 10.1145/1089014.1089020.

[7] H. Akaike, "A new look at the statistical model identification," *IEEE Transactions on Automatic Control*, vol. 19, no. 6, pp. 716–723, 1974, doi: 10.1109/TAC.1974.1100705.

[8] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978, doi: 10.1214/aos/1176344136.

[9] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 327, no. 8476, pp. 307–310, 1986, doi: 10.1016/S0140-6736(86)90837-8.

[10] A. Raue *et al.*, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, no. 15, pp. 1923–1929, 2009, doi: 10.1093/bioinformatics/btp358.

[11] C. Kreutz, A. Raue, D. Kaschek, and J. Timmer, "Profile likelihood in systems biology," *The FEBS Journal*, vol. 280, no. 11, pp. 2564–2571, 2013, doi: 10.1111/febs.12276.

[12] F.-G. Wieland, A. L. Hauber, M. Rosenblatt, C. Tönsing, and J. Timmer, "On structural and practical identifiability," *Current Opinion in Systems Biology*, vol. 25, pp. 60–69, 2021, doi: 10.1016/j.coisb.2021.03.005.

[13] C. R. Harris *et al.*, "Array programming with NumPy," *Nature*, vol. 585, pp. 357–362, 2020, doi: 10.1038/s41586-020-2649-2. Version 2.5.3.