Dapagliflozin PBPK/PD model: fit PD

1 Methods: Dapagliflozin PBPK/PD model: fit PD

1.1 Parameters and data

4 parameters were optimized: EC50 reduction in RTG (`KI__RTG_E50`), hill coefficient reduction in RTG (`KI__RTG_gamma`), FPG effect on RTG (`KI__RTG_m_fpg`) and maximal relative reduction of the RTG (`KI__RTG_max_inhibition`). The parameters of the adopted fit PK_20261020_002954__89bab were fixed at their fitted values.

A fit mapping is a curve of a study, the data and the observable of the model it is compared with. The fit has 31 fit mappings of training data with 254 data points, whose residuals enter the objective; and 15 fit mappings of validation data with 47 data points, which were simulated and evaluated with the fitted parameters but not fitted. The fit was set up and run with sbmlsim [1].

1.2 Objective function

The parameters were estimated by minimizing the objective function

$$F(\vec{p}) = \frac{1}{2} \sum_k \sum_i \rho\left(w_{i,k} r_{i,k}(\vec{p})^2\right) \tag{1}$$

where k runs over the fit mappings of the training data and i over their data points, $r_{i,k}$ is the residual, $w_{i,k} = w_k w^p_{i,k}$ the weight of the data point and ρ the loss function, applied to the squared weighted residual $z = w_{i,k} r^2_{i,k}$. The loss function is linear, $\rho(z) = z$, i.e. a standard least squares problem.

The residuals are normalized by the mean $\overline{y}_k$ of the data of their curve, which makes curves of different quantities and magnitudes comparable:

$$r_{i,k} = \frac{y_{i,k} - f_{k(x_{i,k},\vec{p})}}{\overline{y}_k} \tag{2}$$

The weight of curve k is $w_k = \frac{n_k}{N_k}$, the weight n_k of its fit mapping (the weight its fit mapping collection assigns, 1 by default, or, for a collection which uses the weights of its mappings, the weight of the mapping, by default the number of subjects of its data) divided by the number N_k of its data points.

A data point is weighted with the inverse of its coefficient of variation, $w^p_{i,k} = |y_{i,k} \frac{1}{\sigma_{i,k}}$, with $\sigma_{i,k}$ the standard deviation of the data point, or its standard error for data without a standard deviation. An error of zero is missing, a missing error of a curve with errors is replaced by the largest error of the curve, and the data points of a curve without errors have the weight $w^p_{i,k} = 2$, the weight of a coefficient of variation of 0.5.

1.3 Optimization

The optimizer searched the parameters in log10 space within their bounds; bounds, values and intervals are given in the units of the fit parameters, the units of the table of the parameters.

The objective was minimized in 3 runs with differential evolution [2] of SciPy [3] within the bounds of the parameters, with the strategy `best1bin`, a population of 15 times the number of parameters, at most 1000 generations, the relative tolerance 0.01 of convergence, the recombination constant 0.7 and a mutation dithered between 0.5 and 1. The initial population of a run was drawn by latin hypercube sampling [4], with a seed of every run derived from the seed 1234 of the fit. The best member of the final population was polished with the L-BFGS-B method of SciPy.

The convergence of the runs was assessed from the traces of their cost and the waterfall plot of their final costs, and the parameter set with the lowest cost was used.

The models were simulated with libRoadRunner [5] and its integrator CVODE [6] with the relative tolerance 10^{-6} and the absolute tolerances 10^{-12} of an amount, 10^{-12} of a concentration and 10^{-6} of the other states, which for a species in a concentration is multiplied by the initial volume of its compartment. It ran with the variable step size of libRoadRunner, i.e. with an output at every step of the integrator. A data point at the time of a change after the start of a simulation, e.g. a dose or the reset of a collection, was compared with the state before the change, except the first data point of a curve with data at later times, which starts a collection and was compared with the state after the change.

1.4 Goodness of fit

The goodness of fit is given per kind of data (training data and validation data) by metrics over its n data points with the residuals $e_i = y_i - f\left(x_i, \hat{\vec{p}}\right)$ at the fitted parameters $\hat{\vec{p}}$, the mean squared error, its root and the root mean squared error normalized by the mean $\overline{y}_{k(i)}$ of the data of the curve of a point:

$$\mathrm{MSE} = \frac{1}{n}\sum_{i=1}^{n} e_i^2, \quad \mathrm{RMSE} = \sqrt{\mathrm{MSE}}, \quad \mathrm{NRMSE} = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{e_i}{\overline{y}_{k(i)}}\right)^2} \tag{3}$$

The root mean square of the residuals of the objective, i.e. what the fit minimizes, and the coefficient of determination:

$$\mathrm{RMSE}_w = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\rho(w_i r_i^2)}, \quad R^2 = 1 - \frac{\sum_i e_i^2}{\sum_i \left(y_i - \overline{y}\right)^2} \tag{4}$$

The predictions of a non-linear model are no linear regression of the data, so R^2 can be negative: the prediction is worse than the mean of the data. The information criteria of Akaike [7] and Schwarz [8] with the number n_p of fitted parameters are those of a least squares fit with normally distributed residuals, up to an additive constant, so only differences between models fitted to the same data are meaningful:

$$\mathrm{AIC} = n\ln(\mathrm{MSE}) + 2n_p, \quad \mathrm{BIC} = n\ln(\mathrm{MSE}) + n_p\ln(n) \tag{5}$$

The percentage of the data points whose prediction $f_i = f\left(x_i, \hat{\vec{p}}\right)$ lies within twofold of the data is taken over the n_+ data points with positive data $y_i > 0$ and a positive prediction $f_i > 0$; the others have no ratio and are not counted, and a kind of data without such a point has no percentage:

$$P_2 = \frac{100}{n_+}\left|\left\{i : \frac{1}{2} \le \frac{f_i}{y_i} \le 2\right\}\right| \tag{6}$$

The goodness of fit plot shows the predictions against the data on logarithmic axes, with the lines of twofold $f = 2y$ and $f = \frac{y}{2}$. The Bland-Altman plot [9] shows the ratio of prediction and data as $d_i = \log_{10}\left(\frac{f_i}{y_i}\right)$ against the geometric mean $\sqrt{f_i y_i}$ for the data points with positive values, with the bias $\overline{d}$ and the limits of agreement $\overline{d} \pm 1.96 s_d$, with the sample standard deviation s_d of d_i, of the training data; they are read as the fold factors $10^{\overline{d}}$ and $10^{\overline{d} \pm 1.96 s_d}$.

1.5 Identifiability

The identifiability of the parameters was analysed with the profile likelihood [10], [11]. The profile of a parameter is the lowest objective with the parameter fixed:

$$\mathrm{PL}(p_j) = \min_{p_{l \ne j}} F(\vec{p}) \tag{7}$$

At every point of a scan the other parameters were re-optimized from the previous point with the least squares optimizer of SciPy with the relative step 0.05. A scan ran from the optimum in both directions in the space of the optimizer, with steps in decades of the parameter, starting at 0.1. A step which raised the objective by more than 0.2 of the distance between the objective at the optimum and the threshold was reduced by the factor 2 and repeated, down to 0.01, and the step after one which raised it by less than that divided by the factor was enlarged by the factor, up to 1. A scan stopped at the threshold, at the bound of the parameter, at a point which failed or after 50 points.

The confidence intervals are the values at which the profile crosses the threshold of the likelihood ratio test

$$F_{\mathrm{thr}} = F_{\mathrm{min}} + \chi^2_\alpha \frac{\mathrm{df}}{2} \tag{8}$$

with F_{min} the lowest objective of the fit and of all profiles and $\chi^2_\alpha(\mathrm{df})$ the quantile of the χ^2 distribution at the confidence level $\alpha = 0.95$ with df = 1 degrees of freedom, i.e. 1.92 above the lowest objective. The threshold has this confidence level if F is the negative log-likelihood of the data up to a constant, i.e. if the weighted residuals are independent and normally distributed with unit variance; otherwise the level is approximate. A crossing is interpolated between two points of the profile, on the logarithm of a parameter on a logarithmic scale. A parameter is identifiable if its profile reaches the threshold within the scan on both sides of the optimum, i.e. its confidence interval is finite; practically non-identifiable towards the lower bound if its profile reaches the threshold above the optimum but not below it within the scan, which ends at the lower bound or after 50 points; practically non-

identifiable towards the upper bound if its profile reaches the threshold below the optimum but not above it within the scan, which ends at the upper bound or after 50 points; practically non-identifiable if its profile reaches the threshold on neither side within the scans, which end at the bounds or after 50 points, and is not flat; and structurally non-identifiable if its profile is flat on both sides. A side of a profile is flat if it rises by less than 0.05 of the distance between the lowest objective and the threshold, and a profile with a point which failed or did not converge is not classified. The profile likelihood is preferred over the intervals of the Fisher information, its intervals are invariant under a transformation of the parameters and can be asymmetric [12].

The Fisher information is the local alternative to the profiles: it is the Gauss-Newton approximation of the curvature of the objective at the optimum, from the Jacobian J of the weighted residuals of the training data with respect to the parameters $\vec{p}$ in the space of the optimizer, computed by central finite differences with the step $h_j = \varepsilon \max(|p_j|, 1),\ \varepsilon = 10^{-6}$ of the parameter p_j:

$$I = J^{\top} J, \quad \mathrm{cov} = \sigma^2 I^{-1}, \quad \sigma^2 = \frac{2F}{n - n_p} \tag{9}$$

with the number n of data points of the training data and the number n_p of parameters. The information has rank 4 of 4, the number of its eigenvalues above 10^{-8} of the largest one, and the condition number 799; the covariance of an information of a lower rank is computed with the pseudo-inverse. The Wald intervals are $\hat{p}_j \pm t_{\frac{1+\alpha}{2}, n-n_p} \mathrm{SE}_j$ with the standard errors $\mathrm{SE}_j = \sqrt{\mathrm{cov}_{jj}}$ at the confidence level $\alpha = 0.95$, transformed back into the units of the fit parameters.

1.6 Provenance

The fit PD_20261020_014612__f2826 was run on 2026-10-19 23:46:14 UTC with sbmlsim 0.10.0 [1], libRoadRunner 2.10.0 [5], SciPy 1.18.1 [3], NumPy 2.5.3 [13] and Python 3.14.4 (seed 1234). The models and data were those of commit `266a01e9f9840563604534c348eeeae5ab4e3a46` of `/home/mkoenig/git/pkdb_models-dapagliflozin` their SHA-256 checksums are listed in Table 1. The fit kept the values `ftissue_dap` = 0.411373, `Kp_dap` = 1.20018, `GU__Ka_dis_dap` = 1.09176, `GU__DAPABS_k` = 0.0444919, `GU__FED_f_absorption` = 0.189289, `DAPMET_k` = 0.0658713, `KI__DAPEX_f` = 0.460206 and `KI__D3GEX_f` = 1.80546 of the model fixed.

Table 1: SHA-256 checksums of the models and data of the fit PD_20261020_014612__f2826, their first 12 characters; the full checksums are in the provenance of the result (optimization_result.json).

File	SHA-256
pkdb_models/models/dapagliflozin/data/canagliflozin/Sha2015/.Sha2015_Fig2.tsv	9888c59afb2b...
pkdb_models/models/dapagliflozin/data/canagliflozin/Sha2015/.Sha2015_Fig3.tsv	5058aa68df88...
pkdb_models/models/dapagliflozin/data/canagliflozin/Sha2015/.Sha2015_TabGroups.tsv	1c17a90fc63e...
pkdb_models/models/dapagliflozin/data/canagliflozin/Sha2015/.Sha2015_TabInterventions.tsv	866a70d1f6a7...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Boulton2013/.Boulton2013_Fig1.tsv	cc06471f1d18...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Boulton2013/.Boulton2013_Tab1.tsv	e10b748372b2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Boulton2013/.Boulton2013_TabGroups.tsv	baa2677dd9a0...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Boulton2013/.Boulton2013_TabInterventions.tsv	48ee8e7259f2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Cho2021/.Cho2021_Fig6.tsv	c9f1da565072...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Cho2021/.Cho2021_Tab5.tsv	5f956c0bd354...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Cho2021/.Cho2021_TabGroups.tsv	445946a31a8f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Cho2021/.Cho2021_TabInterventions.tsv	4691fd37ca19...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102002/.FDAMB102002_Fig12.tsv	8d98a3fae7b5...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102002/.FDAMB102002_Fig17.tsv	19868e2823ba...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102002/.FDAMB102002_Tab7.tsv	84d3d77c0d82...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102002/.FDAMB102002_TabGroups.tsv	e563991e0003...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102002/.FDAMB102002_TabInterventions.tsv	9efae56172a1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_Fig13.tsv	3c6bc562b764...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_Fig18.tsv	80954c4abc97...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_Tab12.tsv	4ffcf8e24586...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_Tab8.tsv	9ffc161ed251...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_Tab8A.tsv	2bcefc3176ac...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_TabGroups.tsv	510acbf2acf2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102003/.FDAMB102003_TabInterventions.tsv	a4843e12731a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102006/.FDAMB102006_Tab11.tsv	49e3c6565c1e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102006/.FDAMB102006_Tab11A.tsv	e77578783a53...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102006/.FDAMB102006_TabGroups.tsv	da4ae5a1af69...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102006/.FDAMB102006_TabInterventions.tsv	8f894f9264c1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_Fig3.tsv	fdcd54d1d32f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_Tab13.tsv	8d5ee0f0a8d9...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_Tab13A.tsv	b159e24becb1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_Tab14.tsv	45869ab42ce7...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_Tab14A.tsv	2bd4db1a6c9a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_TabGroups.tsv	02e13ad2de3f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/FDAMB102007/.FDAMB102007_TabInterventions.tsv	ba4ef3395beb...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_Fig1.tsv	d93ae8e1b647...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_Fig1a.tsv	32928f5f3415...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_Tab2.tsv	66064be308d4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_Tab3.tsv	b30cb721cae8...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_TabGroups.tsv	09ff53622f61...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_TabIndividuals.tsv	fbf7f132fbe3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Gould2013/.Gould2013_TabInterventions.tsv	029143b476b3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_Fig1.tsv	c53a3388d323...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_Fig2.tsv	6cfbefd16762...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_Fig4.tsv	488d6e524645...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_Tab2.tsv	3cf62d65a1e7...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_Tab3.tsv	ecbdbc0b2ec6...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_TabGroups.tsv	163e159de102...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_TabInterventions.tsv	0b60f83b6227...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Hwang2022a/.Hwang2022a_TabS2.tsv	0fa6cfc2ac58...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Imamura2013/.Imamura2013_Fig2.tsv	9918b3cea6cf...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Imamura2013/.Imamura2013_Tab2.tsv	2b5e6bba934f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Imamura2013/.Imamura2013_TabGroups.tsv	53304764f2ed...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Imamura2013/.Imamura2013_TabInterventions.tsv	8f05f9649ad7...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Jang2020/.Jang2020_Fig2.tsv	e15c3d846b03...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Jang2020/.Jang2020_Fig3.tsv	513acd3f936e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Jang2020/.Jang2020_Tab1.tsv	221d4fee7fe7...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Jang2020/.Jang2020_TabGroups.tsv	e0a5aae17ce1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Jang2020/.Jang2020_TabInterventions.tsv	814ff82fc5b1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011/.Kasichayanula2011_Fig1.tsv	c2fb3ae6b27a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011/.Kasichayanula2011_Fig2.tsv	0d782a7fc79f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011/.Kasichayanula2011_Tab2.tsv	3e01e5c21939...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011/.Kasichayanula2011_TabGroups.tsv	5a3c03b34db1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011/.Kasichayanula2011_TabInterventions.tsv	1364c9b89ca1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Fig2.tsv	aaed0eb4f448...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Fig3.tsv	5021152af0f7...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Fig4.tsv	3a331ea3638e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Fig5.tsv	6aa1c457948f...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Fig6.tsv	4179d88569fc...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Tab2.tsv	87ba4fae0e9e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_Tab2A.tsv	f7ddb2f91cdf...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_TabGroups.tsv	2ff07e921fec...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_TabIndividuals.tsv	f6a71c1fefe2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011a/.Kasichayanula2011a_TabInterventions.tsv	7fa2cf646d6e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011b/.Kasichayanula2011b_Fig1.tsv	82715bbd1639...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011b/.Kasichayanula2011b_Tab1.tsv	0061412006ad...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011b/.Kasichayanula2011b_TabGroups.tsv	ae27615b4f10...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011b/.Kasichayanula2011b_TabInterventions.tsv	1a96cfb372e9...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_Fig3.tsv	4bfd473b9826...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_Fig4.tsv	4ca9bb12bcde...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_Tab2.tsv	8de2215e981b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_Tab3.tsv	dc71602be922...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_TabGroups.tsv	6e18181424d4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2011c/.Kasichayanula2011c_TabInterventions.tsv	3c7697f1a970...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Fig2.tsv	46c04d6f66d0...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Fig3.tsv	bc3e9f9769ff...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Fig4.tsv	41cf946ad3d2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Tab1.tsv	319ebfd08432...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Tab2.tsv	15ad4982152d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_Tab3.tsv	2cf49f341ed4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_TabGroups.tsv	31142b4b2c9b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2012/.Kasichayanula2012_TabInterventions.tsv	958efe0f56cc...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Fig1.tsv	3a9f535f38e4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Fig2.tsv	6ec70b4384e8...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Tab2.tsv	ace08984344a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Tab2A.tsv	4bedeeed70b3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Tab4.tsv	6f5ec42b18ce...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_Tab4A.tsv	30d888b00017...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_TabGroups.tsv	228678add7c5...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_TabIndividuals.tsv	d70cb20af140...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013/.Kasichayanula2013_TabInterventions.tsv	289df1c46c8a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013a/.Kasichayanula2013a_Fig1.tsv	28e0585cb639...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013a/.Kasichayanula2013a_Tab1.tsv	55aa172ac000...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013a/.Kasichayanula2013a_Tab1A.tsv	99f03406f7fe...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013a/.Kasichayanula2013a_TabGroups.tsv	876c83b13b51...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kasichayanula2013a/.Kasichayanula2013a_TabInterventions.tsv	cb9f4725f8db...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Khomitskaya2018/.Khomitskaya2018_Fig2.tsv	9141b687145c...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Khomitskaya2018/.Khomitskaya2018_Tab2.tsv	eb18af4cf03e...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Khomitskaya2018/.Khomitskaya2018_TabGroups.tsv	39a6b7a2d1f1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Khomitskaya2018/.Khomitskaya2018_TabInterventions.tsv	eb07aa78649d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Fig1.tsv	faf8a67de950...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Fig2.tsv	7419518c5dc2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Tab2.tsv	e4b13d62892b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Tab3.tsv	ea1325423a66...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabGroups.tsv	e6e46d776a7a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabInterventions.tsv	00fa9049fea8...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabS2.tsv	aa635231e391...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_Fig2.tsv	37c777bb108d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_Fig3.tsv	fe3ab763b290...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_Tab1.tsv	375b50f8f0d6...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_Tab2.tsv	c9a0ac803166...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_TabGroups.tsv	ae4dbf307be3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Kim2023a/.Kim2023a_TabInterventions.tsv	a5e38ea82c77...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Fig3.tsv	7d55baedd71d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Fig4.tsv	b5d0cee0bb14...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Fig5.tsv	0dfa0f66b56a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Fig6.tsv	392a6f0e226b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Tab1.tsv	bb02e819a644...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Tab1A.tsv	a1bf6a459427...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Tab2.tsv	a17bfdd6c41d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_Tab6.tsv	a17bfdd6c41d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_TabGroups.tsv	9149127333d9...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Komoroski2009/.Komoroski2009_TabInterventions.tsv	1ee03a62d1a3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/LaCreta2016/.LaCreta2016_Fig2.tsv	f3eda6a27bd4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/LaCreta2016/.LaCreta2016_Tab2.tsv	49e1da53b16b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/LaCreta2016/.LaCreta2016_Tab3.tsv	d1bb4ac6905d...
pkdb_models/models/dapagliflozin/data/dapagliflozin/LaCreta2016/.LaCreta2016_TabGroups.tsv	afe1bf6b7615...
pkdb_models/models/dapagliflozin/data/dapagliflozin/LaCreta2016/.LaCreta2016_TabInterventions.tsv	57ff50e647a3...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Obermeier2010/.Obermeier2010_Fig6.tsv	fb6a24ac0629...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Obermeier2010/.Obermeier2010_Fig7.tsv	cf66e56ba7a4...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Obermeier2010/.Obermeier2010_Tab1.tsv	78bb4acc8c9b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Obermeier2010/.Obermeier2010_TabGroups.tsv	1be4244cd62a...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Obermeier2010/.Obermeier2010_TabInterventions.tsv	b3e812b9a452...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Shah2019a/.Shah2019a_Fig3.tsv	39ad37ab950b...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Shah2019a/.Shah2019a_Tab7.tsv	6ea3f122d095...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Shah2019a/.Shah2019a_TabGroups.tsv	a4a374c11800...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Shah2019a/.Shah2019a_TabInterventions.tsv	e078d756ce4f...

File	SHA-256
pkdb_models/models/dapagliflozin/data/dapagliflozin/Watada2019/.Watada2019_Fig1.tsv	17914be7adc2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Watada2019/.Watada2019_Tab2.tsv	d101ef8d2e69...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Watada2019/.Watada2019_Tab3.tsv	13496a64ebfd...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Watada2019/.Watada2019_TabGroups.tsv	c81302866175...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Watada2019/.Watada2019_TabInterventions.tsv	9978adbc0973...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_Fig.tsv	904258727851...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_Fig1.tsv	b061926dcce1...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_Tab2.tsv	f0ccf3ba01cc...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_Tab3.tsv	b620b4c92760...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_TabGroups.tsv	5783bc234e30...
pkdb_models/models/dapagliflozin/data/dapagliflozin/Yang2013/.Yang2013_TabInterventions.tsv	f422800044b2...
pkdb_models/models/dapagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_Fig3.tsv	03719e965463...
pkdb_models/models/dapagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabGroups.tsv	033515791883...
pkdb_models/models/dapagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabIndividuals.tsv	77b6cfffbdb8...
pkdb_models/models/dapagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabInterventions.tsv	9daa3d0a8674...
pkdb_models/models/dapagliflozin/models/output/models/dapagliflozin_body_flat.xml	32faeb4412ab...

2 Results: Dapagliflozin PBPK/PD model: fit PD

2.1 Parameters

Table 2: Parameters of the fit PD_20261020_014612__f2826, the parameter set PD_20261020_014612__f2826_0, cost 55.5, with their bounds in the optimization. The 95% confidence intervals are those of the profile likelihood, LB and UB an interval which is open towards the lower or the upper bound, and the Wald intervals of the Fisher information for a parameter without a profile. Identifiable is the classification of the profile: yes, no (LB) or no (UB) if the profile does not reach the threshold towards the lower or the upper bound, no if it reaches it on neither side, structural if it is flat.

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
KI__RTG_E50	EC50 reduction in RTG	5.76e-06 [4.88e-06, 6.83e-06]	mmol/l	[1e-07, 0.0001]	yes
KI__RTG_gamma	hill coefficient reduction in RTG	2.8 [1.68, UB]	dimensionless	[1, 4]	no (UB)
KI__RTG_m_fpg	FPG effect on RTG	1.99 [1.55, 2.44]	dimensionless	[0.1, 5]	yes
KI__RTG_max_inhibition	maximal relative reduction of the RTG	0.64 [0.618, 0.67]	dimensionless	[0.3, 0.95]	yes

2.2 Data

Table 3: Data of the fit PD_20261020_014612__f2826: the number of fit mappings and, in parentheses, of data points per study and kind. The training data enters the cost, the validation data and the outliers are evaluated only.

Study	Training	Validation	Outlier	Total
FDAMB102002	5 (60)	-	-	5 (60)
FDAMB102003	6 (36)	-	-	6 (36)
FDAMB102007	-	4 (8)	-	4 (8)
Gould2013	3 (6)	4 (8)	-	7 (14)
Kasichayanula2011a	3 (6)	1 (2)	-	4 (8)
Kim2023	1 (1)	1 (1)	-	2 (2)
Komoroski2009	8 (132)	1 (18)	-	9 (150)
Sha2015	1 (5)	1 (4)	-	2 (9)
Watada2019	-	3 (6)	-	3 (6)
Yang2013	4 (8)	-	-	4 (8)
Total	31 (254)	15 (47)	-	46 (301)

2.3 Goodness of fit

Table 4: Goodness of fit of the fit PD_20261020_014612__f2826 per kind of data, the parameter set PD_20261020_014612__f2826_0. The cost is defined on the training data only, 2-fold is the percentage of the data points with a positive value and prediction whose prediction is within twofold of the data.

Kind	n	cost	MSE	RMSE	NRMSE	RMSE_w	R^2	2-fold [%]	AIC	BIC
training	254	55.5	389	19.7	0.31	0.661	0.868	95.6	1522.6	1536.8
validation	47	-	153	12.4	1.16	3.06	0.805	34.3	244.3	251.7

2.4 Figures




Figure 1: Cost of the optimizers over their steps, fit PD_20261020_014612__f2826. The cost of every optimization run over its steps. Runs which end at the same cost found the same optimum; a run which stops much higher is stuck in a local minimum.




Figure 2: Cost of the optimization runs, ordered, fit PD_20261020_014612__f2826. The final cost of the runs, ordered. A flat plateau at the left is the global optimum found repeatedly, which is the evidence that the multistart converged; steps to the right are local minima.




Figure 3: Prediction against the measured data points, per kind of fit mapping, with the agreement of the training data, the lines of twofold and the metrics of the panel, fit PD_20261020_014612__f2826. Within twofold of the data are 237 of the 248 training data points (95.6%) and 12 of the 35 validation data points (34.3%); only the data points with a positive value and prediction are counted. Prediction against measurement, one point per data point, with a panel per kind of fit mapping. The points scatter around the diagonal when the model describes the data; a systematic deviation from it is a systematic error of the model. The band is the agreement of the training data, i.e. the same band the Bland-Altman plot draws, here as lines parallel to the diagonal. The dark red dotted lines are twofold, prediction = 2 x measurement and prediction = measurement / 2; the caption states the points between them of every kind. A point which is zero or negative is not drawn on the logarithmic axes and not counted, the title of a panel then states the points drawn of all. The validation panel shows how the fit describes data it was not fitted on, the outlier panel the data the fit dropped. The box of a panel carries its R², its normalized RMSE, the RMSE of the residuals of the cost and the percentage of its points within twofold.




Figure 4: Agreement of prediction and measurement as a ratio, per kind of fit mapping, with the agreement of the training data and the agreement of the panel, fit PD_20261020_014612__f2826. The ratio of prediction and measurement over the geometric mean of the two, i.e. the goodness of fit with the diagonal turned into the horizontal. The band is the same, the bias and the limits of agreement `bias ± 1.96 SD` of the training data as fold factors. A bias away from 1 is a systematic over- or underprediction, wide limits are a large scatter, and a trend over the mean is a model which describes the large or the small values better. The box of a panel carries the bias and the SD of its own points as fold factors and the share of them inside the limits of agreement of the training data.




Figure 5: Profile likelihood of every parameter: the cost with the parameter fixed and the other parameters optimized, the threshold of the confidence level and the confidence interval, fit PD_20261020_014612__f2826. One profile per parameter. A profile which crosses the dashed threshold on both sides gives a finite confidence interval; one which stays below it up to a bound is practically non-identifiable; a flat profile is structurally non-identifiable, the parameter is compensated by the others.

2.5 Provenance

Table 5: Provenance of the fit PD_20261020_014612__f2826: the versions it ran with and the revision of its models and data; the SHA-256 checksums of the models and data are listed in Table 1.

Item	Value
Fit	`PD_20261020_014612__f2826`
Created	2026-10-19 23:46:14 UTC
Seed	1234
Python	3.14.4
sbmlsim	0.10.0
libRoadRunner	2.10.0
SciPy	1.18.1
NumPy	2.5.3
Repository	`/home/mkoenig/git/pkdb_models-dapagliflozin`
Commit	`266a01e9f9840563604534c348eeeae5ab4e3a46`
Uncommitted changes	no

Bibliography

[1] M. König, "sbmlsim: SBML simulation made easy." Zenodo, 2026. doi: 10.5281/zenodo.3597770. Version 0.10.0.

[2] R. Storn and K. Price, "Differential evolution - a simple and efficient heuristic for global optimization over continuous spaces," *Journal of Global Optimization*, vol. 11, pp. 341–359, 1997, doi: 10.1023/A:1008202821328.

[3] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python," *Nature Methods*, vol. 17, pp. 261–272, 2020, doi: 10.1038/s41592-019-0686-2. Version 1.18.1.

[4] M. D. McKay, R. J. Beckman, and W. J. Conover, "A comparison of three methods for selecting values of input variables in the analysis of output from a computer code," *Technometrics*, vol. 21, no. 2, pp. 239–245, 1979, doi: 10.2307/1268522.

[5] C. Welsh, J. Xu, L. Smith, M. König, K. Choi, and H. M. Sauro, "libRoadRunner 2.0: a high performance SBML simulation and analysis library," *Bioinformatics*, vol. 39, no. 1, p. btac770, 2023, doi: 10.1093/bioinformatics/btac770. Version 2.10.0.

[6] A. C. Hindmarsh *et al.*, "SUNDIALS: suite of nonlinear and differential/algebraic equation solvers," *ACM Transactions on Mathematical Software*, vol. 31, no. 3, pp. 363–396, 2005, doi: 10.1145/1089014.1089020.

[7] H. Akaike, "A new look at the statistical model identification," *IEEE Transactions on Automatic Control*, vol. 19, no. 6, pp. 716–723, 1974, doi: 10.1109/TAC.1974.1100705.

[8] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978, doi: 10.1214/aos/1176344136.

[9] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 327, no. 8476, pp. 307–310, 1986, doi: 10.1016/S0140-6736(86)90837-8.

[10] A. Raue *et al.*, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, no. 15, pp. 1923–1929, 2009, doi: 10.1093/bioinformatics/btp358.

[11] C. Kreutz, A. Raue, D. Kaschek, and J. Timmer, "Profile likelihood in systems biology," *The FEBS Journal*, vol. 280, no. 11, pp. 2564–2571, 2013, doi: 10.1111/febs.12276.

[12] F.-G. Wieland, A. L. Hauber, M. Rosenblatt, C. Tönsing, and J. Timmer, "On structural and practical identifiability," *Current Opinion in Systems Biology*, vol. 25, pp. 60–69, 2021, doi: 10.1016/j.coisb.2021.03.005.

[13] C. R. Harris *et al.*, "Array programming with NumPy," *Nature*, vol. 585, pp. 357–362, 2020, doi: 10.1038/s41586-020-2649-2. Version 2.5.3.