Dulaglutide PBPK/PD model: fit PD

1 Methods: Dulaglutide PBPK/PD model: fit PD

1.1 Parameters and data

8 parameters were optimized: maximal effect of the GLP-1 agonist on the excess FPG (fraction) (`Emax_glp1_fpg`), EC50 of the GLP-1 agonist on the excess FPG (`EC50_glp1_fpg`), rate normalization FPG with GLP-1 agonist (`k_fpg`), trial effect on the excess FPG (fraction) (`f_trial_fpg`), hb degradation rate (`k_hb_deg`), maximal effect of the GLP-1 agonist on the subcutaneous fat (fraction) (`Emax_SAT_glp1`), EC50 for GLP-1 agonist on fat loss (`EC50_SAT_glp1`) and rate of the change of the subcutaneous fat (`k_SAT_glp1`). The parameters of the adopted fit PK_20261018_102441__faa2d were fixed at their fitted values.

A fit mapping is a curve of a study, the data and the observable of the model it is compared with. The fit has 112 fit mappings of training data with 525 data points, whose residuals enter the objective; 31 fit mappings of validation data with 184 data points, which were simulated and evaluated with the fitted parameters but not fitted; and 3 fit mappings of outliers with 8 data points, data the fit dropped as not usable, which were evaluated like the validation data so that the decision to drop them can be checked. The fit was set up and run with sbmlsim [1].

1.2 Objective function

The parameters were estimated by minimizing the objective function

$$F(\vec{p}) = \frac{1}{2} \sum_k \sum_i \rho\big(w_{i,k} r_{i,k}(\vec{p})^2\big) \tag{1}$$

where k runs over the fit mappings of the training data and i over their data points, $r_{i,k}$ is the residual, $w_{i,k} = w_k w^p_{i,k}$ the weight of the data point and ρ the loss function, applied to the squared weighted residual $z = w_{i,k} r^2_{i,k}$. The loss function is linear, $\rho(z) = z$, i.e. a standard least squares problem.

The residuals are normalized by the mean $\overline{y}_k$ of the data of their curve, which makes curves of different quantities and magnitudes comparable:

$$r_{i,k} = \frac{y_{i,k} - f_{k(x_{i,k},\vec{p})}}{\overline{y}_k} \tag{2}$$

The weight of curve k is $w_k = \frac{n_k}{N_k}$, the weight n_k of its fit mapping (the weight its fit mapping collection assigns, 1 by default, or, for a collection which uses the weights of its mappings, the weight of the mapping, by default the number of subjects of its data) divided by the number N_k of its data points.

A data point is weighted with the inverse of its coefficient of variation, $w^p_{i,k} = |y_{i,k} \frac{|}{\sigma_{i,k}}$, with $\sigma_{i,k}$ the standard deviation of the data point, or its standard error for data without a standard deviation. An error of zero is missing, a missing error of a curve with errors is replaced by the largest error of the curve, and the data points of a curve without errors have the weight $w^p_{i,k} = 2$, the weight of a coefficient of variation of 0.5.

1.3 Optimization

The optimizer searched the parameters in log10 space within their bounds; bounds, values and intervals are given in the units of the fit parameters, the units of the table of the parameters.

The objective was minimized in 3 runs with differential evolution [2] of SciPy [3] within the bounds of the parameters, with the strategy `best1bin`, a population of 15 times the number of parameters, at most 1000 generations, the relative tolerance 0.01 of convergence, the recombination constant 0.7 and a mutation dithered between 0.5 and 1. The initial population of a run was drawn by latin hypercube sampling [4], with a seed of every run derived from the seed 1234 of the fit. The best member of the final population was polished with the L-BFGS-B method of SciPy.

The convergence of the runs was assessed from the traces of their cost and the waterfall plot of their final costs, and the parameter set with the lowest cost was used.

The models were simulated with libRoadRunner [5] and its integrator CVODE [6] with the relative tolerance 10^{-6} and the absolute tolerance 10^{-6}, which for a species in a concentration is multiplied by the initial volume of its compartment, and 18 states of its own. It ran with the variable step size of libRoadRunner, i.e. with an output at every step of the integrator. A data point at the time of a change after the start of a simulation, e.g. a dose or the reset of a

collection, was compared with the state before the change, except the first data point of a curve with data at later times, which starts a collection and was compared with the state after the change.

1.4 Goodness of fit

The goodness of fit is given per kind of data (training data, validation data and outliers) by metrics over its n data points with the residuals $e_i = y_i - f\left(x_i, \hat{\vec{p}}\right)$ at the fitted parameters $\hat{\vec{p}}$, the mean squared error, its root and the root mean squared error normalized by the mean $\overline{y}_{k(i)}$ of the data of the curve of a point:

$$\mathrm{MSE} = \frac{1}{n}\sum_{i=1}^{n} e_i^2, \quad \mathrm{RMSE} = \sqrt{\mathrm{MSE}}, \quad \mathrm{NRMSE} = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{e_i}{\overline{y}_{k(i)}}\right)^2} \tag{3}$$

The root mean square of the residuals of the objective, i.e. what the fit minimizes, and the coefficient of determination:

$$\mathrm{RMSE}_w = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\rho(w_i r_i^2)}, \quad R^2 = 1 - \frac{\sum_i e_i^2}{\sum_i \left(y_i - \overline{y}\right)^2} \tag{4}$$

The predictions of a non-linear model are no linear regression of the data, so R^2 can be negative: the prediction is worse than the mean of the data. The information criteria of Akaike [7] and Schwarz [8] with the number n_p of fitted parameters are those of a least squares fit with normally distributed residuals, up to an additive constant, so only differences between models fitted to the same data are meaningful:

$$\mathrm{AIC} = n\ln(\mathrm{MSE}) + 2n_p, \quad \mathrm{BIC} = n\ln(\mathrm{MSE}) + n_p\ln(n) \tag{5}$$

The percentage of the data points whose prediction $f_i = f\left(x_i, \hat{\vec{p}}\right)$ lies within twofold of the data is taken over the n_+ data points with positive data $y_i > 0$ and a positive prediction $f_i > 0$; the others have no ratio and are not counted, and a kind of data without such a point has no percentage:

$$P_2 = \frac{100}{n_+}\left|\left\{i : \frac{1}{2} \le \frac{f_i}{y_i} \le 2\right\}\right| \tag{6}$$

The goodness of fit plot shows the predictions against the data on logarithmic axes, with the lines of twofold $f = 2y$ and $f = \frac{y}{2}$. The Bland-Altman plot [9] shows the ratio of prediction and data as $d_i = \log_{10}\left(\frac{f_i}{y_i}\right)$ against the geometric mean $\sqrt{f_i y_i}$ for the data points with positive values, with the bias $\overline{d}$ and the limits of agreement $\overline{d} \pm 1.96 s_d$, with the sample standard deviation s_d of d_i, of the training data; they are read as the fold factors $10^{\overline{d}}$ and $10^{\overline{d} \pm 1.96 s_d}$.

1.5 Identifiability

The identifiability of the parameters was analysed with the profile likelihood [10], [11]. The profile of a parameter is the lowest objective with the parameter fixed:

$$\mathrm{PL}(p_j) = \min_{p_{i \ne j}} F(\vec{p}) \tag{7}$$

At every point of a scan the other parameters were re-optimized from the previous point with the least squares optimizer of SciPy with the relative step 0.05. A scan ran from the optimum in both directions in the space of the optimizer, with steps in decades of the parameter, starting at 0.1. A step which raised the objective by more than 0.2 of the distance between the objective at the optimum and the threshold was reduced by the factor 2 and repeated, down to 0.01, and the step after one which raised it by less than that divided by the factor was enlarged by the factor, up to 1. A scan stopped at the threshold, at the bound of the parameter, at a point which failed or after 50 points.

The confidence intervals are the values at which the profile crosses the threshold of the likelihood ratio test

$$F_{\mathrm{thr}} = F_{\mathrm{min}} + \chi^2_\alpha \frac{\mathrm{df}}{2} \tag{8}$$

with F_{min} the lowest objective of the fit and of all profiles and $\chi^2_\alpha(\mathrm{df})$ the quantile of the χ^2 distribution at the confidence level $\alpha = 0.95$ with df = 1 degrees of freedom, i.e. 1.92 above the lowest objective. The threshold has this confidence level if F is the negative log-likelihood of the data up to a constant, i.e. if the weighted residuals are independent and normally distributed with unit variance; otherwise the level is approximate. A crossing is interpolated between two points of the profile, on the logarithm of a parameter on a logarithmic scale. A parameter is identifiable if its profile reaches the threshold within the scan on both sides of the optimum, i.e. its confidence

interval is finite; practically non-identifiable towards the lower bound if its profile reaches the threshold above the optimum but not below it within the scan, which ends at the lower bound or after 50 points; practically non-identifiable towards the upper bound if its profile reaches the threshold below the optimum but not above it within the scan, which ends at the upper bound or after 50 points; practically non-identifiable if its profile reaches the threshold on neither side within the scans, which end at the bounds or after 50 points, and is not flat; and structurally non-identifiable if its profile is flat on both sides. A side of a profile is flat if it rises by less than 0.05 of the distance between the lowest objective and the threshold, and a profile with a point which failed or did not converge is not classified. The profile likelihood is preferred over the intervals of the Fisher information, its intervals are invariant under a transformation of the parameters and can be asymmetric [12].

The Fisher information is the local alternative to the profiles: it is the Gauss-Newton approximation of the curvature of the objective at the optimum, from the Jacobian J of the weighted residuals of the training data with respect to the parameters $\vec{p}$ in the space of the optimizer, computed by central finite differences with the step $h_j = \varepsilon \max(|p_j|, 1),\ \varepsilon = 10^{-6}$ of the parameter p_j:

$$I = J^\top J, \quad \mathrm{cov} = \sigma^2 I^{-1}, \quad \sigma^2 = \frac{2F}{n - n_p} \tag{9}$$

with the number n of data points of the training data and the number n_p of parameters. The information has rank 8 of 8, the number of its eigenvalues above 10^{-8} of the largest one, and the condition number 9580; the covariance of an information of a lower rank is computed with the pseudo-inverse. The Wald intervals are $\hat{p}_j \pm t_{\frac{1+\alpha}{2}, n-n_p} \mathrm{SE}_j$ with the standard errors $\mathrm{SE}_j = \sqrt{\mathrm{cov}_{jj}}$ at the confidence level $\alpha = 0.95$, transformed back into the units of the fit parameters.

1.6 Provenance

The fit PD_20261019_103700__c0378 was run on 2026-10-19 08:37:26 UTC with sbmlsim 0.10.0 [1], libRoadRunner 2.10.0 [5], SciPy 1.18.1 [3], NumPy 2.5.3 [13] and Python 3.14.4 (seed 1234). The models and data were those of commit `a7d90200465ba23560ee963d70986d203c8b41df` of `/home/mkoenig/git/pkdb_models-dulaglutide` their SHA-256 checksums are listed in Table 1. The fit kept the values `DULCAT_k` = 0.00160903, `Kp_dul` = 0.0870081, `ftissue_dul` = 0.0282682, `Ksc_dul` = $4.81 \cdot 10^{-4}$ and `Fsc_dul` = 0.478929 of the model fixed.

Table 1: SHA-256 checksums of the models and data of the fit PD_20261019_103700__c0378, their first 12 characters; the full checksums are in the provenance of the result (optimization_result.json).

File	SHA-256
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011/.Barrington2011_Fig1.tsv`	`20176599267d...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011/.Barrington2011_Fig2.tsv`	`798431c8090b...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011/.Barrington2011_Tab1.tsv`	`3da28f3a8cdc...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011/.Barrington2011_TabGroups.tsv`	`61afc5ec2f7a...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011/.Barrington2011_TabInterventions.tsv`	`b758eaf2cf07...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011a/.Barrington2011a_Fig1.tsv`	`09fa713e7de1...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011a/.Barrington2011a_Fig2.tsv`	`df6839feea0e...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011a/.Barrington2011a_Fig3.tsv`	`8771df0bebe1...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011a/.Barrington2011a_TabGroups.tsv`	`7e478c9c801f...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Barrington2011a/.Barrington2011a_TabInterventions.tsv`	`7d9c6e22cb0a...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Blonde2015/.Blonde2015_Fig2a.tsv`	`d4622077f93a...`
`pkdb_models/models/dulaglutide/data/dulaglutide/Blonde2015/.Blonde2015_Fig2b.tsv`	`032ee105b6af...`

File	SHA-256
pkdb_models/models/dulaglutide/data/dulaglutide/Blonde2015/.Blonde2015_Fig2f.tsv	a82edde842a2...
pkdb_models/models/dulaglutide/data/dulaglutide/Blonde2015/.Blonde2015_TabGroups.tsv	1a4626420876...
pkdb_models/models/dulaglutide/data/dulaglutide/Blonde2015/.Blonde2015_TabInterventions.tsv	57a4f0f0f29d...
pkdb_models/models/dulaglutide/data/dulaglutide/Chen2018/.Chen2018_Fig2b.tsv	e5fd14cbaa36...
pkdb_models/models/dulaglutide/data/dulaglutide/Chen2018/.Chen2018_Fig2d.tsv	d971be80b426...
pkdb_models/models/dulaglutide/data/dulaglutide/Chen2018/.Chen2018_Fig2f.tsv	8a930bd27822...
pkdb_models/models/dulaglutide/data/dulaglutide/Chen2018/.Chen2018_TabGroups.tsv	8fbb984726c8...
pkdb_models/models/dulaglutide/data/dulaglutide/Chen2018/.Chen2018_TabInterventions.tsv	b678d0d7a3e4...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2014/.Dungan2014_Fig2b.tsv	ea390d6166d4...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2014/.Dungan2014_Fig2f.tsv	cdcd76e6f85f...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2014/.Dungan2014_Tab2.tsv	7706661da967...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2014/.Dungan2014_TabGroups.tsv	4645ab15523b...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2014/.Dungan2014_TabInterventions.tsv	689b59ded114...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2016/.Dungan2016_Fig2B.tsv	ae476ac9144c...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2016/.Dungan2016_Fig2D.tsv	7e8d3f39d353...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2016/.Dungan2016_Fig2F.tsv	f604a834bd86...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2016/.Dungan2016_TabGroups.tsv	01050e94d86d...
pkdb_models/models/dulaglutide/data/dulaglutide/Dungan2016/.Dungan2016_TabInterventions.tsv	1e5b2ecf6f42...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_Fig21.tsv	8f92f5065408...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_Fig42.tsv	432830781e6e...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_Tab46.tsv	d91880b0f8db...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_TabGroups.tsv	b5769ed331cd...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_TabIndividuals.tsv	f2170e3a8033...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCM/.FDAGBCM_TabInterventions.tsv	d07ee46a3cfb...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_Fig34.tsv	1b46a7b09bb5...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_Tab11.tsv	c9c0e85c034b...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_Tab33.tsv	87b87b18ed5c...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_Tab34.tsv	1dbd0e08c5fd...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_Tab35.tsv	8b0d2a59637f...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_TabGroups.tsv	71bb1f02817a...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBCN/.FDAGBCN_TabInterventions.tsv	b4fd613fb426...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_Fig22.tsv	7f4f61591983...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_Fig45.tsv	515d77f4f750...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_Tab48.tsv	859fa47c6d6e...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_Tab49.tsv	eebe319efc5e...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_TabGroups.tsv	5ec7dfdfefaf...

File	SHA-256
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDO/.FDAGBDO_TabInterventions.tsv	f6ed64016f63...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_Fig15.tsv	ffe4af50f67d...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_Fig32.tsv	4322da8a2afe...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_Tab10.tsv	26ae65643f48...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_Tab31.tsv	0cef62866325...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_TabGroups.tsv	ce78f59397cc...
pkdb_models/models/dulaglutide/data/dulaglutide/FDAGBDR/.FDAGBDR_TabInterventions.tsv	018babac593f...
pkdb_models/models/dulaglutide/data/dulaglutide/Gao2024/.Gao2024_Fig2.tsv	b7191b7c34af...
pkdb_models/models/dulaglutide/data/dulaglutide/Gao2024/.Gao2024_Tab2.tsv	b4b2664aa260...
pkdb_models/models/dulaglutide/data/dulaglutide/Gao2024/.Gao2024_Tab3.tsv	0a5d7b49ac60...
pkdb_models/models/dulaglutide/data/dulaglutide/Gao2024/.Gao2024_TabGroups.tsv	a13a104a1b2f...
pkdb_models/models/dulaglutide/data/dulaglutide/Gao2024/.Gao2024_TabInterventions.tsv	f9f0eef97a37...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_Fig4a.tsv	a2a59f9f392a...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_Fig4b.tsv	94a7ca31df3d...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_Fig4c.tsv	2f06fbf4df83...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_Fig4d.tsv	04f4dd8414e4...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_TabGroups.tsv	31234340ecbb...
pkdb_models/models/dulaglutide/data/dulaglutide/Gerstein2019/.Gerstein2019_TabInterventions.tsv	3351fe8fd309...
pkdb_models/models/dulaglutide/data/dulaglutide/Giorgino2015/.Giorgino2015_Fig2b.tsv	15cb98a0622d...
pkdb_models/models/dulaglutide/data/dulaglutide/Giorgino2015/.Giorgino2015_Fig2d.tsv	63ee3e9f215e...
pkdb_models/models/dulaglutide/data/dulaglutide/Giorgino2015/.Giorgino2015_Fig2f.tsv	fe262c512f91...
pkdb_models/models/dulaglutide/data/dulaglutide/Giorgino2015/.Giorgino2015_TabGroups.tsv	21c35d0044d6...
pkdb_models/models/dulaglutide/data/dulaglutide/Giorgino2015/.Giorgino2015_TabInterventions.tsv	88cbe904b7c1...
pkdb_models/models/dulaglutide/data/dulaglutide/Grunberger2012/.Grunberger2012_Fig3ab.tsv	b85b8f3637b0...
pkdb_models/models/dulaglutide/data/dulaglutide/Grunberger2012/.Grunberger2012_Fig3d.tsv	785f129af4d7...
pkdb_models/models/dulaglutide/data/dulaglutide/Grunberger2012/.Grunberger2012_TabGroups.tsv	bc3642f53dbd...
pkdb_models/models/dulaglutide/data/dulaglutide/Grunberger2012/.Grunberger2012_TabInterventions.tsv	1fa3e8174272...
pkdb_models/models/dulaglutide/data/dulaglutide/Liu2025/.Liu2025_Fig3.tsv	61233a8b2406...
pkdb_models/models/dulaglutide/data/dulaglutide/Liu2025/.Liu2025_Tab2.tsv	21cff5b77e80...
pkdb_models/models/dulaglutide/data/dulaglutide/Liu2025/.Liu2025_Tab6.tsv	d8087c723671...
pkdb_models/models/dulaglutide/data/dulaglutide/Liu2025/.Liu2025_TabGroups.tsv	c4d2f24be315...

File	SHA-256
pkdb_models/models/dulaglutide/data/dulaglutide/Liu2025/.Liu2025_TabInterventions.tsv	ecc9420ee466...
pkdb_models/models/dulaglutide/data/dulaglutide/Morioka2024/.Morioka2024_Fig1.tsv	d3ceec1fd00e...
pkdb_models/models/dulaglutide/data/dulaglutide/Morioka2024/.Morioka2024_Fig2.tsv	af81b9691c88...
pkdb_models/models/dulaglutide/data/dulaglutide/Morioka2024/.Morioka2024_Fig3.tsv	657b2c5c5a0f...
pkdb_models/models/dulaglutide/data/dulaglutide/Morioka2024/.Morioka2024_TabGroups.tsv	1764ea1e3e19...
pkdb_models/models/dulaglutide/data/dulaglutide/Morioka2024/.Morioka2024_TabInterventions.tsv	076e5063bd79...
pkdb_models/models/dulaglutide/data/dulaglutide/Nauck2014/.Nauck2014_Fig2b.tsv	ed1fb8a72cc8...
pkdb_models/models/dulaglutide/data/dulaglutide/Nauck2014/.Nauck2014_Fig2d.tsv	3a54eceee18f...
pkdb_models/models/dulaglutide/data/dulaglutide/Nauck2014/.Nauck2014_Fig2e.tsv	51eefc1ce59f...
pkdb_models/models/dulaglutide/data/dulaglutide/Nauck2014/.Nauck2014_TabGroups.tsv	5710b4229816...
pkdb_models/models/dulaglutide/data/dulaglutide/Nauck2014/.Nauck2014_TabInterventions.tsv	e45792cfd1ef...
pkdb_models/models/dulaglutide/data/dulaglutide/Pratley2018/.Pratley2018_Fig2a.tsv	1197112fd480...
pkdb_models/models/dulaglutide/data/dulaglutide/Pratley2018/.Pratley2018_Fig2c.tsv	561d8bcff847...
pkdb_models/models/dulaglutide/data/dulaglutide/Pratley2018/.Pratley2018_Fig2g.tsv	3fd86ea36797...
pkdb_models/models/dulaglutide/data/dulaglutide/Pratley2018/.Pratley2018_TabGroups.tsv	cab9507b18a4...
pkdb_models/models/dulaglutide/data/dulaglutide/Pratley2018/.Pratley2018_TabInterventions.tsv	c783944be36e...
pkdb_models/models/dulaglutide/data/dulaglutide/Teraruchi2014/.Teraruchi2014_Fig1b.tsv	ac307c905788...
pkdb_models/models/dulaglutide/data/dulaglutide/Teraruchi2014/.Teraruchi2014_Fig1d.tsv	deac7c758644...
pkdb_models/models/dulaglutide/data/dulaglutide/Teraruchi2014/.Teraruchi2014_Tab2.tsv	2ca9ac22fc1c...
pkdb_models/models/dulaglutide/data/dulaglutide/Teraruchi2014/.Teraruchi2014_TabGroups.tsv	c0b57e510943...
pkdb_models/models/dulaglutide/data/dulaglutide/Teraruchi2014/.Teraruchi2014_TabInterventions.tsv	18be784d3ca8...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_Fig2a.tsv	868b6781bcad...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_Fig2b.tsv	54a14d2d6967...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_Fig3.tsv	55126e83087f...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_Fig5.tsv	77829da4a269...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_TabGroups.tsv	e28c868f2c85...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2011/.Umpierrez2011_TabInterventions.tsv	0c9dcc0f001c...

File	SHA-256
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_Fig2a.tsv	c61e2b7e520d...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_Fig2b.tsv	e42105b88fd0...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_Fig2d.tsv	f63e30025739...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_Fig2f.tsv	72fc96327a03...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_TabGroups.tsv	93c9e2722961...
pkdb_models/models/dulaglutide/data/dulaglutide/Umpierrez2014/.Umpierrez2014_TabInterventions.tsv	2fa22bc84d0b...
pkdb_models/models/dulaglutide/data/dulaglutide/Wang2019/.Wang2019_Fig2b.tsv	d4c5b18f27be...
pkdb_models/models/dulaglutide/data/dulaglutide/Wang2019/.Wang2019_Fig2d.tsv	87cbfed38324...
pkdb_models/models/dulaglutide/data/dulaglutide/Wang2019/.Wang2019_Fig2f.tsv	c9065414eba2...
pkdb_models/models/dulaglutide/data/dulaglutide/Wang2019/.Wang2019_TabGroups.tsv	9a9eae9857e5...
pkdb_models/models/dulaglutide/data/dulaglutide/Wang2019/.Wang2019_TabInterventions.tsv	89b8b02610a6...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_Fig2a.tsv	8d8dd02e55e8...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_Fig2b.tsv	f694ffafada0...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_Fig2d.tsv	498aa4ea5d20...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_Fig2e.tsv	68472152b896...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_TabGroups.tsv	0b3f1dbebd49...
pkdb_models/models/dulaglutide/data/dulaglutide/Weinstock2015/.Weinstock2015_TabInterventions.tsv	5f724b71862b...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_Fig2a.tsv	3b4f56a34fe6...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_Fig2b.tsv	e254a530934c...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_Fig2d.tsv	1dc6478fc989...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_Fig2f.tsv	fcb0e2add93e...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_TabGroups.tsv	612203818bbe...
pkdb_models/models/dulaglutide/data/dulaglutide/Wysham2014/.Wysham2014_TabInterventions.tsv	c65e0986a04d...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_Fig1.tsv	803f3b2df214...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_Fig2.tsv	6d89d7f868c2...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_Fig3.tsv	24624b622386...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_Tab2.tsv	cba84b8464c8...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_Tab4.tsv	6d6cfcd0eaae...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_TabGroups.tsv	76a2c9af3d07...
pkdb_models/models/dulaglutide/data/dulaglutide/Xu2022/.Xu2022_TabInterventions.tsv	553ed212059d...
pkdb_models/models/dulaglutide/data/dulaglutide/Zhang2023/.Zhang2023_Fig2.tsv	7e6f1424e581...
pkdb_models/models/dulaglutide/data/dulaglutide/Zhang2023/.Zhang2023_Tab3.tsv	ada4dad3f25b...
pkdb_models/models/dulaglutide/data/dulaglutide/Zhang2023/.Zhang2023_Tab4.tsv	66478386bf01...

File	SHA-256
pkdb_models/models/dulaglutide/data/dulaglutide/Zhang2023/.Zhang2023_TabGroups.tsv	77f60a5d9ca6...
pkdb_models/models/dulaglutide/data/dulaglutide/Zhang2023/.Zhang2023_TabInterventions.tsv	6f4e6dd8dc90...
pkdb_models/models/dulaglutide/models/output/models/dulaglutide_body.xml	8b128f2ffd5c...

2 Results: Dulaglutide PBPK/PD model: fit PD

2.1 Parameters

Table 2: Parameters of the fit PD_20261019_103700__c0378, the parameter set PD_20261019_103700__c0378_0, cost 33.1, with their bounds in the optimization. The 95% confidence intervals are those of the profile likelihood, LB and UB an interval which is open towards the lower or the upper bound, and the Wald intervals of the Fisher information for a parameter without a profile. Identifiable is the classification of the profile: yes, no (LB) or no (UB) if the profile does not reach the threshold towards the lower or the upper bound, no if it reaches it on neither side, structural if it is flat.

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
Emax_glp1_fpg	maximal effect of the GLP-1 agonist on the excess FPG (fraction)	0.614 [0.509, 0.737]	dimensionless	[0.05, 1]	yes
EC50_glp1_fpg	EC50 of the GLP-1 agonist on the excess FPG	3.4e-07 [1.39e-07, 6.81e-07]	mmol/l	[1e-08, 0.0001]	yes
k_fpg	rate normalization FPG with GLP-1 agonist	0.00031 [0.000138, 0.00188]	l/min	[1e-05, 0.01]	yes
f_trial_fpg	trial effect on the excess FPG (fraction)	0.0424 [LB, 0.118]	dimensionless	[0.001, 0.5]	no (LB)
k_hb_deg	hb degradation rate	2.43e-05 [1.71e-05, 3.41e-05]	mmol/min	[5e-06, 5e-05]	yes
Emax_SAT_glp1	maximal effect of the GLP-1 agonist on the subcutaneous fat (fraction)	0.204 [0.101, UB]	dimensionless	[0.01, 0.8]	no (UB)
EC50_SAT_glp1	EC50 for GLP-1 agonist on fat loss	4.25e-07 [LB, 7.28e-06]	mmol/l	[1e-08, 0.0001]	no (LB)
k_SAT_glp1	rate of the change of the subcutaneous fat	2.11e-05 [3.74e-06, UB]	1/min	[5e-07, 0.0001]	no (UB)

2.2 Data

Table 3: Data of the fit PD_20261019_103700__c0378: the number of fit mappings and, in parentheses, of data points per study and kind. The training data enters the cost, the validation data and the outliers are evaluated only.

Study	Training	Validation	Outlier	Total
Barrington2011a	18 (48)	-	3 (8)	21 (56)
Blonde2015	-	4 (28)	-	4 (28)
Chen2018	6 (26)	-	-	6 (26)
Dungan2014	3 (14)	-	-	3 (14)
Dungan2016	6 (28)	-	-	6 (28)
Gerstein2019	-	4 (26)	-	4 (26)
Giorgino2015	-	6 (64)	-	6 (64)

Study	Training	Validation	Outlier	Total
Grunberger2012	10 (45)	-	-	10 (45)
Liu2025	3 (9)	3 (9)	-	6 (18)
Morioka2024	6 (36)	-	-	6 (36)
Nauck2014	9 (63)	-	-	9 (63)
Pratley2018	6 (42)	-	-	6 (42)
Terauchi2014	12 (44)	-	-	12 (44)
Umpierrez2011	6 (20)	6 (20)	-	12 (40)
Umpierrez2014	6 (40)	-	-	6 (40)
Wang2019	6 (38)	-	-	6 (38)
Weinstock2015	6 (24)	-	-	6 (24)
Wysham2014	5 (32)	4 (25)	-	9 (57)
Xu2022	4 (16)	4 (12)	-	8 (28)
Total	112 (525)	31 (184)	3 (8)	146 (717)

2.3 Goodness of fit

Table 4: Goodness of fit of the fit PD_20261019_103700__c0378 per kind of data, the parameter set PD_20261019_103700__c0378_0. The cost is defined on the training data only, 2-fold is the percentage of the data points with a positive value and prediction whose prediction is within twofold of the data.

Kind	n	cost	MSE	RMSE	NRMSE	RMSE_w	R^2	2-fold [%]	AIC	BIC
training	525	33.1	0.148	0.385	0.0347	0.355	1	100.0	−986.9	−952.8
validation	184	-	0.658	0.811	0.0356	0.731	1	100.0	−60.9	−35.2
outlier	8	-	0.42	0.648	0.0703	0.181	1	100.0	9.1	9.7

2.4 Figures




Figure 1: Cost of the optimizers over their steps, fit PD_20261019_103700__c0378. The cost of every optimization run over its steps. Runs which end at the same cost found the same optimum; a run which stops much higher is stuck in a local minimum.




Figure 2: Cost of the optimization runs, ordered, fit PD_20261019_103700__c0378. The final cost of the runs, ordered. A flat plateau at the left is the global optimum found repeatedly, which is the evidence that the multistart converged; steps to the right are local minima.




Figure 3: Prediction against the measured data points, per kind of fit mapping, with the agreement of the training data, the lines of twofold and the metrics of the panel, fit PD_20261019_103700__c0378. Within twofold of the data are 525 of the 525 training data points (100.0%), 184 of the 184 validation data points (100.0%) and 8 of the 8 outlier data points (100.0%); only the data points with a positive value and prediction are counted. Prediction against measurement, one point per data point, with a panel per kind of fit mapping. The points scatter around the diagonal when the model describes the data; a systematic deviation from it is a systematic error of the model. The band is the agreement of the training data, i.e. the same band the Bland-Altman plot draws, here as lines parallel to the diagonal. The dark red dotted lines are twofold, prediction = 2 x measurement and prediction = measurement / 2; the caption states the points between them of every kind. A point which is zero or negative is not drawn on the logarithmic axes and not counted, the title of a panel then states the points drawn of all. The validation panel shows how the fit describes data it was not fitted on, the outlier panel the data the fit dropped. The box of a panel carries its R^2, its normalized RMSE, the RMSE of the residuals of the cost, its fold errors GMFE and AFE and the percentages of its points within twofold and within two standard deviations of the data.




Figure 4: Agreement of prediction and measurement as a ratio, per kind of fit mapping, with the agreement of the training data and the agreement of the panel, fit PD_20261019_103700__c0378. The ratio of prediction and measurement over the geometric mean of the two, i.e. the goodness of fit with the diagonal turned into the horizontal. The band is the same, the bias and the limits of agreement `bias ± 1.96 SD` of the training data as fold factors. A bias away from 1 is a systematic over- or underprediction, wide limits are a large scatter, and a trend over the mean is a model which describes the large or the small values better. The box of a panel carries the bias and the SD of its own points as fold factors and the share of them inside the limits of agreement of the training data.




Figure 5: Profile likelihood of every parameter: the cost with the parameter fixed and the other parameters optimized, the threshold of the confidence level and the confidence interval, fit PD_20261019_103700__c0378. One profile per parameter. A profile which crosses the dashed threshold on both sides gives a finite confidence interval; one which stays below it up to a bound is practically non-identifiable; a flat profile is structurally non-identifiable, the parameter is compensated by the others.

2.5 Provenance

Table 5: Provenance of the fit PD_20261019_103700__c0378: the versions it ran with and the revision of its models and data; the SHA-256 checksums of the models and data are listed in Table 1.

Item	Value
Fit	`PD_20261019_103700__c0378`
Created	2026-10-19 08:37:26 UTC
Seed	1234
Python	3.14.4
sbmlsim	0.10.0
libRoadRunner	2.10.0
SciPy	1.18.1
NumPy	2.5.3
Repository	`/home/mkoenig/git/pkdb_models-dulaglutide`
Commit	`a7d90200465ba23560ee963d70986d203c8b41df`
Uncommitted changes	no

Bibliography

[1] M. König, "sbmlsim: SBML simulation made easy." Zenodo, 2026. doi: 10.5281/zenodo.3597770. Version 0.10.0.

[2] R. Storn and K. Price, "Differential evolution - a simple and efficient heuristic for global optimization over continuous spaces," *Journal of Global Optimization*, vol. 11, pp. 341–359, 1997, doi: 10.1023/A:1008202821328.

[3] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python," *Nature Methods*, vol. 17, pp. 261–272, 2020, doi: 10.1038/s41592-019-0686-2. Version 1.18.1.

[4] M. D. McKay, R. J. Beckman, and W. J. Conover, "A comparison of three methods for selecting values of input variables in the analysis of output from a computer code," *Technometrics*, vol. 21, no. 2, pp. 239–245, 1979, doi: 10.2307/1268522.

[5] C. Welsh, J. Xu, L. Smith, M. König, K. Choi, and H. M. Sauro, "libRoadRunner 2.0: a high performance SBML simulation and analysis library," *Bioinformatics*, vol. 39, no. 1, p. btac770, 2023, doi: 10.1093/bioinformatics/btac770. Version 2.10.0.

[6] A. C. Hindmarsh *et al.*, "SUNDIALS: suite of nonlinear and differential/algebraic equation solvers," *ACM Transactions on Mathematical Software*, vol. 31, no. 3, pp. 363–396, 2005, doi: 10.1145/1089014.1089020.

[7] H. Akaike, "A new look at the statistical model identification," *IEEE Transactions on Automatic Control*, vol. 19, no. 6, pp. 716–723, 1974, doi: 10.1109/TAC.1974.1100705.

[8] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978, doi: 10.1214/aos/1176344136.

[9] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 327, no. 8476, pp. 307–310, 1986, doi: 10.1016/S0140-6736(86)90837-8.

[10] A. Raue *et al.*, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, no. 15, pp. 1923–1929, 2009, doi: 10.1093/bioinformatics/btp358.

[11] C. Kreutz, A. Raue, D. Kaschek, and J. Timmer, "Profile likelihood in systems biology," *The FEBS Journal*, vol. 280, no. 11, pp. 2564–2571, 2013, doi: 10.1111/febs.12276.

[12] F.-G. Wieland, A. L. Hauber, M. Rosenblatt, C. Tönsing, and J. Timmer, "On structural and practical identifiability," *Current Opinion in Systems Biology*, vol. 25, pp. 60–69, 2021, doi: 10.1016/j.coisb.2021.03.005.

[13] C. R. Harris *et al.*, "Array programming with NumPy," *Nature*, vol. 585, pp. 357–362, 2020, doi: 10.1038/s41586-020-2649-2. Version 2.5.3.